Published in final edited form as:

Am J Psychiatry. 2013 November 1; 170(11): 1367. doi:10.1176/appi.ajp.2013.13060813r.

Response to Keating and Rossell

GUIDO K. FRANK, M.D., MEGAN E. SHOTT, B.S., JENNIFER O. HAGMAN, M.D., and VIJAY A. MITTAL, PH.D.

To the Editor: The correlation between gyrus rectus volume and sweet taste pleasantness in the comparison group is striking as it supports the previous literature and is consistent with an important concept established by the works of Rolls, Kringelbach, and others (1). All eating disorder groups in our sample showed increased gyrus rectus volume relative to individuals in the comparison group, suggesting comparable pathophysiology. Groups were combined for that reason, and we did find a positive correlation between sweet taste pleasantness and volume in the eating disorder case subjects. In addition, within eating disorder subgroups, each individual group showed a positive correlation (anorexia nervosa, $r=0.120$; bulimia nervosa, $r=0.287$; recovered anorexia nervosa, $r=0.345$) between sweet taste pleasantness and gyrus rectus volume, supporting the idea that analyzing the groups as a whole was valid. The correlation coefficients were modest. Despite a significant amount of research, we still know little about the connection between brain structure and function and these results are a step toward identifying target areas that could contribute to altered food intake. Taste pleasantness in individuals with eating disorder could be less well mapped (yet still positively) onto gyrus rectus neurons due to the altered volume, and this could interfere with correct value computation.

It is important to view eating disorders as complex problems with a variety of factors that most likely contribute to development and then perpetuation of the illnesses. Expecting direct correlations between all those measures would be overly simplistic, and we believe that dissecting eating disorders by brain circuits that may contribute to the psychopathology will be more fruitful. Trait alterations seem to exist that cause heightened sensitivity to various (food- and nonfood-related) rewarding or punishing stimuli, contributing to a sense of instability across individuals with anorexia and bulimia nervosa (2, 3). This instability may contribute to heightened anxiety and intolerance of uncertainty (4). Additionally, as suggested in this study, there may be developmental factors such as altered brain volume in the orbitofrontal cortex that could interfere with the processing of value of food stimuli. We recently published a similarly designed study in adolescents with anorexia nervosa and found similarly larger gyrus rectus volumes, further supporting that this structure could be involved in the pathophysiology of eating disorders (5).

All eating disorder groups share an enlarged gyrus rectus, which could impair a healthy sensory satiety processing. In fact individuals with eating disorders share many similar behaviors but on different scales, and there is overlap in symptoms not only across subgroups but even depending on stage of recovery. For instance, individuals with bulimia nervosa are often able to restrict food intake in-between binge episodes. Furthermore, many individuals who meet criteria for one category of eating disorder will often shift into another

eating disorder during the course of illness, which is consistent with a common vulnerability.

References

1. Kringelbach ML, O'Doherty J, Rolls ET, Andrews C. Activation of the human orbitofrontal cortex to a liquid food stimulus is correlated with its subjective pleasantness. *Cereb Cortex*. 2003; 13:1064–1071. [PubMed: 12967923]
2. Frank GK, Reynolds JR, Shott ME, O'Reilly RC. Altered temporal difference learning in bulimia nervosa. *Biol Psychiatry*. 2011; 70:728–735. [PubMed: 21718969]
3. Jappe LM, Frank GK, Shott ME, Rollin MD, Pryor T, Hagman JO, Yang TT, Davis E. Heightened sensitivity to reward and punishment in anorexia nervosa. *Int J Eat Disord*. 2011; 44:317–324. [PubMed: 21472750]
4. Frank GK, Roblek T, Shott ME, Jappe LM, Rollin MD, Hagman JO, Pryor T. Heightened fear of uncertainty in anorexia and bulimia nervosa. *Int J Eat Disord*. 2012; 45:227–232. [PubMed: 21495057]
5. Frank GK, Shott ME, Hagman JO, Yang TT. Localized brain volume and white matter integrity alterations in adolescent anorexia nervosa. *J Am Acad Child Adolesc Psychiatry*. 2013 in press.