Published in final edited form as:

Bipolar Disord. 2014 May ; 16(3): 230–240. doi:10.1111/bdi.12159.

Seroreactive marker for inflammatory bowel disease and associations with antibodies to dietary proteins in bipolar disorder

Emily G. Severance^a, Kristin L. Gressitt^a, Shuoja Yang^a, Cassie R. Stallings^b, Andrea E. Origoni^b, Crystal Vaughan^b, Sunil Khushalani^b, Armin Alaedini^c, Faith B. Dickerson^b, and Robert H. Yolken^a

^aStanley Division of Developmental Neurovirology, Department of Pediatrics, Johns Hopkins University School of Medicine, 600 N. Wolfe Street, Blalock 1105, Baltimore, MD 21287-4933 U.S.A

^bStanley Research Program, Sheppard Pratt Health System, Baltimore, MD, 21285 U.S.A

^cDepartment of Medicine, Columbia University Medical Center, NY, NY, 10032 U.S.A

Abstract

Objectives—Immune sensitivity to wheat gluteins and bovine milk caseins may affect a subset of individuals with bipolar disorder. Digested byproducts of these foods are exorphins that have the potential to impact brain physiology through action at opioid receptors. Inflammation in the gastrointestinal (GI) tract might accelerate exposure of food antigens to systemic circulation and help explain elevated gluten and casein antibody levels in individuals with bipolar disorder.

Methods—We measured a marker of GI inflammation, anti-*Saccharomyces cerevisiae* antibodies (ASCA), in non-psychiatric controls (n=207), bipolar disorder without a recent onset of psychosis (n=226), and bipolar disorder with a recent onset of psychosis (n=38). We compared ASCA levels to antibodies against gluten, casein, EBV, HSV-1, Influenza A, Influenza B, measles, and *Toxoplasma gondii*.

Results—Elevated ASCA conferred a 3.5 to 4.4-fold increased odds ratio of disease association (age-, race- and gender-corrected multinomial logistic regressions, p 0.00001) that was independent of type of medication received. ASCA correlated with food antibodies in both bipolar groups ($R^2=0.29-0.59$, p 0.0005), and with measles and *T. gondii* IgG in the recent onset psychosis bipolar disorder group ($R^2=0.31-0.36$, p 0.004–0.01).

Conclusions—Elevated seropositivity of a GI-related marker and its association with antibodies to food-derived proteins and self-reported GI symptoms suggests a GI comorbidity in at least a subgroup of individuals with bipolar disorder. Marker seroreactivity may also represent part of an overall heightened activated immune state inherent to this mood disorder.

Correspondence: Emily G. Severance, eseverance@jhmi.edu, tel: 410-614-3918, fax: 410-955-3723.

Disclosures

Robert Yolken is a member of the Stanley Medical Research Institute Board of Directors and Scientific Advisory Board. The terms of this arrangement are being managed by the Johns Hopkins University in accordance with its conflict of interest policies.

None of the other authors report any potential conflicts of interest.

Keywords

Gastrointestinal; environment; infection; immunology; mood disorder; autoimmunity

Introduction

A renewed interest in the role of wheat gluteins and bovine milk caseins in the etiology and pathophysiology of psychiatric disorders is currently underway (1–14). F. Curtis Dohan originally proposed that these food antigens might impact schizophrenia, based on observations of celiac disease in individuals with schizophrenia and on correlations between wartime wheat deficits and diminished schizophrenia hospitalization rates (15–20). Several recent studies indicate that individuals with mood disorders may also be affected by food-related immune activation, particularly during the mania and psychosis-associated phases of the disease (3, 5, 8). The mechanism governing how food antigen peptides might impact the brain and behavior is not known, but is thought to involve the digestion of gluteins and caseins into small peptides that are opioid receptor ligands (21–28). For these peptides to access the brain, they must first be absorbed across gastrointestinal (GI) epithelial and endothelial barriers and enter into systemic circulation.

We recently reported that a marker of GI inflammation was elevated and correlated with levels of antibodies to food antigens in individuals with schizophrenia compared to controls (6). In particular, individuals who were in the early stages of disease and/or who were medication-naïve had the highest levels of GI inflammation. Data from serological, gene expression, and imaging studies implicate that inflammation may be an important pathology in individuals with affective disorders (29–38). Defining a direct relationship between mood disorder symptoms and cytokine dysregulation has been difficult, however, given the fluctuating nature of mood cycles, the anti-inflammatory effects of mood disorder drugs, and the natural dynamic process of inflammatory responses (30). Detection of a localized source of inflammation in the GI tract could help to explain disease-associated systemic inflammation, because a permeable GI tract can lead to translocation of gut microbiota into circulation, thus causing a more generalized inflammatory state (39).

To determine if inflammation present in the GI tract might be a comorbid risk factor for bipolar disorder, we measured anti-*Saccharomyces cerevisiae* antibodies (ASCA) in individuals with bipolar disorder without a recent onset of psychosis, bipolar disorder with a recent onset psychosis and in controls with no history of psychiatric disorders. Measures of serological ASCA are used clinically to aid and differentiate the diagnosis of inflammatory bowel diseases including ulcerative colitis and Crohn's disease (40–44). Our objectives were to evaluate if this marker of GI inflammation was (1) elevated in disease groups compared to controls, (2) accompanied by an anti-food antigen immune response, (3) accompanied by an anti-microbial pathogen immune response, (4) associated with recent onset or lifetime psychosis, (5) prevalent during a particular phase of the bipolar disorder cycle, and (6) correlated with the presence of self-reported GI symptoms.

Materials and methods

Study participants

Study participants were recruited from the Baltimore, Maryland, U.S.A. metro area. One objective of our study design was to enable comparisons between recently diagnosed individuals and those individuals who have had their disease for greater amounts of time. For measures of inflammatory indices that are known to fluctuate over time, the ability to compare individuals in different stages of disease is especially pertinent (6, 8, 9, 32, 45). Toward this end, we recruited 226 individuals diagnosed with bipolar disorder without a recent onset of psychosis, 38 individuals with bipolar disorder and a recent onset of psychosis, and 207 individuals who did not have a history of psychiatric disorders from inpatient and outpatient treatment sites and published announcements. These individuals comprise a cohort that has been previously evaluated in a number of studies (3, 4, 8). The methods for identifying and characterizing individuals of the diagnostic groups according to criteria defined by DSM-IV also have been previously described (3, 4, 8). The DSM-IV diagnoses and sample sizes for the two bipolar disorder groups are listed in Table 1.

Basic demographic data of the control and bipolar disorder groups are shown in Table 2. Diagnostic groups differed significantly in age and race and these variables were included in the multivariate analyses described below.

For the 226 individuals with bipolar disorder without a recent onset of psychosis, inclusion criteria were: diagnosis of type I or type II bipolar disorder or bipolar disorder not otherwise specified according to DSM-IV criteria (46); age between 18 and 65, inclusive; absence of primary diagnosis of substance abuse or dependence over the past three months; absence of any history of intravenous substance abuse; absence of mental retardation; absence of clinically significant medical disorder that would affect cognitive performance such as history of encephalitis or serious head trauma, or any other significant neurological disorder of the central nervous system. While these individuals did not have a recent onset of psychosis defined as an onset of psychotic symptoms for the first time within the past two years, 75.2% had a history of past psychosis, as noted in Table 2.

Additional inclusion criteria for the 38 individuals with bipolar disorder and a recent onset of psychosis were the onset of psychotic symptoms for the first time within the past 24 months, defined as the presence of a positive psychotic symptom of at least moderate severity and lasted through the day for several days or that occurred several times a week; age between 18 and 45, inclusive.

The 207 individuals without a history of psychiatric disorder were screened to rule out current or past psychiatric disorders with the Structured Clinical Interview for DSM-IV Axis I Disorders (47). Control participants were between the ages of 20 and 60, inclusive. Exclusion criteria included the following: any history of intravenous substance abuse; mental retardation; clinically significant medical disorder that would affect cognitive performance; current substance abuse that occurred within the last three months. Immunoassay data from this control group have been previously reported (6).

At the time of interview, cognitive function was evaluated with the Repeatable Battery for the Assessment of Neuropsychological Status (RBANS) Form A (48) and psychiatric symptoms rated according to the Positive and Negative Syndrome Scale (PANSS) (49). For the main bipolar disorder group, symptom severity was assessed with the Hamilton Depression Scale and the Young Mania Rating Scale (50, 51). Mean scores for these tests are shown in Table 2. Hamilton Depression Scale and Young Mania Rating Scale data were not available for the recent onset psychosis bipolar disorder group.

During the interview, clinical information was also gathered, and individuals were asked to report any conditions related to their current GI health. Relevant conditions included constipation, Crohn's disease, diarrhea, Gastroesophageal Reflux Disease, Irritable Bowel Syndrome, lactose intolerance, Norwalk virus, pancreatitis, ulcers, any abdominal surgery, and any abdominal pain. For both bipolar disorder groups, Gastroesophageal Reflux Disease was the most frequently reported GI condition, and these data are shown in Table 3.

Blood samples were obtained by venipuncture, and sera were separated and assessed for antibodies to the antigens in the assays described below.

The studies were approved by the Institutional Review Boards (IRB) of the Sheppard Pratt Health System and the Johns Hopkins Medical Institution following established guidelines. All participants provided written informed consent after study procedures were explained. The work described was carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans.

Laboratory procedures

Anti-*Saccharomyces cerevisiae* IgG antibodies (ASCA) were measured according to the manufacturer's protocol using a commercially available kit (Orgentec, Mainz, Germany). IgG antibodies to bovine milk casein and wheat gluten were measured by ELISAs using previously described methods (6, 7). Whole casein was purchased from Sigma-Aldrich (St. Louis, MO, U.S.A.). Whole gluten was extracted from the wheat cultivar Cheyenne as previously described (12). In brief, for both the casein and gluten immunoassays, plate wells were incubated with 100ng protein in 50µl carbonate buffer (0.05M carbonate-bicarbonate, pH 9.6; Sigma-Aldrich, St. Louis, MO, U.S.A.) overnight at 4°C, and plates were blocked for 1 h at 37°C with 1% (wt/vol) human serum albumin (Sigma-Aldrich, St. Louis, MO, U.S.A.) in PBS. Plates were then incubated with samples diluted 1:200 in PBST for 2h at 37°C. Plates were washed and incubated with peroxidase-conjugated goat-anti-human IgG secondary antibodies for 30min at 37°C (Southern Biotech, Birmingham, AL, U.S.A.). A 2,2'-azino-di-(3-ethylbenzthiazoline-6-sulfonate) and 0.02% hydrogen peroxide solution (KPL Protein Research Products, Gaithersburg, MD, U.S.A.) was added for color development, and absorbance was measured at 405 nm, with a reference wavelength of 490 nm, in an automated microtiter plate reader (Molecular Devices, Menlo Park, CA, U.S.A.).

Commercially available ELISA kits for measuring EBV IgG, Influenza A IgG, Influenza B IgG, Measles IgG, and *T. gondii* IgG were purchased from IBL America (Minneapolis, MN, U.S.A.) and/or IBL International GmbH (Hamburg, Germany). HSV-1 kits were purchased from Focus Diagnostics (Cypress, CA, U.S.A.). IgG levels to these infectious disease agents

had been previously measured for a series of studies at the Stanley Division at Johns Hopkins. IgG levels from these assays have also been previously analyzed for the control individuals in a comparison of these subjects with individuals who had schizophrenia (6).

Statistical analyses

Plate-to-plate variation was corrected by control mean-normalizing each plate so that the control individuals on any particular plate equaled a value of “1”, as previously described (6, 7). Quantitative antibody levels to ASCA were compared among groups with ANOVAs, followed by Bonferroni post-hoc analyses and t-tests. ASCA seropositivity levels were defined based on 90% control values. Multinomial logistic regressions were used to establish age-, sex- and race-corrected odds ratios of ASCA seropositivity for disease association. Multiple linear regressions corrected for age, sex and race were implemented to test for inter-correlations of ASCA, food antigen and infectious disease antigen IgG antibody levels. Age, RBANS, PANSS, Young Mania and Hamilton Depression scores were also evaluated separately for correlations with IgG levels in these multiple linear regressions. Each bipolar disorder group was further stratified according to sex, race, the presence of self-reported GI symptoms, the most recent phasic cycle (manic, mixed, depressive), and the presence or absence of past psychotic features; and the same statistical analyses were applied. Regression coefficient values that equaled or exceeded 0.15 and p values less than 0.05 were considered significant. Statistical analyses were performed with STATA version 12 (STATA Corp LP, College Station, Texas, U.S.A.).

Results

Both bipolar disorder groups had significantly higher ASCA levels than controls (Figure 1, panel A; ANOVA, $F=18.51$, $p = 0.00001$; post-hoc Bonferroni p -value range 0.0001–0.001). ASCA seropositivity was associated with an odds ratio for disease of 4.40 in bipolar disorder without a recent onset of psychosis (95% CI 2.54–7.62, $p = 0.00001$) and of 3.52 in bipolar disorder with recent onset psychosis (95% CI 1.48–8.39, $p = 0.00001$) as compared to controls. ASCA levels within diagnostic groups were not significantly different between males and females or between racial groups (African American and Caucasian/other).

To evaluate if ASCA levels were affected by psychiatric medications, we further subdivided the bipolar disorder groups according to current receipt of lithium, valproate or antipsychotic medication (Figure 1, panel B). No significant differences in ASCA measures were found between receipt or not of each of these medication types (Figure 1, panel B). Of note, all individuals in the bipolar disorder recent onset psychosis group were taking antipsychotic medications at the time of the study visit.

When individuals with bipolar disorder without a recent onset of psychosis were broken down according to the absence or presence of past psychotic features, no significant differences in ASCA levels were detected. There were also no statistically significant differences associated with ASCA with respect to lifetime psychosis when both bipolar disorder groups were pooled.

To determine if there were correlations between ASCA levels and antibodies to food and microbial antigens, multiple linear regression models incorporating age, sex and race were implemented. Anti-casein IgG levels have previously been reported for only a subset of this entire cohort (8), so we first measured IgG to casein and to gluten in the entire set. IgG to food antigens were significantly increased in the two bipolar groups compared to controls (casein: ANOVA, $F=16.03$, $p=0.00001$; post-hoc Bonferroni $p=0.0001$; gluten: ANOVA, $F=18.35$, $p=0.00001$; post-hoc Bonferroni $p=0.0001$). In regression models, ASCA levels were highly correlated to IgG of casein and gluten food antigens in both bipolar groups, but not in controls (Figure 2; p -value range 0.00001–0.0005). Casein and gluten IgG were highly inter-correlated in all three diagnostic groups (Table 4). ASCA was generally not correlated with IgG antibodies directed at four of the microbial pathogens (EBV, HSV-1, Influenza A, Influenza B), with the exception of in the recent onset psychosis bipolar disorder group where ASCA levels were significantly associated with IgG to measles and to *T. gondii* (Table 4). When these groups were broken down according to sex, we found that it was only in females where these microbial pathogen correlations were significant (p -value range 0.004–0.0008).

ASCA was not correlated to age, cognitive scores, or symptom severity scores on the RBANS, PANSS, Young Mania or Hamilton Depression assessments.

The bipolar groups were also subdivided according to type of most recent episode: manic, depressed, or mixed. In bipolar disorder without a recent onset of psychosis, ASCA levels were correlated to food antigen IgG in all three phase subgroups (Table 5; p -value range 0.00001–0.001). In the bipolar disorder with recent onset psychosis group, however, it was only in the manic subgroup where ASCA and the food antigen IgG were correlated. We also further investigated IgG to the microbial pathogens, measles and *T. gondii*, and found that ASCA was well-correlated to measles IgG in the manic and mixed subgroups of the bipolar disorder without recent onset psychosis group and with the manic subgroup of the bipolar disorder with recent onset psychosis group (Table 5; p -value range 0.00001–0.04). ASCA and *T. gondii* IgG were not correlated in any of these subgroup comparisons.

We then examined ASCA levels and correlations to food and microbial antigens according to self-reported GI-related symptomatology. Levels of ASCA were not significantly different between those who reported GI-related conditions compared to those who did not, although a trend for higher ASCA in GI-symptom positive individuals was apparent (bipolar disorder without a recent onset of psychosis: t -test, $t=-1.3587$, $p=0.08$, $\text{mean} \pm \text{SEM} = 1.59 \pm 0.28$ for GI-positive ($n=84$) vs. 1.17 ± 0.13 for GI-negative ($n=81$); bipolar disorder with a recent onset of psychosis, $t=-0.93$, $p=0.17$, 3.56 ± 1.31 for GI-positive ($n=24$), 2.32 ± 0.67 for GI-negative ($n=13$)). In bipolar disorder without a recent onset of psychosis, only in the subgroup reporting GI symptoms were significant correlations of ASCA with casein and gluten IgG maintained (Figure 3; casein: $R^2=0.39$, $p=0.00001$; gluten $R^2=0.59$, $p=0.00001$). In the bipolar disorder with recent onset psychosis group, ASCA levels were correlated with food antigen IgG in both those reporting GI symptoms and those who did not (R^2 range 0.58–0.73, p -value range 0.001–0.05).

Discussion

In this study, we found that a marker of inflammatory bowel disease, ASCA, was elevated in bipolar disorder with and without a recent onset of psychosis compared to non-psychiatric controls. An ASCA seropositive status conferred a 3.5- to 4.4-fold increased odds ratio for disease association. Medication status did not appear to impact measures of this GI marker, as no significant differences in ASCA levels between those receiving and not receiving lithium, valproate or antipsychotics were detected. Bipolar disorder-related ASCA levels were significantly greater than those found previously in schizophrenia (data not shown, but ANOVA $p = 0.001$) (6). Further studies are required to fully disentangle if the GI locale represents a source of inflammation or if it is a pathology stemming from a generalized state of immune activation intrinsic to bipolar disorder.

We demonstrated a significant disease association of the ASCA marker with immune reactivity to wheat glutes and bovine milk caseins. In particular, these correlations were significant in individuals of the main bipolar disorder group who reported GI symptoms compared to those who did not have GI symptoms. Together, these findings may support hypotheses related to pre-existing, compromised epithelial barriers; however, other explanations are also possible including a directed interaction of food exorphins with tight junction proteins or epithelial cell transcytosis (52–59). Incomplete digestion of these foods could also generate novel antigens that prime the immune system. We found previously that sera from individuals with bipolar disorder and schizophrenia did not recognize the same epitopes of casein and gluten that were recognized by controls (7, 8, 12). Differential processing of these food proteins may result from alterations in the resident gut microbiota or dysfunction of digestive peptidases. We can speculate that if GI inflammation does indeed cause GI permeability, not only would partially digested gluten and casein peptides enter into the bloodstream, but resident gut microbiota would translocate into systemic circulation. Maes and colleagues have repeatedly demonstrated an association of bacterial translocation with major depressive disorder (60–62), and we are exploring this hypothesis in schizophrenia (63).

Because individuals in our control group may not be afflicted with the same degree of GI inflammation as cases, the constricted quantitative range of control ASCA values may make it difficult to statistically detect its correlations with other variables such as antibodies to food peptides; however, inter-correlations of the food-related antibodies were quite strong in all diagnostic groups. As such, the lack of association of ASCA markers with antibodies to dietary proteins in the controls may reflect mechanisms of immune activation in cases which are alternative to the leaky gut hypothesis. For example, autoimmune conditions might contribute to a heightened immune response and to inter-relatedness between the food-related antibodies, even in the absence of immune activation by specific antigens. Such a pattern may ensue due to an early developmental assault or genetic predisposition affecting immune dysfunction. Currently, there is some evidence supporting an autoimmune component of mood disorders, but results thus far are fairly mixed (63–67). Interestingly, although ASCA is predominately a marker for GI-related diseases, this marker was found to be elevated in patients with a variety of non-GI-related autoimmune conditions such as systemic lupus erythematosus, Graves' disease, cryoglobulinemia and vasculitis (68).

Our cohort included a group of individuals with bipolar disorder who also had a recent onset of psychosis. The small sample size of this group (n=38) limits the extent to which comparisons can be interpreted, but some interesting trends are noteworthy for future research. In contrast to our findings in the main bipolar disorder group, ASCA was correlated with antibodies to food antigens in both those who reported GI symptoms and those who did not, a possible indication of some level of non-specificity of the immune response in the recent onset bipolar disorder group. Furthermore, ASCA levels exhibited correlations with several non-food-related antigens as well, measles and *T. gondii*, but the positive IgG response was not uniform across the antigens tested. For these comparisons, however, statistical power issues might impact our sensitivity in detecting seropositivity. Of note, *T. gondii* and measles have been implicated in bipolar disorder or recent onset psychosis previously, and both have been linked, sometimes controversially, to inflammation processes that affect the GI tract in humans and in experimental animal models (6, 32, 69–82). Although our findings require further scrutiny and larger sample sizes, the possibility of a more generalized inflammatory nature inherent to earlier stages of disease can not be discounted.

In addition to the small sample sizes of our bipolar disorder with recent onset of psychosis group, our study has a number of limitations. The study design is cross-sectional and only allows for speculation regarding mechanisms that might underly our results. With respect to age, the recent onset psychosis group is significantly younger and the established bipolar disorder group significantly older than controls, yet both groups exhibited increased ASCA levels compared to controls, suggesting that age here may not be a strong confounding factor. Furthermore, our GI measures, the ASCA marker and self-reported GI disturbances, would benefit from an additional biological index such as a stool sample or histopathological examination of the GI tract. Blood draws were not standardized nor taken under fasting conditions, although our measures were predominately IgG-based, the levels of which are relatively stable over time and especially over the course of one day. In sub-analyses, we rely on stratifications according to most recent episode (mania, depression, mixed) which results in grouping of asymptomatic with symptomatic individuals. We attempted to address this by examining correlations of ASCA with symptom severity scale scores (Young Mania and Hamilton Depression) with the rationale that symptomatic individuals would score at the polar ranges of these scales. As indicated earlier, no correlations of ASCA with symptom severity scales were observed. Similarly, we tried to further evaluate an effect of lifetime psychosis on ASCA levels by combining the bipolar disorder groups, but no significant differences in marker levels or inter-correlations were detected.

In conclusion, this study offers strong preliminary evidence for a role of the GI tract in the inflammatory pathology of bipolar disorder. The serological biomarkers and their inter-associations may help to identify a subgroup of individuals with bipolar disorder who may be afflicted with GI- and food-based sensitivities as well as those who may be prone to a generalized inflammatory state. Treatment strategies that involve dietary modifications, anti-inflammatory agents and/or microbiota modulations should be evaluated for effects on symptom improvement in carefully controlled clinical trials.

Acknowledgments

We thank Dr. Donald D. Kasarda of the U.S. Department of Agriculture for providing us with the wheat flour used for gluten extraction. This work was supported by the Brain and Behavior Research Foundation where Dr. Severance is a Scott-Gentle Foundation Young Investigator, and by the Stanley Medical Research Institute.

References

1. Cascella NG, Kryszak D, Bhatti B, Gregory P, Kelly DL, Mc Evoy JP, et al. Prevalence of celiac disease and gluten sensitivity in the United States clinical antipsychotic trials of intervention effectiveness study population. *Schizophr Bull.* 2011; 37:94–100. [PubMed: 19494248]
2. Cascella NG, Santora D, Gregory P, Kelly DL, Fasano A, Eaton WW. Increased Prevalence of Transglutaminase 6 Antibodies in Sera From Schizophrenia Patients. *Schizophr Bull.* 2012 Epub ahead of print.
3. Dickerson F, Stallings C, Origoni A, Vaughan C, Khushalani S, Alaedini A, et al. Markers of gluten sensitivity and celiac disease in bipolar disorder. *Bipolar Disord.* 2011; 13:52–8. [PubMed: 21320252]
4. Dickerson F, Stallings C, Origoni A, Vaughan C, Khushalani S, Leister F, et al. Markers of gluten sensitivity and celiac disease in recent-onset psychosis and multi-episode schizophrenia. *Biol Psychiatry.* 2010; 68:100–4. [PubMed: 20471632]
5. Dickerson F, Stallings C, Origoni A, Vaughan C, Khushalani S, Yolken R. Markers of gluten sensitivity in acute mania: A longitudinal study. *Psychiatry Res.* 2012; 196:68–71. [PubMed: 22386570]
6. Severance EG, Alaedini A, Yang S, Halling M, Gressitt KL, Stallings CR, et al. Gastrointestinal inflammation and associated immune activation in schizophrenia. *Schizophr Res.* 2012; 138:48–53. [PubMed: 22446142]
7. Severance EG, Dickerson FB, Halling M, Krivogorsky B, Haile L, Yang S, et al. Subunit and whole molecule specificity of the anti-bovine casein immune response in recent onset psychosis and schizophrenia. *Schizophr Res.* 2010; 118:240–7. [PubMed: 20071146]
8. Severance EG, Dupont D, Dickerson FB, Stallings CR, Origoni AE, Krivogorsky B, et al. Immune activation by casein dietary antigens in bipolar disorder. *Bipolar Disord.* 2010; 12:834–42. [PubMed: 21176030]
9. Severance EG, Gressitt KL, Halling M, Stallings CR, Origoni AE, Vaughan C, et al. Complement C1q formation of immune complexes with milk caseins and wheat gluteins in schizophrenia. *Neurobiol Dis.* 2012; 48:447–53. [PubMed: 22801085]
10. Karlsson H, Blomstrom A, Wicks S, Yang S, Yolken RH, Dalman C. Maternal Antibodies to Dietary Antigens and Risk for Non-affective Psychosis in Offspring. *Am J Psychiatry.* 2012; 169:625–32. [PubMed: 22535227]
11. Niebuhr DW, Li Y, Cowan DN, Weber NS, Fisher JA, Ford GM, et al. Association between bovine casein antibody and new onset schizophrenia among US military personnel. *Schizophr Res.* 2011; 128:51–5. [PubMed: 21376538]
12. Samaroo D, Dickerson F, Kasarda DD, Green PH, Briani C, Yolken RH, et al. Novel immune response to gluten in individuals with schizophrenia. *Schizophr Res.* 2010; 118:248–55. [PubMed: 19748229]
13. Jackson J, Eaton W, Cascella N, Fasano A, Warfel D, Feldman S, et al. A gluten-free diet in people with schizophrenia and anti-tissue transglutaminase or anti-gliadin antibodies. *Schizophr Res.* 2012; 140:262–3. [PubMed: 22771303]
14. Whiteley P, Shattock P, Knivsberg AM, Seim A, Reichelt KL, Todd L, et al. Gluten- and casein-free dietary intervention for autism spectrum conditions. *Front Hum Neurosci.* 2012; 6:344. [PubMed: 23316152]
15. Dohan FC. Cereals and schizophrenia data and hypothesis. *Acta Psychiatr Scand.* 1966; 42:125–52. [PubMed: 5335008]
16. Dohan FC. Wheat “consumption” and hospital admissions for schizophrenia during World War II. A preliminary report. *Am J Clin Nutr.* 1966; 18:7–10. [PubMed: 5900428]

17. Dohan FC. Wartime changes in hospital admissions for schizophrenia. A comparison of admission for schizophrenia and other psychoses in six countries during World War II. *Acta Psychiatr Scand.* 1966; 42:1–23. [PubMed: 5959639]
18. Dohan FC. Is celiac disease a clue to the pathogenesis of schizophrenia? *Ment Hyg.* 1969; 53:525–9. [PubMed: 5344577]
19. Dohan FC, Grasberger JC, Lowell FM, Johnston HT Jr, Arbogast AW. Relapsed schizophrenics: more rapid improvement on a milk- and cereal-free diet. *Br J Psychiatry.* 1969; 115:595–6. [PubMed: 5820122]
20. Dohan FC. Coeliac disease and schizophrenia. *Lancet.* 1970; 1:897–8. [PubMed: 4191543]
21. Dohan FC. Schizophrenia and neuroactive peptides from food. *Lancet.* 1979; 1:1031. [PubMed: 86744]
22. Dohan FC. Schizophrenia, celiac disease, gluten antibodies, and the importance of beta. *Biol Psychiatry.* 1981; 16:1115–7. [PubMed: 7349623]
23. Dohan FC. Genetic hypothesis of idiopathic schizophrenia: its exorphin connection. *Schizophr Bull.* 1988; 14:489–94. [PubMed: 2851166]
24. Reichelt KL, Reichelt WH, Stensrud M. The Role of Peptides in Schizophrenia. *J Neurochem.* 1995; 65:S43-S.
25. Reichelt KL, Seim AR, Reichelt WH. Could schizophrenia be reasonably explained by Dohan's hypothesis on genetic interaction with a dietary peptide overload? *Prog Neuropsychopharmacol Biol Psychiatry.* 1996; 20:1083–114. [PubMed: 8938813]
26. Reichelt KL, Stensrud M. Increase in urinary peptides prior to the diagnosis of schizophrenia. *Schizophr Res.* 1998; 34:211–3. [PubMed: 9850988]
27. Reichelt KL. Exorphins in Schizophrenia and Autism. *J Neurochem.* 1994; 63:S86-S.
28. Kaminski S, Cieslinska A, Kostyra E. Polymorphism of bovine beta-casein and its potential effect on human health. *J Appl Genet.* 2007; 48:189–98. [PubMed: 17666771]
29. Brietzke E, Stertz L, Fernandes BS, Kauer-Sant'anna M, Mascarenhas M, Escosteguy Vargas A, et al. Comparison of cytokine levels in depressed, manic and euthymic patients with bipolar disorder. *J Affect Disord.* 2009; 116:214–7. [PubMed: 19251324]
30. Goldstein BI, Kemp DE, Soczynska JK, McIntyre RS. Inflammation and the phenomenology, pathophysiology, comorbidity, and treatment of bipolar disorder: a systematic review of the literature. *J Clin Psychiatry.* 2009; 70:1078–90. [PubMed: 19497250]
31. Hope S, Dieset I, Agartz I, Steen NE, Ueland T, Melle I, et al. Affective symptoms are associated with markers of inflammation and immune activation in bipolar disorders but not in schizophrenia. *J Psychiatr Res.* 2011; 45:1608–16. [PubMed: 21889167]
32. Leboyer M, Soreca I, Scott J, Frye M, Henry C, Tamouza R, et al. Can bipolar disorder be viewed as a multi-system inflammatory disease? *J Affect Disord.* 2012; 141:1–10. [PubMed: 22497876]
33. Tsai SY, Chung KH, Wu JY, Kuo CJ, Lee HC, Huang SH. Inflammatory markers and their relationships with leptin and insulin from acute mania to full remission in bipolar disorder. *J Affect Disord.* 2012; 136:110–6. [PubMed: 21962564]
34. Padmos RC, Van Baal GC, Vonk R, Wijkhuijs AJ, Kahn RS, Nolen WA, et al. Genetic and environmental influences on pro-inflammatory monocytes in bipolar disorder: a twin study. *Arch Gen Psychiatry.* 2009; 66:957–65. [PubMed: 19736352]
35. Savitz JB, Drevets WC. Neuroreceptor imaging in depression. *Neurobiol Dis.* 2012
36. Stertz L, Magalhaes PV, Kapczinski F. Is bipolar disorder an inflammatory condition? The relevance of microglial activation. *Curr Opin Psychiatry.* 2013; 26:19–26. [PubMed: 23196997]
37. Drexhage RC, Hoogenboezem TH, Versnel MA, Berghout A, Nolen WA, Drexhage HA. The activation of monocyte and T cell networks in patients with bipolar disorder. *Brain Behav Immun.* 2011; 25:1206–13. [PubMed: 21443944]
38. Drexhage RC, Weigelt K, van Beveren N, Cohen D, Versnel MA, Nolen WA, et al. Immune and neuroimmune alterations in mood disorders and schizophrenia. *Int Rev Neurobiol.* 2011; 101:169–201. [PubMed: 22050852]

39. Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med*. 2006; 12:1365–71. [PubMed: 17115046]
40. Ashorn S, Valineva T, Kaukinen K, Ashorn M, Braun J, Raukola H, et al. Serological responses to microbial antigens in celiac disease patients during a gluten-free diet. *J Clin Immunol*. 2009; 29:190–5. [PubMed: 18987962]
41. Desplat-Jego S, Johanet C, Escande A, Goetz J, Fabien N, Olsson N, et al. Update on Anti-Saccharomyces cerevisiae antibodies, anti-nuclear associated anti-neutrophil antibodies and antibodies to exocrine pancreas detected by indirect immunofluorescence as biomarkers in chronic inflammatory bowel diseases: results of a multicenter study. *World J Gastroenterol*. 2007; 13:2312–8. [PubMed: 17511029]
42. Kotze LM, Nisihara RM, Utiyama SR, Kotze PG, Theiss PM, Olandoski M. Antibodies anti-Saccharomyces cerevisiae (ASCA) do not differentiate Crohn's disease from celiac disease. *Arq Gastroenterol*. 2010; 47:242–5. [PubMed: 21140083]
43. Mallant-Hent R, Mary B, von Blomberg E, Yuksel Z, Wahab PJ, Gundy C, et al. Disappearance of anti-Saccharomyces cerevisiae antibodies in coeliac disease during a gluten-free diet. *Eur J Gastroenterol Hepatol*. 2006; 18:75–8. [PubMed: 16357623]
44. Oshitani N, Hato F, Matsumoto T, Jinno Y, Sawa Y, Hara J, et al. Decreased anti-Saccharomyces cerevisiae antibody titer by mesalazine in patients with Crohn's disease. *J Gastroenterol Hepatol*. 2000; 15:1400–3. [PubMed: 11197050]
45. Suvisaari J, Mantere O. Inflammation Theories In Psychotic Disorders: A Critical Review. *Infect Disord Drug Targets*. 2013 Epub ahead of print.
46. DSM-IV. Diagnostic and statistical manual of mental disorders : DSM-IV. 4. Washington, DC: American Psychiatric Association; 1994.
47. First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. Structured Clinical Interview for DSM-IV Axis I Disorders - Non-patient Edition (SCID I/NP). New York: Biometrics Research, New York State Psychiatric Institute; 1998.
48. Randolph, C. RBANS Manual - Repeatable Battery for the Assessment of Neuropsychological Status. San Antonio: Psychological Corporation; 1998.
49. Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull*. 1987; 13:261–76. [PubMed: 3616518]
50. Hamilton M. Development of a rating scale for primary depressive illness. *Br J Soc Clin Psychol*. 1967; 6:278–96. [PubMed: 6080235]
51. Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: reliability, validity and sensitivity. *Br J Psychiatry*. 1978; 133:429–35. [PubMed: 728692]
52. Menard S, Lebreton C, Schumann M, Matysiak-Budnik T, Dugave C, Bouhnik Y, et al. Paracellular versus transcellular intestinal permeability to gliadin peptides in active celiac disease. *Am J Pathol*. 2012; 180:608–15. [PubMed: 22119716]
53. Matysiak-Budnik T, Candalh C, Dugave C, Namane A, Cellier C, Cerf-Bensussan N, et al. Alterations of the intestinal transport and processing of gliadin peptides in celiac disease. *Gastroenterology*. 2003; 125:696–707. [PubMed: 12949716]
54. Heyman M, Abed J, Lebreton C, Cerf-Bensussan N. Intestinal permeability in coeliac disease: insight into mechanisms and relevance to pathogenesis. *Gut*. 2012; 61:1355–64. [PubMed: 21890812]
55. Lammers KM, Lu R, Brownley J, Lu B, Gerard C, Thomas K, et al. Gliadin induces an increase in intestinal permeability and zonulin release by binding to the chemokine receptor CXCR3. *Gastroenterology*. 2008; 135:194–204. e3. [PubMed: 18485912]
56. Tripathi A, Lammers KM, Goldblum S, Shea-Donohue T, Netzel-Arnett S, Buzza MS, et al. Identification of human zonulin, a physiological modulator of tight junctions, as prehaptoglobin-2. *Proc Natl Acad Sci U S A*. 2009; 106:16799–804. [PubMed: 19805376]
57. Drago S, El Asmar R, Di Pierro M, Grazia Clemente M, Tripathi A, Sapone A, et al. Gliadin, zonulin and gut permeability: Effects on celiac and non-celiac intestinal mucosa and intestinal cell lines. *Scand J Gastroenterol*. 2006; 41:408–19. [PubMed: 16635908]

58. Sander GR, Cummins AG, Henshall T, Powell BC. Rapid disruption of intestinal barrier function by gliadin involves altered expression of apical junctional proteins. *FEBS Lett.* 2005; 579:4851–5. [PubMed: 16099460]
59. Rallabhandi P. Gluten and celiac disease--an immunological perspective. *J AOAC Int.* 2012; 95:349–55. [PubMed: 22649918]
60. Maes M, Kubera M, Leunis JC, Berk M, Geffard M, Bosmans E. In depression, bacterial translocation may drive inflammatory responses, oxidative and nitrosative stress (O&NS), and autoimmune responses directed against O&NS-damaged neopeptides. *Acta Psychiatr Scand.* 2013; 127:344–54. [PubMed: 22900942]
61. Maes M, Kubera M, Leunis JC, Berk M. Increased IgA and IgM responses against gut commensals in chronic depression: Further evidence for increased bacterial translocation or leaky gut. *J Affect Disord.* 2012; 141:55–62. [PubMed: 22410503]
62. Maes M, Kubera M, Leunis JC. The gut-brain barrier in major depression: intestinal mucosal dysfunction with an increased translocation of LPS from gram negative enterobacteria (leaky gut) plays a role in the inflammatory pathophysiology of depression. *Neuro Endocrinol Lett.* 2008; 29:117–24. [PubMed: 18283240]
63. Severance EG, Gressitt KL, Stallings CR, Origoni AE, Khushalani S, Leweke FM, et al. Discordant patterns of bacterial translocation markers and implications for innate immune imbalances in schizophrenia. *Schizophr Res.* 2013 Epub ahead of print.
64. Sidhom O, Laadhar L, Zitouni M, Ben Alaya N, Raftafi R, Kallel-Sellami M, et al. Spectrum of autoantibodies in Tunisian psychiatric inpatients. *Immunol Invest.* 2012; 41:538–49. [PubMed: 22612859]
65. Eaton WW, Pedersen MG, Nielsen PR, Mortensen PB. Autoimmune diseases, bipolar disorder, and non-affective psychosis. *Bipolar Disord.* 2010; 12:638–46. [PubMed: 20868462]
66. Vonk R, van der Schot AC, Kahn RS, Nolen WA, Drexhage HA. Is autoimmune thyroiditis part of the genetic vulnerability (or an endophenotype) for bipolar disorder? *Biol Psychiatry.* 2007; 62:135–40. [PubMed: 17141745]
67. Padmos RC, Bekris L, Knijff EM, Tiemeier H, Kupka RW, Cohen D, et al. A high prevalence of organ-specific autoimmunity in patients with bipolar disorder. *Biol Psychiatry.* 2004; 56:476–82. [PubMed: 15450782]
68. Ben-Ami Shor D, Orbach H, Boaz M, Altman A, Anaya J, Bizzaro N, et al. Gastrointestinal-associated autoantibodies in different autoimmune diseases. *American Journal of clinical and experimental immunology.* 2012; 1:49–55. [PubMed: 23885314]
69. Hamdani N, Daban-Huard C, Lajnef M, Richard JR, Delavest M, Godin O, et al. Relationship between *Toxoplasma gondii* infection and bipolar disorder in a French sample. *J Affect Disord.* 2013; 148:444–8. [PubMed: 23273549]
70. Bereswill S, Munoz M, Fischer A, Plickert R, Haag LM, Otto B, et al. Anti-inflammatory effects of resveratrol, curcumin and simvastatin in acute small intestinal inflammation. *PLoS One.* 2010; 5:e15099. [PubMed: 21151942]
71. Bernstein CN, Rawsthorne P, Blanchard JF. Population-based case-control study of measles, mumps, and rubella and inflammatory bowel disease. *Inflamm Bowel Dis.* 2007; 13:759–62. [PubMed: 17230540]
72. D'Souza Y, Dionne S, Seidman EG, Bitton A, Ward BJ. No evidence of persisting measles virus in the intestinal tissues of patients with inflammatory bowel disease. *Gut.* 2007; 56:886–8. [PubMed: 17519494]
73. Dickerson F, Stallings C, Origoni A, Copp C, Khushalani S, Yolken R. Antibodies to measles in individuals with recent onset psychosis. *Schizophr Res.* 2010; 119:89–94. [PubMed: 20051313]
74. Craven M, Egan CE, Dowd SE, McDonough SP, Dogan B, Denkers EY, et al. Inflammation drives dysbiosis and bacterial invasion in murine models of ileal Crohn's disease. *PLoS One.* 2012; 7:e41594. [PubMed: 22848538]
75. Erridge C, Duncan SH, Bereswill S, Heimesaat MM. The induction of colitis and ileitis in mice is associated with marked increases in intestinal concentrations of stimulants of TLRs 2, 4, and 5. *PLoS One.* 2010; 5:e9125. [PubMed: 20161736]

76. Hansen TS, Jess T, Vind I, Elkjaer M, Nielsen MF, Gomborg M, et al. Environmental factors in inflammatory bowel disease: a case-control study based on a Danish inception cohort. *J Crohns Colitis*. 2011; 5:577–84. [PubMed: 22115378]
77. Lidar M, Langevitz P, Shoenfeld Y. The role of infection in inflammatory bowel disease: initiation, exacerbation and protection. *Isr Med Assoc J*. 2009; 11:558–63. [PubMed: 19960852]
78. Liesenfeld O. Immune responses to *Toxoplasma gondii* in the gut. *Immunobiology*. 1999; 201:229–39. [PubMed: 10631572]
79. Munoz M, Heimesaat MM, Danker K, Struck D, Lohmann U, Plickert R, et al. Interleukin (IL)-23 mediates *Toxoplasma gondii*-induced immunopathology in the gut via matrixmetalloproteinase-2 and IL-22 but independent of IL-17. *J Exp Med*. 2009; 206:3047–59. [PubMed: 19995958]
80. Schreiner M, Liesenfeld O. Small intestinal inflammation following oral infection with *Toxoplasma gondii* does not occur exclusively in C57BL/6 mice: review of 70 reports from the literature. *Mem Inst Oswaldo Cruz*. 2009; 104:221–33. [PubMed: 19430647]
81. Wakefield AJ, Pittilo RM, Sim R, Cosby SL, Stephenson JR, Dhillon AP, et al. Evidence of persistent measles virus infection in Crohn's disease. *J Med Virol*. 1993; 39:345–53. [PubMed: 8492105]
82. Carter CJ. Toxoplasmosis and Polygenic Disease Susceptibility Genes: Extensive *Toxoplasma gondii* Host/Pathogen Interactome Enrichment in Nine Psychiatric or Neurological Disorders. *J Pathog*. 2013; 2013:965046. [PubMed: 23533776]

Figure 1.

Quantitative ASCA IgG levels according to diagnosis and medication status.

Panel A. ASCA IgG levels were significantly elevated in bipolar disorder (BP) without a recent onset of psychosis (ROP) and BP with ROP compared to controls (CO). Panel B. ASCA IgG levels between individuals who were medication-positive (+) and medication-negative (–) were not significantly different. LITH refers to lithium; VAL refers to valproate; AP refers to antipsychotic.

Figure 2.

ASCA IgG correlates with food antigen IgG in bipolar disorder groups, but not controls. Gluten data points are shown with an “x” and a solid regression line; casein data points are shown with a closed “o” and a dashed regression line. Multiple linear regressions corrected for age, sex and race were used to generate the listed p-values. “NS” refers to not significant.

Figure 3.

Correlations of ASCA and food antigen IgG levels according to self-reported symptoms of gastrointestinal discomfort. IgG against food antigens is correlated to ASCA in individuals with bipolar disorder without a recent onset of psychosis who reported GI symptoms (+GI symptoms), but not in those who did not report these symptoms. In bipolar disorder with a recent onset of psychosis, ASCA and food antigen IgG were correlated in both those who reported GI symptoms and those who did not. Gluten data points are shown with an “x” and a solid regression line; casein data points are shown with a closed “o” and dashed regression

line. Multiple linear regressions corrected for age, sex and race were used to generate the listed p-values. “NS” refers to not significant.

Table 1

Diagnostic subtypes of bipolar disorder in the study population

Diagnosis	Bipolar disorder w/o ROP ^a		Bipolar disorder w/ ROP	
	n	%	n	%
Bipolar I Disorder, Most Recent Episode Manic	89	39.4%	20	52.6%
Bipolar I Disorder, Most Recent Episode Depressed	53	23.5%	10	26.3%
Bipolar I Disorder, Most Recent Episode Mixed	59	26.1%	6	15.8%
Bipolar Disorder, Not Otherwise Specified (NOS)	1	0.4%	0	0.0%
Bipolar II Disorder	24	10.6%	2	5.3%
	226		38	

^aROP refers to recent onset psychosis

Table 2

Demographic information

		Controls n=207	Bipolar disorder w/o ROP ^a n=226	Bipolar disorder w/ ROP n=38
Duration of illness	Mean years±SEM ^b	-	20.03±0.82	0.90±0.42
Age	Mean years±SEM	32.07±0.80	37.81±0.84 ^c	26.69±1.31 ^d
Males	n (%)	56 (27.1%)	68 (30.1%)	12 (31.6%)
African American	n (%)	66 (31.9%)	48 (21.2%) ^e	9 (23.7%)
RBANS ^f	Mean score±SEM	88.80±0.82	79.23±0.95 ^g	78.63±2.02 ^h
PANSS ⁱ	Mean score±SEM	-	69.35±0.95	70.71±2.47
Y Mania ^j	Mean score±SEM	-	13.40±0.64	-
Hamilton Depression	Mean score±SEM	-	19.97±0.71	-
History of Psychotic Features	n (%)	0 (0%)	170 (75.2%)	38 (100%)

^aROP refers to recent onset psychosis^bSEM refers to standard error of the mean^ct=-4.94, p 0.00001^dt=2.77, p 0.006^echi²=6.31, p 0.012^fRBANS refers to Repeatable Battery for the Assessment of Neuropsychological Status^gt=7.54, p 0.00001^ht=4.8, p 0.00001ⁱPANSS refers to Positive and Negative Syndrome Scale^jY mania refers to Young Mania Rating Scale

Table 3

Gastrointestinal conditions self-reported by individuals with bipolar disorder

	Bipolar disorder w/o ROP ^a	Bipolar disorder w/ ROP
	n=84 ^b	n=13 ^c
Constipation	11 (13.1%)	2 (15.4%)
Diarrhea	3 (3.6%)	1 (7.7%)
Gastroesophageal reflux disease	35 (41.7%)	6 (46.2%)
Crohn's disease or Irritable bowel syndrome	5 (6.0%)	1 (7.7%)
Ulcers	1 (1.2%)	0 (0.0%)
Any abdominal surgery	19 (22.6%)	1 (7.7%)
Any abdominal pain	4 (4.8%)	0 (0.0%)
Other GI conditions	6 (7.1%)	2 (15.4%)

^aROP refers to recent onset psychosis^bInformation regarding absence or presence of GI conditions was collected for n=165^cInformation regarding absence or presence of GI conditions was collected for n=37

Table 4

Correlations of food antigen and microbial pathogen IgG with GI inflammation

		Controls	Bipolar disorder w/o ROP ^a	Bipolar disorder w/ ROP
Food antigens				
ASCA:Casein	n	207	226	38
	R ²	0.04	0.29	0.59
	p	0.12	0.00001	0.00001
ASCA:Gluten	n	207	226	38
	R ²	0.07	0.44	0.45
	p	0.005	0.00001	0.0005
Casein:Gluten	n	207	226	38
	R ²	0.71	0.51	0.77
	p	0.00001	0.00001	0.00001
Microbial pathogens				
EBV	n	207	226	38
	R ²	0.008	0.03	0.03
	p	0.81	0.13	0.91
HSV-1	n	207	220	12
	R ²	0.008	0.04	0.22
	p	0.81	0.09	0.74
Influenza A	n	207	226	38
	R ²	0.01	0.04	0.13
	p	0.67	0.04	0.30
Influenza B	n	207	226	38
	R ²	0.008	0.04	0.16
	p	0.82	0.08	0.21
Measles	n	207	225	38
	R ²	0.01	0.11	0.31^b
	p	0.73	0.0001	0.01
<i>T. gondii</i>	n	207	222	38
	R ²	0.008	0.08	0.36^c
	p	0.8	0.002	0.004

^aROP refers to recent onset psychosis^bFemales: R²=0.45, p 0.004; males NS^cFemales: R²=0.53, p 0.0008; males NS

Table 5

Inter-correlations of GI inflammation, food antigen IgG and microbial pathogen IgG according to bipolar disorder cycle phase

		Bipolar disorder w/o ROP ^a			Bipolar disorder w/ ROP		
		Mania	Depressed	Mixed	Mania	Depressed	Mixed
Food antigens	n ^b	89	53	59	20	10	6
ASCA:Casein	R ²	0.35	0.31	0.47	0.75	0.28	0.99
	p	0.00001	0.001	0.00001	0.0002	0.74	0.16
ASCA:Gluten	R ²	0.44	0.60	0.35	0.80	0.42	0.87
	p	0.00001	0.00001	0.0001	0.00001	0.52	0.52
Casein:Gluten	R ²	0.33	0.72	0.55	0.88	0.88	0.97
	p	0.00001	0.00001	0.00001	0.00001	0.02	0.27
Microbial pathogens							
Measles	R ²	0.25	0.14	0.33	0.48	0.17	0.73
	p	0.0001	0.14	0.0002	0.04	0.89	0.71
<i>T. gondii</i>	R ²	0.06	0.15	0.15	0.35	0.63	0.64
	p	0.25	0.12	0.07	0.14	0.22	0.79

^aROP refers to recent onset psychosis

^bIndividuals with a diagnosis of Bipolar II Disorder or NOS were not included in these analyses.