Published in final edited form as:

Clin Neurophysiol. 2014 August ; 125(8): 1682–1688. doi:10.1016/j.clinph.2013.12.102.

Deficits in startle-evoked arm movements increase with impairment following stroke

Claire Fletcher Honeycutt¹ and Eric Jon Perreault^{1,2,3}

¹Sensory Motor Performance Program, Rehabilitation Institute of Chicago, Chicago, IL, USA

²Department of Biomedical Engineering, Northwestern University, Evanston, IL, USA

³Department of Physical Medicine and Rehabilitation, Northwestern University, Chicago, IL, USA

Abstract

Objective—The startle reflex elicits involuntary release of planned movements (startReact). Following stroke, startReact flexion movements are intact but startReact extension movements are impaired by task-inappropriate flexor activity impeding arm extension. Our objective was to quantify deficits in startReact elbow extension movements, particularly how these deficits are influenced by impairment.

Methods—Data were collected in 8 stroke survivors performing elbow extension following two non-startling acoustic stimuli representing “get ready” and “go” respectively. Randomly, the “go” was replaced with a startling acoustic stimulus. We hypothesized that task-inappropriate flexor activity originates from unsuppressed classic startle reflex. We expected that increasing damage to the cortex (increasing impairment) would relate to increasing task-inappropriate flexor activity causing poor elbow extension movement and target acquisition.

Results—Task-inappropriate flexor activity increased with impairment resulting in larger flexion deflections away from the subjects’ intended target corresponding to decreased target acquisition.

Conclusions—We conclude that the task-inappropriate flexor activity likely results from cortical or corticospinal damage leading to an unsuppressed or hypermetric classic startle reflex that interrupts startReact elbow extension.

Significance—Given startReact’s functional role in compensation during environmental disturbances, our results may have important implications for our understanding deficits in stroke survivor’s response to unexpected environmental disturbances.

Keywords

Startle; stroke; startReact; reaching

© 2013 International Federation of Clinical Neurophysiology. Published by Elsevier Ireland Ltd. All rights reserved.

Corresponding Author: Claire Honeycutt, Rehabilitation Institute of Chicago, 345 E. Superior Street, SMPP, Suite 1406, Chicago, IL 60611, USA, Tel: +1-312.238.1404, FAX: +1-312.238.2208, claire.honeycutt@gmail.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

A unique property of the startle reflex is its ability to involuntarily elicit pre-planned movements throughout the entire arm (Carlsen et al., 2004b, Carlsen et al., 2011, Honeycutt et al., 2013, Rothwell et al., 2002, Valls-Solé et al., 2008, Valls-Solé et al., 1999). When a startling acoustic stimulus is presented in the absence of a movement plan, the classic startle reflex triggers brief co-contraction of muscles resulting in the individual assuming a protective stance – arm flexion in the upper limb. However, when a subject is in a state of movement preparation, a startling acoustic stimulus involuntarily elicits the prepared movement (Carlsen et al., 2004b, Rothwell et al., 2002, Valls-Solé et al., 1999, Valls-Solé et al., 1995). This phenomenon has been called startReact (Valls-Solé et al., 1999). Different from the classic startle response, startReact movements are not significantly different from voluntarily executed movements and reflect the sophistication of voluntarily planned movements in terms of acceleration and target accuracy (Carlsen et al., 2004a, Carlsen et al., 2004b). While the classic startle response results in the generation of a protective, crouched posture, the startReact response appears to be more functionally relevant; specifically, it has been implicated in the ability to actively resist perturbations of the arm and whole-body. We recently demonstrated that arm perturbations, like startling acoustic stimuli, elicit startReact movements (Ravichandran et al., 2013) indicating that the functional role of this reflex likely participates in the effective and efficient response to an environmental perturbations. The startle reflex is also triggered during whole-body perturbations indicating startReact movements may also be functional during balance challenges (Blouin et al., 2006, Campbell et al., 2012, Oude Nijhuis et al., 2010, Siegmund et al., 2008).

The startReact phenomenon was recently shown to improve elbow flexion movements in stroke survivors (Honeycutt and Perreault, 2012) but startReact extension movements were impaired. While voluntary movements were slower with impaired muscle activity patterns, startReact elbow flexion movements were not statistically different from age-matched unimpaired individuals opening a discussion about its potential use in therapy. However, confounding results were found during elbow extension. While there was evidence that startReact extension movements were present, elbow extension was interrupted by task-inappropriate flexor activity causing either delay in elbow extension or elbow flexion away from a subject's intended target.

The specific mechanisms driving this inappropriate flexor activity are unknown making it challenging to properly develop startReact as a therapy tool. However, evaluating the impact of impairment level on task-inappropriate flexor activity following cortical stroke could shed some insight. Impairment level is linked to both lesion size and damage to the corticospinal tract (Ciccarelli et al., 2008, Mohr et al., 1993, Rogers et al., 1997, Saver et al., 1999, Zhu et al., 2010). By investigating changes in task-inappropriate flexor activity with impairment level, we can gain insight into the role of the cortex and the corticospinal tract in its expression. Therefore, our objective was to quantify deficits in startReact extension movements, particularly how these deficits are affected by impairment. These results will give insight not only to the mechanisms driving deficits in startReact extension but given startReact's functional role in responding to environmental perturbations, may also provide

key insights into the mechanisms driving a stroke survivor's impaired responses to arm and balance perturbations.

We have previously hypothesized that the task-inappropriate flexor activity originates from unsuppressed classic startle reflex. This hypothesis was driven by the knowledge that the 1) classic startle response is dominated by flexor activity, 2) amplitude of the classic startle response is modulated by the cortex (Alibiglou and MacKinnon, 2012, Davis et al., 1982, Davis and Gendelman, 1977, Groves et al., 1974), and 3) the classic startle reflex is hypermetric or enlarged following stroke (Jankelowitz and Colebatch, 2004). Damage to the cortex following cortical stroke should diminish the capacity of the cortex to suppress the classic startle reflex during startReact movements. Therefore, we expect that increasing damage to the cortex (increasing impairment) will result in increasing task-inappropriate flexor activity corresponding to increasing deficits in elbow extension movement and decreasing target acquisition. Further, as classic startle is known to adapt (diminish in amplitude over time), we expect that the task-inappropriate flexor activity will diminish over time leading to increasingly appropriate elbow extension movement. A correct hypothesis would indicate that the task-inappropriate flexor activity likely arises from unsuppressed classic startle. Functionally, this result would highlight that more severely impaired individuals will have increased movement trajectory errors that move then away from their intended extension targets during external disturbances.

METHODS

Subjects

Data were collected from 8 chronic stroke subjects ranging in age from 47–81 (mean: 68 $\pm$ 9.8) (Table 1). Stroke subjects with a range of impairment levels were recruited. Impairment was assessed using the upper extremity Fugl-Meyer (UEFM) score, which ranged from 12–59. Inclusion criteria for the stroke subjects included: a unilateral cortical brain lesion from a stroke at least one year prior to the study, an ability to understand the task, lack of aphasia, and a stroke that affected the arm that was dominant prior to injury. We evaluated the dominant arm of all subjects as startReact movements have been studied largely in the dominant arm (Carlsen, Maslovat, 2011). All protocols and recruitment procedures were approved by Northwestern's Institutional Review Board (IRB).

Equipment

In all experiments, the arm was positioned at approximately 70 degrees of shoulder abduction and 25 degrees shoulder flexion; the elbow was positioned at 90 degrees. All subjects were fitted with a custom-made thermoplastic cast that immobilized the wrist and held the arm in the prone position. The top of cast was attached to a force sensor (45E15A4; JR3 Inc, Woodland, CA), which was coupled to a one degree of freedom rotary motor (BSM90N; Baldor Electric Company, WV) through a 10:1 planetary gear (AD140-010-PO; Apex Dynamics, Taiwan). The center of rotation was fixed just above the elbow joint. The rotary motor was used to ensure a repeatable and measured trajectory for evaluation of muscle activity patterns and position, and to support the weight of the arm against gravity. The rotary motor did not assist or perturb the elbow in any way. Rather, it was configured as

an admittance controller set to mimic the properties of a passive inertial load ($0.2 \text{ kg}\cdot\text{m}^2/\text{rad}$) in the flexion/extension axis.

Bipolar Ag/AgCl electromyography (EMG) electrodes (Noraxon Dual Electrodes, #272, Noraxon USA Inc., AZ) were used to record muscle activity from the brachioradialis (Br), triceps long head (TriLo), and the left and right sternocleidomastoid (SCM) muscles. EMG signals were amplified and conditioned using a Bortec AMT-8 (Bortec Biomedical Ltd., Canada), with a band-pass filter of 10–1,000 Hz. The resulting signals were anti-alias filtered using 5th order Bessel filters with a 500 Hz cut-off frequency and sampled at 2500 Hz (PCI-DAS1602/16; Measurement Computing, MA). Elbow position was recorded by an encoder with an effective resolution of 0.0036 degrees. EMG and position data were sampled synchronously. All data collection was synchronized to the GO signal using a pre-trigger collection time of 1s and post-trigger collection time of 5s.

Protocol

Each subject was trained to perform ballistic elbow extension movements of 25 degrees. A computer screen displayed two circles: HOME (located centrally) and TARGET (located at 25 degrees elbow extension). Subjects were instructed to move into the HOME circle and wait for two non-startling (80dB) auditory cues. The first cue (WARNING) signaled the start of the trial, and indicated that the subject should prepare to move. The second cue (GO) was the prompt to initiate the intended movement as rapidly as possible. The GO occurred randomly 2.2 – 2.5 seconds following the WARNING. Subjects were asked to move as quickly as possible from the HOME to TARGET circle. Subjects were instructed to reach to the target as fast as possible. They were given 2–3 seconds to complete the task, but no specific instructions were provided with respect to accuracy. A successful reach was defined as one that exceeded the 25 degree target on the first attempt. Subjects were first trained in the task until they responded reliably and consistently to the GO signal; this typically occurred after approximately 30 trials. After training, a startling auditory stimulus of 128dB was randomly delivered in place of the GO. During a block of 15 trials, startling acoustic stimuli were randomly delivered 3–4 times at the GO. Subjects performed between 2–3 blocks resulting in 6–8 startReact trials collected in each subject. Subjects were given no instruction on how to respond to the startling acoustic stimulus.

Data analysis

Position and EMG traces were visually inspected to eliminate trials in which the subject moved out of HOME prior to the GO or any especially slow trials in which subjects did not move at the GO. Next, SCM muscle activity was evaluated in all trials (voluntary and startReact). Activity in the SCM muscle is known to indicate the presence of startle (Carlsen, Maslovat, 2011, Valls-Solé, Rothwell, 1999). We considered activity in either the left or right SCM muscles within 120 ms of the acoustic stimulus to indicate the presence of startle (Blumenthal et al., 2005). Using this criterion, all trials with SCM activity present in either left or right SCM were classified as SCM+ (startle occurred). Those without SCM activity were classified as SCM- (startle was not detected). Only SCM- voluntary reaching trials and SCM+ startReact reaching trials were analyzed further, which yielded an inclusion rate of 89% voluntary reach and 80% startReact trials in stroke subjects. As SCM+

voluntary reaching trials were elicited during a different acoustic-intensity, which can impact onset latency (Kohfeld, 1969), these trials were excluded from analysis of SCM+ trials.

The latency of muscle activity onset was calculated from the rectified EMG recorded in each trial. The average background activity and standard deviation prior to the stimulus were calculated. Next an automated program identified the time at which the processed EMG increased above 2.5 times the standard deviation of the background activity for a period of 15ms. Following the automatic detection of EMG onset, each trial was evaluated visually to ensure accuracy. The amplitude of the EMG was computed as the average rectified response for a window of 70 ms following the detected onset. A previously reported impairment during startReact extension movements following stroke is abnormal elbow flexion activity prior the planned extension movement. This was quantified by the maximum elbow flexion during the planned elbow extension trials. In addition in order to quantify the time course of the observed trajectory, the elbow position was calculated at 150, 300, and 450ms following movement initiation. The subject's ability to reach the intended target was quantified as the trials in which the 25 degree target was reached on the first attempt and was represented as a probability.

Statistical analysis

First, we quantified the startReact response in comparison to voluntary movement. Muscle onset latency was compared between voluntary and startReact trials. An ANOVA was completed using a linear mixed-effect model. Muscle onset latency was treated as the dependent variable, while trial type (voluntary vs startReact) and muscle type (Br and TriLo) were treated as the independent variables. Subjects were treated as a random effect and a Tukey post-hoc test, which corrects for multiple comparisons, utilized. Our first hypothesis was that increasing damage to the cortex (increasing impairment) would relate to an increase in task-inappropriate flexor activity and consequently to increasing deficits in elbow extension movement and target acquisition. In order to assess the impact of impairment on the inappropriate flexor activity and resulting target acquisition, a linear regression was performed at a confidence interval of 95% with impairment (Upper-Extremity Fugl-Meyer or UEFM score) as the independent variable and the maximum elbow flexion deflection, startReact target acquisition, and average muscle onset latency as the dependent variables. Our second hypothesis was that the task-inappropriate flexor activity would diminish over time leading to increasingly appropriate elbow extension movement over time. In order to assess if the flexor activity diminished in amplitude over time, an ANOVA was performed with elbow position at three time points (150, 300, 450ms), EMG amplitude, and EMG latency as the dependent variables and trial number as the independent variable. Subjects were treated as random effects and trial number was treated as a continuous variable.

RESULTS

Task-inappropriate flexor activity influenced startReact extension movements across all impairment levels (Figure 1). Appropriate agonist (TriLo) activity preceding antagonist

(BR) activity was present during voluntary elbow extension movements (gray). However during startReact elbow extension movements (black) the antagonist BR muscle activity preceded or occurred at the same latency as agonist BR activation. This was confirmed by group results. Onset latency was influenced by trial type ($F_{1,368} = 379$; $p < 0.0001$) and muscle type ($F_{1,368} = 195$; $p < 0.0001$) with a significant interaction ($F_{1,368} = 80$; $p < 0.0001$). Therefore, the average onset latency of the TriLo (211 ± 110 ms) and Br (387 ± 147 ms) muscles during voluntary movements was different ($p \approx 0$), while the onset latencies of TriLo (98 ± 33 ms) and BR (83 ± 23 ms) were not different during startReact extension ($p = 0.25$).

While task-inappropriate flexor activity was present in all subjects evaluated, it led to the most significant deviation from the extension target in more severely impaired individuals. Individuals with mild impairment (Figure 1A) demonstrated a delay in elbow movement towards extension, while individuals with moderate and severe impairment demonstrated large elbow flexion movements prior to elbow extension (Figure 1 B,C). Quantification of group results established that maximum elbow flexion deflection and startReact extension target acquisition were related to impairment level. Maximum elbow flexion deflection (Figure 2A: $p = 0.016$) increased with increasing impairment (decreasing UEFM). Similarly, the probability of startReact extension movements reaching the target (Figure 2B: $p = 0.0009$) decreased with increasing impairment, even though the probability of reaching the extension target during voluntary movements was not related to impairment ($p = 0.12$). Still, changes in elbow flexion deflection and target acquisition were not related to changes in muscle onset latency. BR ($p = 0.79$) and TriLo ($p = 0.47$) muscle onset latencies were not related to changes in impairment (Figure 2C).

Regardless of impairment, elbow position showed decreasing movement in the flexion direction in successive trials (Figure 3) and coincided with decreasing BR muscle amplitude (Figure 4). All subjects showed improved startReact extension movements over time. In mild impairment, the inappropriate flexor activity led to a delay of movement in the extension direction that dissipated in subsequent trials leading to faster target acquisition (Figure 3A). Moderately and severely impaired individuals showed movement trajectories that did not deflect towards flexion in later trials. Group results confirmed that improvements in elbow trajectory were seen in all subjects. Specifically while elbow position was not affected by trial number at 150ms ($F_{(1,46)} = 0.97$; $p = 0.34$), it was affected at 300ms ($F_{(1,46)} = 8.52$; $p = 0.006$) and 450ms ($F_{(1,46)} = 22.6$; $p = 0.0001$) (Figure 4A). Improvements in elbow position trajectory were accompanied by decreased BR muscle amplitude, which was influenced trial number ($F_{(1,43)} = 13.3$; $p = 0.0021$). Conversely, TriLo muscle amplitude remained constant ($F_{(1,43)} = 0$; $p = 0.99$) (Figure 3B) indicating that changes in elbow position were related to changes in BR muscle amplitude. While the amplitude of the BR muscle activity was diminished, the onset latency of BR ($F_{(1,42)} \approx 0$; $p = 0.98$) and TriLo ($F_{(1,45)} \approx 0$; $p = 0.96$) were not impacted by trial number (Figure 3C) indicating that while the task-inappropriate flexor activity diminished, it was not eliminated over the course of the experiment.

DISCUSSION

We hypothesized that the task-inappropriate flexor activity during startReact elbow extension originates from unsuppressed classic startle reflex. Therefore, we expected that 1) increasing damage to the cortex (increasing impairment) would relate to an increase in task-inappropriate flexor activity and consequently to increasing deficits in elbow extension movement and target acquisition and 2) that task-inappropriate flexor activity would diminish over time leading to increasingly appropriate elbow extension movement over time. We indeed found that task-inappropriate flexor activity increased with impairment leading to larger elbow flexion deflections away from the subjects' intended extension target and corresponded to decreased target acquisition. Moreover, severely impaired individuals UEFM < 25 never achieved the extension target during startReact trials. While this task-inappropriate flexor activity diminished leading to improved elbow trajectories over subsequent trials, the inappropriate activity was not eliminated over the course of these experiments. As functional motor impairment is linked to both cortical lesion size (Mohr, Foulkes, 1993, Rogers, Campbell, 1997, Saver, Johnston, 1999) and damage to the corticospinal tract (Ciccarelli, Catani, 2008, Zhu, Lindenberg, 2010), the task-inappropriate flexor activity is likely driven by damage to the cortex or corticospinal tract. Interestingly, adaptation (or diminishing of the flexor activity over time) remained intact in all impairment levels suggesting it is mediated by alternative neural pathways. Finally, as startReact responses are triggered during perturbations of the arm and whole-body, more severely impaired individuals likely will have increased movement trajectory errors that specifically move then away from their intended extension targets during external disturbances.

Mechanisms driving deficits in startReact following stroke

There is growing evidence that classic startle and startReact are separate phenomena mediated by overlapping but distinctive neural structures (Alibiglou and MacKinnon, 2012, Kumru et al., 2006, Maslovat et al., 2012). In unimpaired individuals, classic startle results in brief, synchronous activity of upper limb muscles which is dominated by flexion (Brown et al., 1991). During startReact movements, a startling stimulus releases a planned movement that is not different in muscle activation patterns and target accuracy from a voluntarily elicited planned movement (Carlsen et al., 2011, Maslovat et al., 2011, Valls-Solé, Rothwell, 1999, Valls-Solé, Solé, 1995). In unimpaired individuals, only the planned movement is elicited and the classic startle response is not quantitatively observed in the upper limb muscles (Carlsen, Maslovat, 2011). Animal work demonstrated that the cortex modulates the amplitude of the classic startle reflex (Davis and Gendelman, 1977). Similar results occur in humans as cortical damage leads to hypermetric classic startle responses (Jankelowitz and Colebatch, 2004, Rothwell, 2006). Therefore following a stroke, it is probable that cortical damage impairs the ability to suppress the classic startle reflex during startReact leading to a simultaneous release of both classic startle and startReact. This phenomenon has been observed in other subcortical reflexes e.g. hypermetric stretch reflexes or spasticity (Krakauer, 2005, Levin, 1996, Zackowski et al., 2004) and a resurgence of the typically dormant asymmetric tonic neck reflexes following stroke (Ellis et al., 2012, Lee et al., 2009, Yamshon et al., 1949). Further, the task-inappropriate flexor activity shares several common features with the classic startle reflex. The task-

inappropriate flexor activity shows adaptation over time which has been observed in classic startle (Valls-Solé et al., 1997) but not startReact (Carlsen, Maslovat, 2011). Further the task-inappropriate flexor activity is not statistically different in onset latency from the agonist (TriLo) muscle, which also closely resembles the synchronous agonist/antagonist firing of the classic startle reflex. These observations suggest that the task-inappropriate flexor activity is the result of unsuppressed or hypermetric classic startle resulting from cortical damage.

It is important to note that an intact startReact extension movement is likely present. First, there is a significant decrease in onset latency in the agonist, TriLo muscle during startReact extension (98ms) from voluntary (211ms). TriLo muscle activation showed consistent activation over the duration of the experiment, distinctive from the diminishing activity in the BR muscle, indicating that the mechanisms driving BR and TriLo muscle activation are distinctive. Second, startReact extension resulted in elbow extension. If the classic startle was the only response, we would expect to see only movement in the flexion direction (Brown, Rothwell, 1991, Honeycutt and Perreault, 2012, Jankelowitz and Colebatch, 2004, Yeomans and Frankland, 1996). Finally, we have previously reported that startReact elbow flexion movements are intact and statistically indistinguishable from neurologically healthy age-matched individuals. (Honeycutt and Perreault, 2012) indicating that when task-inappropriate flexor activity does not interrupt movement an appropriate movement can be elicited. This is particularly interesting given that voluntary executed elbow flexion movements are significantly impaired in stroke survivors in comparison to neurologically healthy individuals. StartReact elbow flexion movements are likely not impaired by the unsuppressed startle because this activity assists the flexion movement while it opposes elbow extension. Therefore, we conclude that the task-inappropriate flexor muscle activation most likely results from unsuppressed classic startle.

An alternative mechanism that could drive the task-inappropriate flexor activity is spasticity that often leads to tonic co-contraction during movement in stroke survivors (Dietz and Sinkjaer, 2007, Dietz et al., 1991, Krakauer, 2005, Levin, 1996). However, this is not a probable explanation as spasticity occurs in the response to stretch and the onset latency of the BR muscle is concurrent with the TriLo muscle indicating that it does not result from stretch initiated during movement. Further, the BR activity diminishes in amplitude which would be unexpected if the cause was spasticity.

Functional consequences

The ability to quickly and effectively compensate during environmental perturbations is essential to everyday tasks and injury prevention. It is well-established that following stroke individuals have important deficits resisting perturbations of the arm and ankle (Dietz and Berger, 1984, Dietz, Trippel, 1991, Finley et al., 2008, Lum et al., 2004, Sangani et al., 2007, Trumbower et al., 2013, Trumbower et al., 2010). Perturbation-specific, short and long latency stretch reflexes, are significantly impaired following stroke disrupting their ability to resist arm perturbation (Dietz and Berger, 1984, Dietz, Trippel, 1991, Lum, Patten, 2004, Trumbower, Finley, 2013, Trumbower, Ravichandran, 2010) leading to a reduced torque capacity. Recent evidence demonstrates that this ability is driven not only by

perturbation-specific stretch reflexes but also by task-specific early release of planned movement (Hammond, 1956, Lewis et al., 2006, Ravichandran, Honeycutt, 2013). StartReact movements are identical in complexity and muscle activation patterns as voluntarily planned movements but they are elicited 30–40ms faster – a time that overlaps with the long-latency stretch response to environmental perturbations (Crago et al., 1976, Gottlieb and Agarwal, 1979). Recent work from our laboratory demonstrates that startReact movements are triggered during perturbations of the arm (Ravichandran, Honeycutt, 2013). When startReact is not present, the subject has a significantly diminished and delayed reaction to perturbation indicating that this reflex plays an important role in quickly and effectively resisting perturbations of the arm (Smith et al., 2011).

Furthermore, our results demonstrate that more severely impaired individuals have more severe deficits in startReact extension. Thus, more severely impaired individuals will have increased movement trajectory errors that move them away from their intended extension targets during external disturbances. When coupled with the decreased muscle strength, spasticity, and loss of functional control over the limb, places these individuals at significant risk of injury.

While the role of startReact during whole-body perturbations remains less certain, startle has been recorded during perturbations of the whole body. Furthermore, the startle reflex is mediated in part by the brainstem (Davis, Gendelman, 1982, Groves, Wilson, 1974, Hammond, 1973, Rothwell, 2006, Yeomans and Frankland, 1996), which has known importance during corrective responses during balance challenges (Deliagina et al., 2008, Honeycutt et al., 2009, Honeycutt and Nichols, 2010, Mori, 1987, Mori et al., 1989, Musienko et al., 2008, Schepens et al., 2008, Stapley and Drew, 2009). Thus, it is possible that task-inappropriate flexor activity could lead to movement away from a subject's intended target during balance challenges. Further research is needed to establish a link between the deficits described here and those that occur during active resistance of environmental perturbations.

Limitations

Our results are limited by a lack of detailed lesion data. More thorough imaging data could provide important information about the role of the cortex in these types of responses. Furthermore, tractography (imaging technique quantifying impairment to neural tracts) of the corticospinal tract would provide more definitive evidence that the task-inappropriate flexor activity results from damage of this neural tract.

Finally in this study the arm is supported against gravity and movement restricted to the elbow. Therefore in more functional and less restricted situations, the impact of task-inappropriate flexor activity may be more significant. Furthermore, spasticity and abnormal muscle activation patterns (synergies) would likely also more significantly disrupt movement during these conditions.

Acknowledgments

The authors would like to thank Tim Goetz-Haswell his technical and scientific expertise. In addition, we would like to thank Dr. Jungwha “Julia” Lee for her statistical expertise. This work was supported by the National Institutes of Health grants K12 GM088020 and R01 NS053813 K99 HD073240.

References

- Alibiglou L, MacKinnon CD. The early release of planned movement by acoustic startle can be delayed by transcranial magnetic stimulation over the motor cortex. *J Physiol.* 2012; 590:919–36. [PubMed: 22124142]
- Blouin JS, Inglis JT, Siegmund GP. Startle responses elicited by whiplash perturbations. *J Physiol.* 2006; 573:857–67. [PubMed: 16581859]
- Blumenthal TD, Cuthbert BN, Filion DL, Hackley S, Lipp OV, van Boxtel A. Committee report: Guidelines for human startle eyeblink electromyographic studies. *Psychophysiol.* 2005; 42:1–15.
- Brown P, Rothwell JC, Thompson PD, Britton TC, Day BL, Marsden CD. New observations on the normal auditory startle reflex in man. *Brain.* 1991; 114 :1891–902. [PubMed: 1884184]
- Campbell AD, Chua R, Inglis JT, Carpenter MG. Startle induces early initiation of classically conditioned postural responses. *J Neurophysiol.* 2012; 108:2946–56. [PubMed: 22972964]
- Carlsen AN, Chua R, Inglis JT, Sanderson DJ, Franks IM. Can prepared responses be stored subcortically? *Exp Brain Res.* 2004a; 159:301–9. [PubMed: 15480608]
- Carlsen AN, Chua R, Inglis JT, Sanderson DJ, Franks IM. Prepared movements are elicited early by startle. *J Motor Behav.* 2004b; 36:253–64.
- Carlsen AN, Maslovat D, Lam MY, Chua R, Franks IM. Considerations for the use of a startling acoustic stimulus in studies of motor preparation in humans. *Neurosci Biobehav Rev.* 2011; 35:366–76. [PubMed: 20466020]
- Ciccarelli O, Catani M, Johansen-Berg H, Clark C, Thompson A. Diffusion-based tractography in neurological disorders: concepts, applications, and future developments. *Lancet Neurol.* 2008; 7:715–27. [PubMed: 18635020]
- Crago PE, Houk JC, Hasan Z. Regulatory actions of human stretch reflex. *J Neurophysiol.* 1976; 39:925–35. [PubMed: 978238]
- Davis M, Gendelman DS, Tischler MD, Gendelman PM. A primary acoustic startle circuit: lesion and stimulation studies. *J Neurosci.* 1982; 2:791–805. [PubMed: 7086484]
- Davis M, Gendelman PM. Plasticity of the acoustic startle response in the acutely decerebrate rat. *J Comp Physiol Psychol.* 1977; 91:549–63. [PubMed: 874121]
- Deliagina TG, Beloozerova IN, Zelenin PV, Orlovsky GN. Spinal and supraspinal postural networks. *Brain Res Rev.* 2008; 57:212–21. [PubMed: 17822773]
- Dietz V, Berger W. Interlimb coordination of posture in patients with spastic paresis. Impaired function of spinal reflexes. *Brain.* 1984; 107:965–78. [PubMed: 6478185]
- Dietz V, Sinkjaer T. Spastic movement disorder: impaired reflex function and altered muscle mechanics. *Lancet Neurol.* 2007; 6:725–33. [PubMed: 17638613]
- Dietz V, Trippel M, Berger W. Reflex activity and muscle tone during elbow movements in patients with spastic paresis. *Ann Neurol.* 1991; 30:767–79. [PubMed: 1789693]
- Ellis MD, Drogos J, Carmona C, Keller T, Dewald JP. Neck rotation modulates flexion synergy torques, indicating an ipsilateral reticulospinal source for impairment in stroke. *J Neurophysiol.* 2012; 108:3096–104. [PubMed: 22956793]
- Finley JM, Perreault EJ, Dhaher YY. Stretch reflex coupling between the hip and knee: implications for impaired gait following stroke. *Exp Brain Res.* 2008; 188:529–40. [PubMed: 18446331]
- Gottlieb GL, Agarwal GC. Response to sudden torques about ankle in man: myotatic reflex. *J Neurophysiol.* 1979; 42:91–106. [PubMed: 430116]
- Groves PM, Wilson CJ, Boyle RD. Brain stem pathways, cortical modulation, and habituation of the acoustic startle response. *Behav Biol.* 1974; 10:391–418. [PubMed: 4832941]

- Hammond GR. Lesions of pontine and medullary reticular formation and prestimulus inhibition of the acoustic startle reaction in rats. *Physiol Behav.* 1973; 10:239–43. [PubMed: 4575307]
- Hammond PH. The influence of prior instruction to the subject on an apparently involuntary neuromuscular response. *J Physiol.* 1956; 132:17–8P. [PubMed: 13320395]
- Honeycutt CF, Gottschall JS, Nichols TR. Electromyographic responses from the hindlimb muscles of the decerebrate cat to horizontal support surface perturbations. *J Neurophysiol.* 2009; 101:2751–61. [PubMed: 19321638]
- Honeycutt CF, Kharouta M, Perreault EJ. Evidence for reticulospinal contributions to coordinated finger movements in humans. *J Neurophysiol.* 2013; 110:1476–83. [PubMed: 23825395]
- Honeycutt CF, Nichols TR. The decerebrate cat generates the essential features of the force constraint strategy. *J Neurophysiol.* 2010; 103:3266–73. [PubMed: 20089811]
- Honeycutt CF, Perreault EJ. Planning of Ballistic Movement following Stroke: Insights from the Startle Reflex. *PloS one.* 2012; 7:e43097. [PubMed: 22952634]
- Jankelowitz SK, Colebatch JG. The acoustic startle reflex in ischemic stroke. *Neurology.* 2004; 62:114–6. [PubMed: 14718710]
- Kohfeld DL. Effects of the intensity of auditory and visual ready signals on simple reaction time. *J Exp Psychol.* 1969; 82:88–95. [PubMed: 5371997]
- Krakauer JW. Arm function after stroke: from physiology to recovery. *Semin Neurol.* 2005; 25:384–95. [PubMed: 16341995]
- Kumru H, Valls-Solé J, Kofler M, Castellote J, Sanegre MT. The effects of a prepulse on the startReact phenomenon. *Suppl Clin Neurophysiol.* 2006; 58:101–9. [PubMed: 16623325]
- Lee S, Yao J, Acosta aM, Dewald Ja. The effects of asymmetric tonic neck reflex during reaching movement following stroke: preliminary results. Conference proceedings : Annual International Conference of the IEEE Eng Med Society IEEE Eng Med Society Conference. 2009; 2009:1581–4.
- Levin MF. Interjoint coordination during pointing movements is disrupted in spastic hemiparesis. *Brain.* 1996; 119:281–93. [PubMed: 8624689]
- Lewis GN, MacKinnon CD, Perreault EJ. The effect of task instruction on the excitability of spinal and supraspinal reflex pathways projecting to the biceps muscle. *Exp Brain Res.* 2006; 174:413–25. [PubMed: 16676166]
- Lum PS, Patten C, Kothari D, Yap R. Effects of velocity on maximal torque production in poststroke hemiparesis. *Muscle Nerve.* 2004; 30:732–42. [PubMed: 15468340]
- Maslovat D, Hodges NJ, Chua R, Franks IM. Motor preparation of spatially and temporally defined movements: evidence from startle. *J Neurophysiol.* 2011; 106:885–94. [PubMed: 21613586]
- Maslovat D, Kennedy PM, Forgaard CJ, Chua R, Franks IM. The effects of prepulse inhibition timing on the startle reflex and reaction time. *Neurosci Lett.* 2012; 513:243–7. [PubMed: 22387455]
- Mohr JP, Foulkes MA, Polis AT, Hier DB, Kase CS, Price TR, et al. Infarct topography and hemiparesis profiles with cerebral convexity infarction: the Stroke Data Bank. *J Neurol Neurosurg Psychiatry.* 1993; 56:344–51. [PubMed: 8482953]
- Mori S. Integration of posture and locomotion in acute decerebrate cats and in awake, freely moving cats to initiate and terminate locomotor movements both in the biped and in the quadruped, a smooth transition from and to standing is necessary, standing being. *Prog Neurobiol.* 1987; 28:161–95. [PubMed: 3544055]
- Mori S, Sakamoto T, Ohta Y, Takakusaki K, Matsuyama K. Site-specific postural and locomotor changes evoked in awake, freely moving intact cats by stimulating the brainstem. *Brain Res.* 1989; 505:66–74. [PubMed: 2611678]
- Musienko PE, Zelenin PV, Lyalka VF, Orlovsky GN, Deliagina TG. Postural performance in decerebrated rabbit. *Behav Brain Res.* 2008; 190:124–34. [PubMed: 18359100]
- Oude Nijhuis LB, Allum JHJ, Valls-Sole J, Overeem S, Bloem BR. First trial postural reactions to unexpected balance disturbances: a comparison with the acoustic startle reaction. *J Neurophysiol.* 2010; 104:2704–12. [PubMed: 20810688]
- Ravichandran VJ, Honeycutt CF, Shemmell J, Perreault EJ. Instruction-dependent modulation of the long-latency stretch reflex is associated with indicators of startle. *Exp Brain Res.* 2013; 230:59–69. [PubMed: 23811739]

- Rogers DC, Campbell CA, Stretton JL, Mackay KB. Correlation between motor impairment and infarct volume after permanent and transient middle cerebral artery occlusion in the rat. *Stroke*. 1997; 28:2060–5. discussion 6. [PubMed: 9341719]
- Rothwell JC. The startle reflex, voluntary movement, and the reticulospinal tract. *Suppl Clin Neurophysiol*. 2006; 58:223–31. [PubMed: 16623334]
- Rothwell JC, MacKinnon CD, Valls-Solé J. Role of brainstem-spinal projections in voluntary movement. *Mov Disord*. 2002; 17 (Suppl 2):S27–9. [PubMed: 11836749]
- Sangani SG, Starsky AJ, McGuire JR, Schmit BD. Multijoint reflexes of the stroke arm: neural coupling of the elbow and shoulder. *Muscle Nerve*. 2007; 36:694–703. [PubMed: 17628498]
- Saver JL, Johnston KC, Homer D, Wityk R, Koroshetz W, Truskowski LL, et al. The RANTTAS Investigators. Infarct volume as a surrogate or auxiliary outcome measure in ischemic stroke clinical trials. *Stroke*. 1999; 30:293–8. [PubMed: 9933262]
- Schepens B, Drew T, Rogers MW. Independent and Convergent Signals From the Pontomedullary Reticular Formation Contribute to the Control of Posture and Movement During Reaching in the Cat. *J Neurophysiol*. 2008:2217–38. [PubMed: 18650311]
- Siegmund GP, Blouin J-S, Inglis JT. Does startle explain the exaggerated first response to a transient perturbation? *Exerc Sport Sci Rev*. 2008; 36:76–82. [PubMed: 18362689]
- Smith, RL.; Honeycutt, CF.; Ravichandran, VJ.; Perreault, EJ. Influence of perturbation intensity on the rapid release of triggered reactions. Society for Neuroscience; San Deigo: 2011.
- Stapley PJ, Drew T. The pontomedullary reticular formation contributes to the compensatory postural responses observed following removal of the support surface in the standing cat. *J Neurophysiol*. 2009; 101:1334–50. [PubMed: 19118108]
- Trumbower RD, Finley JM, Shemmell JB, Honeycutt CF, Perreault EJ. Bilateral impairments in task-dependent modulation of the long-latency stretch reflex following stroke. *Clin Neurophysiol*. 2013; 124:1373–80. [PubMed: 23453250]
- Trumbower RD, Ravichandran VJ, Krutky Ma, Perreault EJ. Contributions of altered stretch reflex coordination to arm impairments following stroke. *J Neurophysiol*. 2010; 104:3612–24. [PubMed: 20962072]
- Valls-Solé J, Kumru H, Kofler M. Interaction between startle and voluntary reactions in humans. *Exp Brain Res*. 2008; 187:497–507. [PubMed: 18458890]
- Valls-Solé J, Rothwell JC, Goulart F, Cossu G, Muñoz E. Patterned ballistic movements triggered by a startle in healthy humans. *J Physiol*. 1999; 516:931–8. [PubMed: 10200438]
- Valls-Solé J, Solé A, Valdeoriola F, Muñoz E, Gonzalez LE, Tolosa ES. Reaction time and acoustic startle in normal human subjects. *Neurosci Lett*. 1995; 195:97–100. [PubMed: 7478277]
- Valls-Solé J, Valdeoriola F, Tolosa E, Nobbe F. Habituation of the auditory startle reaction is reduced during preparation for execution of a motor task in normal human subjects. *Brain Res*. 1997; 751:155–9. [PubMed: 9098581]
- Yamshon L, Machek O, Covalt D. The tonic neck reflex in the hemiplegic; an objective study of its therapeutic implication. *Arch Phys Med Rehabil*. 1949; 11:706–11. [PubMed: 15391233]
- Yeomans JS, Frankland PW. REVIEWS Review article The acoustic startle reflex : neurons and connections. *Brain Res Rev*. 1996; 21:301–14. [PubMed: 8806018]
- Zackowski KM, Dromerick AW, Sahrman Sa, Thach WT, Bastian AJ. How do strength, sensation, spasticity and joint individuation relate to the reaching deficits of people with chronic hemiparesis? *Brain*. 2004; 127:1035–46. [PubMed: 14976070]
- Zhu LL, Lindenberg R, Alexander MP, Schlaug G. Lesion load of the corticospinal tract predicts motor impairment in chronic stroke. *Stroke*. 2010; 41:910–5. [PubMed: 20378864]

HIGHLIGHTS

1. Startle-evoked movements remain intact following stroke, but there are deficits specific to extension movements that increase with impairment level.
2. These extension-related deficits appear to originate from a hypermetric classic startle reflex likely resulting from damage to cortical pathways.
3. These results may have important implications for our understanding of deficits in stroke survivor's response to unexpected environmental disturbances.

Figure 1.

EMG responses from the BR and TriLo muscles and elbow position during voluntary (gray) and startReact (black) trials. A) Mild impairment, UEFM = 56 B) Moderate impairment, UEFM = 34 C) Severe impairment, UEFM = 15. Black line indicates extension target.

Figure 2.

Quantification of startReact deficits across impairment level. A) The maximum flexion deflection position during startReact trials for each subject graphed against Upper Extremity Fugl-Meyer (UEFM) scores. B) The probability of target acquisition during startReact trials graphed against UEFM. C) Onset latency for BR (black) and TriLo (gray) muscles against impairment.

Figure 3.

Elbow position presented across early, middle, and late startReact trials to depict changes in elbow position over time. A) Mild Impairment, UEFM = 59 B) Moderate Impairment: UEFM = 40. C) Severe Impairment: UEFM = 24. Black line represents extension target.

Figure 4.

Change in position, EMG amplitude, and onset latency over successive trials. A) Arm position during startReact at 150ms (light gray), 300ms (dark gray), and 450ms (black) across trials. B) Average EMG amplitude during startReact for the BR (black) and TriLo (gray) muscles across trials. C) Onset latencies during startReact for the BR (black) and TriLo (gray) muscles across trials.

Table 1

Subject Characteristics

Subject #	Sex	Age	Paretic Limb *	Years since Stroke	UEFM	Chedoke Arm
1	M	68	L	10	34	4
2	M	81	R	17	24	3
3	M	61	R	8	54	5
4	F	70	R	22	40	4
5	M	69	R	6	12	2
6	M	74	R	23	15	--
7	M	65	R	11	22	3
8	M	47	R	36	59	5

* Paretic limb was dominant hand prior to stroke

-- Scores were not available