Published in final edited form as:

Sleep Med. 2014 April ; 15(4): 444–450. doi:10.1016/j.sleep.2014.01.003.

Sleep Disturbance and Cardiometabolic Risk Factors in Early Pregnancy: A Preliminary Study

Alyssa HANEY^a [Medical Student], Daniel J. BUYSSE^a [Professor], Bedda L. ROSARIO^b [Visiting Assistant Professor of Epidemiology], Yi-Fan CHEN^b [Graduate Student Researcher], and Michele L. OKUN^{a,*} [Assistant Professor]

^aUniversity of Pittsburgh School of Medicine, Department of Psychiatry Pittsburgh, PA 15213

^bUniversity of Pittsburgh, Graduate School of Public Health, Epidemiology Data Center Pittsburgh, PA 15213

Abstract

Background: Cardiometabolic (CM) risk factors are linked to increased morbidity. Disturbed sleep is associated with CM risk factors in late pregnancy, but little is known about sleep in early pregnancy and CM risk factors.

Methods: Diary and actigraphy-assessed sleep information, as well as CM outcomes (blood pressure (BP) and BMI), were collected thrice from pregnant women (N = 161) in early pregnancy: T1 (10-12 weeks), T2 (14-16 weeks), and T3 (18-20 weeks). The sleep variables evaluated included sleep onset latency (SOL), wake after sleep onset (WASO) and total sleep time (TST). Sleep variables were dichotomized using established clinical cutoffs.

Results: BMI and BP significantly changed across time. Women with persistent SOL ≥ 20 minutes had greater BMI than women without persistent SOL ≥ 20 minutes prior to covariate adjustment at T1 and T2, but at T3 BMI values converged. Similar results were observed for persistent WASO ≥ 30 minutes. Persistently long WASO, as measured by actigraphy, was associated with elevated SBP, after controlling for covariates.

Conclusions: Consistent with anecdotal evidence, it appears as if a subset of women report substantial difficulty initiating and maintaining sleep during early pregnancy and this may augment risk for higher blood pressure and BMI. Understanding these relationships is important since CM risk factors are linked to maternal and infant morbidity. Assessing sleep in early pregnancy may bestow time necessary for appropriate intervention.

Keywords

Pregnancy; sleep disturbance; blood pressure; BMI; sleep latency; weight; actigraphy

© 2014 Elsevier B.V. All rights reserved

*Corresponding author: Michele Okun, Ph.D. University of Pittsburgh Western Psychiatric Institute and Clinic 3811 O'Hara St. E1124 Pittsburgh, PA 15213 T: 412-586-9434 mlo34@pitt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

Cardiometabolic disturbances during pregnancy, such as hypertension and obesity, have become increasingly prevalent problems in the United States. Hypertension impacts 6-8% of all pregnancies, while obesity rates have reached upwards of 50% in reproductive aged women (1). Hypertensive disorders in pregnancy increase the risk of abruptio placentae, disseminated intravascular coagulation, cerebral hemorrhage, hepatic failure, and acute renal failure, (2) all of which are potentially deadly disorders to both mother and fetus. Elevated blood pressure in pregnancy can also be part of pre-eclampsia and eclampsia, which carry maternal mortality rates of up to 10-15% (3). In addition, hypertensive disorders that develop in pregnancy impart risk for maternal cardiovascular disease later in life (4). Obesity in pregnancy is also associated with maternal morbidity, including gestational and later life diabetes, hypertensive disorders, and higher rate of cesarean delivery (5;6). Likewise, maternal obesity is associated with fetal morbidity, such as fetal macrosomia and fetal anomalies, including neural tube defects and cardiac malformations (7).

Sleep is a common behavioral pathway associated with both hypertension and obesity. Sleep disturbances including short/long sleep, poor sleep quality and poor continuity, are associated with hypertension and obesity in non-pregnant cohorts, even after controlling for traditional risk factors such as age, smoking, and race (8). Recent studies have observed similar associations in pregnant women. For example, Williams et al (9) reported that both long and short sleep duration in early pregnancy were associated with elevated blood pressure in later pregnancy. Likewise, Althuisen et al (10) identified short sleep as a correlate of excessive weight gain in pregnancy. However, they only assessed sleep in later pregnancy (30 weeks'). We speculate that poor sleep in early gestation (operationalized as less than 20 weeks) may contribute to dysregulation in the CM pathways resulting in hypertension and obesity; thereby increasing risk for adverse pregnancy outcomes (9).

Sleep can be measured subjectively or objectively using various methods. Subjective methods include questionnaires and sleep diaries, whereas objective methods include actigraphy and polysomnography (PSG) (11). Actigraphy is a convenient and reliable adjunct to questionnaire data. At the present time, few studies have ascertained both subjective and objective sleep in pregnant women. Previous studies using actigraphy in pregnancy have been limited by short collection periods (<72 hours) and small cohorts (<50) (12). Thus, the purpose of this study was to 1) examine whether women with poor sleep, defined as short sleep duration, longer wake after sleep onset, or longer sleep onset latency at 10-12 weeks', measured by both sleep diary and actigraphy, would have higher blood pressure or higher BMI across early pregnancy (10-20 weeks); and 2) whether women with persistent sleep disturbance (i.e. poor sleep between 10-20 weeks) would have greater increases in BP or BMI compared to women with intermittent sleep disturbance or no sleep disturbance.

METHODS

Participants

Participants assessed for this secondary analysis were drawn from a prospective, observational study (Sleep in Pregnancy (SLIP)) (13), in which sleep disturbance and inflammation in early gestation are evaluated for their role in adverse pregnancy outcomes. Women in general good health were recruited at approximately 10 weeks gestation by self-referral, physician referral, local advertising or participation in University research registries. Exclusion criteria included a current diagnosis of depression, bipolar disorder or anxiety, self-reported sleep disorders (e.g., sleep apnea, narcolepsy, hypersomnia or insomnia), current use of an antidepressant, antipsychotic, or anti-inflammatory medication, gynecologic anatomical abnormality, hypertension, diabetes, HIV or other major chronic disease. All participants provided written informed consent and the study was approved by the University of Pittsburgh Internal Review Board. Participants received monetary compensation for their participation.

Study Protocol and Procedures

At enrollment participants provided information on current health, health behaviors, and socio-demographic information. Sleep data were collected T1 (10-12), T2 (14-16) and T3 (18-20 weeks' gestation). Eligible participants completed the Pittsburgh sleep diary (14) daily and wore a wrist actigraph (Phillips/Respironics, Murrysville, PA) for 14 consecutive days, 24 hours a day during each assessment period. On Day 15 during each measurement cycle, participants presented for a health assessment, including blood pressure and weight measurements.

Sleep

Daily sleep diaries (14) were completed by participants for three 2-week periods. Participants recorded information about nocturnal sleep upon awakening and about daytime information prior to retiring for the day. Participants concurrently wore a wrist actigraph (Actiwatch64, Philips Respironics, Bend, OR) on the non-dominant wrist. The actigraph continuously monitors movement and activity in 1-minute epochs via an accelerometer and storage chip. Actigraphy data were scored using Actiware-Sleep 5.0 analysis software (Actiwatch64, Philips Respironics, Bend, OR).

Measures of sleep included subjective and actigraphy-assessed indices of (a) *sleep onset latency* (SOL), (the amount of time between reported bedtime and sleep onset time); (b) *wake after sleep onset* (WASO), (the amount of time awake after sleep onset); and (c) *total nocturnal sleep time* (TST), (defined as the amount of time between sleep onset and sleep offset). All measures were averaged across all available nights for each period (up to 14 days). Sleep variables were examined as continuous or categorical as appropriate. The three sleep variables were categorized to depict the presence or absence of sleep disturbance using published data from pregnant women in early pregnancy (9;15). SOL was considered disturbed if the diary or actigraphy 2-week average was ≥ 20 minutes. WASO was considered disturbed if the diary or actigraphy 2-week average ≥ 30 minutes. TST was considered disturbed if the diary 2-week average was < 7 hours or if the actigraphy 2-weeks

average was < 6 hours. We chose these two different TST cutoffs to maintain similar cell sizes, since approximately equal percentages were represented from each method (24% from diary-assessed < 7hrs and 20% from actigraphy-assessed < 6 hrs). In addition to the cross-sectional groupings, we further categorized participants into “persistent” (had sleep disturbance at all time points), “intermittent” (has sleep disturbance at 1 or 2 time points), and “no” (has no disturbed sleep) sleep disturbance groups.

Outcomes measures

Blood Pressure (BP) assessments were taken using a WelchAllyn® automated sphygmomanometer while participants were in a seated position. Sitting values of systolic (SBP) and diastolic blood pressure (DBP) were collected by the study nurse. *Weight* was assessed using a Detecto® electronic scale and is reported in pounds. *Height* was assessed using a Seca® electronic measuring rod. Both measures were taken with no shoes on. *BMI* was calculated as kg/m². All measures were collected thrice during the study period: 12, 16, and 20 weeks.

Covariates

Potential covariates included sociodemographics and indices of health behaviors previously associated with blood pressure or weight gain/BMI during pregnancy. *Age*, *race* (Caucasian or African-American/Other), *marital status*, *exercise* (yes or no), and *number of children < 18 years at home* (0 or > 1) were established by self-report. *Pre-pregnancy weight* was collected by a nurse at the first clinical visit. Self-reported symptoms of *stress* were measured with the 10-item Perceived Stress Scale (PSS 10) (Cohen *et al.*, 1983) was calculated as a continuous variable. Self-reported symptoms of depression were measured with the Inventory of Depressive Symptoms (IDS) questionnaire (16). The IDS, minus sleep items, was calculated as a continuous variable.

Statistical Analyses

Descriptive statistics were examined to characterize the demographics for the total cohort and by diary and actigraphy sleep measures. Means and standard deviations of all continuous measures, by time point, for the total sample are reported. Examination of normal distribution assumption for continuous data was determined by q-q plots, histograms and Shapiro Wilk test. Counts and percentages for categorical data are reported. Pearson’s or Spearman’s rank correlation analyses were conducted to assess the degree of linear relationship between continuous variables, as applicable. Partial correlations adjusted by age, race, marital status, number of children, stress and depression were also calculated. Point biserial and polyserial correlation analyses were conducted to assess degree of relationship between dichotomous and continuous measures, or ordinal and continuous measures, respectively. Sleep measures were dichotomized to define “good” and “poor” sleep. Diary measures poor sleep were defined as follows: SOL (> 20 min), WASO (> 30 min) and TST (<7 hrs). Poor sleep for actigraphy measures were defined as: SOL (> 20 min), WASO (> 30 min) and TST (< 6 hrs). Frequencies of “good” and “poor” are reported to further describe the degree of poor sleep of each sleep variable at each time point and across time. Mixed modeling techniques were used to examine whether binary measures of sleep

were associated with BMI and BP. SAS procedure MIXED was used for modeling the main effects of time and sleep, and time by sleep interactions, and to account for within subject correlation. For these models, logarithm base 10 transformations were used for BMI. These models were also adjusted for potential confounders including age, race, marital status whether there was a child at home, IDS with sleep item removed, PSS, pre-pregnancy weight and exercise. All analyses were conducted using SAS, version 9.3 statistical software (SAS Institute Inc., Cary, NC).

RESULTS

Participant characteristics are presented in Table 1. Participants ($n = 161$) were 29 ± 5 years of age and 70.2% of participants self-identified as Caucasian. Over 80% were married or living with a partner and over 37% had at least one child < 18 years of age living at home. Blood pressures of these women reflected the healthy status. Only two women had high BP at 12 weeks ($>140/90$). Table 2 presents the sleep parameters at baseline (10-12 weeks), T2 (14-16 weeks) and T3 (18-20 weeks). Evaluation of the covariates indicated that pre-pregnancy weight was correlated with T1 SBP ($r(93) = 0.46$, $p < 0.0001$) and DBP ($r(93) = 0.39$, $p = 0.0002$) after adjusting for age, race, marital status, number of children, stress and depressive symptoms. Women who were married had shorter diary and actigraphy-assessed sleep onset latency ($r(159) = -0.28$, $p = 0.0003$; $r(146) = -0.24$, $p = 0.0038$), actigraphy-assessed WASO ($r(146) = -0.33$, $p < 0.0001$), and longer actigraphy-assessed sleep duration ($r(146) = 0.26$, $p = 0.0012$). Having more children at home was also associated with longer diary-assessed SOL (T1: $r(153) = 0.22$, $p = 0.0138$), greater weight (T1: $r(148) = 0.23$, $p = 0.0119$), and higher BMI (T1: $r(145) = 0.24$, $p = 0.0102$), but not SBP or DBP. Caucasian women had shorter SOL (diary and actigraphy) (T1: $r(159) = -0.27$, $p = 0.0005$; $r(146) = -0.24$, $p = 0.0035$), less actigraphy-assessed WASO (T1: $r(146) = -0.29$, $p = 0.0004$), and longer actigraphy-assessed sleep duration (T1: $r(146) = 0.32$, $p < 0.0001$).

The next set of correlations assessed whether sleep at T1 (10-12 weeks') was associated with any of the CM factors at each time point (T1, T2 or T3). Diary-assessed SOL at T1 was associated with SBP at T2 after adjusting for covariates ($r(132) = 0.18$, $p = 0.0329$). However, this trend disappeared after correcting for multiple comparisons. Neither diary- nor actigraphy-assessed WASO and TST at T1 were associated with any of the CM factors.

We further evaluated the frequency of disturbed sleep at each time point, and whether sleep disturbance was present, absent or intermittent across all time points. Table 3 presents the frequencies of women who met 1 criterion at each time point, as well as the number and percentage of women who met criteria for persistent, intermittent or no sleep disturbance for each sleep variable. Considering SOL, for instance, we identified 22 (17%) women who had persistent SOL ≥ 20 minutes as measured by diary, yet only 6 (6%) who met similar criteria based on actigraphy. Twenty five (19%) women had long SOL intermittently as measured by diary, with 12 (11%) meeting similar criteria as measured by actigraphy. In contrast to previous reports (15;17), over half of the women 71 (54.6%) had SOL less than 20 minutes based on diary, as well as for actigraphy-assessed SOL, 72 (68%) indicating no concerns with sleep initiation. Table 4 presents the frequencies at each time point of women who met 2 or 3 criteria.

Finally, we examined whether persistent sleep disturbance (present at all time points) was differentially associated with the CM factors. For diary-assessed SOL and WASO, there was a statistically significant interaction between time and sleep group (SOL: $F(2,128) = 3.67$, $p = 0.028$; WASO: $F(2,128) = 3.12$, $p = 0.047$) and a significant within-subjects main effect of time (SOL: $F(2,128) = 164.16$, $p < 0.0001$; WASO: $F(2,128) = 180.16$, $p < 0.0001$) for unadjusted models, revealing that changes in BMI over time are not the same for all groups. These effects remained statistically significant following covariate adjustment for age, race, marital status, exercise, having children at home, the IDS with the sleep item removed, the PSS, and pre-pregnancy weight. Only pre-pregnancy weight was a statistically significant correlate of BMI or BP ($P_s < 0.001$). Figures 1 and 2 show that BMI increases over time for both groups (persistent SOL/WASO and no persistent SOL/WASO). Interestingly, women with intermittent long WASO showed the greater increase in BMI over time. Lastly for actigraphy-assessed WASO and systolic BP, there was a significant within-subjects main effect of time ($F(2,112) = 4.79$, $p = 0.0101$) and a significant interaction between time and sleep group ($F(2,112) = 3.33$, $p = 0.0394$) after controlling for covariates (Figure 3). When evaluating sleep measures as continuous data, diary-assessed WASO and TST, and actigraphy-assessed SOL, WASO, and TST were not associated with BMI.

DISCUSSION

This study evaluated sleep collected concurrently via diary and actigraphy in early pregnancy and two particularly relevant cardiometabolic risk factors, BP and BMI. We found that issues with diary-assessed SOL and WASO (two measures of sleep continuity) were associated with higher BP and/or BMI, which remained significant after controlling for covariates. However, given the cross-sectional nature of the BP and BMI data, we, encourage the reader to interpret the findings with caution. Total sleep time was not associated with either of the CM risk factors. This is in contrast to a report by Williams et al (9). However, our BP assessments were in early pregnancy when BP is typically lower than pre-pregnancy values or in later pregnancy (18). To our surprise, we observed no significant associations among the actigraphy-assessed variables and BP or BMI. While actigraphy has been validated in the general population, it has yet to be thoroughly studied in pregnant women. Actigraphy determines sleep or wake by measuring nocturnal movement. In pregnancy there are multiple factors that may cause a woman to have more movement during sleep without actually awakening (19). Further studies are needed in order to better interpret actigraphy data in the pregnant population.

Several studies have noted an association between sleep disturbance, especially poor sleep quality, and higher blood pressure (8;20). We noted a positive relationship between (T1) actigraphy assessed WASO > 30 minutes and blood pressure across all time points, and this association remained significant after adjusting for covariates. Furthermore, we found that women with longer WASO did not exhibit the typical decrease and gradual rise in blood pressure that is often observed in the later part of the second trimester. Blood pressure fluctuates in women entering the second trimester due to a series of normal vascular adaptations (18). Our findings indicate that women with longer WASO show a blunted change in their blood pressure when compared to those with average WASO (Figure 3). While suggestive, further studies, including 24-hour ambulatory BP measurements, are

needed to better understand the patterns and mechanisms associated with sleep disturbance and changes in blood pressure, and how this may impact health in later pregnancy. Moreover, the criteria used to determine irregular blood pressure values may not be sufficient to detect a signal. Future studies are needed to determine optimal cutoff values for defining abnormal sleep as well as critical BP values in early pregnancy.

The relationship between SOL and BMI was diminished to a trend after controlling for the covariates of age, race, marital status, having children at home, stress, depressive symptoms and prepregnancy weight. This finding is consistent with the published literature noting that stressful life events are associated with specific aspects of sleep disturbance, like long SOL (21). Additionally, several studies have shown that long WASO, as well as short sleep duration, are linked to higher BMI (22;23). Our findings do not support previous findings. However, this may be a reflection of the population under study. Women are expected to gain weight during pregnancy (24) and report of substantial sleep disturbance is considered the norm (25). The association between SOL and BMI is in line with the hypothesis that sleep disturbance is associated with increased BMI (8;26). We speculate that women who report disturbances in several domains of sleep, including difficulty initiating and maintaining sleep, as well as shorter sleep duration and poor quality, may be at the greatest risk for gaining excessive weight or developing high BP. The cell sizes were insufficient to adequately address this question. While all women gain weight during pregnancy, it is the degree of change in relation to pre-pregnancy BMI that is particularly relevant to risk for adverse outcomes (27;28). When analyzing the degree of change in BMI across time, women with longer SOL had higher BMI at each time point. Our data also suggest that women who have longer SOL also experienced more stress than their counterparts. Therefore, it is possible that stress via sleep disturbance may account for the increased BMI. Not surprisingly, pre-pregnancy weight was a significant correlate in both models examining BMI. The degree to which women gain weight following conception is a strong determinant of future maternal morbidity (28). Women who have low net gestational weight gain are also at risk of adverse pregnancy outcomes. Further evaluation of these bidirectional relationships is warranted.

Lastly, given the evidence that chronic sleep disturbance, rather than cross-sectionally measured sleep, is associated with greater morbidity (29), we identified those women who had persistent sleep disturbance and compared them to women with no sleep disturbance. Although the cell sizes were not powered to detect a difference, we did observe that women with persistent SOL ≥ 20 minutes had higher BMI values at all time points. Additional studies that follow women throughout pregnancy are needed to determine whether chronic sleep disturbance is necessary to substantially increase risk for morbidity.

These findings add to the growing evidence that sleep disturbance in pregnancy may play a role in adverse pregnancy outcomes (30). In accordance with previously published studies, our data suggest that disturbed sleep as early as 10-12 weeks' gestation may affect CM risk factors known to adversely affect health and outcomes (9). We do however acknowledge some limitations in the sample population and measures which temper our interpretations. While some may argue that PSG (polysomnography) is the gold standard with which to record sleep information, we contend that the use of actigraphy in this study is a better

measurement of sleep. Unlike PSG, actigraphy affords the opportunity to capture change over time and evaluate variability in pregnant sleep. Another limitation is that we only assessed BP once per visit rather than the recommended two or more readings separated by 2 minutes (31). We also did not have data after 20 weeks. It is feasible that we would see other associations if we had data from across all of pregnancy. Moreover, a single BP measurement does not allow for the examination of diurnal variation. However, few studies have collected 24-hour ambulatory BP in pregnant women due to high participant burden (32). Future studies need to utilize this method to account for nocturnal blood pressure and associated dipping. Nevertheless, all physiologic measurements were taken between 7:00am and 10:00am to control for time effects. Finally, we sought to recruit healthy, low risk women in order to mitigate the effects of various comorbidities. However, we were limited to self-report of sleep disorders, psychopathology, other illnesses and medication use. Reliance on self-report can result in some participants entering the study with an undiagnosed condition, which could influence the findings. Particularly relevant here is the absence of data regarding snoring or sleep disordered breathing (SDB) at each of the time points. While only approximately 10% of pregnant women have SDB in early pregnancy (33), estimates of frequent snoring are around 20% (30). Hence, future studies need to assess snoring and/or SDB in early pregnancy studies on sleep.

This study lays ground work for future endeavors assessing sleep in pregnancy and sheds light on the importance of clinical evaluation of sleep patterns in early pregnancy. The significant morbidity associated with elevated maternal blood pressure and BMI make early detection important to avoid subsequent morbidity. One pertinent aspect of sleep is that it is a modifiable behavior. A growing literature confirms the utility of behavioral interventions to ameliorate sleep disturbances (34). While this has not been evaluated in pregnant women, we speculate that this option may be especially pertinent given that there are few interventions that can be implemented in pregnancy. A better understanding of the role sleep may play in the development and/or progression of maternal hypertension and unhealthy BMI during pregnancy may prove beneficial for the incorporation of future screening, evaluation, and treatment of these conditions.

Acknowledgments

Supported by NIH NR010813 and CTSI UL1 RR024153 and UL1 TR000005. The National Institutes of Health specifically disclaim responsibility for any analyses, interpretations, or conclusions. M.L.O. has had full access to the data, and takes full responsibility for the integrity of the data and accuracy of the analysis.

Reference List

1. Kopelman PG. Obesity as a medical problem. *Nature*. Apr 6; 2000 404(6778):635–43. [PubMed: 10766250]
2. Roberts, JM. Pregnancy-Related hypertension. In: Creasey, RK.; Resnik, R., editors. *Maternal-Fetal Medicine*. 5th. Saunders; Philadelphia: 2004. p. 859-99.
3. Hibbard JU, Shroff SG, Lang RM. Cardiovascular changes in preeclampsia. *Semin Nephrol*. Nov; 2004 24(6):580–7. [PubMed: 15529293]
4. Evans CS, Gooch L, Flotta D, Lykins D, Powers RW, Landsittel D, et al. Cardiovascular system during the postpartum state in women with a history of preeclampsia. *Hypertension*. Jul; 2011 58(1):57–62. [PubMed: 21606389]

5. Guelinckx I, Devlieger R, Beckers K, Vansant G. Maternal obesity: pregnancy complications, gestational weight gain and nutrition. *Obes Rev.* Mar; 2008 9(2):140–50. [PubMed: 18221480]
6. Sohlberg S, Stephansson O, Cnattingius S, Wikstrom AK. Maternal body mass index, height, and risks of preeclampsia. *Am J Hypertens.* Jan; 2012 25(1):120–5. [PubMed: 21976280]
7. Watkins ML, Botto LD. Maternal prepregnancy weight and congenital heart defects in offspring. *Epidemiology.* Jul; 2001 12(4):439–46. [PubMed: 11428386]
8. Altman NG, Izci-Balserak B, Schopfer E, Jackson N, Rattanaumpawan P, Gehrman PR, et al. Sleep duration versus sleep insufficiency as predictors of cardiometabolic health outcomes. *Sleep Med.* Dec; 2012 13(10):1261–70. [PubMed: 23141932]
9. Williams MA, Miller RS, Qiu C, Cripe SM, Gelaye B, Enquobahrie D. Associations of early pregnancy sleep duration with trimester-specific blood pressures and hypertensive disorders in pregnancy. *Sleep.* Oct 1; 2010 33(10):1363–71. [PubMed: 21061859]
10. Althuisen E, van Poppel MN, Seidell JC, van MW. Correlates of absolute and excessive weight gain during pregnancy. *J Womens Health.* Oct; 2009 18(10):1559–66.
11. Hall, M.; Okun, ML.; Atwood, CW.; Buysse, DJ.; Strollo, PJ. Measurement of sleep by polysomnography. In: Luecken, LL.; Gallo, LC., editors. *Handbook of Physiological Research Methods in Health Psychology.* Sage Publications; 2008. p. 341–67.
12. Beebe KR, Lee KA. Sleep disturbance in late pregnancy and early labor. *J Perinat Neonatal Nurs.* Apr; 2007 21(2):103–8. [PubMed: 17505229]
13. Okun ML, Kline CE, Roberts JM, Wettlaufer B, Glover K, Hall M. Prevalence of Sleep Deficiency in Early Gestation and its Associations with Stress and Depressive Symptoms. *J Womens Health (Larchmt).* Oct 12.2013
14. Monk TH, Reynolds CF, Kupfer DJ, Buysse DJ, Coble PA, Hayes AJ, et al. The Pittsburgh Sleep Diary. *J Sleep Res.* 1994; 3:111–20.
15. Okun ML, Coussons-Read ME. Sleep disruption during pregnancy: how does it influence serum cytokines? *J Reprod Immunol.* Apr; 2007 73(2):158–65. [PubMed: 17074396]
16. Rush AJ, Gullion CM, Basco MR, Jarrett RB, Trivedi MH. The Inventory of Depressive Symptomatology (IDS): psychometric properties. *Psychol Med.* May; 1996 26(3):477–86. [PubMed: 8733206]
17. Lee KA, Zaffke ME, McEnany G. Parity and sleep patterns during and after pregnancy. *Obstet Gynecol.* Jan; 2000 95(1):14–8. [PubMed: 10636494]
18. Grindheim G, Estensen ME, Langesaeter E, Rosseland LA, Toska K. Changes in blood pressure during healthy pregnancy: a longitudinal cohort study. *J Hypertens.* Feb; 2012 30(2):342–50. [PubMed: 22179091]
19. Sahota PK, Jain SS, Dhand R. Sleep disorders in pregnancy. *Curr Opin Pulm Med.* Nov; 2003 9(6):477–83. [PubMed: 14534398]
20. Knutson KL, Van CE, Rathouz PJ, Yan LL, Hulley SB, Liu K, et al. Association between sleep and blood pressure in midlife: the CARDIA sleep study. *Arch Intern Med.* Jun 8; 2009 169(11):1055–61. [PubMed: 19506175]
21. Schafer V, Bader K. Relationship Between Early-life Stress Load and Sleep in Psychiatric Outpatients: A Sleep Diary and Actigraphy Study. *Stress Health.* Jun 15.2012
22. Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB. Inadequate sleep as a risk factor for obesity: analyses of the NHANES I. *Sleep.* Oct 1; 2005 28(10):1289–96. [PubMed: 16295214]
23. Spiegel K, Knutson K, Leproult R, Tasali E, Van Cauter E. Sleep loss: a novel risk factor for insulin resistance and Type 2 diabetes. *J Appl Psychol.* Nov; 2005 99(5):2008–19.
24. American Congress of Obstetricians and Gynecologists (ACOG). *Obesity in Pregnancy.* Committee on Obstetric Practice; 2013.
25. Okun, ML. Sleep in Pregnancy and the Postpartum. In: Kushida, CA., editor. *Encyclopedia of Sleep.* Academic Press; Waltham: 2013. p. 674–9.
26. Grandner MA, Perlis ML. Insomnia as a cardiometabolic risk factor. *Sleep.* Jan; 2013 36(1):11–2. [PubMed: 23288965]

27. Guendelman S, Pearl M, Kosa JL, Graham S, Abrams B, Kharrazi M. Association Between Preterm Delivery and Pre-pregnancy Body Mass (BMI), Exercise and Sleep During Pregnancy Among Working Women in Southern California. *Matern Child Health J.* Jul 11.2012
28. Heude B, Thiebaugeorges O, Goua V, Forhan A, Kaminski M, Foliguet B, et al. Pre-pregnancy body mass index and weight gain during pregnancy: relations with gestational diabetes and hypertension, and birth outcomes. *Matern Child Health J.* Feb; 2012 16(2):355–63. [PubMed: 21258962]
29. Patel SR, Ayas NT, Malhotra MR, White DP, Schernhammer ES, Speizer FE, et al. A prospective study of sleep duration and mortality risk in women. *Sleep.* May 1; 2004 27(3):440–4. [PubMed: 15164896]
30. Facco FL, Kramer J, Ho KH, Zee PC, Grobman WA. Sleep disturbances in pregnancy. *Obstet Gynecol.* Jan; 2010 115(1):77–83. [PubMed: 20027038]
31. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation.* Feb 8; 2005 111(5):697–716. [PubMed: 15699287]
32. Booker CJ, Dodson WC, Kunselman AR, Repke JT, Legro RS. Twenty-four-hour ambulatory blood pressure monitor heart rate: a potential marker for gestational hypertension in at-risk women. *Am J Perinatol.* May; 2012 29(5):339–46. [PubMed: 22147639]
33. Frederick IO, Qiu C, Sorensen TK, Enquobahrie DA, Williams MA. The prevalence and correlates of habitual snoring during pregnancy. *Sleep Breath.* May; 2013 17(2):541–7. [PubMed: 22907649]
34. Davidson JR. Insomnia treatment options for women. *Obstet Gynecol Clin North Am.* Dec; 2009 36(4):831–xi. [PubMed: 19944303]

Highlights

- Pre-pregnancy weight is correlated with blood pressure at 20 weeks
- Difficulty initiating sleep in early pregnancy is associated with higher BMI
- Difficulty initiating sleep in early pregnancy is associated with higher blood pressure
- Stress and difficulty initiating sleep manifests into greater weight gain and higher blood pressure

Figure 1.

Estimated logarithm body mass index (log BMI) values for women with persistent long diary-assessed SOL (≥ 20 minutes, solid line) and without persistent long diary-assessed SOL (< 20 minutes, dash line) are expected means from the linear mixed-effects model with covariate adjustment for age, race, marital status, whether she has a child at home, IDS with sleep item removed, PSS, pre-pregnancy weight and exercise. Error bars indicate 95% confidence intervals. Linear mixed-effects model analysis showed significant main effect for time ($F(2,69)=127.09$, $p<0.0001$). Results show that BMI is increasing over time for both sleep groups (persistent SOL and no persistent SOL)..

Figure 2.

Estimated logarithm body mass index (log BMI) values for women with persistent long diary-assessed WASO (≥ 30 minutes, solid line) and without persistent long diary-assessed WASO (< 30 minutes, dash line) are expected means from the linear mixed-effects model with covariate adjustment for age, race, marital status, whether she has a child at home, IDS with sleep item removed, PSS, pre-pregnancy weight and exercise. Error bars indicate 95% confidence intervals. Linear mixed-effects model analysis showed significant main effect for time ($F(2,69)=138.62$, $p<0.0001$). Results show that BMI is increasing over time for both sleep groups (persistent WASO and no persistent WASO)..

Figure 3.

Estimated systolic blood pressure (Systolic BP) values for women with persistent long actigraphy-assessed WASO (≥ 30 minutes, solid line) and without persistent long actigraphy-assessed WASO (< 30 minutes, dash line) are expected means from the linear mixed-effects model with covariate adjustment for age, race, marital status, whether she has a child at home, IDS with sleep item removed, PSS, pre-pregnancy weight and exercise. Error bars indicate 95% confidence intervals. Linear mixed-effects model analysis showed significant main effect for time ($F(2,112)=4.79$, $p<0.0101$) and significant interaction between sleep group and time ($F(2,112)=3.33$, $p=0.0394$). Results show that Systolic BP decreases over time for the no persistent actigraphy-assessed WASO group, and remains constant across time for the persistent actigraphy-assessed WASO group.

Table 1

Subject Demographics

	Mean $\pm$ SD	N
Age (yrs)	29 $\pm$ 5 Range (19-40)	161
Pre-pregnancy weight (lbs)	159.5 $\pm$ 40.0 Range 97.0 - 332)	119
Weight at T1 (lbs)	159.9 $\pm$ 40.4 Range (88.4-343.5)	154
BMI at T1 (kg/m ²)	26.6 $\pm$ 6.2 Range (15.2 – 55.2)	151
Blood Pressure at T1 Diastolic	67.6 $\pm$ 8.5 Range (44 -92)	151
Systolic	108.0 $\pm$ 12.6 Range (80-150)	151
Perceived Stress Scale	13.9 (6.3) Range (0-33)	
Inventory of Depressive Symptoms (sleep item removed)	4.9 (3.0) Range (0-19)	
	N (%)	
Race	113 (70.2)	
Caucasian	48 (29.8)	
African-American/Other		
Marital Status	104 (65.8)	
Married	27 (17.1)	
Living with Partner	1 (0.6)	
Divorced	26 (16.5)	
Never Married		
Number of Children	96 (61.9)	
0	44 (28.4)	
1	10 (6.5)	
2	5 (3.2)	
3+		

Table 2

Sleep Characteristics at All Time Points

Sleep Measure (minutes)	N	T1 (10-12 weeks) Mean (SD)	N	T2 (14-16 weeks) Mean (SD)	N	T3 (18-20 weeks) Mean (SD)
Diary						
SOL	161	20.0 (14.9)	138	15.7 (12.6)	133	16.8 (12.8)
WASO	161	24.2 (18.2)	138	20.5 (16.6)	133	20.1 (14.3)
TST	161	460.9 (52.0)	138	459.3 (55.7)	133	454.1 (53.8)
Actigraphy						
SOL	148	14.7 (11.4)	124	12.5 (9.0)	122	12.9 (10.3)
WASO	148	90.8 (56.6)	124	78.5 (42.6)	122	71.5 (37.6)
TST	148	390.0 (63.5)	124	390.5 (57.9)	122	395.9 (50.8)

SOL = Sleep onset latency; WASO = Wake after sleep Onset; TST = total sleep duration

Table 3

Frequency of women who met individual criteria for sleep disturbance

Sleep Measure	T1 N (%)	T2 N (%)	T3 N (%)	Persistent Sleep Disturbance N (%)	Intermittent Sleep Disturbance N (%)	No Sleep Disturbance N(%)
Diary						
SOL 20 mins	60 (37.3)	34 (24.6)	39 (29.3)	22 (17.0)	25 (19.2)	71 (54.6)
WASO 30 mins	44 (27.3)	34 (24.6)	23 (17.3)	14 (11.0)	28 (21.5)	79 (60.8)
TST < 7 hrs	39 (24.2)	28 (20.3)	35 (26.3)	15 (12.0)	18 (13.9)	83 (63.9)
Actigraphy						
SOL 20 mins	31 (21.0)	20 (16.1)	22 (18.0)	6 (6.0)	12 (11.3)	72 (67.9)
WASO 30 mins	145 (98.0)	121 (97.6)	116 (95.0)	96 (91.0)	1 (.94)	0 (0)
TST < 7 hrs	38 (25.7)	36 (29.0)	25 (20.5)	13 (12.0)	16 (15.1)	67 (63.2)

Persistent sleep disturbance = poor sleep at T1, T2 and T3.

No sleep disturbance = no poor sleep at T1, T2 and T3.

Intermittent sleep disturbance = poor at T1 but not T2 and T3, or poor at T2 but not T1 and T3, or poor at T3 but not T1 and T2

Table 4

Frequency of women who met two or more criteria for sleep disturbance

Sleep Measure	T1 N (%)	T2 N (%)	T3 N (%)
Diary			
SOL 20 mins + TST < 7 hrs	18 (11.2)	11 (8.0)	16 (12.0)
WASO 30 mins + TST < 7 hrs	15 (9.3)	11 (8.0)	12 (9.0)
SOL 20 mins + WASO 30 mins	23 (14.3)	14 (10.1)	14 (10.5)
Actigraphy			
SOL 20 mins + TST < 6 hrs	15 (10.1)	10 (8.1)	7 (5.7)
WASO 30 mins + TST < 6 hrs	38 (25.7)	35 (28.2)	23 (18.9)
SOL 20 mins + WASO 30 mins	31 (21.0)	20 (16.1)	22 (18.0)
All Diary-assessed criteria	9 (5.6)	7 (5.1)	8 (6.0)
All Actigraphy-assessed criteria	15 (10.1)	10 (8.1)	7 (5.7)