Published in final edited form as:

Chem Commun (Camb). 2011 September 28; 47(36): 10025–10027. doi:10.1039/c1cc13597j.

Characterization and Application of the Fe(II) and α -Ketoglutarate Dependent Hydroxylase FrbJ

Matthew A. DeSieno^{a,c}, Wilfred A. van der Donk^{b,c}, and Huimin Zhao^{a,b,c,d}

Huimin Zhao: zhao5@illinois.edu

^aDepartment of Chemical and Biomolecular Engineering, University of Illinois at Urbana-Champaign, 600 S. Mathews Avenue, Urbana, IL 61801 (USA), Fax: +1-217- 333-5052 Tel: +1-217- 333-2631

^bDepartments of Chemistry and Biochemistry, University of Illinois at Urbana-Champaign

^cInstitute for Genomic Biology, University of Illinois at Urbana-Champaign

^dDepartment of Bioengineering, Center for Biophysics, and Computational Biology, University of Illinois at Urbana-Champaign

Abstract

The Fe(II) and α -ketoglutarate-dependent hydroxylase FrbJ was previously demonstrated to utilize FR-900098 synthesizing a second phosphonate FR-33289. Here we assessed its ability to hydroxylate other possible substrates, generating a library of potential antimalarial compounds. Through a series of bioassays and *in vitro* experiments, we identified two new antimalarials.

The gene *frbJ* was identified during the cloning and sequencing of the biosynthetic gene cluster for the phosphonate antibiotic FR-900098 from *Streptomyces rubellomurinus*.¹ The enzyme is homologous to Fe(II) and α -ketoglutarate (α KG)-dependent hydroxylases, which catalyze substrate hydroxylation with concomitant oxidative decomposition of α KG to succinate and carbon dioxide.² Enzymes within this family play a pivotal role in a diverse set of reactions, including antibiotic synthesis where they generate pathway precursors, intermediates, and final products.^{3–4} For example, clavaminic synthase (CAS) catalyzes three independent oxidative reactions in the biosynthesis of clavulanic acid.⁵ Interestingly, another enzyme from this family has demonstrated epoxidase activity in the semisynthetic pathway for the phosphonate antibiotic fosfomicin. The enzyme encoded by *epoA* from *Penicillium decumbens* converts the substrate *cis*-propenylphosphonic acid to the final product fosfomicin.⁶ Although the natural function of this enzyme in the native host is unknown, the epoxidase activity signifies the broad range of functions of Fe(II) and α KG-dependent enzymes.

Deletion studies of *frbJ* within the heterologous hosts *Streptomyces lividans* and *Escherichia coli* indicated that the enzyme was not required for FR-900098 biosynthesis, leading to some

uncertainty as to its function.^{1, 7} It was originally postulated that FrbJ might be responsible for the *N*-hydroxylation step; however, catalysis of this nature by this class of enzyme would have been exceedingly rare. Fe(II) and α KG-dependent enzymes can efficiently catalyze the hydroxylation of inactivated C-H bonds, so we investigated whether FrbJ might be involved in the synthesis of another phosphonate antibiotic FR-33289 ((*R*)-2-hydroxy-3-(*N*-acetyl-*N*-hydroxyamino)propylphosphonic acid).⁸⁻⁹ Production of this compound was observed in both the native producer *S. rubellomurinus* producer strain and a heterologous *S. lividans* 4G7 host. We have recently shown that FrbJ is indeed capable of catalyzing the hydroxylation of FR-900098 at the β -position (Fig. 1).⁷ This study focused on the expression, purification, and further biochemical characterization of FrbJ and the determination of its role in FR-900098 biosynthesis. We also demonstrated its ability to catalyze the hydroxylation of additional substrates as part of the creation of a new library of potential antimalarial drugs, all of which were screened for inhibitory activity against a malarial drug target.

The full-length *frbJ* gene was cloned from fosmid 4G7.⁷ *E. coli* BL21(DE3) cells were transformed with this construct and IPTG-induced cultures were purified to homogeneity. As previously described, the final steps in the pathway contain a series of intermediates conjugated to the nucleotide cytidine-5'-monophosphate (CMP).⁷ If hydroxylation by FrbJ were to occur before cleavage of the CMP by FrbI, CMP-5'-FR-900098 could serve as the true substrate for the enzyme. FrbJ showed activity towards the natural substrate FR-900098 but not the nucleotide containing substrate, indicating that nucleotide cleavage must occur before hydroxylation in the biosynthetic pathway (Table 1). We also checked for enzymatic activity with another known phosphonate antibiotic, fosmidomycin (FR-31564). This substrate contains an *N*-formyl group instead of the *N*-acetyl group on FR-900098. FrbJ demonstrated activity towards this structurally similar substrate, but displayed a 6-fold preference for the natural substrate. As a result of fosmidomycin hydroxylation by FrbJ, a novel phosphonate was able to be synthesized, (*R*)-2-hydroxy-3-(*N*-formyl-*N*-hydroxyamino)propylphosphonic acid, hereby called hydroxyfosmidomycin. To the best of our knowledge, this compound has never been detected from a biological source or produced synthetically.

Heterologous production of FR-33289 in *E. coli* was attempted by expressing all of the genes in the biosynthetic cluster, excluding *frbI* (pETDuet-*frbABCD*, pRSFDuet-*frbHFG*, pACYCDuet-*frbEJ-dxrB*). Although FR-900098 production was evident in the LC-MS analysis, no detectable amounts of FR-33289 were detected. Cell-free lysate of *E. coli* strains that only expressed FrbJ was incubated with FR-900098. The lysate only demonstrated conversion to FR-33289 upon addition of exogenous α -ketoglutarate, suggesting that FrbJ may be regulated by the intracellular concentration of this cofactor, thus preventing *in vivo* production in *E. coli*.

The biosynthesis of secondary metabolites can be effectively regulated via the availability of precursors, cofactors, and cellular energy originating from primary metabolism. Rate-limiting enzymes control the flux of these critical metabolites to ensure that secondary metabolism does not become favored over primary metabolism. Khetan *et al.* described the function of lysine-6-aminotransferase (LAT), which was determined to be the rate-limiting

enzyme in cephamycin biosynthesis from *Streptomyces clavuligerus*.¹⁰ This enzyme, which utilizes α KG as a cofactor in the conversion of lysine to α -aminoadipic acid, is the first step in the biosynthetic pathway. The authors measured the K_M for α KG to be 8.6 mM, substantially higher than its intracellular concentration ranging from 0.3–1.3 mM.¹⁰ The K_M value for LAT is also higher than that of other characterized Fe(II) and α KG-dependent hydroxylases, which is typically on the order of 10s of micromolar.^{11–13} In the same way that LAT regulates the cephamycin biosynthetic pathway, FrbJ may regulate the ratio of the two phosphonates FR-900098 and FR-33289 using α -ketoglutarate. The K_M of α KG for the oxidation of FR-900098 by FrbJ was determined to be 1.3 ± 0.19 mM, comparable to the high K_M of LAT and close to the value of the intracellular level of α KG. One crucial difference between the two pathways is that the entire cephamycin biosynthesis can be regulated by the intracellular concentration of α KG since LAT is the first enzyme in the pathway, making the high K_M of the enzyme ideal for switching the biosynthesis of cephamycin entirely on or off. The lower K_M of FrbJ is a much weaker control valve, which can regulate the conversion of FR-900098 to FR-33289.

The dianionic charge of phosphonates at physiological pH poses problems for their efficacy; in particular, they suffer from poor bioavailability as a result of their high polarity. The oral bioavailability of FR-900098 and fosmidomycin is moderate with a resorption rate of approximately 30%.¹⁴ Recent efforts have focused on masking the charged moiety of the phosphonate group through esterification. Such chemically synthesized prodrugs cross the gastrointestinal tract into the blood stream with higher efficiency, where they are cleaved by broad specificity plasma esterases, thus releasing the free and active phosphonate.^{15–17} We sought a method for the preparation of similar esterified FR-900098 prodrugs through biosynthetic means. The gene cluster for dehydrophos from *Streptomyces luridus* contains an *S*-adenosyl-*L*-methionine (SAM)-dependent *O*-methyltransferase encoded by *dhpl*.¹⁸ Dhpl has very low substrate specificity,¹⁹ and is able to *O*-methylate the phosphonate of a wide range of substrates, making it an excellent candidate for further engineering.

As shown in Table 1, FrbJ is capable of catalyzing the hydroxylation of FR-900098 and fosmidomycin to produce FR-33289 and hydroxyfosmidomycin, respectively. These four phosphonates were used as the basis for the creation of a prospective antimalarial library; each was methylated by Dhpl (Fig. 2). FR-900098 and fosmidomycin were commercially available and the subsequent six compounds were synthesized by *in vitro* reactions with purified FrbJ, Dhpl, and Pfs. The phosphonate library was then purified to remove any unreacted substrate or cofactors. Although not identical to the di-ester prodrugs prepared by chemical synthesis, these methylated compounds may enhance the bioavailability of the final active phosphonate drugs.

The library was first screened by bioassays of an *E. coli* phosphonate uptake strain.¹ Bioassay plates of this FR-900098-sensitive strain showed clear inhibition zones around discs impregnated with FR-900098, FR-33289, fosmidomycin, and hydroxyfosmidomycin; all of which were lost with a resistant *E. coli* strain. However, there was no detectable growth inhibition for any of the corresponding methylated compounds with the FR-900098-sensitive strain. To ensure the lack of bioactivity on the plate was not specific for the Dxr target, we utilized a second phosphonate antibiotic with a completely different target.

Fosfomycin, a broad-range antibacterial used to treat urinary tract infections,²⁰ and the corresponding fosfomycin-OMe was prepared in a similar method as described for the other phosphonates. Similar to the prospective methylated antimalarials, there was no growth inhibition of the sensitive *E. coli* strain by fosfomycin-OMe. The methyl group is either not cleaved in these *in vivo* bioassays, preventing the release of the active phosphonate, or the methyl ester hampers uptake.

The library was also assessed using purified *Plasmodium falciparum* Dxr (pfDxr). The recombinant enzyme was codon-optimized for heterologous expression in *E. coli* and was designed to lack the N-terminal signal peptide.^{21–22} Enzymatic activity was monitored by NADPH oxidation at 340 nm, and subsequent inhibition values for the IC₅₀ and K_I were obtained for each of the prospective antimalarial compounds. Similar to the *in vivo* bioassays, only the free, unmethylated phosphonate compounds demonstrated *in vitro* inhibition of pfDxr. There was no apparent decrease in the enzymatic activity upon addition of the methylated compounds to any of the reaction mixtures. The approximately two-fold difference between the inhibition of FR-900098 and fosmidomycin is similar to published data²¹. Also, the inhibition constants for FR-900098 and fosmidomycin with their hydroxylated counterparts, FR-33289 and hydroxyfosmidomycin, respectively, were determined to be nearly identical to each other (Table 2). Even though the natural pfDxr substrate also contains a secondary hydroxyl group similar to FR-33289 and hydroxyfosmidomycin, albeit at different positions on the carbon backbone, there was no apparent increase in inhibition compared to FR-900098 and fosmidomycin.

The combinatorial biosynthesis with FrbJ was successful in producing two new antimalarial target compounds, FR-33289 and hydroxyfosmidomycin. The *E. coli* bioassays showed some improvement for these hydroxylated compounds as the zones of inhibition were slightly larger. Although the inhibition of the Dxr enzyme itself does not change, the bioavailability of these hydroxylated compounds appears to increase. In order to more accurately assess these compounds, we will look to expand the scope of the inhibition assays to include *Plasmodium* species.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by a grant from the National Institute of General Medical Sciences GM077596 (H.Z. and W.A.V). M.A.D. acknowledges support from the US National Institutes of Health under Ruth L. Kirschstein National Research Award 5 T32 GM070421 from the National Institute of General Medical Sciences.

Notes and references

1. Eliot AC, Griffin BM, Thomas PM, Johannes TW, Kelleher NL, Zhao H, Metcalf WW. Chem Biol. 2008; 15:765–770. [PubMed: 18721747]
2. Hausinger RP. Crit Rev Biochem Mol Biol. 2004; 39:21–68. [PubMed: 15121720]
3. Kovaleva EG, Lipscomb JD. Nat Chem Biol. 2008; 4:186–193. [PubMed: 18277980]
4. Schofield CJ, Zhang Z. Curr Opin Struct Biol. 1999; 9:722–731. [PubMed: 10607676]

5. Lloyd MD, Merritt KD, Lee V, Sewell TJ, Wha-Son B, Baldwin JE, Schofield CJ, Elson SW, Baggaley KH, Nicholson NH. *Tetrahedron*. 1999; 55:10201–10220.
6. Watanabe M, Sumida N, Murakami S, Anzai H, Thompson CJ, Tateno Y, Murakami T. *Appl Environ Microbiol*. 1999; 65:1036–1044. [PubMed: 10049860]
7. Johannes TW, DeSieno MA, Griffin BM, Thomas PM, Kelleher NL, Metcalf WW, Zhao H. *Chem Biol*. 2010; 17:57–64. [PubMed: 20142041]
8. Okuhara M, Kuroda Y, Goto T, Okamoto M, Terano H, Kohsaka M, Aoki H, Imanaka H. *J Antibiot (Tokyo)*. 1980; 33:24–28. [PubMed: 6768705]
9. Kuroda Y, Okuhara M, Goto T, Okamoto M, Terano H, Kohsaka M, Aoki H, Imanaka H. *J Antibiot (Tokyo)*. 1980; 33:29–35. [PubMed: 7372547]
10. Khetan A, Malmberg LH, Kyung YS, Sherman DH, Hu WS. *Biotechnol Prog*. 1999; 15:1020–1027. [PubMed: 10585184]
11. White AK, Metcalf WW. *J Biol Chem*. 2002; 277:38262–38271. [PubMed: 12161433]
12. Eichhorn E, van der Ploeg JR, Kertesz MA, Leisinger T. *J Biol Chem*. 1997; 272:23031–23036. [PubMed: 9287300]
13. Vaz FM, Ofman R, Westinga K, Back JW, Wanders RJ. *J Biol Chem*. 2001; 276:33512–33517. [PubMed: 11431483]
14. Tsuchiya T, Ishibashi K, Terakawa M, Nishiyama M, Itoh N, Noguchi H. *Eur J Drug Metab Pharmacokinet*. 1982; 7:59–64. [PubMed: 7067725]
15. Reichenberg A, Wiesner J, Weidemeyer C, Dreiseidler E, Sanderbrand S, Altincicek B, Beck E, Schlitzer M, Jomaa H. *Bioorg Med Chem Lett*. 2001; 11:833–835. [PubMed: 11277531]
16. Ortmann R, Wiesner J, Reichenberg A, Henschker D, Beck E, Jomaa H, Schlitzer M. *Bioorg Med Chem Lett*. 2003; 13:2163–2166. [PubMed: 12798327]
17. Wiesner J, Ortmann R, Jomaa H, Schlitzer M. *Arch Pharm (Weinheim)*. 2007; 340:667–669. [PubMed: 17994601]
18. Circello BT, Eliot AC, Lee JH, van der Donk WA, Metcalf WW. *Chem Biol*. 2010; 17:402–411. [PubMed: 20416511]
19. Lee JH, Bae B, Kuemin M, Circello BT, Metcalf WW, Nair SK, van der Donk WA. *Proc Natl Acad Sci U S A*. 2010; 107:17557–17562. [PubMed: 20876132]
20. Hendlin D, Stapley EO, Jackson M, Wallick H, Miller AK, Wolf FJ, Miller TW, Chaiet L, Kahan FM, Foltz EL, Woodruff HB, Mata JM, Hernandez S, Mochales S. *Science*. 1969; 166:122–123. [PubMed: 5809587]
21. Jomaa H, Wiesner J, Sanderbrand S, Altincicek B, Weidemeyer C, Hintz M, Turbachova I, Eberl M, Zeidler J, Lichtenthaler HK, Soldati D, Beck E. *Science*. 1999; 285:1573–1576. [PubMed: 10477522]
22. Bae B, Cobb RE, Li Z, DeSieno MA, Zhao H, Nair SK. submitted.

Fig. 1.

FrbJ catalyzes the hydroxylation of FR-900098 to FR-33289 with concomitant oxidative decomposition of α KG to succinate and carbon dioxide.

Fig. 2. Prospective antimalarial library created using FrbJ and Dhpl (with Pfs). Due to the low substrate specificity of Dhpl, hydroxylation by FrbJ was carried out before *O*-methylation.

Table 1

Kinetic parameters of FrbJ.

Substrate	k_{cat} (min^{-1}) (mean $\pm$ SD)	K_{M} (μM) (mean $\pm$ SD)	$k_{\text{cat}}/K_{\text{M}}$ ($\text{mM}^{-1} \text{min}^{-1}$)
FR-900098	147 $\pm$ 31	289 $\pm$ 48	509
Fosmidomycin	57.3 $\pm$ 8.1	700 $\pm$ 137	81.9
CMP-5'-FR-900098			NR

NR: no reaction detected; SD: standard deviation

Table 2

Inhibition of purified Dxr from *Plasmodium falciparum* by members of the phosphonate library.

Substrate	IC ₅₀ (nM) (mean ± SD)	K _I (nM) (mean ± SD)
FR-900098	15 ± 3	3.2 ± 1.0
FR-33289	16 ± 3	3.0 ± 0.8
Fosmidomycin	36 ± 6	6.8 ± 1.6
Hydroxyfosmidomycin	41 ± 5	7.1 ± 1.4
FR-900098-OMe	NI	NI
FR-33289-OMe	NI	NI
Fosmidomycin-OMe	NI	NI
Hydroxyfosmidomycin-OMe	NI	NI

NI: no enzyme inhibition detected; SD: standard deviation