Published in final edited form as:

Methods Mol Biol. 2013 ; 1032: 271–286. doi:10.1007/978-1-62703-496-8_21.

Use of the Cockroach Antigen Model of Acute Asthma to Determine the Immunomodulatory Role of Early Exposure to Gastrointestinal Infection

Carolyn G. Durham,

Departments of Cell, Developmental and Integrative Biology and Pathology, University of Alabama at Birmingham

Lisa M. Schwiebert, and

Departments of Cell, Developmental and Integrative Biology, University of Alabama at Birmingham

Robin G. Lorenz*

Departments of Cell, Developmental and Integrative Biology and Pathology, University of Alabama at Birmingham

Summary

The increased incidence of asthma over the last 50 years in developed countries has been associated with a decrease in infections acquired early in childhood. These early infections are thought to shape subsequent immune responses. Although there have been multiple clinical associations between gastrointestinal infections and decreased asthma incidence, it has been difficult to move beyond a simple correlation when studying human patients. This section describes an acute asthma model in C57BL/6 mice designed to specifically evaluate the effect of prior gastric *Helicobacter* colonization and inflammation in a murine model of cockroach allergen-induced asthma.

Keywords

asthma; helicobacter; gastritis; murine; cockroach antigen; hygiene hypothesis; toll-like receptors; neonatal

1. Introduction

In 1989, Strachan proposed the hygiene hypothesis to answer the growing epidemiological trend of the rise of industrialization, decline in infectious diseases, and subsequent increase in diseases associated with a T-helper 2 (Th2) cell response, including asthma.[1] Despite the growing evidence for this hypothesis, the immunological mechanism by which decreasing infection rates and increasing Th2 diseases occurs has not been elucidated.[2]

*Address correspondence and reprint requests to: Dr. Robin G. Lorenz, Department of Pathology, University of Alabama at Birmingham, 1825 University Blvd., SHEL 602, Birmingham, AL 35294-2182. rlorenz@uab.edu. Ph: 205-934-0676. Fax: 205-996-9113.

CGD Current Address: Department of Medicine, University of North Carolina, Chapel Hill, NC 27599

Numerous studies have shown a role for viruses in asthma incidence. In fact, Strachan, et al. showed that newborns who were exposed to respiratory viruses had an increased incidence of asthma.[3] However, there is a growing body of research that suggests that immune system development, spurred by early colonization with bacteria, decreases the rate of asthma development later in life.[2] For example, studies have shown that children reared in a household with a dog or whose bedrooms had high endotoxin levels were at a lower risk for developing asthma.[4,5] Additionally, numerous studies have shown that living on a farm dramatically reduces the risk for asthma, presumably because of the constant levels of bacterial exposure.[6] This clearly demonstrates the benefit of exposing young children to bacteria and bacterial components in reducing asthma incidence.

In developing countries, many of the infectious diseases are known to elicit a Th1 phenotype. One of these bacteria is *Helicobacter pylori*, which is endemic in many developing countries. As countries become more industrialized, antibiotic therapy is more widespread and these infectious diseases decline. This is especially noted when the incidence of infection is higher in adults, since they were young when the bacterial exposure was greater, than in children who were reared under more newly established standards of infection control.[7] Interestingly, the timing of bacterial exposure is critical in whether or not the microbial acquisition helps in shaping the immune system or causing a pathogenic problem. A study done with *Helicobacter pylori* infection in children and adults indicated that in children less than 10 years old, IL10 and IFN γ were significantly upregulated when compared to older children and adults.[8] The IL10 increase persisted past 10 years old and was higher than that of non-infected people, indicating that this early exposure promotes a Th1 immune response and that regulation of this response persists.

The bacterial exposure component of the hygiene hypothesis has gained acknowledgment and prompted members of the American Thoracic Society to set forth the notion that bacterial manipulation is a key factor in asthma prevention. Furthermore, they charge that bacterial exposure, especially through the gastrointestinal tract, is fundamental to properly developing a Th1-skewed immune system.[2] A key component of this immune system development is the body being able to detect the presence of the bacteria. Our lab has previously shown that, in C57BL/6 mice, toll-like receptors (TLRs) 2 and 4 are significantly upregulated two weeks after birth.[9] TLRs are pathogen-associated pattern receptors. TLRs 2 and 4 are specific for Gram positive and Gram negative bacterial ligands, respectively. However, in mice born to dams on a broad spectrum antibiotic cocktail and who are then weaned onto this cocktail, this upregulation does not occur.[9] This indicates that the ability to sense bacteria is critical in immune system development. Our lab has also shown that mice who were born to parents that had been on broad spectrum antibiotics and were, themselves, maintained on this regimen, had delayed Th1/17 development.[9] These data demonstrate the need for bacterial recognition and bacterial exposure in immune system development. To evaluate the role of bacteria in the hygiene hypothesis, we have developed a model where gastritis is induced by infection with the gastrointestinal bacteria *Helicobacter felis*; and subsequently asthma is induced using cockroach antigen (CRA).

Helicobacter pylori infection is endemic in most developing countries, affecting at least 70% of the population and colonizing about 20% of the United States population.

Physiologically, the presence of *H. pylori* in the gut has been shown to elicit a strong IFN γ response, which has been shown to downregulate Th2 cells.[10] Specifically, administering the neutrophil-activating protein of *H. pylori*, both systemically and mucosally, reduces characteristic Th2 cytokines, IgE, and eosinophilia by activating the Th1 pathway.[11] These studies show that this bacterial infection causes an overwhelming Th1 response, one that epidemiologically and physiologically have been shown to directly affect the response of the Th2 pathway. In recent years, the role of Th17 cells in *Helicobacter* infections has been shown to be critical in pathogenesis. Shi, et al. demonstrated that the Th17 cells are critical in acute infection, where it works with IL8 to recruit neutrophils and decrease bacterial burden.[12] Depleting these cells leads to proliferation of bacteria. Interestingly, if there is no control mechanism in place, such as regulatory T cells, IL17 is upregulated to the point where it is ineffective in pathogen clearance.[13] Data from our lab shows that in germfree mice, IL17 levels continue to climb after 8 weeks, while *H. felis* is never cleared. With conventionally reared C57BL/6 mice, IL17 increases initially, but decreases after 8 weeks.[14] Th1 and Th17 adaptive immunity plays a significant role in *Helicobacter* infections, though both must be tightly regulated.

Human studies demonstrate that polymorphisms in the TLR2 and 4 genes affect the pathogenesis of *Helicobacter* gastritis.[15,16] However, there have been conflicting *in vitro* reports about which of these TLRs are needed to recognize the bacteria. Many epidemiological studies have shown an inverse correlation with this pathogen and asthma susceptibility. Chen and Blaser demonstrated that asthma onset in children younger than 5 years old was inversely associated with seropositivity for *H. pylori* (OR, 0.58; 95% CI, 0.38-0.88). In the same study, seropositivity for *H. pylori* in children 3-19 years old was significantly inversely correlated with having a current case of asthma (OR, 0.41; 95% CI, 0.24-0.69).[17] Interestingly, with the rise of asthma, there has also been an increase in administering broad spectrum antibiotics in small children. This illustrates that, not only is there a dysregulation in global bacteria, but also that the bacteria, such as *H. pylori*, that were acquired during childhood that could potentially shape the immune system are being eliminated before their physiological effects can be properly achieved.[18] Additionally, *H. pylori* is associated with decreasing the severity of gastroesophageal reflux disorder, which is a positive correlate of asthma.[10] Immunologically, asthma is characterized as a Th2 disease, as asthmatic patients have increased serum IgE, production of which is known to be initiated by IL-4, 5, and/or 13.[10] Interestingly, IFN γ production has been shown to hinder Th2 cytokine production; IFN γ is a downstream target of NF κ B, the transcription factor induced by TLR2 and 4 activation. Thus, activating TLR2 and 4 by environmental exposure to bacterial ligands could elicit a more Th1- and Th17-skewed adaptive response and result in the downregulation of the Th2 phenotype in asthma patients.[19]

Because asthma is a complex disease, various mouse models are used to elucidate various facets of the disease. Since mice do not spontaneously get asthma, the mice must be sensitized with a specific allergen and subsequently challenged with that allergen. Many labs use BALB/c mice for their studies because these mice have a Th2-skewed immune system and a robust asthmatic response upon asthma induction. However, this model does not parallel typical human disease since most people do not have a strongly Th2 skewed

immune system. Therefore, the use of C57BL/6 mice is a good model because it has a Th1 bias, which is more like the human immune system. Using C57BL/6 mice is also beneficial when studying the bacterial colonization and/or sensing in asthma, as these mice respond more strongly to bacteria than the BALB/c mice.[20] Likewise, the allergens used in asthma studies vary greatly. Historically, ovalbumin has been used to induce the asthmatic phenotype because of its ability to produce strong Th2 responses.[21,22] However, this allergen does not have clinical application, as ovalbumin is not a common allergen in humans. Therefore, more clinically relevant allergens, such as house dust mite and cockroach extracts, are becoming more frequently used.[23,24] There are two categories of asthma models: acute and chronic. The acute models are used to study the onset and beginning stages of asthma, whereas the chronic models are used to study the long term effect of asthma, such as airway remodeling. The chronic model is also helpful in studying therapies for asthma patients, since most patients already have remodeling taking place in their lungs.[24]

In this model, we chose cockroach antigen (CRA) because the cockroach is a common household pest. However, this model has only previously been evaluated in adult mice, whereas we needed a model of childhood asthma induction if the effect of early gastrointestinal infections is to be adequately tested. Therefore, we utilized 2, 4, and 6 week old mice to determine the induction of acute asthma by CRA. We concluded that, at 2 weeks, the newly weaned pups did not have a mature enough immune response to mount a Th2 acute asthma response. At 6 weeks, their immune reactions had switched to a more Th1/17 phenotype and the results were more variable. However, 4 week old C57BL/6 mice had both a robust and consistent response to CRA and can now be established as an excellent model for acute sensitization and challenge with cockroach antigen (CRA) to study the development and early stages of asthma (Figures 1 and 2).

2. Materials

2.1. *Helicobacter* Components

1. C57BL/6 mice (Jackson Laboratory, Bar Harbor, ME)
2. *Helicobacter felis* ATCC 49179
3. ATCC Medium 260 Plates: trypticase soy agar, defibrinated calf blood (5% v/v) (Colorado serum Company, Denver, CO), Trimethoprim (1 mg/ml), Vancomycin (10 mg/ml), Fungizone (1% v/v). Resuspend 20 grams of Trypticase Soy Agar in 500 ml of ultrapure water (e.g. Milli-Q or equivalent) and heat with frequent agitation to boiling for one minute to completely dissolve the powder. This should be autoclaved for 15 minutes at 121°C and then cooled to 55°C in a water bath. When cooled, add 25 ml of the defibrinated calf blood using sterile technique and stir slowly to mix (*see* Note 1). Add 5 ml of Trimethoprim, 1.5 ml of Vancomycin, and 5 ml of Fungizone (all previously sterilized) to the mix. Immediately pour plates in a sterile hood to approximately ½ full and flame the tops to get rid of any

---

¹Because the defibrinated calf blood is frozen, it must be thawed in a 37°C water bath.

air bubbles. Allow the medium solidify, and then store at 4°C in a sealed container for no more than one month. This makes 20-25 plates.

4. Brain Heart Infusion (BHI) Broth: Add 37 g of BHI broth to 1 L of deionized distilled water. Autoclave for 30 minutes on the liquid cycle. Let cool to 55°C in a water bath and then add the following reagents: 3 µg/ml Vancomycin (3.0 ml of a 10 mg/ml stock); 10 µg/ml Trimethoprim (10.0 ml of a 1 mg/ml stock); 1% Fungizone (10 ml); and 5% defibrinated fetal calf serum (50 ml).
5. BBL™ CampyPak™ Plus Microaerophilic System Envelopes with Palladium Catalyst (BD, San Jose, CA)
6. Histology sponges and cassettes
7. Citrosolv (Fisher)
8. Pepsin (0.25% in PBS)
9. Rabbit anti-*H. felis*/*H. pylori* antibody (Covance, Emeryville, CA)
10. Cy3 donkey anti-rabbit antibody (1:200 dilution, Jackson ImmunoResearch)
11. FITC-labeled lectin N-acetyl-D-glucoaminy-specific *Griffonia simplicifolia* II (5 µg/mL, Invitrogen, Eugene, OR)
12. Hoechst 33258 (0.5 ug/ml; bis-Benzimide, Sigma, St. Louis, MO)

2.2. CRA Asthma Model Components

1. Cockroach antigen (Hollister-Stier, Spokane, WA) (*see* Note 2)
2. Incomplete Freund's Adjuvant
3. Phosphate buffered saline (PBS)
4. 2-glass syringes (5 ml each)
5. Three-way stopcock
6. 1 ml syringe with needle (32G)
7. Eppendorf tubes (1.5 ml, one per mouse)
8. Isoflurane
9. P20 & P200 pipette
10. Plastic disposable pipette dropper (one per mouse)
11. 10% buffered formalin
12. Bouin's fixative solution: This is a picric acid-formalin-acetic acid mixture that can either be made within the lab (300 ml saturated picric acid, 100 ml formaldehyde, 20 ml acetic acid) or purchased. This fixative allows better and crisper nuclear staining than 10% neutral-buffered formalin. As picric acid is extremely explosive

---

²CRA is not stable after dilution. Therefore, all dilutions must be made fresh and used immediately.

if allowed to dry out, it is usually safer to just purchase the Bouin's fixative solution.

13. RNAlater RNA Stabilization Reagent (Ambion, Austin, TX): This is an immediate RNA stabilization and protection reagent. It allows tissue archiving without the risk of RNA degradation.
14. Methacholine: Used in the methacholine challenge, in which the subject inhales aerosolized methacholine to determine the level of bronchial hyperreactivity.
15. Ketamine: Used for the induction and maintenance of general anesthesia
16. Flexivent (Scireq, Montreal, Canada): This is a computer-controlled precision pump that controls mechanical ventilation while also obtaining measurements of respiratory mechanics.

2.3. ELISA Components

1. Immulon 96 well plates (Thermo Fisher Scientific, Waltham, MA)
2. Goat Anti-Mouse IgE-UNLB (10 µg/ml; Southern Biotech, Birmingham, AL): For use in coating the ELISA plate.
3. Wash buffer (PBS, 0.5% Tween-20)
4. Blocking buffer (PBS, 5% bovine serum albumin (BSA))
5. Diluent buffer (PBS, 1% BSA)
6. Mouse IgE Standard (Southern Biotech)
7. Goat Anti-Mouse IgE-AP (Southern Biotech) (1:2000 in diluent buffer): For detection of IgE in serum.
8. 3N NaOH: 120g in 1000 ml of water.
9. SIGMAFAST™ p-Nitrophenyl phosphate Tablets (Sigma)
10. VERSAmax microplate reader (Molecular Devices, Sunnyvale, CA).

2.4. RNA and QPCR Components

1. Applied Biosystems Assays-On-Demand primer/probe sets.
2. TaqMan Universal PCR Mix (PE Applied Biosystems; Foster City, CA).
3. Trizol® (Life Technologies, Grand Island, NY)
4. Turbo DNase Kit (Ambion, Austin, TX).
5. Roche Transcriptor First Strand cDNA Synthesis Kit (Roche, Painsberg, Germany)
6. Stratagene MX3000P Real-Time Cycler (Agilent Technologies)

3. Methods

3-1. Growing *Helicobacter felis*

1. One vial of *H. felis* (ATCC 49179) is inoculated onto an ATCC Medium 260 plate (see Note 3).
2. Incubate at 37°C for 2 days in an anaerobic jar with a CampyPak (activate CampyPak per instructions with water). Make sure the container has an airtight seal (see Note 4).
3. Check *H. felis* viability by dropping one drop of the bacterial suspension onto a microscope slide and covering with a standard coverslip. Using a 20x or 40x objective lens on a light microscope, focus on the bacteria. Make sure the majority of them are motile by watching them swim through multiple viewing fields.
4. If motile, using the broth from the plate, inoculate BHI broth (100 ml) in a flask. Secure in an anaerobic jar with a CampyPak. (see Note 5).
5. Incubate at 37°C for 18-24 h, with gentle shaking.
6. Check viability, as outlined in 3-1.3.
7. Determine the optical density of the bacteria culture (OD₄₅₀; 1 OD₄₅₀ ≈ 10⁹ bacteria) (see Note 6).
8. Harvest bacteria by centrifuging at 3000 g for 10 minutes, and then resuspend the pellet in glycerol: BHI freezing media (31 ml glycerol: 69 ml BHI). Store aliquots of bacteria (2 × 10⁹ CFU/ml) at -80°C. The frozen stock should remain viable for ~4 months.

3-2. *H. felis* Innoculation

1. Day 0 and 3: Inoculate mice orally (per os (p.o.)) using a 200 µl pipette with 0.05 ml of *H. felis* (2 × 10⁹ CFU/ml in glycerol: BHI) using a frozen aliquot.
2. Day 7: Inoculate mice p.o. as described in 3-2.1 using freshly grown *H. felis*. Use the same culture technique as described above in 3-1, with the exception that a vial of frozen stock of *H. felis* can be used to start the culture instead of a new vial from ATCC (see Note 7).

3-3. Asthma Sensitization (Day 0)

1. Cockroach antigen (CRA) is resuspended in PBS to a final concentration of 4 mg/ml (see Note 2).

³The frozen aliquots of *H. felis* must be thawed at 50°C.

⁴When incubating the cultures, incubate with the solid agar on the bottom. Because the *H. felis* grows at the solid/liquid interface, turning the plate upside down will cause the liquid to spill and not be in contact with the solid media. This will result in no bacterial growth.

⁵Because *H. felis* does not grow in colonies, remove the broth culture from the plate using a 5 ml pipette. Using a clean pipette, draw up 2 ml of BHI and put it on the used plate. Swirl the plate on a flat surface and then tilt it at a 45° angle to remove the liquid. Place this liquid onto the freshly inoculated plate. Repeat with a clean pipette and 2 ml of additional clean BHI.

⁶If frequently taking aliquots for OD readings, replace the CampyPak microaerophilic packets every time you open the container.

⁷In our lab, this infection scheme results in a 100% infection rate.

2. It is then put into one glass syringe. A second glass syringe is filled with an equal amount (v/v) of Incomplete Freund's Adjuvant (IFA).
3. Connect these 2 glass syringes to a 3-way stopcock, and then emulsify the IFA and CRA by syringe-extrusion (alternatively pushing the solution in each syringe through progressively smaller pore sizes in the stopcock) for 10-15 minutes (*see* Note 8).
4. The solution is finished emulsifying when a drop of the solution does not disperse on the top of ice-cold water. Use the solution immediately (*see* Note 9).
5. Using a non-tuberculin syringe and a 32G needle, inject the mice intraperitoneally (50 μ l) and subcutaneously (50 μ l).

3-4. Asthma Induction via Intranasal Challenge (Days 14, 18, 22, 26)(*see* Note 10)

1. Mix CRA with PBS (0.075 mg/ml) (*see* Note 2).
2. Anaesthetize mice individually using isoflurane (*see* Note 11).
3. Using a P20 pipette, drop PBS: CRA solution (10 μ l, 0.75 μ g) onto nostril. Once the mouse inhales the solution, drop another 10 μ l onto the nostril (*see* Note 12).

3-5. Asthma Induction via Intratracheal Challenge (Day 28)

1. Mix CRA with PBS (0.12 μ g/ml) (*see* Note 2).
2. Anaesthetize mice individually using isoflurane (*see* Note 13).
3. When the mouse begins to awaken and gasp using its diaphragm, instill PBS: CRA (50 μ l, 6.0 μ g) solution into the mouse's throat between gasps with a P200 pipette (*see* Note 14).

⁸During the emulsification process, the pore size in the 3-way stopcock must be made progressively smaller by closing the pore incrementally. This solution will be progressively more difficult to mix. This means that the emulsification is occurring.

⁹If the solution is not completely emulsified, the antigen will disperse immediately after injection and will not induce the appropriate immune response.

¹⁰The intranasal challenge causes proliferation of the CRA-specific T cell and elicits their migration into the airways, causing an asthmatic phenotype. Performing this procedure several days apart gradually builds up the asthmatic response in the mouse, similar to the development of asthma in children. The intratracheal challenge is designed to elicit a maximum number of T cells into the airways without causing the animal respiratory distress. The intratracheal challenge is much harder for the mouse to endure. Therefore, this route of administration is only conducted at the end of the procedure.

¹¹When performing the intranasal challenge, the mouse should only be exposed to the isoflurane for about 15 sec, which should induce a low level of anesthesia.

¹²The procedure must be done very quickly to get the "sniff" response to the drop of CRA, which ensures that it goes into their lungs and is not swallowed. During this procedure, the mouse will begin recovering from unconsciousness and the muscles will begin to tighten. Brace the lower jaw of the mouse with your thumb and the top of the mouse's head with your forefinger, wrapping your other fingers around its torso. In the event that the mouse awakens from the anesthesia before the procedure is complete, you will have a firm grip on the mouse.

¹³The mouse should be exposed to the isoflurane for 30 seconds until completely limp with barely detectable breath movements in its chest. Wait to administer the challenge until the mouse begins to awaken and its diaphragm spasms.

¹⁴The mouse should be held as previously mentioned upon being removed from the isoflurane through the entire procedure until the mouse is fully awake. Often, if the mouse is not held it will die, possibly due to drop in body temperature. The pipette should be poised at the back of the throat, depressing the mouse's tongue. Ensure that the pipette is ready to deposit the antigen when the mouse awakens. The procedure goes very quickly and the mouse can awaken very rapidly, so care must be taken continuously restrain the mouse while anesthetized and release the mouse if it awakens.

3-6. Sacrifice (Day 29)

1. Anaesthetize mice using isoflurane.
2. Perform cervical dislocation.

3-7. Tissue Collection

1. Mice are euthanized using 5 ml of isoflurane for approximately 30 sec and followed by cervical dislocation.
2. Blood is collected via heart puncture, allowed to clot for 20 min at room temperature (r.t.), centrifuged for 10 minutes and 1300 rpm, and the serum removed and stored at -20°C until analysis.
3. The stomach is removed and quartered. Each quarter is either fixed in Bouin's Fixative for histology, frozen at -80°C (2 quarters) for protein analysis, or stored in RNAlater for RNA analysis, as per protocol.
4. Place 1 stomach quarter flat between 2 thin histology sponges in a histology cassette. Store the cassette in Bouin's fixative for 24 hours at 4°C. Bouin's fixative is replaced with 70% ethanol every 24 hours for 48 hours (*see* Note 15). The tissue can be embedded in paraffin along its long, cut edge in order to get a cross section of the epithelium. One 5 µm slide is stained with hemotoxylin and eosin and scored for pathology. Other slides can be stained with various antibodies to determine helicobacter infection rates, presence of inflammatory cells, differentiated epithelial cells, proliferation, and/or apoptosis, as needed.
5. Freeze stomach quarters 2 and 3 at -80°C for protein extraction, if needed.
6. Store stomach quarter 4 in RNAlater and process as directed by the manufacture's protocol for future quantitative real-time RT-PCR (qRT-PCR).
7. The lungs are removed above the main stem branch directly below the larynx. The lungs are then separated directly below the main stem branch. Using a 25G needle and a 3 ml syringe, the left lung is perfused with 1-2 ml of formalin through the main bronchus until it is expanded but not over extended. Pressure must be maintained on the fluid in the lungs by gripping the forceps around the needle inserted into the bronchus, while gently applying pressure to the syringe. This is held for 1-2 min. Then, the lung is removed from the needle and placed into a histology cassette and stored in formalin.
8. The tissue in the cassettes is stored in formalin for 24 hours. The formalin is then replaced with 70% ethanol once every 24 hours for 2 days. The tissue can be embedded in paraffin and stained with hemotoxylin and eosin. Lung sections are then scored for pathology.
9. The right lung is stored in RNAlater as directed by the manufacture's protocol for future qRT-PCR.

¹⁵Tissue should not be fixed in Bouin's fixative for more than 24 hours or pigments can begin to form. Excess fixative should be washed out of the tissue using the alcohol/water washes.

3-8. Airway Hyperresponsiveness

1. Mice are anesthetized with 450 mg/kg of ketamine, and a tracheotomy tube (18G) is inserted and connected to the inspiratory and expiratory ports of a ventilator (Felixvent). Mice are ventilated at a rate of 160 breaths per minute at a tidal volume of 0.2 ml with a positive end-expiratory pressure of 2-4 cm water.
2. Increasing concentrations of methacholine (0, 10, 20, 30, 40, and 50 mg/ml) are administered via aerosolization. From 20 seconds to 3 minutes after each aerosol challenge, detailed measurements should be recorded continuously. The measurements should include resistance (R), compliance (C), and elastance (E) (*see* Note 16).

3-9. IgE ELISAs

1. Immulon 96 well plates are coated with Goat Anti-Mouse IgE-UNLB (10 µg/ml) in PBS, overnight at 4°C.
2. The next day, the plate is washed with wash buffer five times. The non-specific binding sites are blocked for >1 hour using blocking buffer.
3. After washing the plates five times with wash buffer, the serum samples are added. The serum from mice who received CRA is diluted 1:2 with diluent buffer, and the mock-treated mice serum samples are used neat. Mouse IgE-UNL Bis diluted to a start concentration of 2000 pg/ml, and then diluted 1:2 seven more times for the complete standard curve. The standards and samples are incubated overnight at 4°C.
4. The next day, the plate is washed 5 times with wash buffer. Goat Anti-Mouse IgE-AP (1:2000 in diluent buffer) is added and incubated at r.t. for 2 hours.
5. After washing the plate 5 times with wash buffer, SIGMAFAST™ p-Nitrophenyl phosphate Tablets are dissolved in 20 ml of deionized water and added to the plate. Plates are incubated for 30 minutes at r.t. in the dark.
6. 3N NaOH is added to stop the reaction, and the plate is read at 405 nm on a microplate reader.

3-10. RNA and RT-PCR

1. RNA Isolation: The Trizol® (phenol and guanidine isothiocyanate) method can be used to isolate the total RNA from one quarter of each stomach and the left lung. [25]
2. Before making cDNA, the RNA is processed to remove contaminating DNA using the Turbo DNase Kit. cDNA is made using the Roche Transcriptor First Strand cDNA Synthesis Kit with mRNA (2 µg) from each sample. Quantitative real-time reverse transcription polymerase chain reaction (QPCR) is performed on each

¹⁶During this procedure, several measurements (including compliance and resistance) are taken; however, we only report the resistance measurement because this particular asthma induction protocol is not designed to dramatically affect other parameters, such as compliance. The acute nature of this model that does not result in significant, long-term airway remodeling.

sample. Primers/probes used are from Applied Biosystems Assays-On-Demand. These are used with TaqMan Universal PCR Mix. All RNA data is analyzed using the 2^{-Ct} relative quantitation method, described in the Applied Biosystems manufacturer's protocol (*see* Note 17).[25-27]

3-11. Pathology Scoring

1. Stomach: One quarter stomach from each mouse that was stained using hemotoxylin and eosin should be scored on a scale of 0-3 in each of 3 categories. The scores from the 3 categories are then added together for a total score, with 0 being the lowest and 9 being the highest possible scores.[28]

Score	Longitudinal Extent of Inflammation	Vertical Extent of Inflammation	Histological Changes
0	none	none	none
1	patchy	basal lamina propria only	mild loss of differentiated epithelial cells
2	<50%	transmural	moderate loss of differentiated epithelial cells
3	>50%	both mucosa and submucosa involved	severe loss of differentiated epithelial cells

2. Lung: Each section should be scored using a method derived from Curtis, et. al. in which the inflammation around the vasculature and the bronchial is evaluated and added together for a total inflammation score, with 0 being the lowest and 6 being the maximum.[29]

Score	Vascular	Bronchial
0	no inflammation	no inflammation
1	occasional cuffing	occasional cuffing
2	most vessels surrounded by a thin layer (1-5) of inflammatory cells	most vessels surrounded by a thin layer (1-5) of inflammatory cells
3	most vessels surrounded by a thick layer (>5) of inflammatory cells	most vessels surrounded by a thick layer (>5) of inflammatory cells

3-12. *H. felis* Staining and Quantification (*see* Note 18).[30]

1. Deparaffinization an unstained tissue section as follows: wash the slides two times for 10 mins per wash with Citrosolv; next, wash the slides three times for 10 mins with isopropyl alcohol; and finally, rinse the slides for 5 mins with running deionized water.
2. After rehydrating the tissue in phosphate-buffered saline (PBS), pepsin (0.25% in PBS) is incubated on the slides for 10 mins at r.t..

¹⁷The housekeeping gene for comparison used in these experiments was the 18S gene because this gene has been determined to be relatively stable, even under inflammatory conditions.[26,31,32] This method uses the difference of the average crossing threshold (Ct) of the 18S gene from the average Ct of the target gene to determine the relative expression of the target gene within each group of animals (Ct). Next, the Ct is calculated determining the difference of the experimental Ct (*H. felis*-infected mice) from the control Ct (mock-infected mice). Finally, the average fold change of the gene is calculated with the following formula: 2^{-Ct} . Using the standard deviation of the Ct of the experimental group in the average fold change formula, the upper and lower limits are calculated.

¹⁸A quarter of the stomach with the squamo-columnar junction and antrum from each mouse is deparaffinized, stained, and quantitated.

3. After rinsing the slides in PBS, to block non-specific binding sites and to permeabilize the tissue, add PBS blocking buffer (1% bovine serum albumin, 0.3% Triton X-100) to each slide and incubated for 1 hour at r.t.
4. The slides are then washed in PBS and the tissue stained with undiluted rabbit anti-*H. pylori* antibody to semi-quantitate *H. felis* colonization. This antibody is known to cross-react with *H. felis*.
5. After washing the slides in PBS, Cy3 donkey anti-rabbit antibody (1:200 dilution) and FITC-labeled lectin N-acetyl-D-glucoaminy-specific *Griffonia simplicifolia* II (5 µg/mL) are added to the tissue and incubated for 1 hour at r.t. for detection of *H. felis* and mucous neck cells, respectively.
6. To counterstain the nuclei, the slides are incubated for 20 minutes at r.t. with Hoechst 33258 (0.5 µg/ml).
7. Colonization of the antrum with *H. felis* is evaluated on a scale of 0 to 4, where 0 = no bacteria per gland; 1 = 1 – 2 bacteria per gland; 2 = 3 – 10 bacteria per gland; 3 = 11 – 20 bacteria per gland; and 4 = >20 bacteria per gland.(33)

Acknowledgments

The authors would like to thank Kim Estell for assistance with airway hyperresponsiveness analysis and Ben Christmann for helping with the lung inflammation technique. We would also like to thank J. McNaught for slide preparation and M. Harris for animal husbandry, and members of the Lorenz lab for valuable advice. This study was supported in part by NIH grants R01 DK059911; P01 DK071176; the American Asthma Foundation grant 06-0167; and University of Alabama at Birmingham Digestive Diseases Research Development Center grant P30 DK064400. CGD is supported by the Howard Hughes Medical Institute Med into Grad Fellowship. Aspects of this project were conducted in biomedical research space that was constructed with funds supported in part by NIH grant C06RR020136.

References

1. Strachan DP. Family size, infection and atopy: the first decade of the “hygiene hypothesis”. *Thorax*. 2000; 55(Suppl 1):S2–10. [PubMed: 10943631]
2. Yoo J, Tcheurekdjian H, Lynch SV, Cabana M, Boushey HA. Microbial manipulation of immune function for asthma prevention: inferences from clinical trials. *Proc Am Thorac Soc*. 2007; 4(3): 277–282. [PubMed: 17607013]
3. Strachan DP, Seagroatt V, Cook DG. Chest illness in infancy and chronic respiratory disease in later life: an analysis by month of birth. *International journal of epidemiology*. 1994; 23(5):1060–1068. [PubMed: 7860157]
4. Ownby DR, Johnson CC, Peterson EL. Exposure to dogs and cats in the first year of life and risk of allergic sensitization at 6 to 7 years of age. *JAMA : the journal of the American Medical Association*. 2002; 288(8):963–972. [PubMed: 12190366]
5. Braun-Fahrlander C, Riedler J, Herz U, Eder W, Waser M, Grize L, Maisch S, Carr D, Gerlach F, Bufe A, Lauener RP, Schierl R, Renz H, Nowak D, von Mutius E. Environmental exposure to endotoxin and its relation to asthma in school-age children. *The New England journal of medicine*. 2002; 347(12):869–877. [PubMed: 12239255]
6. Riedler J, Braun-Fahrlander C, Eder W, Schreuer M, Waser M, Maisch S, Carr D, Schierl R, Nowak D, von Mutius E. Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet*. 2001; 358(9288):1129–1133. [PubMed: 11597666]
7. Kusters JG, van Vliet AH, Kuipers EJ. Pathogenesis of *Helicobacter pylori* infection. *Clin Microbiol Rev*. 2006; 19(3):449–490. [PubMed: 16847081]

8. Harris PR, Wright SW, Serrano C, Riera F, Duarte I, Torres J, Pena A, Rollan A, Viviani P, Guiraldes E, Schmitz JM, Lorenz RG, Novak L, Smythies LE, Smith PD. Helicobacter pylori gastritis in children is associated with a regulatory T-cell response. *Gastroenterology*. 2008; 134(2): 491–499. [PubMed: 18242215]
9. Dimmitt RA, Staley EM, Chuang G, Tanner SM, Soltau TD, Lorenz RG. Role of postnatal acquisition of the intestinal microbiome in the early development of immune function. *J Pediatr Gastroenterol Nutr*. 2010; 51(3):262–273. [PubMed: 20639773]
10. Blaser MJ, Chen Y, Reibman J. Does Helicobacter pylori protect against asthma and allergy? *Gut*. 2008; 57(5):561–567. [PubMed: 18194986]
11. Amedei A, Cappon A, Codolo G, Cabrelle A, Polenghi A, Benagiano M, Tasca E, Azzurri A, D'Elia MM, Del Prete G, de Bernard M. The neutrophil-activating protein of Helicobacter pylori promotes Th1 immune responses. *The Journal of clinical investigation*. 2006; 116(4):1092–1101. [PubMed: 16543949]
12. Shi Y, Liu XF, Zhuang Y, Zhang JY, Liu T, Yin Z, Wu C, Mao XH, Jia KR, Wang FJ, Guo H, Flavell RA, Zhao Z, Liu KY, Xiao B, Guo Y, Zhang WJ, Zhou WY, Guo G, Zou QM. Helicobacter pylori-induced Th17 responses modulate Th1 cell responses, benefit bacterial growth, and contribute to pathology in mice. *J Immunol*. 2010; 184(9):5121–5129. [PubMed: 20351183]
13. Kao JY, Zhang M, Miller MJ, Mills JC, Wang B, Liu M, Eaton KA, Zou W, Berndt BE, Cole TS, Takeuchi T, Owyang SY, Luther J. Helicobacter pylori immune escape is mediated by dendritic cell-induced Treg skewing and Th17 suppression in mice. *Gastroenterology*. 2010; 138(3):1046–1054. [PubMed: 19931266]
14. Schmitz JM, Durham CG, Ho SB, Lorenz RG. Gastric mucus alterations associated with murine Helicobacter infection. *J Histochem Cytochem*. 2009; 57(5):457–467. [PubMed: 19153195]
15. Tahara T, Arisawa T, Wang F, Shibata T, Nakamura M, Sakata M, Hirata I, Nakano H. Toll-like receptor 2 (TLR) -196 to 174del polymorphism in gastro-duodenal diseases in Japanese population. *Digestive diseases and sciences*. 2008; 53(4):919–924. [PubMed: 17934843]
16. Trejo-de la OA, Torres J, Perez-Rodriguez M, Camorlinga-Ponce M, Luna LF, Abdo-Francis JM, Lazcano E, Maldonado-Bernal C. TLR4 single-nucleotide polymorphisms alter mucosal cytokine and chemokine patterns in Mexican patients with Helicobacter pylori-associated gastroduodenal diseases. *Clinical immunology*. 2008; 129(2):333–340. [PubMed: 18755634]
17. Chen Y, Blaser MJ. Helicobacter pylori colonization is inversely associated with childhood asthma. *The Journal of infectious diseases*. 2008; 198(4):553–560. [PubMed: 18598192]
18. Johnson CC, Ownby DR, Alford SH, Havstad SL, Williams LK, Zoratti EM, Peterson EL, Joseph CL. Antibiotic exposure in early infancy and risk for childhood atopy. *J Allergy Clin Immunol*. 2005; 115(6):1218–1224. [PubMed: 15940137]
19. Schaub B, Lauener R, von Mutius E. The many faces of the hygiene hypothesis. *J Allergy Clin Immunol*. 2006; 117(5):969–977. quiz 978. [PubMed: 16675321]
20. Gueders MM, Paulissen G, Crahay C, Quesada-Calvo F, Hacha J, Van Hove C, Tournoy K, Louis R, Foidart JM, Noel A, Cataldo DD. Mouse models of asthma: a comparison between C57BL/6 and BALB/c strains regarding bronchial responsiveness, inflammation, and cytokine production. *Inflamm Res*. 2009; 58(12):845–854. [PubMed: 19506803]
21. Pastva A, Estell K, Schoeb TR, Atkinson TP, Schwiebert LM. Aerobic exercise attenuates airway inflammatory responses in a mouse model of atopic asthma. *Journal of immunology*. 2004; 172(7): 4520–4526.
22. Hewitt M, Estell K, Davis IC, Schwiebert LM. Repeated bouts of moderate-intensity aerobic exercise reduce airway reactivity in a murine asthma model. *Am J Respir Cell Mol Biol*. 2010; 42(2):243–249. [PubMed: 19423772]
23. Epstein MM. Do mouse models of allergic asthma mimic clinical disease? *Int Arch Allergy Immunol*. 2004; 133(1):84–100. [PubMed: 14726635]
24. Nials AT, Uddin S. Mouse models of allergic asthma: acute and chronic allergen challenge. *Dis Model Mech*. 2008; 1(4-5):213–220. [PubMed: 19093027]
25. Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem*. 1987; 162(1):156–159. [PubMed: 2440339]

26. Bas A, Forsberg G, Hammarstrom S, Hammarstrom ML. Utility of the housekeeping genes 18S rRNA, beta-actin and glyceraldehyde-3-phosphate-dehydrogenase for normalization in real-time quantitative reverse transcriptase-polymerase chain reaction analysis of gene expression in human T lymphocytes. *Scand J Immunol.* 2004; 59(6):566–573. [PubMed: 15182252]
27. Biosystems A. A 4371095 Rev A.
28. Roth K, Kapadia S, Martin S, Lorenz R. Cellular immune responses are essential for the development of *Helicobacter felis*-associated gastric pathology. *J Immunol.* 1999; 163(3):1490–1497. [PubMed: 10415051]
29. Curtis JL, Warnock ML, Arraj SM, Kaltreider HB. Histologic analysis of an immune response in the lung parenchyma of mice. Angiopathy accompanies inflammatory cell influx. *The American journal of pathology.* 1990; 137(3):689–699. [PubMed: 2399937]
30. Brown JK, Pemberton AD, Wright SH, Miller HR. Primary antibody-Fab fragment complexes: a flexible alternative to traditional direct and indirect immunolabeling techniques. *J Histochem Cytochem.* 2004; 52(9):1219–1230. [PubMed: 15314089]
31. Ropenga A, Chapel A, Vandamme M, Griffiths NM. Use of reference gene expression in rat distal colon after radiation exposure: a caveat. *Radiat Res.* 2004; 161(5):597–602. [PubMed: 15161363]
32. Rubie C, Kempf K, Hans J, Su T, Tilton B, Georg T, Brittner B, Ludwig B, Schilling M. Housekeeping gene variability in normal and cancerous colorectal, pancreatic, esophageal, gastric and hepatic tissues. *Mol Cell Probes.* 2005; 19(2):101–109. [PubMed: 15680211]

Figure 1.

Four week old mice have more robust pathology and airway resistance than 2 or 6 week old mice. **(A)** Four week old mice developed more inflammation within their airways and vasculature. Although 6 week old CRA-treated mice have moderate perivascular and peribronchial inflammation, the younger 4 week old mice have much greater perivascular and peribronchial inflammation after asthma induction. The 2 week old mice have only mild perivascular and peribronchial inflammation after CRA treatment. **(B-D)** Representative H&E images of the 2 week old mice, 4 week old mice, and 6 week old mice, respectively. **(E)** Resistance (R) in the airways was consistently higher in the 4 week old mice when compared with the mock-treated, as well as **(F)** the treated 2 week old mice and **(G)** the treated 6 week old mice.

Figure 2.

Four week old mice developed higher serum asthma markers. **(A)** Serum IgE was consistently greater in 4 week old mice than the other two groups. **(B-D)** The disease in the 4 week old mice **(C)** appeared to be IL13 mediated, though all of the Th2 cytokines were elevated in these mice, as opposed to **(B)** the 2 week old mice and **(D)** the 6 week old mice.