Published in final edited form as:

Schizophr Res. 2014 August ; 157(0): 238–243. doi:10.1016/j.schres.2014.05.005.

A fundamental role for hippocampal parvalbumin in the dopamine hyperfunction associated with schizophrenia

Angela M. Boley, Stephanie. M. Perez, and Daniel J. Lodge*

Department of Pharmacology & Center for Biomedical Neuroscience, University of Texas Health Science Center at San Antonio, San Antonio, TX, USA

Abstract

Postmortem studies in schizophrenia patients have demonstrated robust alterations in GABAergic markers throughout the neuraxis. It has been suggested that these alterations are restricted to subpopulations of interneurons, such as those containing the calcium binding protein parvalbumin. Indeed, a reduction in parvalbumin expression is a consistent observation in human postmortem studies, as well as, in a wide and diverse variety of animal models. However, it still remains to be determined whether this decrease in parvalbumin expression contributes to, or is a consequence of the disease. Here we utilize lentiviral delivered shRNA and demonstrate that a selective reduction in parvalbumin mRNA expression induces hyperactivity within the ventral hippocampus. In addition, we observe downstream increases in dopamine neuron population activity without changes in average firing rate or percent burst firing. These changes in dopamine neuron activity were associated with an enhanced locomotor response to amphetamine administration. These data therefore demonstrate that a reduction in ventral hippocampal parvalbumin expression is sufficient, in and of itself, to induce an augmented dopamine system function and behavioral hyper-responsivity to amphetamine, implicating a potential key role for parvalbumin in the pathophysiology of schizophrenia.

Keywords

Parvalbumin; Hippocampus; Dopamine; Schizophrenia

© 2014 Elsevier B.V. All rights reserved.

*Correspondence should be addressed to: Dr. Daniel Lodge, Department of Pharmacology & Center for Biomedical Neuroscience, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Dve, MC 7764, San Antonio, TX, USA. Ph: 210-567-4188. LodgeD@uthscsa.edu.

Contributors

Dr. Lodge designed the study and assisted with manuscript preparation. Drs. Boley and Perez, assisted with study design and performed all of the experiments, analyzed data and participated in writing the manuscript. All authors contributed to and have approved the final manuscript.

Conflict of Interest

Dr. Lodge reports receiving consulting fees from Dey Pharmaceuticals, while Boley and Perez have no disclosures nor conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Schizophrenia is a neuropsychiatric disease affecting up to 1% of the population (Bhugra, 2005; Saha et al., 2005). It is known to be multifactorial with both genetic and environmental factors contributing to the disease (Agid et al., 1999; Purcell et al., 2009; Sawa and Snyder, 2002). This heterogeneity has led to a number of distinct hypotheses of schizophrenia, however as yet, the specific neuropathology underlying this disease has not been conclusively determined. Nonetheless, clinical data from post mortem studies have provided some consistent observations, including an altered expression of GABAergic markers throughout both cortical and hippocampal regions (Konradi et al., 2011; Lewis et al., 2005). Specifically, decreases in glutamic acid decarboxylase (GAD)-1 mRNA and GAD-67 protein are observed throughout the cortex of schizophrenia patients (Hashimoto et al., 2003). It should be noted that these alterations in GABAergic markers are not global; rather, they are more prevalent among distinct subclasses of interneurons, including those that express the calcium binding protein parvalbumin (PV) (Lewis et al., 2005). Indeed, a decreased expression of PV is a consistent observation not only in postmortem human schizophrenia patients (Konradi et al., 2011; Lewis et al., 2005), but also in a diverse number of rodent models of the disease (Abdul-Monim et al., 2007; Behrens et al., 2007; Cunningham et al., 2006; Francois et al., 2009; Harte et al., 2007; Lodge et al., 2009). However, whether this decrease in PV expression is the cause, or a consequence, of the illness is not currently known.

Decreases in PV expression are observed in both the prefrontal cortex (Lewis et al., 2012) and hippocampus (Konradi et al., 2011), two brain regions that are consistently implicated in schizophrenia. Deficits in prefrontal cortical function likely contribute to cognitive impairments and working memory deficits (Goldman-Rakic, 1995; Weinberger et al., 1986), whereas aberrant hippocampal function is associated with positive symptom severity (Schobel et al., 2009). We have recently demonstrated, in the methylazoxymethanol acetate (MAM) rodent model of schizophrenia (for review see - (Lodge and Grace, 2009)), that the dopamine hyperfunction and associated behavioral hyper-responsivity to psychomotor stimulants are driven by aberrant activity within the ventral hippocampus (vHipp) (Lodge and Grace, 2007; Perez and Lodge, 2013; Perez et al., 2013). Given that PV interneurons are perisomatic targeting (Benes and Berretta, 2001; Gonzalez-Burgos and Lewis, 2012; Kawaguchi and Hama, 1987), we posit that a loss of PV interneuron function would result in an increased pyramidal cell firing (in the vHipp) that is sufficient to induce a schizophrenia-like phenotype. Here we utilize lentiviral delivered short hairpin RNA (shRNA) to examine the consequence of vHipp PV knockdown, as it pertains to the dopamine dysfunction in schizophrenia.

Materials and Methods

All experiments were performed in accordance with the guidelines outlined in the USPHS Guide for the Care and Use of Laboratory Animals and were approved by the Institutional Animal Care and Use Committee of the University of Texas Health Science Center.

Lentivirus Administration

All survival surgical procedures were performed under general anesthesia in a semi-sterile environment. Adult male Sprague Dawley rats obtained from Harlan Laboratories were anesthetized with sodium pentobarbital (60mg/kg, i.p.) and placed in a stereotaxic apparatus. Anesthesia was maintained by supplemental administration of sodium pentobarbital as required to maintain suppression of limb compression withdrawal reflex. A core body temperature of 37°C was sustained by a thermostatically controlled heating pad. The skin was reflected and bore holes drilled bilaterally overlying the vHipp (A/P +5.3, M/L +5.2, D/V -7.0 mm from bregma). Rats were administered (0.75µl/side) commercially available (Thermo Scientific) high-titer lentivirus particles containing GIPZ vectors expressing shRNA targeting either parvalbumin (mature antisense: TAGCAGACAAGTCTCTGGC) or a non-silencing control. Rats were sutured and housed under ABSL 2 conditions for 72 hours before being transferring to standard housing conditions. Rats were housed for a period of 6 weeks prior to behavioral and electrophysiological examinations to ensure stable transgene expression.

Amphetamine-Induced Locomotion

Rats were acclimatized to the behavioral facility for at least 1 hour prior to being placed in an open field arena (Med Associates) where spontaneous locomotor activity in the X-Y plane was determined for 45 mins by beam breaks and recorded with Open Field Activity Software (Med Associates). Following the baseline period, all rats were injected with D-amphetamine sulfate (0.5mg/kg, i.p.) and locomotor activity recorded for 45 minutes. A subsequent dose (2.0mg/kg, i.p.) was then administered and an additional 45 minutes of locomotor activity was recorded.

Extracellular Recordings

Rats were anesthetized with chloral hydrate (400 mg/kg, i.p.), as this anesthetic does not significantly depress dopamine neuron activity (Hyland et al., 2002), and placed in a stereotaxic apparatus. Anesthesia was maintained by supplemental administration of chloral hydrate as required to maintain suppression of limb compression withdrawal reflex. A core body temperature of 37°C was sustained by a thermostatically controlled heating pad. Extracellular glass microelectrodes (impedance 6–14MΩ) were lowered into the right ventral hippocampus (A/P +5.3, M/L +5.0, D/V -5.0 to -8mm from bregma) using a hydraulic micropositioner (KOPF - Model 640). Putative pyramidal neurons were defined as those with firing frequencies less than 2Hz as reported previously (Shah and Lodge, 2013; van der Meer and Redish, 2011). Electrodes were also inserted into the right ventral tegmental area (VTA; A/P -5.3, M/L +0.6 mm from bregma and -6.5 to -9.0 mm ventral of brain surface) and the activity of the population of dopamine neurons was determined by counting the number of spontaneously active dopamine neurons encountered while making multiple vertical passes (typically 6), separated by 200µm, in a predetermined pattern to sample equivalent regions of the VTA. Spontaneously active dopamine neurons were identified with open filter settings (low pass: 30Hz, high pass: 30kHz) using previously established electrophysiological criteria (Grace and Bunney, 1983) and once isolated, activity was recorded for 2–3 mins.

Quantitative PCR

Rats were decapitated following electrophysiological recordings and the brain was separated into two hemispheres. The vHipp was dissected from one hemisphere and homogenized. RNA was precipitated, separated by filtration and the concentration determined by absorbance at 260nm. Single stranded RNA was converted to cDNA using a High Capacity cDNA Reverse Transcription Kit (Ambion). Real time PCR was performed with FAM-labeled TaqMan primers targeting either parvalbumin (Rn00574541_m1), GAD1 (Rn00690300_m1) or GAPDH (Rn01775763_g1). Detection of FAM labeled DNA was performed by a CFX384 Real-Time PCR Detection System (Bio-Rad). C_t was calculated as the number of PCR cycles required for mRNA detection (compared to the control mRNA, GAPDH), while fold-changes were expressed using the 2^{-C_t} method.

Parvalbumin Immunohistochemistry

The hemisphere not used for PCR was post-fixed for at least 24 hours, and cryoprotected (25% w/v sucrose in PBS) until saturated. Hemispheres were coronally sectioned (50 μ m) using a cryostat (Leica). Ventral hippocampal slices were used to detect the expression of PV. Sections were washed three times (10 minutes) in PBS then blocked (2% normal goat serum & 0.3% Triton Tx100) for 30 minutes at room temperature. Primary antibodies [anti-PV 1:1000 (Abcam; ab11427) were applied (in PBS containing 1% normal goat serum and 0.3% Triton Tx100) overnight at 4°C followed by incubation with AlexaFluor® 594 goat anti-rabbit IgG (H+L) for 1 hour at room temperature. Slices were then mounted and cover slipped with ProLong gold anti-fade reagent. Computer assisted estimates of PV positive interneurons were performed using Neurolucida in combination with a Zeiss epifluorescent microscope.

Analysis

Locomotor data were analyzed by three separate 2-way ANOVAs (treatment and time as factors), one for each of the relevant time periods (spontaneous, 0.5 mg/kg, 2.0 mg/kg), followed by a Holm-Sidak post-hoc test, where appropriate. Electrophysiological analysis of single unit neuron activity was performed using commercial computer software (LabChart Pro – ADInstruments), and compared with student's t-test unless data failed test for normality and/or equal variance were a Mann Whitney Rank Sum test was utilized. PCR data was analyzed using commercial computer software (CFX Manager – BioRad). Cell counts were performed using Neurolucida and significance determined by a student's t-test. All data are represented as the mean $\pm$ standard error of the mean (SEM) unless otherwise stated. All statistical analyses were calculated using SigmaPlot (Systat Software Inc).

Materials

High titer lentiviral particles were obtained from Thermo Scientific. Pentobarbital sodium, chloral Hydrate, and D-Amphetamine sulfate were all purchased from Sigma. FAM-labeled TaqMan probes, the secondary AlexaFluor® 594 goat anti-rabbit IgG (H+L) antibody as well as the ProLong gold antifade mountant were obtained from Life Technologies. The anti-parvalbumin antibody was purchased from ABCAM. All other chemicals and reagents were of either analytical or laboratory grade and purchased from various suppliers.

Results

Confirmation of PV knockdown

To recapitulate the postmortem finding of reduced hippocampal PV expression, we performed region specific knockdown of PV by microinjection of commercially available high-titer lentiviral particles expressing microRNA-adapted small hairpin ribonucleic acid (shRNA) targeting PV mRNA in the vHipp. Lentiviral particles expressing a non-silencing shRNA were used as a control. The knockdown of PV expression was determined postmortem by quantitative polymerase chain reaction (qPCR) and immunohistochemistry (Fig 1). Specifically, qPCR demonstrated a decrease in mRNA expression (fold-change; control shRNA: 1.0 ± 0.7 c.f. PV shRNA: 0.2 ± 0.3) without overt changes in GAD mRNA expression (fold-change; control shRNA: 1.0 ± 0.6 c.f. PV shRNA: 1.6 ± 1.1); however, statistical analysis was precluded as PV mRNA expression was below the limit of detection in the majority (75%) of PV shRNA samples analyzed. To examine whether this decrease in PV expression resulted in a decrease in PV immunoreactivity, we performed quantitative cell counts throughout the vHipp (Fig 1). Rats administered PV shRNA displayed a significant reduction in PV positive interneurons throughout the vHipp (control shRNA: 9.11 ± 0.39 c.f. PV shRNA: 6.74 ± 0.48 cells/mm²; t-test, $t = 3.865$, $p < 0.05$, $n = 6$ /group; Fig 1).

Hippocampal Activity

PV neurons are fast-firing and perisomatic targeting which means that they are well situated to regulate pyramidal neuron activity (Benes and Berretta, 2001; Gonzalez-Burgos and Lewis, 2012; Kawaguchi and Hama, 1987). Thus, a decrease in PV interneuron function is likely sufficient to augment pyramidal activity within the vHipp. Indeed, we now demonstrate that a decrease in vHipp PV is sufficient to increase pyramidal cell output as determined by in vivo extracellular electrophysiology in chloral hydrate anesthetized rats. Specifically, putative pyramidal neurons in PV shRNA treated rats fired on average at a rate almost twice that observed in control rats (control shRNA: 0.73 ± 0.09 c.f. PV shRNA: 1.22 ± 0.12 Hz; Mann-Whitney Rank Sum Test, $U = 169.5$, $p < 0.05$, $n = 22-31$ neurons; Fig 2A).

Dopamine System Function

We have previously demonstrated that aberrant vHipp activity results in an augmented dopamine system function (Lodge and Grace, 2007). The vHipp has been demonstrated to selectively regulate the proportion of spontaneously active dopamine neurons in the ventral tegmental area (VTA). This is thought to reflect the gain of the dopamine system (Lodge and Grace, 2007). For this reason, we examined whether vHipp PV knockdown was sufficient to alter dopamine neuron activity. Indeed, we now demonstrate that vHipp PV knockdown significantly increases dopamine neuron population activity (control shRNA: 1.02 ± 0.06 c.f. PV shRNA: 1.59 ± 0.18 cells/track; Mann-Whitney Rank Sum Test, $U = 4.50$, $p < 0.05$, $n = 7-8$ rats; Fig 2B) without altering average firing rate (control shRNA: 4.03 ± 0.36 c.f. PV shRNA: 3.75 ± 0.28 Hz; Mann-Whitney Rank Sum Test, $U = 1597$, $p > 0.05$, $n = 49-69$ neurons; Fig 2C) or average percent burst firing (control shRNA: 43.2 ± 3.8 c.f. PV shRNA: 35.1 ± 3.4 %; Mann-Whitney Rank Sum Test, $U = 1375$, $p > 0.05$, $n = 49-69$ neurons; Fig 2D).

Behavioral Response to Amphetamine

Schizophrenia patients consistently demonstrate exaggerated responses to psychomotor stimulants (Janowsky et al., 1973). A similar paradigm can be used to examine behavioral deficits in putative animal models of schizophrenia (Lodge and Grace, 2007; Perez and Lodge, 2013; Perez et al., 2013). Here we demonstrate that vHipp PV knockdown produces an enhanced locomotor response to amphetamine administration. Specifically, PV shRNA treated rats display a significantly enhanced locomotor response to low dose (0.5mg/kg, i.p.) amphetamine administration when compared to non-silencing shRNA controls (2-way ANOVA of 0.5mg/kg dose; $F(\text{treatment})=9.851$; Holm-Sidak $t=3.139$, $p<0.05$; Fig 3), and also in response to the higher dose (2-way ANOVA of 2.0mg/kg dose; $F(\text{treatment})=4.225$; Holm-Sidak $t=2.056$, $p<0.05$; Fig 3) without changes in baseline activity (2-way ANOVA of baseline; $F(\text{treatment})=2.468$; $p>0.05$; Fig 3), consistent with previous observations in rodent models (Lodge and Grace, 2007; Perez and Lodge, 2013; Perez et al., 2013).

Discussion

Previous studies in postmortem schizophrenia patients, as well as, rodent models have demonstrated a reduction in PV expression throughout the cortex (Lewis et al., 2012; Lodge et al., 2009) and hippocampus (Konradi et al., 2011; Lodge et al., 2009); however, whether this contributes to, or is a consequence of, the disease is not currently known. We now demonstrate that a reduction in vHipp PV expression is sufficient, in and of itself, to increase vHipp activity and induce downstream changes in dopamine neuron activity and a behavior that is analogous to the positive symptoms of schizophrenia.

There is a considerable literature implicating aberrant dopamine signaling in the pathophysiology of schizophrenia (Abi-Dargham, 2004; Grace, 2004; Howes et al., 2012; Howes and Kapur, 2009; Lodge and Grace, 2011). This can be directly examined in rodent models by recording the electrophysiological activity of midbrain dopamine neurons (Lodge and Grace, 2007; Perez and Lodge, 2012, 2013). Such studies have suggested that the increase in dopamine system function is associated with changes in the number of spontaneously active neurons in the absence of changes in firing rate or bursting pattern (Lodge and Grace, 2007; Perez and Lodge, 2012, 2013). Under control conditions, it is thought that about 50% of the midbrain dopamine neurons are quiescent due to a tonic GABAergic drive from the ventral pallidum (VP) (Grace et al., 2007). This provides an opportunity whereby the gain of the dopamine signal can be enhanced or attenuated by modulation of afferent inputs. In the case of schizophrenia, we posit that the ability to appropriately regulate the gain of the system is lost, as most of the dopamine neurons are spontaneously active (Lodge and Grace, 2007, 2011). As a result, it is likely that all stimuli are now being assigned the same salience, and may contribute to the paranoia consistently observed in patients. While a number of regions can regulate pallidal activity, the hippocampus has been consistently implicated in schizophrenia. The ventral regions of the hippocampus regulate the number of spontaneously active dopamine neurons via a multi-synaptic pathway that involves the nucleus accumbens (NAc) (Lodge and Grace, 2007, 2011). Thus, activation of the vHipp will increase GABAergic transmission from the NAc that, in turn, suppresses VP tone to the VTA. The net result is a disinhibition of the VTA and

a selective increase in the number of dopamine neurons firing spontaneously (Lodge and Grace, 2007, 2011).

While hippocampal hyperactivity has been observed in both rodent models (Lodge and Grace, 2007; Perez and Lodge, 2013; Shah and Lodge, 2013), as well as human schizophrenia patients (Heckers, 2004; Kraguljac et al., 2013; Malaspina et al., 1999; Medoff et al., 2001; Schobel et al., 2009), the cause of this aberrant activity has not been conclusively demonstrated. One hypothesis is that a deficit in interneuron function may result in increases in pyramidal cell firing; however, this has not been verified experimentally. Here we demonstrate that selectively decreasing PV mRNA expression in the vHipp of 'normal' Sprague-Dawley rats, is sufficient to recapitulate the hippocampal hyperactivity observed in rodent models, as well as patients (Heckers, 2004; Kraguljac et al., 2013; Lodge and Grace, 2007; Malaspina et al., 1999; Medoff et al., 2001; Perez and Lodge, 2013; Schobel et al., 2009; Shah and Lodge, 2013). The exact mechanism by which a decrease in PV leads to hippocampal hyperactivity has not been conclusively demonstrated. PV is a calcium binding protein with a slow onset that does not significantly affect the amplitude of fast Ca^{2+} transients, but alters the decay of $[\text{Ca}^{2+}]_i$ (Schwaller et al., 2002). In the hippocampus, a loss of PV expression does not alter either single inhibitory postsynaptic responses or paired-pulse modulation of IPSCs (Vreugdenhil et al., 2003). Rather, an enhanced facilitation of GABA release is observed after repetitive high frequency stimulation in PV $-/-$ mice (Vreugdenhil et al., 2003). Thus, the hyperactivity observed following PV shRNA in the current study is not likely attributable to changes in the functionality of the interneuron but may be attributable to neuronal loss. Indeed, the buffering capacity of PV has been suggested to protect from Ca^{2+} -mediated excitotoxic insult (Figueredo-Cardenas et al., 1998; Van Den Bosch et al., 2002).

In addition to increases in vHipp activity, PV shRNA induced downstream changes in VTA dopamine neuron activity. Consistent with the pharmacological studies detailed above, rats with selective vHipp PV knockdown display increases in the number of spontaneously active VTA dopamine neurons, without overt changes in firing rate or bursting pattern. This neurophysiological phenotype is consistent with that observed in rodent models of schizophrenia, such as the MAM-treated rat (Lodge and Grace, 2007; Perez and Lodge, 2012, 2013; Perez et al., 2013). Interestingly, while the increase in dopamine neuron activity is qualitatively similar to that of the MAM-treated rat, it appears to be of a smaller magnitude. This is likely due to differences between the acute PV knockdown performed in the current study and the more sustained neurodevelopmental loss observed in the MAM rat (Chen et al., 2014). Indeed, an increase in dopamine neuron population activity, similar to that observed following vHipp PV shRNA, is present following the degradation of vHipp perineuronal nets in adulthood (Shah and Lodge, 2013).

Symptoms such as hallucinations, delusions and paranoia obviously cannot be assessed in rodents; however, there are behavioral alterations that can be modeled in both patients and rats. One reliable and reproducible observation is an enhanced response to psychomotor stimulants. Thus, the administration of small doses of amphetamine will precipitate psychosis in schizophrenia patients (Janowsky et al., 1973). Similarly, rodent models display an exaggerated response to psychomotor stimulants, which is quantified by examining the

magnitude of the locomotor response (Lodge and Grace, 2007; Moore et al., 2006; Perez and Lodge, 2013; Perez et al., 2013). As a behavioral correlate of the augmented dopamine neuron activity, we examined the locomotor response to amphetamine in rats treated with vHipp PV shRNA and demonstrate that a reduction in vHipp PV expression produces an enhanced response consistent with that observed in rodent models of the disease.

Taken together, we demonstrate a potential key role for PV in the pathophysiology of schizophrenia that contributes to aberrant hippocampal neuronal activity, as well as, downstream changes in dopamine system function that contribute to positive symptoms. It should be noted that deficits in PV expression in patients and rodent models are not limited to the hippocampus, but are also observed throughout the prefrontal cortex (Lewis et al., 2005). Given that we did not investigate the effects of PV knockdown in regions other than the vHipp, we cannot conclude that the symptoms are exclusively hippocampal. Indeed, deficits in PV interneuron function in the mPFC are likely associated with non-dopaminergic symptoms in schizophrenia patients, such as deficits in cognition (Pantelis et al., 1997). These data are also consistent with recent demonstrations examining the early postnatal interneuron-specific knockdown of the NMDA receptor. Specifically, these mice demonstrate reductions in PV expression and increases in pyramidal neuron activity which are associated with behavioral deficits consistent with a schizophrenia-like phenotype; including deficits in pre-pulse inhibition of startle, novelty-induced hyperlocomotion, a decrease in saccharine preference, and deficits in spatial working memory (Belforte et al., 2010). Collectively, we demonstrate that a knockdown of PV expression in the vHipp is sufficient, in and of itself, to produce the aberrant dopamine neuron activity and behavioral hyper-responsivity to psychomotor stimulants that is thought to contribute to psychosis in schizophrenia patients.

Acknowledgments

Funding body agreements:

This work was supported by an R01 to DJL (MH090067) and F31 to SMP (MH098564) from the NIH.

References

- Abdul-Monim Z, Neill JC, Reynolds GP. Sub-chronic psychotomimetic phencyclidine induces deficits in reversal learning and alterations in parvalbumin-immunoreactive expression in the rat. *Journal of psychopharmacology*. 2007; 21(2):198–205. [PubMed: 17329300]
- Abi-Dargham A. Do we still believe in the dopamine hypothesis? New data bring new evidence. *Int J Neuropsychopharmacol*. 2004; 7(Suppl 1):S1–5. [PubMed: 14972078]
- Agid O, Shapira B, Zislin J, Ritsner M, Hanin B, Murad H, Troudart T, Bloch M, Heresco-Levy U, Lerer B. Environment and vulnerability to major psychiatric illness: A case control study of early parental loss in major depression, bipolar disorder and schizophrenia. *Molecular Psychiatry*. 1999; 4(2):163–172. [PubMed: 10208448]
- Behrens MM, Ali SS, Dao DN, Lucero J, Shekhtman G, Quick KL, Dugan LL. Ketamine-induced loss of phenotype of fast-spiking interneurons is mediated by NADPH-oxidase. *Science*. 2007; 318(5856):1645–1647. [PubMed: 18063801]
- Belforte JE, Zsiros V, Sklar ER, Jiang Z, Yu G, Li Y, Quinlan EM, Nakazawa K. Postnatal NMDA receptor ablation in corticolimbic interneurons confers schizophrenia-like phenotypes. *Nature neuroscience*. 2010; 13(1):76–83.

- Benes FM, Berretta S. GABAergic interneurons: Implications for understanding schizophrenia and bipolar disorder. *Neuropsychopharmacology*. 2001; 25(1):1–27. [PubMed: 11377916]
- Bhugra D. The global prevalence of schizophrenia. *PLoS Med*. 2005; 2(5):e151. quiz e175. [PubMed: 15916460]
- Chen L, Perez SM, Lodge DJ. An augmented dopamine system function is present prior to puberty in the methylazoxymethanol acetate rodent model of schizophrenia. *Developmental neurobiology*. 2014
- Cunningham MO, Hunt J, Middleton S, LeBeau FE, Gillies MJ, Davies CH, Maycox PR, Whittington MA, Racca C. Region-specific reduction in entorhinal gamma oscillations and parvalbumin-immunoreactive neurons in animal models of psychiatric illness. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2006; 26(10):2767–2776. [PubMed: 16525056]
- Figueredo-Cardenas G, Harris CL, Anderson KD, Reiner A. Relative resistance of striatal neurons containing calbindin or parvalbumin to quinolinic acid-mediated excitotoxicity compared to other striatal neuron types. *Experimental neurology*. 1998; 149(2):356–372. [PubMed: 9500958]
- Francois J, Ferrandon A, Koning E, Angst MJ, Sandner G, Nehlig A. Selective reorganization of GABAergic transmission in neonatal ventral hippocampal-lesioned rats. *The international journal of neuropsychopharmacology/official scientific journal of the Collegium Internationale Neuropsychopharmacologicum*. 2009; 12(8):1097–1110.
- Goldman-Rakic PS. Cellular basis of working memory. *Neuron*. 1995; 14(3):477–485. [PubMed: 7695894]
- Gonzalez-Burgos G, Lewis DA. NMDA receptor hypofunction, parvalbumin-positive neurons, and cortical gamma oscillations in schizophrenia. *Schizophr Bull*. 2012; 38(5):950–957. [PubMed: 22355184]
- Grace, AA. Developmental dysregulation of the dopamine system and the pathophysiology of schizophrenia. In: Keshavan, MS.; Kennedy, JL.; Murray, RM., editors. *Neurodevelopment and Schizophrenia*. Cambridge University Press; Cambridge, UK: 2004. p. 273-294.
- Grace AA, Bunney BS. Intracellular and extracellular electrophysiology of nigral dopaminergic neurons--I. Identification and characterization. *Neuroscience*. 1983; 10(2):301–315. [PubMed: 6633863]
- Grace AA, Floresco SB, Goto Y, Lodge DJ. Regulation of firing of dopaminergic neurons and control of goal-directed behaviors. *Trends in Neurosciences*. 2007; 30(5):220–227. [PubMed: 17400299]
- Harte MK, Powell SB, Swerdlow NR, Geyer MA, Reynolds GP. Deficits in parvalbumin and calbindin immunoreactive cells in the hippocampus of isolation reared rats. *Journal of Neural Transmission*. 2007; 114(7):893–898. [PubMed: 17594127]
- Hashimoto T, Volk DW, Eggen SM, Mirnics K, Pierri JN, Sun Z, Sampson AR, Lewis DA. Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *Journal of Neuroscience*. 2003; 23(15):6315–6326. [PubMed: 12867516]
- Heckers S. The hippocampus in schizophrenia. *American Journal of Psychiatry*. 2004; 161(11):2138–2139. [PubMed: 15514431]
- Howes OD, Kambeitz J, Kim E, Stahl D, Slifstein M, Abi-Dargham A, Kapur S. The nature of dopamine dysfunction in schizophrenia and what this means for treatment: Meta-analysis of imaging studies. *Archives of General Psychiatry*. 2012; 69(8):776–786. [PubMed: 22474070]
- Howes OD, Kapur S. The Dopamine Hypothesis of Schizophrenia: Version III—The Final Common Pathway. *Schizophrenia Bulletin*. 2009; 35(3):549–562. [PubMed: 19325164]
- Hyland BI, Reynolds JNJ, Hay J, Perk CG, Miller R. Firing modes of midbrain dopamine cells in the freely moving rat. *Neuroscience*. 2002; 114(2):475–492. [PubMed: 12204216]
- Janowsky DS, el-Yousel MK, Davis JM, Sekerke HJ. Provocation of schizophrenic symptoms by intravenous administration of methylphenidate. *Arch Gen Psychiatry*. 1973; 28(2):185–191. [PubMed: 4630714]
- Kawaguchi Y, Hama K. Fast-spiking non-pyramidal cells in the hippocampal CA3 region, dentate gyrus and subiculum of rats. *Brain Res*. 1987; 425(2):351–355. [PubMed: 3427435]
- Konradi C, Yang CK, Zimmerman EI, Lohmann KM, Gresch P, Pantazopoulos H, Berretta S, Heckers S. Hippocampal interneurons are abnormal in schizophrenia. *Schizophrenia Research*. 2011; 131(1–3):165–173. [PubMed: 21745723]

- Kraguljac NV, White DM, Reid MA, Lahti AC. Increased hippocampal glutamate and volumetric deficits in unmedicated patients with schizophrenia. *JAMA psychiatry*. 2013; 70(12):1294–1302. [PubMed: 24108440]
- Lewis DA, Curley AA, Glausier JR, Volk DW. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci*. 2012; 35(1):57–67. [PubMed: 22154068]
- Lewis DA, Hashimoto T, Volk DW. Cortical inhibitory neurons and schizophrenia. *Nature Reviews Neuroscience*. 2005; 6(4):312–324.
- Lodge DJ, Behrens MM, Grace AA. A loss of parvalbumin-containing interneurons is associated with diminished oscillatory activity in an animal model of schizophrenia. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2009; 29(8):2344–2354. [PubMed: 19244511]
- Lodge DJ, Grace AA. Aberrant hippocampal activity underlies the dopamine dysregulation in an animal model of schizophrenia. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2007; 27:11424–11430. [PubMed: 17942737]
- Lodge DJ, Grace AA. Gestational methylazoxymethanol acetate administration: A developmental disruption model of schizophrenia. *Behav Brain Res*. 2009; 7(204 2):306–312. [PubMed: 19716984]
- Lodge DJ, Grace AA. Hippocampal dysregulation of dopamine system function and the pathophysiology of schizophrenia. *Trends in Pharmacological Sciences*. 2011; 32(9):507–513. [PubMed: 21700346]
- Malaspina D, Storer S, Furman V, Esser P, Printz D, Berman A, Lignelli A, Gorman J, Van Heertum R. SPECT study of visual fixation in schizophrenia and comparison subjects. *Biol Psychiatry*. 1999; 46(1):89–93. [PubMed: 10394477]
- Medoff DR, Holcomb HH, Lahti AC, Tamminga CA. Probing the human hippocampus using rCBF: contrasts in schizophrenia. *Hippocampus*. 2001; 11(5):543–550. [PubMed: 11732707]
- Moore H, Jentsch JD, Ghajarnia M, Geyer MA, Grace AA. A neurobehavioral systems analysis of adult rats exposed to methylazoxymethanol acetate on E17: implications for the neuropathology of schizophrenia. *Biol Psychiatry*. 2006; 60(3):253–264. [PubMed: 16581031]
- Pantelis C, Barnes TRE, Nelson HE, Tanner S, Weatherley L, Owen AM, Robbins TW. Frontal-striatal cognitive deficits in patients with chronic schizophrenia. *Brain*. 1997; 120(10):1823–1843. [PubMed: 9365373]
- Perez SM, Lodge DJ. Aberrant dopamine D2-like receptor function in a rodent model of schizophrenia. *Journal of Pharmacology and Experimental Therapeutics*. 2012; 343(2):288–295. [PubMed: 22859862]
- Perez SM, Lodge DJ. Hippocampal interneuron transplants reverse aberrant dopamine system function and behavior in a rodent model of schizophrenia. *Molecular Psychiatry*. 2013; 18(11):1193–1198. [PubMed: 23979606]
- Perez SM, Shah A, Asher A, Lodge DJ. Hippocampal deep brain stimulation reverses physiological and behavioural deficits in a rodent model of schizophrenia. *Int J Neuropsychopharmacol*. 2013; 16(6):1331–1339. [PubMed: 23190686]
- Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, Ruderfer DM, McQuillin A, Morris DW, Odushlaine CT, Corvin A, Holmans PA, Odonovan MC, MacGregor S, Gurling H, Blackwood DHR, Craddock NJ, Gill M, Hultman CM, Kirov GK, Lichtenstein P, Muir WJ, Owen MJ, Pato CN, Scolnick EM, St Clair D, Williams NM, Georgieva L, Nikolov I, Norton N, Williams H, Toncheva D, Milanova V, Thelander EF, O'Dushlaine CT, Kenny E, Quinn EM, Choudhury K, Datta S, Pimm J, Thirumalai S, Puri V, Krasucki R, Lawrence J, Quested D, Bass N, Crombie C, Fraser G, Leh Kuan S, Walker N, McGhee KA, Pickard B, Malloy P, MacLean AW, Van Beck M, Pato MT, Medeiros H, Middleton F, Carvalho C, Morley C, Fanous A, Conti D, Knowles JA, Paz Ferreira C, MacEdo A, Helena Azevedo M, Kirby AN, Ferreira MAR, Daly MJ, Chambert K, Kuruvilla F, Gabriel SB, Ardlie K, Moran JL, Sklar P. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009; 460(7256):748–752. [PubMed: 19571811]
- Saha S, Chant D, Welham J, McGrath J. A Systematic Review of the Prevalence of Schizophrenia. *PLoS Medicine*. 2005; 2(5):e141. [PubMed: 15916472]

- Sawa A, Snyder SH. Schizophrenia: Diverse approaches to a complex disease. *Science*. 2002; 296(5568):692–695. [PubMed: 11976442]
- Schobel SA, Lewandowski NM, Corcoran CM, Moore H, Brown T, Malaspina D, Small SA. Differential targeting of the CA1 subfield of the hippocampal formation by schizophrenia and related psychotic disorders. *Arch Gen Psychiatry*. 2009; 66(9):938–946. [PubMed: 19736350]
- Schwaller B, Meyer M, Schiffmann S. ‘New’ functions for ‘old’ proteins: The role of the calcium-binding proteins calbindin D-28k, calretinin and parvalbumin, in cerebellar physiology. Studies with knockout mice. *Cerebellum*. 2002; 1(4):241–258. [PubMed: 12879963]
- Shah A, Lodge DJ. A loss of hippocampal perineuronal nets produces deficits in dopamine system function: relevance to the positive symptoms of schizophrenia. *Transl Psychiatry*. 2013; 3:e215. [PubMed: 23321812]
- Van Den Bosch L, Schwaller B, Vleminckx V, Meijers B, Stork S, Ruehlicke T, Van Houtte E, Klaassen H, Celio MR, Missiaen L, Robberecht W, Berchtold MW. Protective effect of parvalbumin on excitotoxic motor neuron death. *Experimental neurology*. 2002; 174(2):150–161. [PubMed: 11922657]
- van der Meer MA, Redish AD. Theta phase precession in rat ventral striatum links place and reward information. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2011; 31(8):2843–2854. [PubMed: 21414906]
- Vreugdenhil M, Jefferys JG, Celio MR, Schwaller B. Parvalbumin-deficiency facilitates repetitive IPSCs and gamma oscillations in the hippocampus. *J Neurophysiol*. 2003; 89(3):1414–1422. [PubMed: 12626620]
- Weinberger DR, Berman KF, Zec RF. Physiologic dysfunction of dorsolateral prefrontal cortex in schizophrenia. I. Regional cerebral blood flow evidence. *Archives of General Psychiatry*. 1986; 43(2):114–124. [PubMed: 3947207]

Figure 1. Lentiviral administration of parvalbumin (PV) shRNA decreases the number of PV positive interneurons in the ventral hippocampus. Images in A & B represent PV positive neurons (red) as determined by immunohistochemistry. The graph in C depicts the quantification of computer assisted cell counts throughout the ventral extent of the hippocampus. † represents $p < 0.05$, Student's t-test.

Figure 2.

Lentiviral administration of parvalbumin (PV) shRNA increases hippocampal and dopamine neuron activity. Putative pyramidal neurons throughout the hippocampus were recorded (A) and demonstrated significantly greater activity in PV shRNA treated rats. Downstream increases in dopamine neuron population activity (B) were also observed without significant changes in average firing rate (C) or burst firing (D). † represents $p < 0.05$, Mann Whitney Rank Sum test.

Figure 3.

Lentiviral administration of parvalbumin (PV) shRNA augments the locomotor response to amphetamine. Baseline locomotor activity was not significantly affected by vHipp PV knockdown (black squares) relative to control (non-silencing shRNA: Grey circles). In contrast, the locomotor response to both low (0.5mg/kg) and high (2.0mg/kg) doses of amphetamine were significantly enhanced demonstrating that a decrease in vHipp PV expression is sufficient to induce behavioral alterations consistent with those observed in schizophrenia. † represents $p < 0.05$, 2-way ANOVA.