Using species distribution models to optimize vector control in the framework of the tsetse eradication campaign in Senegal

Ahmadou H. Dicko^a, Renaud Lancelot^{b,c}, Momar T. Seck^a, Laure Guerrini^{d,e}, Baba Sall^f, Mbargou Lo^f, Marc J. B. Vreysen^g, Thierry Lefrançois^{b,c}, William M. Fonta^h, Steven L. Peckⁱ, and Jérémy Bouyer^{a,b,c,1}

^aLaboratoire National d'Elevage et de Recherches Vétérinaires, Institut Sénégalais de Recherches Agricoles, BP 2057, Hann, Dakar, Sénégal; ^bUnité Mixte de Recherche Contrôle des Maladies Animales Exotiques et Emergentes, Centre de Coopération Internationale en Recherche Agronomique pour le Développement, 34398 Montpellier, France; ^cUnité Mixte de Recherche 1309 Contrôle des Maladies Animales Exotiques et Emergentes, Institut National de la Recherche Agronomique, 34398 Montpellier, France; ^dUnité de Recherche Animal et Gestion Intégrée des Risques, Centre de Coopération Internationale en Recherche Agronomique pour le Développement, 34398 Montpellier, France; ^eDepartment Environment and Societies, University of Zimbabwe, Harare, Zimbabwe; ^fDirection des Services Vétérinaires, BP 45 677, Dakar, Sénégal; ^gInsect Pest Control Laboratory, Joint Food and Agriculture Organization of the United Nations/International Atomic Energy Agency Programme of Nuclear Techniques in Food and Agriculture, A-1400 Vienna, Austria; ^hWest African Science Center for Climate Change and Adapted Land Use, BP 13621, Ouagadougou, Burkina Faso; and ⁱBiology Department, Brigham Young University, Provo, UT 84602

Edited by Fred L. Gould, North Carolina State University, Raleigh, NC, and approved June 6, 2014 (received for review April 29, 2014)

Tsetse flies are vectors of human and animal trypanosomoses in sub-Saharan Africa and are the target of the Pan African Tsetse and Trypanosomiasis Eradication Campaign (PATTEC). *Glossina palpalis gambiensis* (Diptera: Glossinidae) is a riverine species that is still present as an isolated metapopulation in the Niayes area of Senegal. It is targeted by a national eradication campaign combining a population reduction phase based on insecticide-treated targets (ITTs) and cattle and an eradication phase based on the sterile insect technique. In this study, we used species distribution models to optimize control operations. We compared the probability of the presence of *G. p. gambiensis* and habitat suitability using a regularized logistic regression and Maxent, respectively. Both models performed well, with an area under the curve of 0.89 and 0.92, respectively. Only the Maxent model predicted an expert-based classification of landscapes correctly. Maxent predictions were therefore used throughout the eradication campaign in the Niayes to make control operations more efficient in terms of deployment of ITTs, release density of sterile males, and location of monitoring traps used to assess program progress. We discuss how the models' results informed about the particular ecology of tsetse in the target area. Maxent predictions allowed optimizing efficiency and cost within our project, and might be useful for other tsetse control campaigns in the framework of the PATTEC and, more generally, other vector or insect pest control programs.

area-wide integrated pest management | genetic control | aerial release | chilled adult technique

Tsetse are vectors of human African trypanosomosis, a major neglected tropical disease (1), and African animal trypanosomosis (AAT), one of the most important pathological constraints to livestock development in 38 infested African countries (2). The Pan African Tsetse and Trypanosomiasis Eradication Campaign is a political initiative started in 2001 that calls for intensified efforts to reduce the tsetse and trypanosomosis problem (3). As part of this global effort, the government of Senegal embarked in 2007 on a tsetse eradication campaign in a 1,000-km² target area of the Niayes region, neighboring the capital Dakar. In this area, the limits of the distribution of the tsetse target populations were assessed using a stratified entomological sampling frame based on remote sensing indicators (4). The only tsetse species present was *Glossina palpalis gambiensis* Vanderplank, and it was responsible for the cyclical transmission of three trypanosome species, namely *Trypanosoma vivax*, *T. congolense*, and *T. brucei brucei*, listed in order of importance (5). The high prevalence of animal trypanosomosis in the Niayes (serological prevalence of 28.7% for *T. vivax*) hampered

peri-urban intensification of cattle production (particularly dairy cattle). A population genetics study demonstrated that the *G. p. gambiensis* population of the Niayes was completely isolated from the main tsetse belt in the southeastern part of Senegal (6). This comprehensive set of entomological, veterinary, genetic, and other baseline data confirmed the isolated nature of the *G. p. gambiensis* population in the Niayes, which prompted the government of Senegal to select an eradication strategy using area-wide integrated pest management (AW-IPM) principles (7).

The successful implementation of an AW-IPM strategy requires a thorough understanding of the ecology of the target population, particularly its spatial distribution, a study which was undertaken in the Niayes from 2007 to 2011 before the start of the operational eradication efforts. The selected strategy integrates insecticide-treated targets (ITTs) and cattle with the release of sterile insects. As the habitat of *G. p. gambiensis* is very fragmented (4), the targeting of suitable habitats for deployment of the ITTs is crucial to optimize cost efficiency (8) but also to enable the selection of appropriate sites for deployment of the

Significance

Tsetse flies transmit human and animal trypanosomoses in sub-Saharan Africa, respectively a major neglected disease and the most important constraint to cattle production in infested countries. They are the target of the Pan African Tsetse and Trypanosomiasis Eradication Campaign (PATTEC). Here we show how distribution models can be used to optimize a tsetse eradication campaign in Senegal. Our results allow a better understanding of the relationships between tsetse presence and various environmental parameters measured by remote sensing. Furthermore, we argue that the methodology developed should be integrated into future tsetse control efforts that are planned under the umbrella of the PATTEC initiative. The results have a generic value for vector and pest control campaigns, especially when eradication is contemplated.

Author contributions: R.L., M.T.S., B.S., M.L., M.J.B.V., and J.B. designed research; A.H.D. and J.B. performed research; A.H.D., R.L., M.T.S., L.G., T.L., W.M.F., S.L.P., and J.B. contributed new reagents/analytic tools; A.H.D., R.L., and J.B. analyzed data; and A.H.D., R.L., M.T.S., L.G., B.S., M.L., M.J.B.V., T.L., W.M.F., S.L.P., and J.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: bouyer@cirad.fr.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.140773111/-DCSupplemental.

Fig. 2. ROC curves and AUCs of the regularized logistic regression and Maxent models. These results were obtained using the presence-absence validation dataset (Left) and the expert-based landscape classification from aerial images (Right).

Marginal response curves for the different predictors (Fig. S3) showed that the percentage of the area covered by forest was positively correlated with habitat suitability, with a sharp increase after 5% of forests and a plateau after 20%. With respect to average day LST, habitat suitability was stable between 20 and 35 °C and then sharply decreased; whereas average day LST was not low enough in the study area to limit the presence of *G. p. gambiensis*, high temperatures were unsuitable. The correlation between minimum day LST and habitat suitability was bell-shaped, with a maximum between 20 and 25 °C. This could be related to the buffering effect of dense tree cover on temperature drops at night. Indeed, night temperature drops faster in open environments than in the forest. Habitat suitability dropped sharply, with minimum temperatures exceeding 25 °C. The correlation between mean NDVI and habitat suitability was a sigmoid curve that increased sharply after 0.2. Regarding NDVI range, a plateau of habitat suitability was observed until 0.5 and sharply decreased after this threshold, thus illustrating the importance of perennial tree vegetation for this tsetse species. The relationship with habitat suitability was similar for MIR SD, related to the strong negative correlation between NDVI and MIR.

Use of Maxent Predictions in the Tsetse Eradication Project. Before the availability of the Maxent model in the Senegal project, operational choices such as selection of trap sites were made using a vegetation classification obtained from a Landsat 7 Enhanced Thematic Mapper Plus (ETM+) image of April 2001. Suitable habitat for *G. p. gambiensis* was mapped from this classification with high sensitivity but low specificity (4). The availability of the Maxent predictions was used in four ways to optimize the implementation of the eradication effort.

First, boundaries of the target eradication area, shown as black grid cells in Fig. 3, were validated by the model. All of the new suitable habitat areas identified by the Maxent model within a range of 5 km around sites where wild *G. p. gambiensis* had been trapped (at any occasion) were already included in the control strategy (Discussion). The target area was subdivided into four operational blocks, which are being sequentially addressed during the operational program (Fig. 4). Each block was subjected to a 1-y tsetse-density reduction phase, followed by an 18-mo eradication period using the sterile insect technique.

Second, the spatial distribution of the monitoring traps deployed since January 2012 in block 1 and since October 2012 in block 2 was modified through relocating 22 of the 97 monitoring traps (23%) to more suitable sites according to habitat suitability as predicted by the Maxent model (Fig. S4). In block 3, where monitoring had not started yet at the time of writing, the monitoring traps will also be deployed in sites that have a high (predicted) suitability value as indicated by the model.

Third, 1,347 insecticide-impregnated traps used for tsetse-density reduction were deployed in block 2 according to predictions of

Maxent during the period December 2012 to February 2013, from which 661 were renewed during the period January to February 2014. The total surface area covered by block 2 is close to 500 km² but contains only 80.6 km² of predicted suitable habitat, thus giving a final trap density of 16.7 traps per km² of suitable habitat (Fig. 4).

Fourth, predicted suitable habitats were also used to optimize aerial releases of sterile male *G. p. gambiensis*. Polygons with similar surface areas of suitable patches were identified (RL1 and RL2 in block 1, for example; Fig. 4) and the density of released sterile males was adjusted proportionally to the area of suitable habitat in these polygons (0.24 and 0.11 flies per ha in RL1 and RL2, respectively).

Optimization of Tsetse Suppression and Eradication by the Model. In the first block, the apparent density of *G. p. gambiensis* dropped from an average of 0.42 (SD 0.39) flies per trap per d before control to an average of 0.04 (SD 0.11) flies per trap per d and 0.003 (SD 0.01) flies per trap per d during the suppression and eradication phases, respectively (Fig. 5). In the second block, the apparent density dropped from an average of 1.24 (SD 1.23) flies per trap per d before control to an average of 0.005 (SD 0.017) flies per trap per d during the suppression phase.

Apparent fly density in block 2 was higher than in block 1 (generalized linear mixed model, $P = 0.009$; SI Methods) before the start of the suppression, which initially sharply reduced tsetse densities ($P = 0.001$). This effect was limited in time (6 mo), and tsetse density remained stable and even increased with time thereafter ($P < 10^{-3}$), although remaining at a very low level (< 0.04 fly per trap per d) until the eradication phase started. The cumulated reduction of densities with time was higher in block 2 (reduction of 99.6%) than in block 1 (reduction of 90.4%) ($P < 10^{-3}$).

In block 1, the last wild fly was captured on August 9, 2012, ~6 mo after the start of sterile male releases. It was an old female (more than 40 d old) in its fourth larviposition cycle with an empty uterus, and the next follicle was immature and small, indicating an abortion. This female showed a copulation scar and its spermatheca was 85% filled, indicating that its sterility was probably induced by a mate with one sterile male. From the beginning of the eradication phase in block 1 (March 16, 2012) to the date corresponding to the last capture, only three other wild

Fig. 3. Predictions of the regularized logistic regression and Maxent models. In the case of the logistic model, the predictions correspond to probabilities of presence, whereas in the case of the Maxent model, they correspond to a rescaled suitability index (logistic output). The black cells correspond to the target area. Black and red crosses represent absence and presence data, respectively.

Table 1. Prediction qualities of the regularized logistic regression and Maxent models using the validation dataset

Model	Threshold	Sensitivity	Specificity	PCC	AUC
Glmnet	0.5	0.63 (0.07)	0.93 (0.03)	0.78 (0.04)	0.89 (0.03)
Maxent	0.5	0.64 (0.06)	0.96 (0.03)	0.80 (0.04)	0.92 (0.02)
Glmnet	0.11*	0.96 (0.03)	0.50 (0.07)	0.73 (0.04)	—
Maxent	0.13*	0.96 (0.03)	0.57 (0.07)	0.77 (0.04)	—
Glmnet	0.35 [†]	0.82 (0.05)	0.89 (0.04)	0.86 (0.03)	—
Maxent	0.32 [†]	0.86 (0.05)	0.79 (0.06)	0.82 (0.04)	—

These results were obtained with the presence-absence validation dataset. SDs are presented in parentheses. Glmnet, regularized logistic regression.

*Selected to optimize sensitivity.

[†]Selected to optimize PCC.

females could be dissected and all had indications of having mated with a sterile male. The average percentage of sterile males as a proportion of all catches was then 99.2% (SD 1.6%), corresponding to a sterile-to-wild male ratio of 130. The percentage of sterile males remained 100% thereafter (no wild fly was captured for 78 weekly collections with 25 monitoring traps), corresponding to a very likely eradication (probability of not detecting potential remaining flies of 0.002 only).

Discussion

Ecological Niche of *G. p. gambiensis* in the Niayes. Our analysis showed that the ecological niche of *G. p. gambiensis* in the Niayes area corresponded to permanent ligneous vegetation with a tree density sufficient to provide adequate shade and buffer temperature and relative hygrometry variations in comparison with macroclimatic conditions occurring in the surrounding open environments. Air temperature in dense tree vegetation in gallery forests can be 4 °C lower compared with the surroundings, and relative humidity 15% higher. This habitat provides resting sites for *G. p. gambiensis* in contrast to the more open habitat into which they may disperse for short periods (some hours) in search of a blood meal, namely their hunting sites. Suitable *G. p. gambiensis* habitat may be seasonal because of the variations of macroclimatic conditions and of nonpermanent vegetation.

The Maxent model confirmed that permanent dense tree vegetation was important for *G. p. gambiensis*, but also that the larger the area occupied by the flies during the rainy season (corresponding to their dispersal capacity) the more suitable this habitat appears to be for *G. p. gambiensis*. Evidence is provided by the positive correlation of the forest range with the ENFA marginality axis (Fig. 1).

The data layers derived from Landsat images, which have a higher spatial resolution (30 m) than Moderate Resolution

Imaging Spectroradiometer from the National Aeronautics and Space Administration (MODIS) data (250 m), were important predictors of the presence of *G. p. gambiensis*. This was expected, given their ability to survive in very small vegetation patches in the Niayes (4). It would be difficult to obtain such fine-scale data for larger regions of West Africa. Fortunately, a good correlation was found between forest predictors and minimum night LST (Fig. 1). Moreover, the regularized gain is only reduced by 6% when all forest predictors are removed.

Comparison of the Regularized Logistic Regression and Maxent Models.

The nature of predictions differs between regularized logistic regression (probability) and Maxent (index). However, the two predictions were highly correlated, as observed elsewhere (17). Model quality-assessment metrics were similar using the presence-absence validation dataset. Maxent predicted suitable areas better than regularized logistic regression based on the expert-based landscape classification. Tsetse presence data are generally more meaningful than absence data, because all known traps have a very low efficiency with respect to trapping rates (as a percentage of available individuals), namely $\leq 1\%$ per d per km² (4). In the Niayes, observed trapping efficiencies were as low as 0.3% per d per km² following the release of more than 200,000 sterile male flies. In our case, particular conditions (feasibility study of an eradication project) allowed a rigorous selection of false absences (*SI Methods*), including homogeneous trapping protocol and known trap efficiency (4). Under other circumstances (trap efficiency unknown, different trap models, or trapping protocols), trapping may generate false absence data. Using only presence data to assess habitat suitability has the advantage that data derived from different sources (e.g., compilation of published data) can be combined to inform control projects.

Moreover, Maxent predictions were important for the eradication program as all suitable habitat needed to be included in the monitoring and the target area, even if they were not infested at the time of sampling due to possible movement among the patches. In the Niayes area, *G. p. gambiensis* are never present in all suitable patches at the same time. Instead, they form a metapopulation with patches connected through dispersal (18).

Use of Model Predictions for Optimization of the Eradication Project.

To generate a binary presence-absence map to delimitate the infested area, we chose a threshold providing a high sensitivity (96%). Indeed, in the case of an eradication project, it is paramount to reduce false negatives as much as possible, to avoid leaving tsetse-infested pockets not subjected to the control effort. Such areas can act as a source of reinvasion into previously cleared areas. During the entomological baseline data survey, the target area was divided into operational 5 × 5 km grid cells where at least one tsetse had been captured, plus a buffer zone

Fig. 4. Optimization of the integrated control strategy using model predictions. The Maxent model was used with a threshold of 0.13 to predict the suitable habitats for *G. p. gambiensis* (sensitivity of 0.96 and specificity of 0.57). In block 1, the suitable habitats allowed delimitating two polygons for aerial releases (RL1 and RL2) where the minimum numbers of sterile males released per km² were 24 and 11, respectively, based on Maxent predictions. Adult chilled tsetse flies are released with a Mubarqui smart release machine on board a gyrocopter of the Kalahari aerodrome (Upper Left) (20). In block 1, the green and gray lines represent the track flying records of the gyrocopter on April 14, 2014 in RL1 and April 11, 2014 in RL2, respectively. In block 2, 1,347 insecticide-impregnated traps were set from December 2012 to February 2013 in the predicted suitable sites (blue lozenges) to suppress tsetse populations.

removed as soon as a tsetse fly was caught (minimum 1 d). In the absence of capture, the traps were retrieved after a maximum of 42 d.

To homogenize data quality and avoid pseudoreplication (data points in the same pixel), the dataset was simplified in two ways. First, the study area was rasterized into square pixels with a 250-m resolution to match the model predictors (*SI Methods*).

For the presence data, all trap positions located within the same pixel were aggregated and concentrated in the pixel center. Presence data were observed in 68 pixels from 91 presence points (*Fig. S5*).

For the absence data, all trap positions located within a buffer of 500 m around a presence point were removed (to account for tsetse dispersal capacity). Furthermore, we removed absence pixels with $P > 0.01$ that the flies were present despite the absence of trapping using the model described (24) (*SI Methods*). From the initial 592 absence data, 333 were finally retained (*Fig. S5*), 56 of which were used in the validation dataset. Therefore, the training dataset was composed of 68 presence and 269 absence data.

Validation datasets. Trap data were collected independent of the training dataset from April 2009 to February 2013 during different surveys (tsetse dispersal and competitiveness studies, longitudinal monitoring of the demographic structure of tsetse populations, etc.). This dataset included 92 presence and 64 absence data. It was processed as described above, resulting in a dataset of 64 presence and 1 absence data (trapping times were not long enough to ascertain tsetse absence). A subset of 56 absence data was extracted from the 333 absence data described above.

A second validation dataset was created using 182 aerial photos taken from a gyrocopter at an altitude ranging from 100 to 300 m. The environment identified from these pictures was subsequently categorized as suitable or unsuitable for tsetse habitat (4). Picture coordinates were corrected using Google Earth to take the angle and deviation from the ground into account. Overall, 23 suitable and 159 unsuitable habitats were identified, thereafter called “expert-based habitats” (*Fig. S6*).

Predictors Used for Tsetse Suitable Habitat. Climatic and environmental data were derived from a time series of MODIS (version V005). Composite day and night land surface temperature (MOD11A2; 8-d averages), middle infrared (MOD13Q1; 16-d averages), and normalized difference vegetation index (MOD13Q1; 16-d averages) were selected for the analysis (25). Spatial resolution of pixels was 250 × 250 m (*SI Methods*).

A supervised classification of the vegetation was achieved using Landsat 5 Thematic Mapper satellite images with a spatial resolution of 30 × 30 m (*SI Methods*). Four cloud-free and haze-free satellite images were used, from

October 2009, April 2010, June 2010, and December 2010, to take into account the seasonal dynamics of these habitats, named “forests” in the manuscript.

Models. Ecological niche factor analysis was used to characterize the habitat of *G. p. gambiensis* in the target area (*SI Methods*).

Regularized logistic model was used to predict tsetse presence probability. Maxent analysis was used to predict habitat suitability. Approximately 10,000 pixels were sampled from the environmental variables (background) to calibrate the model. Because of the small sample size for presence data (56 points), only linear and quadratic transformations of environmental variables were used.

Model performance and comparison were assessed using the two validation sets (26). Model performance metrics were computed for each predictive model (27). The metrics used were the area under the receiver operating curve, called the area under the curve (28), the percentage of consonants correct (PCC), the specificity, and the sensibility (*SI Methods*).

Regularized gain is an additional metric for Maxent models. Marginal response curves were also computed to assess the relationship between the predicted suitability index and a given environmental datum. These curves were obtained by varying this variable while keeping all others at their average value.

Predicted values from regularized logistic regression and Maxent models were compared using Spearman’s rank correlation coefficient.

Impact of Control Operations. We used a generalized linear mixed model (29) to measure the impact of the suppression on tsetse apparent densities. The response data were tsetse counts in the traps. Time (measured in weeks), treatment (suppression or not), and the block (1 or 2) and their interactions were used as fixed effects, whereas the trap locations were used as random effects (*SI Methods*). Raw data are presented in *Dataset S1*.

The probability that eradication was effective in block 1 was estimated using the same model used to clean the absence dataset (24) (*SI Methods*), considering that at least a couple of flies were necessary to maintain the population.

ACKNOWLEDGMENTS. This work was funded by the US State Department through the Peaceful Uses Initiative, the Joint Food and Agriculture Organization of the United Nations/International Atomic Energy Agency Division of Nuclear Techniques in Food and Agriculture, the Department of Technical Cooperation, the Directorate of Veterinary Services of Sénégal, Institut Sénégalais de Recherches Agricoles, and Centre de Coopération Internationale en Recherche Agronomique pour le Développement.

1. Simarro PP, Jannin J, Cattand P (2008) Eliminating human African trypanosomiasis: Where do we stand and what comes next? *PLoS Med* 5(2):e55.
2. Itard J, Cuisance D, Tacher G (2003) Trypanosomoses: Historique - répartition géographique. *Principales Maladies Infectieuses et Parasitaires du Bétail. Europe et Régions Chaudes*, eds Lefèvre P-C, Blancou J, Chermette R (Lavoisier, Paris), Vol 2, pp 1607–1615.
3. Kabayo JP (2002) Aiming to eliminate tsetse from Africa. *Trends Parasitol* 18(11): 473–475.
4. Bouyer J, et al. (2010) Stratified entomological sampling in preparation for an area-wide integrated pest management program: The example of *Glossina palpalis gambiensis* (Diptera: Glossinidae) in the Niayes of Senegal. *J Med Entomol* 47(4):543–552.
5. Seck MT, Bouyer J, Sall B, Bengaly Z, Vreysen MJB (2010) The prevalence of African animal trypanosomoses and tsetse presence in western Senegal. *Parasite* 17(3): 257–265.
6. Solano P, et al. (2010) Population genetics as a tool to select tsetse control strategies: Suppression or eradication of *Glossina palpalis gambiensis* in the Niayes of Senegal. *PLoS Negl Trop Dis* 4(5):e692.
7. Vreysen MJB (2001) Principles of area-wide integrated tsetse fly control using the sterile insect technique. *Med Trop (Mars)* 61(4-5):397–411.
8. Kagbadouno MS, et al. (2011) Progress towards the eradication of tsetse from the Loos islands, Guinea. *Parasit Vectors* 4:18.
9. Guisan A, Thuiller W (2005) Predicting species distribution: Offering more than simple habitat models. *Ecol Lett* 8(9):993–1009.
10. Brotons L, Thuiller W, Araújo MB, Hirzel AH (2004) Presence-absence versus presence-only modelling methods for predicting bird habitat suitability. *Ecography* 27(4): 437–448.
11. Zou H, Hastie T (2005) Regularization and variable selection via the elastic net. *J R Stat Soc Series B Stat Methodol* 67(2):301–320.
12. Friedman J, Hastie T, Tibshirani R (2010) Regularization paths for generalized linear models via coordinate descent. *J Stat Softw* 33(1):1–22.
13. Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. *Ecol Modell* 190(3-4):231–259.
14. Fithian W, Hastie T (2012) Statistical models for presence-only data: Finite-sample equivalence and addressing observer bias. *arXiv* 1207.6950.
15. Hirzel AH, Haussler J, Chessel D, Perrin N (2002) Ecological-niche factor analysis: How to compute habitat-suitability maps without absence data? *Ecology* 83(7):2027–2036.
16. Hutchinson G (1957) Concluding remarks. *Cold Spring Harb Symp Quant Biol* 22: 415–427.
17. Gormley AM, et al. (2011) Using presence-only and presence-absence data to estimate the current and potential distributions of established invasive species. *J Appl Ecol* 48(1):25–34.
18. Peck SL (2012) Networks of habitat patches in tsetse fly control: Implications of metapopulation structure on assessing local extinction. *Ecol Modell* 246:99–102.
19. Bouyer J, Seck MT, Sall B (2013) Misleading guidance for decision making on tsetse eradication: Response to Shaw et al. (2013). *Prev Vet Med* 112(3-4):443–446.
20. Bouyer J, Lefrançois T (2014) Boosting the sterile insect technique to control mosquitoes. *Trends Parasitol* 30(6):271–273.
21. Mubarqui RL, et al. (2014) The smart aerial release machine, a universal system for applying the sterile insect technique. *PLoS ONE*, in press.
22. Vreysen MJB, et al. (2000) *Glossina austeni* (Diptera: Glossinidae) eradicated on the island of Unguja, Zanzibar, using the sterile insect technique. *J Econ Entomol* 93(1): 123–135.
23. Vreysen MJB, Seck MT, Sall B, Bouyer J (2013) Tsetse flies: Their biology and control using area-wide integrated pest management approaches. *J Invertebr Pathol* 112 (Suppl):S15–S25.
24. Barclay HJ, Hargrove JW (2005) Probability models to facilitate a declaration of pest-free status, with special reference to tsetse (Diptera: Glossinidae). *Bull Entomol Res* 95(1):1–11.
25. Rogers DJ, Hay SI, Packer MJ (1996) Predicting the distribution of tsetse flies in West Africa using temporal Fourier processed meteorological satellite data. *Ann Trop Med Parasitol* 90(3):225–241.
26. Elith J, et al. (2006) Novel methods improve prediction of species’ distributions from occurrence data. *Ecography* 29(2):129–151.
27. Liu C, Berry PM, Dawson TP, Pearson RG (2005) Selecting thresholds of occurrence in the prediction of species distributions. *Ecography* 28(3):385–393.
28. DeLong ER, DeLong DM, Clarke-Pearson DL (1988) Comparing the areas under two or more correlated receiver operating characteristic curves: A nonparametric approach. *Biometrics* 44(3):837–845.
29. Laird NM, Ware JH (1982) Random-effects models for longitudinal data. *Biometrics* 38(4):963–974.