Published in final edited form as:

Org Biomol Chem. 2011 June 7; 9(11): 4070–4078. doi:10.1039/c0ob00938e.

Novel anti-Prelog stereospecific carbonyl reductases from *Candida parapsilosis* for asymmetric reduction of prochiral ketones

Yao Nie^{a,b}, Rong Xiao^a, Yan Xu^{a,c,*}, and Gaetano T. Montelione^{a,*}

^aCenter for Advanced Biotechnology and Medicine, Department of Molecular Biology and Biochemistry, Rutgers University, Piscataway, New Jersey 08854, USA

^bSchool of Biotechnology and Key laboratory of Industrial Biotechnology, Ministry of Education, Jiangnan University, Wuxi, Jiangsu 214122, China

^cState Key Laboratory of Food Science and Technology, Jiangnan University, Wuxi, Jiangsu 214122, China

Abstract

Application of biocatalysis in the synthesis of chiral molecules is one of the greenest technologies for the replacement of chemical routes due to its environmentally benign reaction conditions and unparalleled chemo-, regio- and stereoselectivities. We have been interested in searching for carbonyl reductase enzymes and assessing their substrate specificity and stereoselectivity. We now report a gene cluster identified in *Candida parapsilosis* that consists of four open reading frames including three putative stereospecific carbonyl reductases (*scr1*, *scr2*, and *scr3*) and an alcohol dehydrogenase (*cpadh*). These newly identified three stereospecific carbonyl reductases (SCRs) showed high catalytic activities for producing (*S*)-1-phenyl-1,2-ethanediol from 2-hydroxyacetophenone with NADPH as the coenzyme. Together with CPADH, all four enzymes from this cluster are carbonyl reductases with novel anti-Prelog stereoselectivity. SCR1 and SCR3 exhibited distinct specificities to acetophenone derivatives and chloro-substituted 2-hydroxyacetophenones, and especially very high activities to ethyl 4-chloro-3-oxobutyrates, a β -ketoester with important pharmaceutical potentials. Our study also showed that genomic mining is a powerful tool for the discovery of new enzymes.

Introduction

The NAD(P)H-dependent carbonyl reductases catalyze reduction of a variety of endogenous and xenobiotic carbonyl compounds, including biologically and pharmacologically active substrates.¹ There are considerable interests in the use of carbonyl reductases in the

*Fax: +1-732-235-5633; Tel: +1-732-235-5321; gxy@cabm.rutgers.edu. * Fax: +86-510-85918201; Tel: +86-510-85864112; yxu@jiangnan.edu.cn.

† Electronic Supplementary Information (ESI) available: Experimental details including docking settings and analytic methods for chiral alcohol products, and 1H NMR spectrum of 1-phenyl-1,2-ethanediol. See DOI: 10.1039/b000000x/

‡ Footnotes should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.

pharmaceutical and fine chemicals industries for the production of chiral alcohols, important building blocks for synthesis of chirally pure pharmaceutical agents.² For the production of chiral auxiliaries from their corresponding prochiral ketones, carbonyl reductases has inherent advantages over chemocatalysts in terms of their highly chemo-, enantio-, and regioselectivities. These features make stereospecific carbonyl reductases very interesting from both scientific and industrial perspectives.³

Stereospecific carbonyl reductases are ubiquitous in nature and have been characterized from diverse sources including bacteria,⁴ yeasts,⁵ plants,⁶ and tissues from several mammalian species.⁷ However, despite the diversity of stereospecific oxidoreductases,⁸ their range of applications remains modest. This situation may be attributed to several perceived limitations including the stereospecificity and availability of the enzymes. In addition, researches on molecular mechanisms of oxidoreductases are still in its infancy. So far, most enzymes catalyzing asymmetric reductions generally follow the Prelog's rule in terms of stereochemical outcomes,⁹ while enzymes with anti-Prelog stereospecificity, such as *Pseudomonas* sp. ADH and *Lactobacillus kefir* ADH, are still limited among the carbonyl reductases classified to four stereochemical patterns concerning hydride transfer from reduced cofactor (NAD(P)H) to ketone.¹⁰ In particular, the precise mechanism of enzymatic anti-Prelog stereo-preference in asymmetric reduction is not yet fully understood.

Although most oxidoreductases possessing anti-Prelog selectivity have been identified in bacteria, such as *Geotrichum* sp.,¹¹ *Lactobacillus brevis*,¹² *Lactobacillus kefir*,¹³ and *Pseudomonas* sp.,¹⁴ *Candida* yeast species are attractive as sources of highly-stereospecific oxidoreductases.¹⁵ In a previous study, an NADPH-dependent alcohol dehydrogenase of anti-Prelog type has been discovered from *C. parapsilosis*, which catalyzes asymmetric reduction of 2-hydroxyacetophenone into (*S*)-1-phenyl-1,2-ethanediol (PED),^{16,17} a versatile chiral building block for the synthesis of pharmaceuticals, agrochemicals, and liquid crystals. PED is also a precursor for the production of chiral biphosphines and a chiral initiator for stereoselective polymerization.¹⁸

As a result of recent advances in genomics, proteomics, and bioinformatics, the area of biocatalysis is developed significantly, and the availability of genome sequences from large numbers of microorganisms allows scientists to discover novel enzymes with application potentials.¹⁹ The genome sequence of *C. parapsilosis* has been completed recently by the Wellcome Trust Sanger Institute Pathogen Genomics group (<http://www.sanger.ac.uk/sequencing/Candida/parapsilosis/>), which provided us an opportunity to look into the genome of *C. parapsilosis* in extensive details. We have identified three open reading frames (ORFs) in the 960-kb contig005802 of *C. parapsilosis* coding for putative stereospecific carbonyl reductase genes *scr1*, *scr2*, and *scr3*, respectively. These ORFs have been cloned and expressed, and the encoded proteins were purified to homogeneity and their functions were confirmed as stereospecific carbonyl reductases (SCR1, SCR2, and SCR3), with anti-Prelog selectivity that converts 2-hydroxyacetophenone to (*S*)-PED. These oxidoreductases have unique specificities that are useful for fine biochemical synthesis.

Results and discussion

Identification of putative stereospecific carbonyl reductases-encoding genes

Bioinformatic analysis based on sequence-similarity with CPADH, a known stereospecific alcohol dehydrogenase,¹⁷ in the *C. parapsilosis* genome revealed additional homologous ORFs, named here as *scr1*, *scr2*, and *scr3* coding for putative stereospecific carbonyl reductases (SCRs). As shown in Fig.1, these three ORFs, as well as the *cpadh* gene, locate in the 960-kb contig005802 of the *C. parapsilosis* genome. The *scr1*, *scr2*, and *scr3* genes comprise 846, 840, and 840 bp, encoding polypeptides of 281, 279, and 279 amino acid residues with the calculated molecular masses of 30,061, 29,993, and 30,097 Da, respectively, and thus there is not any intron found in these ORFs. Multiple sequence alignment of these four ORFs (Fig. 2) revealed high sequence identity between CPADH and SCR1 (68%), SCR2 (88%), and SCR3 (84%). From the amino acid sequence and secondary structure prediction, the three putative enzymes exhibit a classic α/β Rossmann-fold structure, the cofactor-binding motif, Gly⁴³-X-X-X-Gly⁴⁷-X-Gly⁴⁹ in SCR1 and Gly⁴¹-X-X-X-Gly⁴⁵-X-Gly⁴⁷ in SCR2 and SCR3, and the catalytic triad, Ser¹⁷⁴-Tyr¹⁸⁹-Lys¹⁹³ in SCR1 and Ser¹⁷²-Tyr¹⁸⁷-Lys¹⁹¹ in SCR2 and SCR3.²⁰

To gain insights into the proposed mechanism of cofactor binding, structure models of SCRs were obtained by homology modeling based on the X-ray structure of CPADH (PDB code: 3ctm). Because of the high degree of sequence identity, it can be expected that these four proteins including CPADH and SCRs are very similar in overall conformation, except for some loop regions due to amino-acid sequence difference. Thus, the cofactor-binding domain is expected to fold into the classic Rossmann-fold structure, a seven-strand parallel β -sheet flanked on both sides by α -helices.²¹ Based on the highly conserved cofactor-binding domain, putative enzyme-cofactor docking was performed with NADPH and NADH, respectively (Fig. 3). In both lowest-energy conformational ensembles, structure comparison with different cofactors suggested that the steric conformation in the cofactor-binding domain is more favorable to NADPH. Additionally the positive charged residue His (His⁷⁰ in SCR1, His⁶⁸ in SCR2 and SCR3) is expected to play a role in the affinity of the enzyme on NADPH, rather than NADH, by forming salt-bridge with the phosphate moiety at the 2'-position of AMP.²²

According to catalytic properties and primary structure information, stereospecific oxidoreductases including alcohol dehydrogenases and carbonyl reductases are mainly classified into three different groups, the zinc-dependent alcohol dehydrogenase, the short-chain dehydrogenase/reductase (SDR), and the aldo-keto reductase (AKR).⁸ All four proteins share sequence motifs characteristic of the SDR superfamily, including the cofactor-binding motif Gly-x-x-x-Gly-x-Gly (x denotes any amino acid) and the catalytic triad of Ser-Tyr-Lys, and further the extended tetrad of Asn-Ser-Tyr-Lys observed in the majority of SDRs.²³ In addition, the SCRs also have the conserved sequence motifs of secondary structure elements and key positions for assignment of coenzyme specificity of the cP2 subfamily in classic SDRs, except that the conserved basic residue Lys/Arg responsible for binding phosphate group in NADPH is replaced by weak basic residue His,²⁴ indicating that the cofactor dependence of the putative carbonyl reductases might be not

rigidly restricted to NADPH and these enzymes still show somewhat low activity with NADH. These highly-conserved characteristic sequence motifs suggested that the SCRs belong to the cP2 subfamily of the classical SDR superfamily, one of the three NADPH-dependent subfamilies.²⁴

Cloning, expression, and purification of SCRs

The sequence and structure information involving the overall protein structure analysis and the conserved SDR structural characteristics revealed that the identified genes and *cpadh* show the absence of intron and the encoded proteins are all not glycoproteins, and then these genes were attempted to be expressed in *Escherichia coli* system. From the nucleotide sequence of ORF, the *scr1*, *scr2*, and *scr3* were amplified by PCR from genomic DNA of *C. parapsilosis* CCTCC M203011, and the PCR products were inserted into pET21c vector by ligation-independent cloning to construct the recombinant plasmids. These three plasmids, pET21-SCR1, pET21-SCR2, and pET21-SCR3, were then transformed into expression host *E. coli* BL21(DE3) pMgK cells, and recombinant SCR1, SCR2, and SCR3 were produced in *E. coli* as fusion proteins containing a C-terminal His₆ tag. All three recombinant enzymes were expressed at very high level. Of them, SCR1 and SCR3 were expressed as soluble form at yields of 50 mg L⁻¹ broth and 46 mg L⁻¹ broth, respectively, while SCR2 has relatively low solubility with a yield of 5 mg L⁻¹ broth (Fig. 4).

The three recombinant enzymes were purified to homogeneity as judged by Coomassie Brilliant Blue staining of SDS-PAGE (Fig. 5) by Ni affinity purification followed by gel filtration chromatography. The relative molecular mass of the SCR1 and SCR3 were estimated to be 124.6 kDa and 123.4 kDa by analytic gel filtration and static light scattering using the same low salt buffer,^{25,26} but SCR2 was detected as aggregated form. Since the relative molecular mass of the monomer of the recombinant enzymes should be around 30 kDa based on their amino acid composition, these results suggested that both SCR1 and SCR3 have tetrameric structures.

Catalytic properties of recombinant SCRs

Because SCR1, SCR2, and SCR3 showed high homology to CPADH, which exhibits catalytic activity to 2-hydroxyacetophenone,¹⁷ the enzymatic activities of these three SCRs were investigated for reduction of 2-hydroxyacetophenone. Under the assay conditions, SCR1 gave the highest specific activity of 5.16 U mg⁻¹, and SCR3 had the catalytic activity of 4.23 U mg⁻¹, while SCR2 has a lower specific activity of 1.55 U mg⁻¹. In addition, corresponding to the results from cofactor-binding model and secondary structure prediction and analysis, the SCRs all displayed catalytic activity with NADPH as the coenzyme, but very low activities with NADH, indicating that these three enzymes, like CPADH,¹⁷ are all NADPH-dependent in oxidoreductases.

Since environmental pH value can have an influence on the stereochemistry of enzymatic reactions,²⁷ the effect of the reaction pH on the activities of SCRs catalyzing 2-hydroxyacetophenone reduction were also investigated. All three enzymes exhibited the highest activity at the pH ranging from 5.0 to 6.0 (Fig. 6). Subsequently, the enzymes were evaluated under their individual optimal pH, and apparent kinetic parameters were further

measured by double reciprocal Lineweaver-Burk plots at various 2-hydroxyacetophenone concentrations with fixed NADPH concentrations. As shown in Table 1, for reduction of 2-hydroxyacetophenone, these three enzymes exhibited different kinetic parameters.

Stereoselectivity to prochiral carbonyl group

Using 2-hydroxyacetophenone as the substrate, optically pure 1-phenyl-1,2-ethanediol (PED) of (*S*)-enantiomer (>99% e.e.) was produced by each of SCR1, SCR2, and SCR3, respectively (Fig. 7). These three enzymes are capable of catalyzing asymmetric reduction of prochiral carbonyl compounds and are all (*S*)-specific carbonyl reductase toward 2-hydroxyacetophenone. Of them, however, SCR2 is not as efficient as the other two enzymes, corresponding to its lower activity. Like the homologous anti-Prelog specific alcohol dehydrogenase, CPADH, these data demonstrate that the paralogous SCR1, SCR2, and SCR3 enzymes are anti-Prelog-type stereospecific carbonyl reductases.^{17,28}

Oxidoreductases perform a wide variety for asymmetric reductions, differing in stereospecificity and substrate specificity, and have been used for producing optically active alcohols from various prochiral ketones, ketoacids, and ketoesters. The SCRs catalyze (*S*)-specific reduction of 2-hydroxyacetophenone, an anti-Prelog type reaction.²⁸ Therefore, these new enzymes complement the stereospecific oxidoreductases described to date for catalysis of the reduction of prochiral carbonyl compounds to the corresponding optically pure alcohols with anti-Prelog stereopreference.

Otherwise, the finding of stereospecific carbonyl reductases capable of catalyzing (*S*)-specific reduction of 2-hydroxyacetophenone from the same host would give us a profound knowledge on the reaction mechanism of *C. parapsilosis* whole-cell mediated stereoinversion, which involves two steps, the oxidation step of (*R*)-PED to the intermediate (2-hydroxyacetophenone) and the reduction step of the intermediate to (*S*)-PED.¹⁶ *C. parapsilosis* whole cells catalyzing deracemization of racemic PED to (*S*)-enantiomer would be updated as a one-pot concurrent tandem process involving at least four stereoselective alcohol dehydrogenases and carbonyl reductases, but not the two-step redox sequentially catalyzed by an alcohol oxidase and a carbonyl reductase (Scheme 1).^{17,29} In addition, the contribution of newly discovered SCRs to the whole-cell mediated PED stereoinversion can be further proved by comparing the kinetic parameters such as the maximum initial reaction rates and Michaelis constants between the two steps involved in the one-pot redox reaction, in which the maximum initial reaction rate of reduction is almost 10 times higher than the oxidation reaction.³⁰ Thus, the reduction step catalyzed by CPADH and SCRs would drive the whole reaction process to the product formation and the oxidation step is the restrictive step in the sequential two-step reaction.

Substrate specificity

Since the three new enzymes showed distinct (*S*)-specific carbonyl-reducing activity, we then further examined substrate specificity of these enzymes to various carbonyl compounds including aryl ketones, aliphatic ketones, α - and β -ketoesters. As shown in Table 2, on the one hand, the enzymes all exhibited higher catalytic activity to ketoesters than to alkyl and aromatic ketones, while compared with the other two enzymes, SCR2 generally showed

lower activities to keto substrates, corresponding to the observation in activity and kinetics evaluation of these enzymes catalyzing 2-hydroxyacetophenone reduction; on the other hand, the enzymes exhibited diversity on specificity to aryl ketones and ketoesters, respectively. For both substituted acetophenone and 2-hydroxyacetophenone derivatives, bearing chloro or methyl at various positions of the phenyl ring, the ortho substituents were poor substrates for these enzymes, indicating that the substitution at ortho position might have steric effect on the hydrogen attack from electron donor NADPH to the carbonyl group and therefore significantly influence the reactivity of the enzymes. However, SCR3 was more specific to *p*-Cl-2-hydroxyacetophenone, while SCR1 had higher activity to *m*-Cl-2-hydroxyacetophenone. Of the aryl ketones, despite its low activities to the substrates, SCR2 was more active to acetophenone and its derivatives than to 2-hydroxyacetophenone, which would be a trend observed different from SCR1 and SCR3. For ketoesters, the enzymes exhibited high activity to those with small groups, but compared with SCR3, SCR1 was more active to bulky ketoesters with phenyl ring and generally showed higher activities to β -ketoesters. For stereoselective reduction of carbonyl compounds with different chemical structures, these three enzymes followed the anti-Prelog rule, as their behaviors in catalyzing asymmetric reduction of 2-hydroxyacetophenone, although chiral alcohol enantiomers were produced in different absolute configurations, affording aryl diols, phenylhydroxyl esters, ethyl 4-trifluoro-3-hydroxybutyrate, and ethyl 4-chloro-3-hydroxybutyrate in (*S*)-configuration, whereas (*R*)-antipode for others due to the switch in CIP priority.

Among the tested β -ketoesters, additionally, SCR1 performed the highest activity to ethyl 4-chloro-3-oxobutyrate, an important chiral building block. The enzyme-cofactor-substrate docking model, as shown in Fig. 8, revealed that, in the lowest-energy conformational ensemble, the substrate-binding domain is favorable to this kind of substrates, and the carbonyl carbon is in a position proximal to the C4 atom of the nicotinamide ring of NADPH. These results suggested that, for enzymatic reactions, the enzyme structure generally has a significant influence on the enzyme activity and affinity to substrates. The steric conformation of active site in catalytic domain determines the substrate recognition, binding, and orientation in enzyme.³¹

It is worthy to note that, as shown above, SCR1 catalyzes the reduction of a broad spectrum of ketones including aryl, aliphatic ketones, α - and β -ketoesters, and shows a particular substrate specificity towards ethyl 4-chloro-3-oxobutyrate, a precursor for the synthesis of ethyl 4-chloro-3-hydroxybutyrate which is an important pharmaceutical intermediate.³² Therefore, the new discovered stereospecific carbonyl reductases would be useful enzymes with application potentials.

Experimental

Materials

C. parapsilosis strain CCTCC M203011 was obtained from the China Center for Type Culture Collection (CCTCC, Wuhan, China). *E. coli* XL-10 gold cells were used for gene cloning and plasmid preparation, and *E. coli* BL21 (DE3) pMgK competent cells, a rare codon-enhanced strain, were used for gene expression. High-fidelity PCR kit including DNA polymerase was purchased from FINNZYMES (Finland). The plasmid pET21c was

obtained from Novagen (USA). (*R*)- and (*S*)-1-phenyl-1,2-ethanediol, all the aliphatic ketones and ketone esters, aryl ketones including acetophenone and its derivatives, 2-hydroxyacetophenone, propiophenone, butyrophenone, valerophenone, hexanophenone, and coenzymes including NAD(P)H and NAD(P)⁺ were purchased from Sigma-Aldrich (USA). All other 2-hydroxyacetophenone derivatives including *o*-Cl-2-hydroxyacetophenone, *m*-Cl-2-hydroxyacetophenone, *p*-Cl-2-hydroxyacetophenone, and *p*-CH₃O-2-hydroxyacetophenone were prepared using the method described by Itsuno.³³ All other chemicals used in this work were of analytical grade and commercially available.

Cloning and expression of genes encoding stereospecific reductases

The genes encoding SCR1, SCR2, and SCR3 were amplified by polymerase chain reaction from *C. parapsilosis* genomic DNA. PCR-amplified DNA products were purified by QIAquick PCR Purification Kit (QIAGEN, USA) and inserted into pET21c expression vector (Novagen, USA) by ligation-independent cloning (LIC) using In-Fusion PCR Cloning Kit (Clontech, USA) for construction of recombinant plasmids. The infusion reaction mixtures were used to transform *E. coli* XL-10 gold cells. The plasmids isolated from these transformants were verified by DNA sequence analysis using BigDye Terminator cycle sequencing kit and an ABI PRISM 310 Genetic Analyzer (Applied Biosystems, USA). The plasmids with the correct inserts, pET21-SCR1, pET21-SCR2, and pET21-SCR3, were transformed into *E. coli* BL21(DE3) pMgK competent cells for the production of SCRs. These plasmids provide SCRs with a six-His tag fused at the C-terminus.

E. coli BL21 (DE3) pMgK transformants were cultivated at 37 °C in Luria broth (LB) medium in the presence of ampicillin (100 µg mL⁻¹) and kanamycin (50 µg mL⁻¹). When the optical density of the culture at 600 nm reached 0.6, the temperature was changed to 17 °C and isopropyl-β-D-thiogalactopyranoside (IPTG) was added to the culture to give a final concentration of 1 mM for induction of gene expression. After an additional incubation of 20 h at 17 °C, cells harvested by centrifugation were disrupted by sonication, and expressions of the recombinant proteins were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). For expression of SCR2, the expression conditions involving lower concentration of IPTG (0.1 mM) and shorter induction time (12 hours) were adopted, besides the above standard expression conditions.

Purification of recombinant enzymes

The cells were suspended in binding buffer (20 mM Tris-HCl, pH7.5, 0.3 M NaCl, 40 mM imidazole, 1 × protease inhibitors, 1 mM Tris (2-carboxyethyl) phosphine (TCEP)) and disrupted on ice by sonication. The supernatant of the cell lysate was collected by centrifugation at 26,000 × *g* for 40 min at 4 °C and purified by an AKTApurify system using HisTrap HP affinity column followed by Superdex 75 gel filtration column (GE Healthcare, USA), and the purified fractions were exchanged into low salt buffer (10 mM Tris-HCl, pH 7.5, 0.1 M NaCl, 0.02% NaN₃, 5 mM D,L-dithiothreitol).^{25,26} The final recombinant enzymes were purified with an apparent homogeneity on sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) with 12% polyacrylamide gels. Their molecular masses were measured by matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry (Applied Biosystems, USA), and their

oligomerization states were determined by analytic gel filtration using Agilent 1200 series HPLC system followed by static light scattering (Wyatt Technology, USA). These final preparations of purified SCRs were used in all of the experiments in this study.

Enzyme assays

Carbonyl reductase activity was measured by a continuous spectrophotometric assay using 2-hydroxyacetophenone as a substrate. One unit of enzyme activity was defined as the amount of enzyme catalyzing the oxidation of 1 μmol NAD(P)H per min under the assay conditions. The standard assay mixture for the enzyme activity comprised of 0.1 M potassium phosphate buffer (pH 6.5), 0.3 mM NAD(P)H, 0.7 mM 2-hydroxyacetophenone and appropriate enzyme in a total volume of 100 μL . The decrease in the amount of the coenzyme was measured spectrophotometrically at 340 nm extinction coefficient $[\epsilon]=6.22 \text{ mM}^{-1} \text{ cm}^{-1}$. Protein concentration was determined using Bradford reagents (Bio-Rad) with bovine serum albumin as a standard. The pH dependence of enzyme activity was determined over a pH range of 4.0 to 8.5 using the following buffers: 0.1 M acetate pH 4.0 to 6.0), 0.1 M sodium phosphate (pH 6.0 to 8.0) and 0.1 M Tris-HCl (pH 8.0 to 8.5).

The substrate specificity of SCRs was investigated under the same conditions as described above. Various carbonyl compounds including aryl ketones, aliphatic ketones, α - and β -ketoesters were used as the substrates with the cofactor of NADPH.

Asymmetric reduction and stereoselectivity assay

Asymmetric reduction of 2-hydroxyacetophenone by the purified enzymes were carried out at 30°C for 6 h with shaking in a reaction mixture comprising 0.1 M potassium phosphate buffer (pH 6.5), 1 g L⁻¹ 2-hydroxyacetophenone, NADPH (7 mM) and 0.5 mg of the purified enzyme in a total volume of 0.5 ml. The reaction products were extracted with ethyl acetate and the organic layer was used for analysis. The optical purity of reaction products were analyzed by HPLC using a Chiralcel OB-H column (4.6 $\times$ 250 mm, Daicel Chemical Ind., Ltd., Japan). Enantiomers were eluted with hexane and 2-propanol (9:1) at a flow rate of 0.5 mL min⁻¹. The effluent was monitored at 215 nm, and the areas under each peak were integrated.¹⁶ To further confirm the product of 1-phenyl-1,2-ethanediol but not byproducts of the reduction, the reaction product was analyzed by ¹H NMR (for assay details see the Supplementary Information).

Asymmetric reduction of various carbonyl compounds, including aryl ketones, aliphatic ketones, α - and β -ketoesters, by the enzymes were carried out at 30°C for 6 h with shaking in a reaction mixture comprising 0.1 M potassium phosphate buffer (pH 6.5), NADPH (7 mM) and 0.5 mg of the purified enzyme in a total volume of 0.5 ml. The reaction products were extracted with ethyl acetate or hexane and the organic layer was used for analysis. The reaction products were analyzed by chiral HPLC (HP 1100, Agilent, USA) equipped with Chiralcel OB-H column (4.6 mm $\times$ 250 mm; Daicel Chemical Ind. Ltd., Japan) or chiral GC (7890A, Agilent, USA) equipped with FID detector and Chrompack Chirasil-Dex CB chiral capillary column (25 m $\times$ 0.25 mm; Varian, USA) (for assay details see the Supplementary Information).

Homology modeling, alignment, and flexible docking

The crystal structure of CPADH (PDB code: 3ctm) was used. The homology modeling of SCRs was carried out by HOMA (Homology Modeling Automatically) (<http://www-nmr.cabm.rutgers.edu/HOMA/>). The structure alignment of CPADH and SCRs was performed with the Deep View Swiss-Pdb Viewer 4.0 program. All docking calculations were accomplished with AutoDock Vina 1.0.³⁴ A docking algorithm that took account of ligand flexibility but kept the protein rigid was employed. Docking runs were carried out using the standard parameters of the program for interactive growing and subsequent scoring (for settings of grid box dimensions and centers see the Supplementary Information).

Nucleotide sequence accession number

The nucleotide sequence for the stereospecific carbonyl reductase genes *scr1*, *scr2*, and *scr3* have been deposited in the GenBank database under accession numbers FJ939565, FJ939563, and FJ939564, respectively.

Conclusion

In this work, we describe a new gene cluster of enantioselective oxidoreductases with unusual stereospecificity in *C. parapsilosis*. We confirmed that these genes encode three unique stereospecific carbonyl reductases through cloning, expression, purification, and characterization of the corresponding gene products. We verified enantiomer configuration of the enzymatic products from asymmetric reduction of prochiral carbonyl groups using multiple substrates. SCR1, SCR2, and SCR3, like CPADH, all exhibit a novel anti-Prelog stereospecificity in reducing prochiral carbonyl groups such as forming (*S*)-1-phenyl-1,2-ethanediol from the corresponding ketone substrate, 2-hydroxyacetophenone. The enzymes are, however, distinct in their catalytic properties, including their pH dependency and substrate specificity spectrum. The genes encoding SCR2, SCR3, CPADH, and SCR1 are sequential in the cluster (Fig. 1) with intervals of several hundred base pairs. Further studies are necessary to understand the metabolic roles of these homologues and potential synergies among them.

The discovery of novel stereospecific carbonyl reductases of anti-Prelog selectivity further demonstrates the diversity of stereospecific oxidoreductases in microorganisms. Such enzymes provide a basis for elucidating the molecular mechanism of enzyme-mediated asymmetric reaction involving stereo-recognition between protein and chiral molecule, and the mechanism of electron transfer between functional groups of chiral molecule and key amino acid residues in the enzyme. Apart from their unique value in study on mechanism of stereospecific oxidoreduction reaction, these novel carbonyl reductases with anti-Prelog stereopreference discovered by genomic mining have multiple potentials to generate chiral alcohols useful as intermediates in fine chemical synthesis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the Wellcome Trust Sanger Institute Pathogen Genomics group for *Candida parapsilosis* Sequencing data base (<http://www.sanger.ac.uk/sequencing/Candida/parapsilosis/>). We thank Dr. R. Zhang for useful discussions and comments on the manuscript.

This work was supported by the National Institutes of General Medical Science Protein Structure Initiative program, grant U54 GM074958 (to G.T.M), the National Natural Science Foundation of China (NSFC) (No. 30800017), the National Key Basic Research and Development Program of China (973 Program) (No. 2009CB724706 and 2011CB710802), and the Program of Introducing Talents of Discipline to Universities (111 Project) (111-2-06).

Notes and references

- Forrest GL, Gonzalez B. Chem. Biol. Interact. 2000; 129:21–40. [PubMed: 11154733]
- Panke S, Held M, Wubbolts M. Curr. Opin. Biotechnol. 2004; 15:272–279. [PubMed: 15357999]
- Schmid A, Dordick JS, Hauer B, Kiener A, Wubbolts M, Witholt B. Nature. 2001; 409:258–268. [PubMed: 11196655]
- Schoemaker HE, Mink D, Wubbolts MG. Science. 2003; 299:1694–1697. [PubMed: 12637735]
- Kroutil W, Mang H, Edegger K, Faber K. Curr. Opin. Chem. Biol. 2004; 8:120–126. [PubMed: 15062771]
- Itoh N, Mizuguchi N, Mabuchi M. J. Mol. Catal. B Enzym. 1999; 6:41–50.
- Katz M, Hahn-Hagerdal B, Gorwa-Grauslund MF. Enzyme Microb. Technol. 2003; 33:163–172.
- Müller M, Katzberg M, Bertau M, Hummel W. Org. Biomol. Chem. 2010; 8:1540–1550. [PubMed: 20237665]
- Zhu D, Yang Y, Buynak JD, Hua L. Org. Biomol. Chem. 2006; 4:2690–2695. [PubMed: 16826293]
- Melis LED, Whiteman PH, Stevenson TW. Plant Sci. 1999; 143:173–182.
- Wsol V, Skalova L, Szotakova B, Trejtnar F, Kvasnickova E. Chirality. 1999; 11:505–509. [PubMed: 10368924]
- Kamitori S, Iguchi A, Ohtaki A, Yamada M, Kita K. J. Mol. Biol. 2005; 352:551–558. [PubMed: 16095619]
- Reid MF, Fewson CA. Crit. Rev. Microbiol. 1994; 20:13–56. [PubMed: 8185833]
- Bradshaw CW, Fu H, Shen GJ, Wong CH. J. Org. Chem. 1992; 57:1526–1532.
- Ernst M, Kaup B, Müller M, Bringer-Meyer S, Sahm H. Appl. Microbiol. Biotechnol. 2005; 66:629–634. [PubMed: 15549291]
- Niefind K, Müller J, Riebel B, Hummel W, Schomburg D. J. Mol. Biol. 2003; 327:317–328. [PubMed: 12628239]
- Prelog V. Pure Appl. Chem. 1964; 9:119–130.
- Ankati H, Zhu D, Yang Y, Biehl ER, Hua L. J. Org. Chem. 2009; 74:1658–1662. [PubMed: 19161324]
- De Wildeman SMA, Sonke T, Schoemaker HE, May O. Acc. Chem. Res. 2007; 40:1260–1266. [PubMed: 17941701]
- Wei ZL, Lin GQ, Li ZY. Bioorg. Med. Chem. 2000; 8:1129–1137. [PubMed: 10882023]
- Fantin G, Fogagnolo M, Giovannini PP, Medici A, Pedrini P, Gardini F, Lanciotti R. Tetrahedron. 1996; 52:3547–3552.
- Bradshaw CW, Hummel W, Wong CH. J. Org. Chem. 1992; 57:1532–1536.
- Bradshaw CW, Fu H, Shen GJ, Wong CH. J. Org. Chem. 1992; 57:1526–1532.
- Engelking H, Pfaller R, Wich G, Weuster-Botz D. Tetrahedron: Asymmetry. 2004; 15:3591–3593.
- Gamenara D, de María PD. Biotechnol. Adv. 2009; 27:278–285. [PubMed: 19500548]
- Kamble AL, Soni P, Banerjee UC. J. Mol. Catal. B Enzym. 2005; 35:1–6.
- Kataoka M, Hoshino-Hasegawa A, Thiwhong R, Higuchi N, Ishige T, Shimizu S. Enzyme Microb. Technol. 2006; 38:944–951.
- Kataoka M, Delacruz-Hidalgo ARG, Akond MA, Sakuradani E, Kita K, Shimizu S. Appl. Microbiol. Biotechnol. 2004; 64:359–366. [PubMed: 14593510]
- Nie Y, Xu Y, Mu XQ, Tang Y, Jiang J, Sun ZH. Biotechnol. Lett. 2005; 27:23–26. [PubMed: 15685415]
- Pollard DJ, Telari K, Lane J, Humphrey G, McWilliams C, Nidositko S, Salmon P, Moore J. Biotechnol. Bioeng. 2006; 93:674–686. [PubMed: 16395718]
- Nie Y, Xu Y, Mu XQ. Org. Process Res. Dev. 2004; 8:246–251.
- Nie Y, Xu Y, Mu XQ, Wang HY, Yang M, Xiao R. Appl. Environ. Microbiol. 2007; 73:3759–3764. [PubMed: 17435004]

18. Iwasaki F, Maki T, Nakashima W, Onomura O, Matsumura Y. *Org. Lett.* 1999; 1:969–972. Liese A, Karutz M, Kamphuis J, Wandrey C, Kragl U. *Biotechnol. Bioeng.* 1996; 51:544–550. [PubMed: 18629818]
19. Kaluzna IA, Matsuda T, Sewell AK, Steward JD. *J. Am. Chem. Soc.* 2004; 126:12827–12832. [PubMed: 15469278] Raskin DM, Seshadri R, Pukatzki SU, Mekalanos JJ. *Cell.* 2006; 124:703–714. [PubMed: 16497582]
20. Oppermann U, Filling C, Hult M, Shafqat N, Wu X, Lindh M, Shafqat J, Nordling E, Kallberg Y, Persson B, Jörnvall H. *Chem. Biol. Interact.* 2003; 143–144:247–253.
21. Ghosh D, Weeks CM, Grochulski P, Duax WL, Erman M, Rimsay RL, Orr JC. *Proc. Natl. Acad. Sci.* 1991; 88:10064–10068. [PubMed: 1946424]
22. Zhang R, Zhu G, Zhang W, Cao S, Ou X, Li X, Bartlam M, Xu Y, Zhang XC, Rao Z. *Protein Sci.* 2008; 17:1412–1423. [PubMed: 18566346]
23. Filling C, Berndt KD, Benach J, Knapp S, Prozorovski T, Nordling E, Ladenstein R, Jörnvall H, Oppermann U. *J. Biol. Chem.* 2002; 277:25677–25684. [PubMed: 11976334]
24. Kallberg Y, Oppermann U, Jörnvall H, Persson B. *Eur. J. Biochem.* 2002; 269:4409–4417. [PubMed: 12230552]
25. Acton TB, Gunsalus K, Xiao R, Ma L, Aramini J, Baron MC, Chiang Y, Clement T, Cooper B, Denissova N, Douglas S, Everett JK, Palacios D, Paranjli RH, Shastry R, Wu M, Ho C-H, Shih L, Swapna GVT, Wilson M, Gerstein M, Inouye M, Hunt JF, Montelione GT. *Methods Enzymol.* 2005; 394:210–243. [PubMed: 15808222]
26. Xiao R, Anderson S, Aramini J, Belote R, Buchwald WA, Ciccocanti C, Conover K, Everett JK, Hamilton K, Huang YJ, Janjua H, Jiang M, Kornhaber GJ, Lee DY, Locke JY, Ma L, Maglaqui M, Mao L, Mitra S, Patel D, Rossi P, Sahdev S, Sharma S, Shastry R, Swapna GVT, Tong SN, Wang D, Wang H, Zhao L, Montelione GT, Acton TB. *J. Struct. Biol.* 2010; 172:21–33. [PubMed: 20688167]
27. Secundo F, Phillips RS. *Enzyme Microb. Technol.* 1996; 19:487–492.
28. Manzocchi A, Fiecchi A, Santaniello E. *J. Org. Chem.* 1988; 53:4405–4407. Prelog V. *Pure Appl. Chem.* 1964; 9:119–130.
29. Gruber CC, Lavandera I, Faber K, Kroutil W. *Adv. Synth. Catal.* 2006; 348:1789–1805. Voss CV, Gruber CC, Kroutil W. *Angew. Chem. Int. Ed.* 2008; 47:741–745. Voss CV, Gruber CC, Faber K, Knaus T, Macheroux P, Kroutil W. *J. Am. Chem. Soc.* 2008; 130:13969–13972. [PubMed: 18821754]
30. Mu XQ, Xu Y, Yang M, Sun ZH. *Process Biochem.* 2010
31. Zhu D, Mukherjee C, Rozzell JD, Kambourakis S, Hua L. *Tetrahedron.* 2006; 62:901–905.
32. Thayer AM. *Chem. Eng. News.* 2006; 84:26–27.
33. Itsuno S, Nakano M, Miyazaki K, Masuda H, Ito K, Hirao A, Nakahama S. *J. Chem. Soc. Perkin Trans. I.* 1985:2039–2044.
34. Trott O, Olson AJ. *J. Comput. Chem.* 2010; 31:455–461. [PubMed: 19499576]

contig005802 of *Candida parapsilosis* genome **Fig. 1.**

Map of contig005802 of *Candida parapsilosis* genome including the four open reading frames, *scr1*, *scr2*, *scr3*, and *cpadh*

Fig. 2.

Amino acid sequence alignment of CPADH (GenBank accession number DQ675534), SCR1 (GenBank accession number FJ939565), SCR2 (GenBank accession number FJ939563), and SCR3 (GenBank accession number FJ939564) from *C. parapsilosis*. Gaps in the aligned sequences are indicated by dashes. Identical amino acid residues are enclosed in boxes.

Fig. 3. Stereoview of the overall model structure of enzyme-cofactor docking with NADPH (a) and NADH (b), respectively, illustrated by the PyMOL program.

Fig. 4.

Analysis of the overexpression of SCR1, SCR2, and SCR3. The proteins were separated on a 12% SDS-polyacrylamide gel and stained with Coomassie Brilliant Blue G-250. Lane 1, total protein for SCR1; Lane 2, soluble fraction for SCR1; Lane 3, total protein for SCR2; Lane 4, soluble fraction for SCR2; Lane 5, total protein for SCR3; Lane 6, soluble fraction for SCR3; Lane 7, molecular mass standard.

Fig. 5. SDS-PAGE analysis of purified enzymes. The purified proteins were resolved by SDS-PAGE on a 12% polyacrylamide gel and stained with Coomassie Brilliant Blue G-250. Lane 1, molecular mass standard; Lane 2, purified SCR1; Lane 3, purified SCR2; Lane 4, purified SCR3.

Fig. 6. pH dependence of SCR1, SCR2, and SCR3 catalyzing 2-hydroxyacetophenone reduction. The enzyme activities of SCR1 (■), SCR2 (▲), and SCR3 (●) were measured in 0.1 M acetate buffer (pH 4.0 to 6.0) or 0.1 M sodium phosphate buffer (pH 6.0 to 8.0) or 0.1 M Tris-HCl buffer (pH 8.0 to 8.5) with 2-hydroxyacetophenone as the substrate and NADPH as the cofactor. Maximal enzyme activity observed was set as 100% relative activity for each enzyme.

Fig. 7. Asymmetric reduction of 2-hydroxyacetophenone (2-HAP) to 1-phenyl-1,2-ethanediol (PED) enantiomer by SCR1, SCR2, and SCR3, respectively. (a) Standard sample of (R)-PED. (b) Standard sample of (S)-PED. (c) SCR1 catalyzed asymmetric reduction of 2-HAP. (d) SCR2 catalyzed asymmetric reduction of 2-HAP. (e) SCR3 catalyzed asymmetric reduction of 2-HAP.

Fig. 8.
Stereoview of the model structure of SCR1-NADPH-ethyl 4-chloro-3-oxobuturate (COBE) docking, illustrated by the PyMOL program.

Scheme 1.

One-pot concurrent tandem stereoinversion of 1-phenyl-1,2-ethanediol by *Candida parapsilosis* involving C PADH and SCRs.

Table 1

Activities and kinetic parameters for reduction of 2-hydroxyacetophenone (2-HAP) by SCR1, SCR2, SCR3, and CPADH, using NADPH as cofactor

Enzyme	pH ^a	Specific activity /U mg ⁻¹	K _m /mM	V _{max} /μmol min ⁻¹ mg ⁻¹	K _{cat} /s ⁻¹
SCR1	5.0	5.16	9.83	42.0	21.04
SCR2	5.5	1.55	4.81	8.83	4.41
SCR3	6.0	4.23	4.68	32.3	16.20
CPADH	4.5	2.70	5.83	18.3	9.18

^a Activity assay was carried out at the optimum pH for each enzyme.

Table 2

Substrate specificity and stereoselectivity of SCR1, SCR2, and SCR3 to various carbonyl compounds

Substrate			Relative activity ^d /%			Product configuration
R1	R2		SCR1	SCR2	SCR3	
<div>Aryl ketones</div> <div> </div>						
1	H	CH ₃	1.27	3.36	2.57	<i>R</i>
2	H	CH ₂ OH	2.84	0.78	3.08	<i>S</i>
3	2'-Cl	CH ₂ OH	0.60	1.74	0.19	<i>S</i>
4	3'-Cl	CH ₂ OH	14.61	2.21	5.91	<i>S</i>
5	4'-Cl	CH ₂ OH	2.96	2.97	18.49	<i>S</i>
6	4'-OCH ₃	CH ₂ OH	-	-	-	-
7	H	CH ₂ CH ₃	6.31	0.32	2.08	<i>R</i>
8	H	(CH ₂) ₂ CH ₃	10.90	0.16	0.81	<i>R</i>
9	H	(CH ₂) ₃ CH ₃	1.96	-	0.93	<i>R</i>
10	H	(CH ₂) ₄ CH ₃	-	-	-	-
11	2'-CH ₃	CH ₃	1.65	0.42	0.12	<i>R</i>
12	3'-CH ₃	CH ₃	0.05	0.57	2.50	<i>R</i>
13	4'-CH ₃	CH ₃	1.63	0.51	7.29	<i>R</i>
14	2'-Cl	CH ₃	1.88	1.21	0.36	<i>R</i>
15	3'-Cl	CH ₃	0.88	1.68	2.33	<i>R</i>
16	4'-Br	CH ₃	2.02	3.87	6.54	<i>R</i>
17	4'-OCH ₃	CH ₃	-	-	-	-
<div>Aliphatic ketones</div> <div> </div>						

Substrate			Relative activity ^d /%			Product configuration
R1	R2		SCR1	SCR2	SCR3	
20	CH ₃	(CH ₂) ₃ CH ₃	0.86	0.83	0.40	<i>R</i>
21	CH ₃	(CH ₂) ₄ CH ₃	0.15	0.52	0.28	<i>R</i>
22	CH ₃	(CH ₂) ₅ CH ₃	1.39	-	0.52	<i>R</i>
23	CH ₂ CH ₃	CH(CH ₃) ₂	0.37	-	0.82	-
<div><div>α-ketoesters</div><div> </div></div>						
24	CH ₃	CH ₃	67.12	18.92	74.12	<i>R</i>
25	Phenyl	CH ₃	32.93	5.76	3.82	<i>S</i>
26	CH ₃	CH ₂ CH ₃	65.32	19.89	80.72	<i>R</i>
27	Phenyl	CH ₂ CH ₃	19.15	2.78	0.19	<i>S</i>
<div><div>β-ketoesters</div><div> </div></div>						
28	CF ₃	CH ₂ CH ₃	29.51	15.24	11.86	<i>S</i>
29	CH ₃	CH ₃	29.31	9.88	4.96	<i>R</i>
30	CH ₂ CH ₃	CH ₃	9.44	7.82	5.05	<i>R</i>
31	4-fluorophenyl	CH ₃	14.12	5.62	3.89	<i>S</i>
32	CH ₃	CH ₂ CH ₃	45.43	8.17	21.31	<i>R</i>
33	CH ₂ CH ₃	CH ₂ CH ₃	8.79	7.16	20.85	<i>R</i>
34	Chloromethyl	CH ₂ CH ₃	100.00	17.58	53.20	<i>S</i>
35	Phenyl	CH ₂ CH ₃	7.01	-	0.22	<i>S</i>
36	3,4-dimethoxyphenyl	CH ₂ CH ₃	4.44	-	0.47	<i>S</i>

^aThe enzyme activities of SCRs to various substrates and the configuration of produced alcohols were measured as described in the text. Maximal enzyme activity observed was set as 100% relative activity for the enzymes to various substrates.