Published in final edited form as:

Am J Prev Med. 2014 August ; 47(2): 188–197. doi:10.1016/j.amepre.2014.04.004.

Effects of the Communities That Care System on Cross-Sectional Profiles of Adolescent Substance Use and Delinquency

M. Lee Van Horn, PhD, Abigail A. Fagan, PhD, J. David Hawkins, PhD, and Sabrina Oesterle, PhD

Department of Psychology (Van Horn), University of South Carolina, Columbia, South Carolina; Department of Sociology, Criminology & Law (Fagan), University of Florida, Gainesville, Florida; and Social Development Research Group (Hawkins, Oesterle), University of Washington, Seattle, Washington

Abstract

Introduction—Adolescent substance use and delinquency are major public health problems. Although community-based prevention strategies have been recommended to produce population-level reductions in rates of substance use and delinquency, few models show evidence of effectiveness.

Purpose—To test the efficacy of a community-based prevention system, Communities That Care (CTC), in reducing community rates of problem behaviors, particularly effects on specific profiles of adolescent substance use and delinquency in eighth and tenth graders.

Methods—Twenty-four communities were randomized to CTC intervention or control groups. Data were collected from 14,099 eighth and tenth grade students in these communities using anonymous cross-sectional surveys in 2004 and 2010 and analyzed in 2012. Outcomes were four different profiles of self-reported substance use and delinquency in eighth grade and five profiles in tenth grade.

Results—In the cross-sectional 2010 data, there was no intervention effect on the probability of experimenting with substances or of substance use coupled with delinquent activities for either grade. However, tenth graders in intervention communities were significantly less likely to be alcohol users than those in control communities (OR=0.69, CI=0.48, 1.00).

Conclusions—Cross-sectional population surveys showed evidence of CTC effects in reducing tenth grade alcohol users but not experimenters. A community-wide reduction in adolescent alcohol use is important because alcohol is the most commonly used illicit substance during

© 2014 American Journal of Preventive Medicine. Published by Elsevier Inc. All rights reserved.

Address correspondence to: M. Lee Van Horn, Department of Psychology, Barnwell College, University of South Carolina, Columbia SC 29206. vanhorn@sc.edu.

No financial disclosures were reported by the authors of this paper.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

adolescence, and early initiation of alcohol use has been associated with alcohol-related disorders in adulthood. Failure to find hypothesized effects on experimenters qualifies these results.

Introduction

Adolescence is marked by increasing involvement in risk-seeking behaviors such as substance use and delinquency.^{1,2} Such behaviors can result in negative consequences such as poor academic performance, compromised health, and increased rates of victimization and physical injury.^{3,4} Initiation of alcohol use during adolescence increases risk for alcohol and other drug disorders in adulthood,^{5,6} and early initiation of delinquency predicts continued involvement in severe, violent offending.^{7,8} The high financial costs of these problem behaviors are related to medical care, work loss, drug treatment programs, and correctional systems.^{9–11}

Many interventions have been developed and demonstrated effective in preventing adolescent problem behaviors.¹² However, relatively few community-based preventive interventions have shown effectiveness, however, leading federal agencies to call for more development and evaluation of such approaches.^{10,12,14,15} Compared with single interventions, community-based strategies have the potential to affect multiple outcomes across a larger population, given their focus on simultaneously addressing numerous risk and protective factors via the implementation of multiple, locally coordinated preventive interventions that are likely to affect the majority of residents in a community.^{16,17}

To date, only two community-based prevention systems have been found to impact adolescent delinquency and/or substance use.¹⁸ The first is the Promoting School–Community–University Partnerships to Enhance Resilience (PROSPER) project, which relies on university staff, school district officials, and other community members to implement family-focused and school-based prevention services. This model has demonstrated reductions in the initiation, prevalence, and escalation of substance use during adolescence.¹⁹ The second, Communities That Care (CTC),^{20,21} relies on local coalitions to implement a range of services that reduce risk factors and strengthen protective factors experienced by youth. CTC has been shown to reduce the initiation of tobacco use, alcohol use, delinquency, and violence among a longitudinal panel of students followed from grade five to ten.^{22,23}

Previous tests of community-based prevention systems examine average effects on each behavioral outcome. However, etiologic evidence indicates that risk and protective factors predict numerous problem behaviors that often co-occur.^{24,25} Thus, when evaluating intervention effects, it would be useful to examine the degree to which an intervention affects involvement in multiple problem behaviors, such as substance use **and** violence, as well as combinations of such behaviors. The current study investigates the degree to which CTC affects the probability that adolescents will engage in specific behavioral profiles of substance use, delinquency, and violence.

Although the methodology described in this paper could be applied to the evaluation of any type of preventive intervention, it is particularly relevant for comprehensive approaches that

involve the implementation of multiple services targeting diverse predictors of related outcomes. In the CTC system, community coalitions receive training and technical assistance and follow structured protocols to prevent youth problem behaviors. Based on a public health framework, CTC assists communities in conducting population surveys of public secondary school students to identify locally elevated risk factors and suppressed protective factors and in implementing tested and effective preventive interventions that target these factors. Thus, the interventions are specific to the needs of each community and include one or more types of services such as school-based curricula and organizational change models, afterschool services (e.g., tutoring and mentoring programs), and family-focused interventions such as parenting workshops.

Coalitions are encouraged to deliver an array of preventive interventions, which target to change multiple risk and protective factors in as many participants as possible, so long as the interventions are implemented with fidelity and in a coordinated fashion to minimize duplication of services. Although communities may implement interventions targeting higher-risk youth (i.e., selective or indicated prevention programs), CTC emphasizes the delivery of services to the general population. Therefore, the CTC system can be considered a universal prevention approach intended to reduce community-wide problem behaviors.

The CTC theory of change posits that this system will produce reductions in the **overall prevalence** of health problems with similar effects across individuals. However, the effects of CTC could be specific to certain groups. For example, evaluations of other universal interventions have reported stronger or weaker impacts on problem behaviors according to demographic status (e.g., sex or race/ethnicity) or for higher- versus lower-risk youth.^{26–32} The current study examines outcomes in a different manner. Rather than assessing the degree to which the intervention has produced changes in a single behavior for groups of individuals defined according to demographic characteristics or risk status, we investigate the degree to which the CTC system affects the probability that adolescents engage in specific behavioral profiles of substance use, delinquency, and violence.

This approach is based on prior examination of cross-sectional data from the CTC study, which identified exclusive profiles of multiple problem behaviors.³³ Students in the *abstainer* profile had very low probabilities of reporting any substance use, delinquency, or violence; *experimenters* reported some lifetime substance use but little recent use, and were very unlikely to have engaged in delinquent behaviors; students in the *drug use* profile were very likely to report current drug use but less likely to engage in delinquent behaviors; and *problem students* were very likely to report substance use and delinquent activities. There was also evidence for a fifth profile: a group of students characterized as *alcohol users*, who were very likely to engage in current alcohol use but had limited involvement with other substances or delinquency, and primarily came from the experimenter profile when only four profiles were estimated. Evidence that students engage in different combinations of problem behaviors provides the foundation for this more nuanced evaluation of how CTC can effect changes in communities. Investigating the degree to which CTC or any intervention impacts profiles of co-occurring behaviors can better illuminate the ways in which the intervention achieves change and better specify where the strongest intervention effects occur, although the methods are limited by the assumption that the profiles are stable

across samples and across time. This approach jointly tests intervention effects on the probability that study participants are in each of the identified categories while also allowing tests of specific hypotheses about the impacted groups.

Research Aims

Although effects of CTC have been shown in a longitudinal panel, they have not been demonstrated with cross-sectional data and groups of students most impacted by the intervention have not been identified. The current paper uses repeated cross-sectional data to examine the probability of eighth and tenth grade students in intervention and control communities engaging in different profiles of problem behaviors. Prior analyses have shown that in 2004, before implementation of preventive interventions selected through the CTC process, there was no difference in profiles among eighth grade students.³³ This paper shows evidence for the comparability of these profiles in 2004 surveys of tenth graders. The primary aim of this study was to examine effects of the CTC intervention on cross-sectional profiles of problem behaviors in 2010, 6 years after the intervention communities began targeted prevention services. Given that CTC generally advocates for implementation of universal preventive interventions, we previously hypothesized that the greatest preventive impact would be localized to youth engaging in more common and less serious problem behaviors.³³ We test the hypothesis that in contrast to students in control communities, students in the CTC intervention communities will have a reduced likelihood of being an experimenter or alcohol user compared to being an abstainer, assuming that these classes are replicated.

Methods

Data were obtained from students in the Community Youth Development Study (CYDS),³⁴ an RCT testing the effectiveness of the CTC model in 24 small to moderate-sized communities (12 matched pairs) located in seven states (Figure 1). One community from each pair was randomized to implement CTC and the other to conduct prevention services as usual. From spring 2003 through spring 2008 (Years 1–5, the efficacy phase of the trial), each of the 12 intervention communities was provided with training in the CTC model and proactive technical assistance via weekly phone calls and one to three annual site visits by research staff to ensure faithful delivery of the CTC model and selected preventive interventions. They also received funding for a CTC coordinator and up to \$75,000 in Years 2–5 to implement programs, policies, and practices showing prior evidence of effectiveness in well-conducted trials, as identified for communities in the CTC *Prevention Strategies Guide* (communitiesthatcare.net), and that targeted fifth- to ninth-grade students and their families.

Communities selected interventions that targeted risk factors reported as elevated by local youth on the same surveys used in the current analyses. Priority risk factors and interventions differed between communities. Across communities, 17 different school-, family-, and community-based interventions (examples include: Life Skills Training, Guiding Good Choices, and Big Brothers/Big Sisters) were implemented during Years 2–5, with an average of three programs per community.^{35–38} The sustainability phase of the

CYDS trial began in 2008; from that point, data were collected to evaluate long-term changes in student outcomes, but training and technical assistance and financial resources were no longer provided. Most intervention communities continued to implement CTC and evidence-based programs.³⁹

The present study began by examining profiles of problem behaviors among tenth graders in spring 2004. Although the randomized trial began in spring 2003, intervention communities spent the first 15–18 months receiving training, forming coalitions, and planning. Intervention programming began in fall 2004. No intervention effects were expected in spring 2004, and none were found for eighth graders at that time point.³³ By spring 2010, intervention communities had been implementing CTC with technical and financial support from the study for 5 years and without study support for 2 additional years. Eleven of the original 12 CTC coalitions were still active³⁹ and intervention sites were delivering effective prevention programming at higher rates than control communities.⁴⁰

Procedures

Data were obtained from cross-sectional population surveys of students administered in schools during a classroom period by teachers using standardized anonymous administration procedures. Teachers received specific instructions to assure anonymity of responses and signed an agreement to abide by these instructions. Screening criteria were used to exclude respondents who lied or responded inconsistently. For instance, if students responded *never* to lifetime use of cigarettes and indicated smoking one or more cigarettes per day during the past 30 days on a separate question, their answers were considered inconsistent. Students with over two inconsistent responses or who reported that they were not honest in completing the survey were excluded. Because responses were anonymous, passive consent was obtained from parents and informed assent was obtained from students. The University of Washington's Human Subjects Review Committee approved the protocol.

Participants

Cross-sectional data were collected from students in grades six, eight, ten, and 12. Intervention effects were examined for eighth and tenth grades only. In grade six, reported levels of substance use and delinquency were too low to expect detectable intervention effects, and lower response rates made the 12th grade cross-sectional samples less representative of the population. Across data sets, the sample consisted of between 49% and 53% girls. Between 3% and 6% of students self-identified as African American, and 77% to 80% as Caucasian. In the 2004 sample, 11% of surveyed tenth graders identified as Hispanic. In 2010, when new reporting requirements resulted in a change in the wording of race/ethnicity items such that Hispanic was designated as a separate category, 23% of eighth graders and 19% of tenth graders identified as Hispanic.

Measures

Data were collected using the CTC Youth Survey,^{41,42} which is designed to measure risk and protective factors, substance use, and delinquency in middle and high school samples. Analyses used items asking the frequency of alcohol, tobacco, and marijuana use across the lifetime and in the last 30 days, and the number of times in the last 2 weeks they had more

than five drinks at one time. Analyses included three levels of substance use: no use, lifetime but not current use, and use within the last 30 days. Students also indicated the frequency with which they engaged in eight delinquent behaviors over the last year (Table 1); these items and binge drinking were coded yes/no in analyses.

Three community-level variables were included to reduce community-level variability in outcomes. The percentage of students receiving free or reduced-price lunch and the total number of students in the community in thousands were obtained from the National Center for Educational Statistics for the 2003–2004 school year. Additionally, an indicator for Utah was included because these students were uniformly lower in levels of substance use. When community matching was explicitly modeled, the Utah adjustment was unnecessary.

Data Analysis

In 2012, multi-level latent class analysis (MLCA) estimated in Mplus⁴³ was used to examine the effects of CTC on the probability of students' membership in latent classes. The multi-level component of the model accounted for the fact that communities rather than individual subjects were randomized.^{44,45} The least restrictive, fully random approach³³ was used to account for clustering between communities in the probability that a student would be in each latent class versus the reference class (abstainer was the reference class in this case). Because communities rather than individuals were randomized to condition, fixed and random effects for the probability of membership in each class were included to maintain the nominal type I error rate. Full information maximum likelihood estimation allowed for the inclusion of subjects with missing data, except those missing data on all variables.

Because the meaning of latent classes may change across time and samples, it is important to also verify that classes are replicated with independent data; results of these analyses are only valid in populations where these latent classes exist. Previous analyses of eighth grade students found support for both four and five-class solutions in independent samples collected in communities from the same states.³³ For this study, the five-class solution was also replicated for an independent sample of tenth graders (Appendix A).

Initial analyses found that the differences between communities in the distribution of students across classes were greater than those shown in previous analyses as indicated by intraclass correlation coefficients (ICCs) for latent class means ranging from 0.05 to 0.10. This is problematic because higher ICCs for the outcome(s) lower statistical power to find treatment effects.^{46,47} Community-level covariates were included in the analyses to account for these differences. The meaning of the latent classes was virtually unchanged, but the ICCs for class means decreased to between 0 and 0.03, due primarily to communities in Utah compared to other states. Rates of substance use and problem behaviors were much lower in Utah communities and thus there were far more abstainers (Table 2).

Results

Initial analyses tested the equivalence of intervention and control conditions for tenth grade students in 2004, before program implementation. The five-class model including abstainer, experimenter, alcohol user, drug user, and problem behavior classes found previously in

eighth grade an independent tenth grade sample (Appendix A) was replicated in 2004. No evidence was found for pre-intervention differences between intervention and control communities in profiles of substance use and delinquency.

The first step in assessing intervention effects on latent classes was the replication of the previously identified latent classes. The four-class solution was replicated for eighth graders in 2010. However, the five-class solution did not distinguish an alcohol use profile, but instead contained a small group (6%) of students who reported current marijuana use and no binge drinking (Appendix A). Rather than use this result that was not replicated in any other sample and therefore potentially unreliable, we proceeded by examining intervention effects for eighth grade students using four classes, including abstainers, experimenters, drug users, and problem students, which were substantively the same as the four previously identified classes. For tenth grade students, the five-class solution including the alcohol class was replicated in 2010 and was utilized in subsequent analyses. Response profiles for each grade are shown in Table 1.

Effects of the CTC intervention on class membership in 2010 for both eighth and tenth grades are reported in Table 2. We hypothesized that CTC would reduce the probability of being in the experimenter and alcohol use classes versus the abstainer classes for students in the intervention compared to the control condition. Results showed no intervention effects for the experimenter class for either grade. For tenth grade students, the intervention reduced the probability of being in the alcohol use versus the abstainer class, as hypothesized. The odds of a tenth grade student in an intervention community being in the alcohol use class as opposed to the abstainer class was 0.69 (CI=0.48, 1.00) times those of a student in the control condition. To show the practical implication of these effects, the aforementioned multinomial regression model was used to calculate the predicted probability that a student from a control community not in Utah, at the average level of free and reduced lunch (39%), and of the average student population size (3,417 students) would be an alcohol user was 0.26 versus 0.19 if the student lived in a treatment community, the expected reduction in alcohol users due to CTC is from 26% to 19%. Analyses showed no difference in the probability of belonging to either the drug use or problem behavior classes versus the abstainer class in the CTC intervention communities compared to the control communities in eighth or tenth grade cross-sectional samples (Table 2).

Discussion

CTC was designed to reduce levels of substance use, delinquency, and other problem behaviors using a community-based prevention system that relies on broad-based coalitions to reduce elevated risk factors and strengthen protective factors with effective preventive interventions. Previous studies have shown significant intervention effects in smoking, drinking, and delinquency in a longitudinal panel of students followed from grade five through ten.^{22, 23} Instead of the longitudinal panel, the current study used two cross-sectional samples of eighth and tenth grade students collected 6 years apart: in 2004, before prevention programs had been initiated, and in 2010, 2 years after external support for CTC and prevention program implementation had been withdrawn but while intervention sites were continuing to deliver effective prevention programming.⁴⁰

The current study differed from prior outcome analyses in that it tested the specific hypothesis that CTC would reduce the proportion of students in the experimenter and alcohol user classes in repeated cross-sectional surveys. Because some universal interventions have been shown to have the strongest effects among the highest-risk youth,^{26–30} the study also assessed the degree to which intervention effects were found among youth engaging in more serious or higher levels of problem behaviors. This possibility was tested by default, given that differences in all profiles were assessed with the LCA approach.

Only one of three hypothesized intervention effects was found: a 7–percentage point reduction in alcohol users in tenth grade. Contrary to our hypothesis, no differences in experimenters were found across intervention conditions for either eighth or tenth grade students in the cross-sectional 2010 survey. Analyses showed no evidence of intervention effects on more serious levels of problem behaviors, although these analyses were exploratory and effects were not specifically hypothesized. The inclusion of community-level covariates substantially reduced variability in latent classes between communities because the students in Utah were much more likely to be abstainers than students in any other state. These conclusions assume the stability of the four- and five-class models in the population.

The analyses utilized in this study allowed identification of a group of students for whom the intervention was most effective: tenth graders who displayed a specific profile of being likely to engage in current drinking and binge drinking, but not other substance use or problem behaviors. Although it is unclear why intervention effects would be largest on alcohol users, that effects on alcohol were found in both cross-sectional and longitudinal data^{22,23} is noteworthy. Cross-sectional analyses typically have less power to detect intervention effects than analyses of longitudinal data.⁴⁸ The reduction in rates of alcohol use is also important because alcohol is the most commonly used substance among adolescents.⁴⁹ Moreover, use of alcohol during adolescence has been associated with an increased risk of other negative outcomes not measured in this survey, including victimization, risky sexual behavior, and mental health problems,^{25,50–52} as well as an increased likelihood of alcohol abuse and dependency during adulthood.^{25,53}

An unexpected finding was that the intervention did not affect the proportion of students who were experimenters in either grade and that the alcohol use class was not replicated in the eighth grade. One explanation for these null findings is the systematic decrease in substance use seen both in these data (the percentage of eighth grade abstainers increased from 58% in 2004 to 66% in 2010) and in recent years in national surveys.⁴⁹ These trends could have impacted profiles of substance use such that the alcohol use class was no longer evident for eighth graders. Decreasing percentages of experimenters in eighth and tenth grade also reduces power to find intervention effects for these students. Future cross-sectional surveys of students in the CTC study communities could help inform some of these issues. Although this study did not look at intervention effects in other years, it would be especially informative to examine these effects in 2008 and, given that interventions are sustained, in 2012 data.

This study did not find a reduction in rates of serious drug use or problem behaviors for students in intervention versus control communities. These analyses also had low power to detect effects given the low rates of involvement in these behaviors. Additionally, given that communities largely implemented universal prevention strategies targeting youth who had not yet initiated involvement in problem behaviors, we did not have strong expectations for finding intervention effects in these groups.

Although this paper did not demonstrate the hypothesized intervention effects, the more nuanced evaluation provided by these analyses did allow identification of a specific subgroup of students for whom the intervention was effective. The strength of this LCA approach is that it recognizes that not all substance or alcohol use is the same, and allows for the creation of particular profiles of problem behaviors based on all measured outcomes. In this case, the analyses were able to distinguish between alcohol users, those who experimented with substance use, and those who used alcohol as part of a pattern of more serious problem behaviors. It is possible that students who engage only in alcohol use view this behavior as normative, whereas they consider other illegal drugs or delinquent behaviors as more deviant. The current study suggests that CTC may be effective at altering the attitudes and behaviors of alcohol users more than those of others.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by Research Grant R01DA015183 from the National Institute on Drug Abuse awarded to J. David Hawkins. We are grateful to the co-funding organizations: the National Cancer Institute, Center for Substance Abuse Prevention, National Institute on Alcohol Abuse and Alcoholism, National Institute on Mental Health, and National Institute on Child Health and Development. Thanks are due to John Briney for making 10 years of data usable.

References

1. Sampson RJ, Laub JH. Crime and deviance in the life course. *Annu Rev Sociol.* 1993; 18:63–84.
2. Moffitt TE. Adolescence-limited and life-course persistent anti-social behavior: a developmental taxonomy. *Psychol Rev.* 1993; 100(4):674–701. [PubMed: 8255953]
3. Patton GC, Coffey C, Sawyer SM, et al. Global patterns of mortality in young people: a systematic analysis of population health data. *Lancet.* 2009; 374(9693):881–92. [PubMed: 19748397]
4. Fergusson DM, Lynskey MT, Horwood LJ. Factors associated with continuity and changes in disruptive behavior patterns between childhood and adolescence. *J Abnorm Child Psych.* 1996; 24(5):533–54.
5. Windle M, Spear LP, Fuligni AJ, et al. Transitions into underage and problem drinking: summary of developmental processes and mechanisms: ages 10–15. *Alcohol Res Health.* 2009; 32(1):30–40. [PubMed: 23104445]
6. Hingson RW, Heeren T, Winter MR. Age at drinking onset and alcohol dependence: age at onset, duration, and severity. *Arch Pediatr Adolesc Med.* 2006; 160(7):739–46. [PubMed: 16818840]
7. Farrington DP. Developmental and life-course criminology: key theoretical and empirical issues—the 2002 Sutherland Award address. *Criminology.* 2003; 41(2):221–55.
8. Sampson, RJ.; Laub, JH. Crime in the making: pathways and turning points through life. Cambridge MA: Harvard University Press; 1993.

9. Woolf SH. The big answer: rediscovering prevention at a time of crisis in health care. *Harvard Health P Rev*. 2006; 7(2):5–20.
10. Specter A. Making youth violence prevention a national priority. *Am J Prev Med*. 2008; 34(3S):S3–S4. [PubMed: 18267196]
11. The National Center on Addiction and Substance Abuse. Shoveling up II: the impact of substance abuse on federal, state, and local budgets. New York: Columbia University; 2009.
12. National Research Council and Institute of Medicine. Preventing mental, emotional, and behavioral disorders among young people: progress and possibilities. Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults: research advances and promising interventions. Washington D.C: Board on Children, Youth, and Families, Division of Behavioral and Social Sciences and Education, The National Academies Press; 2009.
13. National Prevention Council. National prevention strategy. Washington D.C: USDHHS, Office of the Surgeon General; 2011.
14. Woolf SH. The power of prevention and what it requires. *JAMA*. 2008; 299(20):2437–9. [PubMed: 18505953]
15. Office of National Drug Control Policy. National drug control strategy 2012. Washington DC: Office of National Drug Control Policy; 2012.
16. Wagenaar AC, Erickson DJ, Harwood EM, O'Malley PM. Effects of state coalitions to reduce underage drinking: a national evaluation. *Am J Prev Med*. 2006; 31(4):307–15. [PubMed: 16979455]
17. David-Ferdon C, Hammond WR. Community mobilization to prevent youth violence and to create safer communities. *Am J Prev Med*. 2008; 34(3S):S1–S2. [PubMed: 18267193]
18. Catalano RF, Fagan AA, Gavin LE, et al. Worldwide application of prevention science in adolescent health. *Lancet*. 2012; 379(9826):1653–64. [PubMed: 22538180]
19. Spoth RL, Redmond C, Clair S, Shin C, Greenberg MT, Feinberg ME. Preventing substance misuse through community–university partnerships randomized controlled trial outcomes 4½ years past baseline. *Am J Prev Med*. 2011; 40(4):440–7. [PubMed: 21406278]
20. Hawkins, JD.; Catalano, RFJ. Communities that care: action for drug abuse prevention. San Francisco CA: Jossey–Bass Publishers; 1992.
21. Hawkins JD, Catalano RF, Arthur MW. Promoting science-based prevention in communities. *Addict Behav*. 2002; 27:951–76. [PubMed: 12369478]
22. Hawkins JD, Oesterle S, Brown EC, et al. Results of a type 2 translational research trial to prevent adolescent drug use and delinquency: a test of communities that care. *Arch Pediatr Adolesc Med*. 2009; 163(9):789–98. [PubMed: 19736331]
23. Hawkins JD, Oesterle S, Brown EC, et al. Sustained decreases in risk exposure and youth problem behaviors after installation of the communities that care prevention system in a randomized trial. *Arch Pediatr Adolesc Med*. 2012; 166(2):141–8. [PubMed: 21969362]
24. Coie JD, Watt NF, West SG, et al. The science of prevention: a conceptual framework and some directions for a national research program. *Am Psychol*. 1993; 48(10):1013–22. [PubMed: 8256874]
25. Hawkins JD, Catalano RF, Miller JY. Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychol Bull*. 1992; 112(1):64–105. [PubMed: 1529040]
26. Allen JP, Philliber S. Who benefits most from a broadly targeted prevention program? Differential efficacy across populations in the teen outreach program. *J Community Psychol*. 2001; 29(6):637–55. [PubMed: 17235364]
27. Tolan PH, Gorman-Smith D, Henry DB. Supporting families in a high-risk setting: proximal effects of the SAFEChildren preventive intervention. *J Consult Clin Psych*. 2004; 72(5):855–69.
28. Kellam SG, Brown CH, Poduska JM, et al. Effects of a universal classroom behavior management program in first and second grades on young adult behavioral, psychiatric, and social outcomes. *Drug Alcohol Depen*. 2008; 95S:S5–S28.
29. Conduct Problems Prevention Research Group. Fast track randomized controlled trial to prevent externalizing psychiatric disorders: findings from grades 3 to 9. *J Am Acad Child Psy*. 2007; 46(10):1250–62.

30. Olds DL, Henderson CR Jr, Cole R. Long-term effects of nurse home visitation on children's criminal and antisocial behavior: 15-year follow-up of a randomized controlled trial. *JAMA*. 1998; 280(14):1238–44. [PubMed: 9786373]
31. Eisen M, Zellman GL, Massett HA, Murray DM. Evaluating the Lion's-Quest "skills for adolescence" drug education program: first-year behavioral outcomes. *Addict Behav*. 2002; 27:619–32. [PubMed: 12188596]
32. Perry CL, Williams CL, Komro KA, et al. Project Northland high school interventions: community action to reduce adolescent alcohol use. *Health Educ Behav*. 2000; 27(1):29–49. [PubMed: 10709791]
33. Van Horn ML, Fagan AA, Jaki T, et al. Using multilevel mixtures to evaluate intervention effects in group randomized trial. *Multivar Behav Res*. 2008; 43(2):289–326.
34. Hawkins JD, Catalano RF, Arthur MW, et al. Testing communities that care: the rationale, design and behavioral baseline equivalence of the community youth development study. *Prev Sci*. 2008; 9(3):178–90. [PubMed: 18516681]
35. Fagan AA, Hanson K, Hawkins JD, Arthur MW. Bridging science to practice: achieving prevention program implementation fidelity in the community youth development study. *Am J Community Psychol*. 2008; 41(3–4):235–49. [PubMed: 18302016]
36. Fagan AA, Hanson K, Hawkins JD, Arthur MW. Implementing effective community-based prevention programs in the community youth development study. *Youth Violence Juv Justice*. 2008; 6(3):256–78.
37. Fagan AA, Hanson K, Hawkins JD, Arthur MW. Translational research in action: implementation of the communities that care prevention system in 12 communities. *J Community Psychol*. 2009; 37(7):809–29. [PubMed: 22121303]
38. Quinby RK, Fagan AA, Hanson K, Brooke-Weiss B, Arthur MW, Hawkins JD. Installing the communities that care prevention system: implementation progress and fidelity in a randomized controlled trial. *J Community Psychol*. 2008; 36(3):313–32.
39. Gloppen KM, Arthur MW, Hawkins JD, Shapiro VB. Sustainability of the communities that care prevention system by coalitions participating in the community youth development study. *J Adolescent Health*. 2012; 51:259–64.
40. Fagan AA, Hanson K, Briney JS, Hawkins JD. Sustaining the utilization and high quality implementation of tested and effective prevention programs using the communities that care prevention system. *Am J Community Psychol*. 2012; 49(3–4):365–77. [PubMed: 21809149]
41. Arthur MW, Hawkins JD, Pollard JA, Catalano RF, Baglioni AJ. Measuring risk and protective factors for substance use, delinquency, and other adolescent problem behaviors—the communities that care youth survey. *Eval Rev*. 2002; 26:575–601. [PubMed: 12465571]
42. Glaser RR, Van Horn ML, Arthur MW, Hawkins JD, Catalano RF. Measurement properties of the communities that care youth survey across demographic groups. *J Quant Criminol*. 2005; 21:73–102.
43. Muthén, LK.; Muthén, BO. *Mplus*. 6. Los Angeles CA: Muthén & Muthén; 2010.
44. Vermunt JK. Multilevel latent class models. *Sociol Methodol*. 2003; 33:213–39.
45. Vermunt JK, Van Dijk L. A nonparametric random-coefficients approach: the latent class regression model. *Multilevel Modeling Newsletter*. 2001; 13:6–13.
46. Murray, DM. *Design and analysis of group-randomized trials*. New York: Oxford University Press; 1998.
47. Murray DM, Van Horn ML, Hawkins JD, Arthur MW. Analysis strategies for a community trial to reduce adolescent ATOD use: a comparison of random coefficient and ANOVA/ANCOVA models. *Contemp Clin Trials*. 2006; 27:188–206. [PubMed: 16324889]
48. Zeger SL, Liang KL. An overview of methods for the analysis of longitudinal data. *Stat Med*. 1992; 11(14–15):1825–39. [PubMed: 1480876]
49. Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE. *Demographic subgroup trends for various licit and illicit drugs, 1975–2010 (Monitoring the Future Occasional Paper No. 74)*. Ann Arbor MI: Institute for Social Research; 2011.
50. Donovan JE. Adolescent alcohol initiation: a review of psychosocial risk factors. *J Adolescent Health*. 2004; 35(6):e7–18.

51. Schinke SP, Fang L, Cole KCA. Substance use among early adolescent girls: risk and protective factors. *J Adolescent Health*. 2008; 43(2):191–4.
52. Begle AM, Hanson RF, Danielson CK, et al. Longitudinal pathways of victimization, substance use, and delinquency: findings from the national survey of adolescents. *Addict Behav*. 2011; 36:682–9. [PubMed: 21377805]
53. Hingson RW, Heeren T, Winter M. Age at drinking onset and alcohol dependence—age at onset, duration, and severity. *Arch Pediatr Adolesc Med*. 2006; 160(7):739–46. [PubMed: 16818840]

Figure 1.

CONSORT diagram for eighth and ninth grade CYDS cross-sectional sample used in latent class analysis

CYDS, Community Youth Development Study

Table 1

Predicted responses probabilities from multilevel unconditional latent class analyses

	8th Grade 2010					10th Grade 2010				
	Abstainer	Experiment	Drug Use	Problem	Abstainer	Experiment	Alcohol	Drug Use	Problem	
Proportion	66%	22%	9%	3%	57%	16%	11%	13%	3%	
Cigarettes										
Never	0.98	0.45	0.11	0.17	0.97	0.30	0.52	0.07	0.24	
Lifetime	0.02	0.48	0.34	0.23	0.03	0.58	0.34	0.30	0.16	
Current	0.00	0.07	0.55	0.60	0.01	0.12	0.14	0.63	0.60	
Alcohol										
Never	0.82	0.17	0.03	0.06	0.75	0.11	0.01	0.00	0.08	
Lifetime	0.14	0.49	0.13	0.10	0.19	0.78	0.10	0.11	0.06	
Current	0.04	0.34	0.83	0.84	0.07	0.11	0.89	0.89	0.85	
Marijuana										
Never	1.00	0.72	0.17	0.12	0.99	0.37	0.54	0.03	0.18	
Lifetime	0.00	0.21	0.21	0.15	0.01	0.46	0.25	0.21	0.16	
Current	0.00	0.07	0.62	0.74	0.01	0.17	0.22	0.76	0.66	
Binge Drinking										
No	1.00	0.86	0.39	0.22	1.00	1.00	0.45	0.34	0.25	
Yes	0.00	0.15	0.62	0.78	0.00	0.00	0.55	0.66	0.75	
Suspended										
No	0.95	0.72	0.54	0.11	0.96	0.78	0.88	0.64	0.30	
Yes	0.05	0.28	0.46	0.89	0.05	0.22	0.12	0.36	0.70	
Carried gun										
No	0.98	0.88	0.89	0.18	0.97	0.94	0.93	0.91	0.20	
Yes	0.02	0.12	0.11	0.83	0.03	0.06	0.07	0.09	0.80	
Sold drugs										
No	1.00	0.99	0.78	0.23	1.00	0.96	0.99	0.60	0.17	
Yes	0.00	0.01	0.22	0.77	0.00	0.04	0.01	0.40	0.83	
Stole Vehicle										
No	1.00	0.95	0.93	0.42	0.99	0.97	0.99	0.94	0.35	
Yes	0.00	0.06	0.07	0.59	0.01	0.04	0.01	0.06	0.65	

		8th Grade 2010				10th Grade 2010				
		Abstainer	Experiment	Drug Use	Problem	Abstainer	Experiment	Alcohol	Drug Use	Problem
Arrested										
No	Yes	0.99	0.88	0.77	0.25	0.98	0.88	0.95	0.70	0.24
Yes		0.01	0.13	0.24	0.75	0.02	0.12	0.05	0.30	0.76
Attacked someone										
No	Yes	0.94	0.71	0.50	0.10	0.96	0.82	0.90	0.63	0.11
Yes		0.06	0.29	0.50	0.90	0.04	0.18	0.10	0.37	0.90
High at school										
No	Yes	1.00	0.88	0.34	0.18	1.00	0.79	0.86	0.18	0.06
Yes		0.00	0.12	0.66	0.82	0.00	0.21	0.14	0.82	0.94
Gun to school										
No	Yes	1.00	0.99	0.98	0.46	1.00	0.99	0.99	1.00	0.43
Yes		0.00	0.01	0.02	0.54	0.00	0.01	0.01	0.01	0.57

Note: Proportion is the proportion of student estimated to be in each class.

Table 2

Effects of the Communities that Care intervention on latent classes

	8th Grade		10th grade	
	Estimate	SE	Estimate	SE
Experimenters versus Abstainer Class				
Intercept	-2.09	0.17	-1.49	0.29
CTC treatment	0.06	0.08	-0.21	0.16
Free/reduced lunch	2.57	0.41	0.64	0.60
Number of students	0.03	0.03	0.12	0.05
Community in Utah	-0.71	0.12	-1.03	0.26
Random intercept	0.00	0.00		0.36
Alcohol Use versus Abstainer Class				
Intercept	NA		-1.32	0.27
CTC treatment	NA		-0.37	0.18
Free/reduced lunch	NA		0.20	0.53
Number of students	NA		0.05	0.07
Community in Utah	NA		-1.44	0.48
Random intercept	NA		0.07	0.04
Drug Use versus Abstainer Class				
Intercept	-2.99	0.30	-1.36	0.30
CTC treatment	0.10	0.08	-0.11	0.15
Free/reduced lunch	2.29	0.56	-0.37	0.59
Number of students	0.04	0.08	0.15	0.03
Community in Utah	-0.80	0.46	-1.62	0.19
Random intercept	0.10	0.04	0.04	0.04
Problem Behavior versus Abstainer Class				
Intercept	-3.94	0.58	-3.62	0.37
CTC treatment	0.40	0.25	1.49	0.15
Free/reduced lunch	0.40	1.16	1.63	0.75
Number of students	0.26	0.09	1.30	0.08
Community in Utah	-1.80	0.45	-1.22	0.29

	8th Grade			10th grade		
	Estimate	SE	OR	Estimate	SE	OR
Random intercept	0.12	0.12		0.00	0.00	

Note: Analyses use data from 2010 included 4,671 8th graders and 4,810 10th graders who were in the cross sectional Communities that Care sample. The alcohol use class was not included in the 8th grade analyses. Number of students was divided by 1,000 in analyses.

Boldface indicates statistical significance ($p<0.05$).

CTC, Communities that Care; NA, not applicable.