Published in final edited form as:

Am J Prev Med. 2014 August ; 47(2): 141–149. doi:10.1016/j.amepre.2014.02.011.

E-Cigarette Awareness and Perceived Harmfulness:

Prevalence and Associations with Smoking Cessation Outcomes

Andy S.L. Tan, MBBS, MPH, MBA, PhD and Cabral A. Bigman, PhD

Center of Excellence in Cancer Communication Research (Tan), Annenberg School for Communication, University of Pennsylvania, Philadelphia, Pennsylvania; and the Department of Communication (Bigman), College of Liberal Arts & Sciences, University of Illinois at Urbana-Champaign, Urbana, IL

Abstract

Background—Electronic cigarettes, or e-cigarettes, are increasingly advertised as replacements for regular cigarettes or cessation aids for smokers.

Purpose—To describe the prevalence and correlates of e-cigarette awareness and perceived harmfulness among U.S. adults and analyze whether these variables are associated with smokers' past year quit attempts and intention to quit.

Methods—Data were obtained from the Health Information National Trends Survey (HINTS 4 Cycle 2), conducted from October 2012 to January 2013. Data analyses were performed from June to August 2013.

Results—Overall, 77% of respondents were aware of e-cigarettes. Of these, 51% believed e-cigarettes were less harmful than cigarettes. Younger, white (compared with Hispanic), more educated respondents, and current or former smokers (compared with non-smokers) were more likely to be aware of e-cigarettes. Among those who were aware of e-cigarettes, younger, more educated respondents and current smokers (compared with former and non-smokers) were more likely to believe that e-cigarettes were less harmful. Awareness and perceived harm were not associated with smokers' past year quit attempts or intention to quit.

Conclusions—Overall e-cigarette awareness increased while smokers' perceived harm of e-cigarettes declined compared with earlier surveys. However, awareness and perceived harm of e-cigarettes did not show evidence of promoting smoking cessation at the population level.

© 2014 American Journal of Preventive Medicine. Published by Elsevier Inc. All rights reserved.

Address correspondence to: Andy S.L. Tan, Annenberg School for Communication, University of Pennsylvania, 3620 Walnut Street, Philadelphia PA 19104. andytan@alumni.upenn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

No financial disclosures were reported by the authors of this paper.

Introduction

Electronic cigarettes, also known as e-cigarettes, are defined as “battery-operated products designed to deliver nicotine, flavor, and other chemicals.”¹ E-cigarettes are marketed through various media channels,² sports or music events,^{3,4} and outdoor or in-store displays.^{2,5} E-cigarettes have been portrayed in the media as competing products to traditional cigarettes and a cessation aid. For instance, some online marketing strategies communicate that e-cigarettes are safer alternatives to regular cigarettes.⁶ Increasingly, e-cigarette use is appearing in news stories and entertainment media.^{2,7} Some YouTube videos, for example, feature e-cigarettes use on entertainment shows by celebrities who claim that e-cigarettes are safer than regular cigarettes.^{8,9} Other videos misrepresent e-cigarettes as medical products and smoking cessation aids.^{8,9} The products are not currently approved by the U.S. Food and Drug Administration (FDA) for smoking cessation.¹⁰

As awareness of e-cigarettes and their potential competitive advantages to traditional cigarettes increases because of marketing and publicity, it is possible that use of traditional cigarettes will decline. Prior research on the marketing of nicotine replacement therapies and other established cessation aids suggest a link between exposure and cessation intentions and behaviors. Avery and colleagues, for example, found that increased exposure to magazine advertisements of cessation products (e.g., Habitrol, Nicoderm patch, Nicorette Gum, Nicotrol, Prostep patch, and Zyban) was associated with an increased likelihood of attempted and successful quitting.¹¹ In a study among smokers, increased recall of cessation methods—the most frequently recalled methods being the nicotine patch, nicotine gum, and Zyban or prescription medications—and perceived effectiveness of cessation aids were significantly associated with intentions to quit.¹² These studies suggest that advertising exposure and recall of cessation products could influence smokers’ quit intentions and successful quitting.

Of course, awareness is a necessary but not always sufficient condition for adoption of new technologies or behaviors.^{13,14} Diffusion of innovation and behavior change theories offer frameworks that help explain how awareness of e-cigarettes and beliefs about their harmfulness could influence cessation-related psychosocial factors (e.g., outcome expectancies, subjective norms, and self-efficacy to use e-cigarettes to quit smoking) and ultimately cessation behavior.^{13–16} For example, adoption of alternatives to traditional cigarette smoking such as e-cigarettes and other cessation products appear to be driven in part by perceptions of reduced harm as a relative advantage.^{17,18} However, there is an ongoing debate within the public health community about whether e-cigarettes are a viable alternative for harm reduction, and whether smokers are merely supplementing or truly replacing their smoking with e-cigarettes and achieving smoking cessation.

Although e-cigarettes are characterized as replacements for regular cigarettes or cessation aids for smokers,² the evidence on the harm-reduction potential of e-cigarettes as cessation products is mixed. For instance, clinical trials showed that e-cigarettes could help some smokers who are trying to quit to achieve abstinence rates comparable to those using nicotine patches.^{19–21} A national survey found that current e-cigarette users were more likely to have attempted to quit smoking in the previous year and tried to quit for longer than

24 hours compared with those who were not current e-cigarette users.²² Other studies suggest that e-cigarette use might interfere with successful quitting. One study reported that current smokers who tried using e-cigarettes did not differ from those who never tried e-cigarettes in their quit intentions or attempts in the past year.²³ In another survey, smokers who used e-cigarettes were more likely to reduce cigarette consumption, but the rates of successful quitting were not different from those who did not use e-cigarettes.²⁴ State quitline clients who had used e-cigarettes were significantly less likely to be tobacco-abstinent at follow-up compared with non-users.²⁵

Study Objectives

If e-cigarettes are in fact being viewed as a competitor or replacement for smoking cigarettes owing to a comparative advantage with regard to reduced harmfulness, then awareness and perceived harmfulness of e-cigarettes may be associated with cessation intentions or quit attempts among current smokers. Several national surveys indicate a steady increase in awareness of e-cigarettes between 2009 and 2012 among the general U.S. population.^{22–24,26–28} Given the rapidly evolving landscape in advertising and media coverage of e-cigarettes, the first objective of this study is to describe the prevalence of awareness and perceived harmfulness of e-cigarettes compared with regular cigarettes using a nationally representative survey of U.S. adults conducted between late 2012 and early 2013. There is also a gap in the literature regarding the impact of increased publicity and awareness about the purported harm-reduction potential of e-cigarettes on cessation outcomes. Therefore, the second objective of this study is to examine whether the awareness and perceived harmfulness of e-cigarettes are associated with quit attempts or intentions to quit among smokers.

Methods

Data Source

Data were obtained from the most recent round of the Health Information National Trends Survey (HINTS 4 Cycle 2) conducted between October 2012 and January 2013 by the National Cancer Institute.²⁹

Sample

The sampling frame included U.S. adults aged 18 years. For the first stage of the stratified sampling procedure, a sample of addresses was randomly selected from three strata: areas with high concentrations of minorities, areas with low concentrations of minorities, and counties comprising Central Appalachia. In the second stage, one adult was selected within each sampled household using the next birthday method. Minority respondents (Hispanics and African Americans) were oversampled. Data was collected using mailed self-administered questionnaires following a modified Dillman approach.³⁰ In this survey, 3,630 respondents completed the questionnaire, yielding an overall response rate of 39.97%, as computed using the RR2 formula from the American Association of Public Opinion Research.³¹ This response rate is comparable with previous HINTS iterations using address-based sampling strategies.³² Additional details of the survey design and administration are available elsewhere.³³

Measures

The survey asked the following question: *New types of cigarettes are now available called electronic cigarettes (also known as e-cigarettes or personal vaporizers). These products deliver nicotine through a vapor. Compared to smoking cigarettes, would you say that electronic cigarettes are...much less harmful/less harmful/just as harmful/more harmful/much more harmful/I've never heard of electronic cigarettes?*

Respondents who answered they have never heard of electronic cigarettes were coded as being unaware of e-cigarettes and all other responses were coded as being aware. Item non-response, multiple responses, and missing values were coded as missing ($n=143$). Perceived harmfulness among those who were aware of e-cigarettes was dichotomized into less harmful versus equally or more harmful because of the skewed distribution of the responses ($<3\%$ of respondents answered more harmful or much more harmful).

Smoking status was based on self-reported smoking, consistent with other research.^{23,26,27} Respondents who have never smoked up to 100 cigarettes in their lives were classified as non-smokers. Those who smoked at least 100 cigarettes in their lives but were currently not smoking at all were classified as former smokers. Those who smoked at least 100 cigarettes in their lives and were smoking daily or on some days were classified as current smokers.

Current smokers were asked if in the past year they have stopped smoking for 1 day or longer because they were trying to quit (past year quit attempt) and if they are seriously considering quitting smoking in the next 6 months (intention to quit).

Potential confounders included demographic variables (i.e., age, gender, race/ethnicity, and education level), health status, Census region, and household income.

Statistical Analyses

Data were analyzed between June and August 2013 with Stata, version 13 (StataCorp LP, College Station TX) using descriptive statistics and multiple logistic regression models. Prevalence and predictors of awareness of e-cigarettes were analyzed among the overall sample. For the distribution and predictors of perceived harmfulness of e-cigarettes, the analysis was restricted to respondents who were aware of e-cigarettes. To evaluate the harm-reduction potential of e-cigarettes, multiple logistic models assessed whether awareness and perceived harmfulness of e-cigarettes were associated with reporting past year quit attempts and the intention to quit among current smokers. All logistic regression models controlled for the aforementioned potential confounders and were weighted to the U.S. general adult population using full sample and replicate weights provided in the HINTS dataset.³³ These weights were necessary to adjust for the stratified sampling design, oversampling of certain subpopulations, and correction for non-response and non-coverage biases in order to ensure valid inferences from the responding sample to the general population.³³ Missing values on dependent variables were excluded from the analyses. To reduce potential bias, missing values for each predictor variable were treated as a separate category in the logistic regression models.³²

Results

Table 1 summarizes the sample characteristics (weighted to the general U.S. adult population). Most respondents were white, had some college or higher education, reported annual household income <\$75,000, were in good to excellent health, and did not smoke currently.

Prevalence and Correlates of E-Cigarette Awareness

Figure 1 shows that a steady increase in the awareness of e-cigarettes based on earlier national surveys among U.S. adults occurred from 16.4% in 2009 to 77.1% in the current study.^{23,26,27}

In the multivariate analysis, older participants aged 65–74 years and those aged ≥75 years had 62% and 82% lower odds of being aware of e-cigarettes, respectively, compared with young adults aged 18–34 years. Hispanic respondents had 62% lower odds of e-cigarette awareness than white respondents. Respondents with a college or higher education level had 1.75-fold increased odds of being aware than those with high school or lower education. Current and former smokers had 3.37- and 1.62-fold higher odds of being aware of e-cigarettes, respectively, than non-smokers (Table 2).

Prevalence and Correlates of Perceptions of E-Cigarette Harmfulness

Figure 2 shows that a slight decline in the proportion of current smokers who were aware of e-cigarettes and believed that e-cigarettes are less harmful than smoking regular cigarettes occurred from 84.7% in 2010 to 65.0% in the current study. Data for non-smokers and former smokers were not available from these earlier surveys for comparison with the current study.

Table 2 presents the multivariate analyses predicting beliefs about e-cigarette harms versus regular cigarettes. Older respondents had 38–72% lower odds of believing that e-cigarettes are less harmful than regular cigarettes compared with those aged 18–34 years. Those who had college or higher education had 2.06-fold higher odds than participants with high school or lower education of believing e-cigarettes were less harmful. Current smokers in the study had 2.53-fold higher odds of holding this belief compared with non-smokers.

Associations with Past Year Quit Attempts and Intention to Quit

Awareness was not associated with current smokers' quit attempts in the past year or their intention to quit in the next 6 months. Similarly, among those who were aware of e-cigarettes, perceived harmfulness of e-cigarettes compared with regular cigarettes was not associated with current smokers' quit attempts or intention to quit (Tables 3 and 4).

Discussion

Compared with earlier national surveys among U.S. adults, this study found a notable increase in public awareness of e-cigarettes since 2009. In the beginning of 2013, more than three in four respondents were aware of this novel product. The rise in awareness of e-cigarettes could reflect sharp increases in advertising expenditures by manufacturers,

availability in retail stores across the country, and presence in popular media in the time between the national surveys.² This trend suggests that e-cigarette awareness is increasingly prevalent and could be almost universal among U.S. adults in the near future. However, it is worth noting that there were differences in awareness across certain socioeconomic factors. Consistent with previous research, older respondents, those with lower education, and some minorities were not as aware of e-cigarettes^{22,23,26,27} These persistent disparities in awareness may be consequential for tobacco control efforts among vulnerable populations depending on whether e-cigarettes turn out to be, on balance, helpful or harmful regarding smoking uptake and cessation efforts.

The implications of increasingly widespread awareness of e-cigarettes in the general population are presently unclear. Despite the high levels of awareness, only 6% of U.S. adults reported ever using e-cigarettes.²⁷ Among smokers, ever use of e-cigarettes was estimated at 32% in 2012, but current e-cigarette use is relatively low at only 6%.²² Although one recent news report predicted that e-cigarette sales would soon reach \$1.7 billion,³⁴ this impressive figure represents only 1% of current regular cigarette sales in the U.S. The low usage levels do not presently support fears that e-cigarettes pose a serious public health threat that could unravel decades of tobacco control programs to de-normalize smoking. At the same time, claims that e-cigarettes are currently a major force in reducing harm from regular cigarette use at the population level also appear premature.

We compared current smokers' perceptions of reduced harm of e-cigarettes with earlier surveys and found a gradual decline in this belief from 84% in 2010 to 65% in 2013.²⁶ A formal analysis of this trend was not possible because of differences in study sampling. This apparent decline in smokers' beliefs about reduced harm of e-cigarettes compared with regular cigarettes is perplexing against the background of advertising and media messages touting e-cigarettes as safer alternatives and cessation aids. One possible explanation is that the increased media attention over the lack of FDA approval and regulation of this emerging tobacco product,^{34–36} injuries arising from e-cigarette-induced fires,^{37,38} and health concerns from toxic chemicals in e-cigarettes^{39–41} in recent years may have conveyed conflicting information about the relative safety of e-cigarette use.

In this study, we found that fewer former (50%) and never smokers (46%) were convinced that e-cigarettes are less harmful than regular cigarettes compared with current smokers (65%). These differences could be due to smokers being more exposed to marketing materials that frame the benefits and reduced harms of e-cigarettes because they are the target audience of e-cigarette marketing, or they could reflect greater attention to the topic among smokers because it has more personal relevance. However, we recommend continued monitoring of perceptions of e-cigarette harms among former and non-smokers for a few reasons. First, a qualitative study suggested that non-smokers who have no intention to initiate smoking may nevertheless be open to trying e-cigarettes, which could serve as a gateway to other tobacco products including cigarettes.⁴² Second, young adult former smokers in a community survey believed that e-cigarettes are less addictive than cigarettes; this may tempt former smokers to resume their smoking habits and result in relapse.⁴³

The current analyses showed that there were no significant associations between e-cigarette awareness or perceived harmfulness and smokers' intentions to quit or past-year quit attempts. These null findings suggest several possibilities. One potential interpretation is that adult smokers have not yet accepted e-cigarettes as a means to quit smoking. Even if they are aware of e-cigarettes and believe that they are less harmful, it is not sufficient to prompt them to consider quitting or try to quit smoking. For example, this relationship may be mediated through greater perceived efficacy of the product and moderated by perceived efficacy to quit smoking. A recent study reported that smokers in Hawaii who used e-cigarettes for quitting were more likely to have higher motivation and self-efficacy to quit and to achieve longer recent quit durations than non-users of e-cigarettes.⁴⁴ An alternative interpretation of the null associations is that it could be an artifact of the cross-sectional analyses. For example, it may be that there was insufficient time lag between smokers becoming aware of e-cigarettes and reduced harm and their decisions to try to quit smoking. It is also possible that there has not been enough exposure to information about e-cigarettes or related persuasive content that would influence smokers' cessation intentions or behaviors at this point in time. The null findings underscore the importance of further research to track smokers' exposure to e-cigarette information and their cessation behaviors over time, which could help to rule out these alternate explanations about insufficient time lag, lack of cumulative exposure to e-cigarette information, or lack of exposure to specific messages to use e-cigarettes for smoking cessation.

Study Limitations

There are several limitations to this study. First, the e-cigarette awareness measure did not directly ask if respondents were aware of e-cigarettes. However, the results from this study are in line with the trend of increasing e-cigarette awareness across various studies over time as shown in Figure 1. Currently, there is no standardized measure of e-cigarette awareness; prior surveys have all measured awareness using different survey items.^{22–24,26,27} Second, owing to the cross-sectional nature of the survey design, it is not possible to draw causal inferences. The study sample may not represent the U.S. adult population because of the low response rate. However, steps were taken to adjust for survey non-response and to weight the study sample to the general population. This study did not assess exposure to e-cigarette information in the media, the content of such exposures, e-cigarette usage, and other quit behaviors (e.g., reducing the number of cigarettes smoked per day) because these were not measured in the HINTS questionnaire. Future rounds of HINTS could consider including these items. We recommend further population-based research to assess smokers' beliefs and attitudes toward e-cigarettes as cessation aid devices, quit attempts, smoking reduction, and successful tobacco cessation among e-cigarette users in the U.S. In particular, it may be that as population-level e-cigarette use increases, a relationship among awareness, perceived harm, e-cigarette use, and smoking cessation could emerge.

Conclusions

This study found that awareness of e-cigarettes among U.S. adults increased over time compared with earlier surveys. It is uncertain whether increased population e-cigarette awareness and perceptions about reduced harm might play a role in encouraging smoking

cessation behaviors. However, public health professionals should systematically scrutinize the nature of marketing activities and media coverage of e-cigarettes, their impact on population awareness and perceptions of e-cigarettes, and how these factors may influence e-cigarette use and smoking prevalence in the U.S. population.

Acknowledgments

Andy Tan conducted this work while a postdoctoral fellow in the Center of Excellence in Cancer Communication Research at the University of Pennsylvania (supported by National Cancer Institute grant P20CA095856). Cabral Bigman conducted this work while a Yerby Postdoctoral Fellow at the Harvard School of Public Health and an assistant professor at the University of Illinois at Urbana-Champaign.

References

1. U.S. Food and Drug Administration Center for Tobacco Products. Public Health Focus - Electronic Cigarettes (e-Cigarettes). fda.gov/NewsEvents/PublicHealthFocus/ucm172906.htm.
2. Noel JK, Rees VW, Connolly GN. Electronic cigarettes: a new “tobacco” industry? *Tob Control*. 2011; 20(1):81. [PubMed: 20930060]
3. National Public Radio. E-Cigarettes Bring Smokers Back Inside, For Now. npr.org/2013/04/27/179490799/e-cigarettes-bring-smokers-back-inside-for-now.
4. Veppo, Inc. Veppo E-Cigarettes at 2013 Billboard Music Awards. veppocig.com/billboard-music-awards/.
5. Internet Movie Database. Demo Reel (Blu E-Cig National TV Commercial). IMDb.imdb.com/video/demo_reel/vi2066392089/.
6. Cobb NK, Brookover J, Cobb CO. Forensic analysis of online marketing for electronic nicotine delivery systems. *Tob Control*. 2013 In press.
7. Grana RA, Glantz SA, Ling PM. Electronic nicotine delivery systems in the hands of Hollywood. *Tob Control*. 2011; 20(6):425–426. [PubMed: 21659450]
8. Luo, C.; Zheng, X.; Zeng, DD., et al. Portrayal of electronic cigarettes on YouTube. In: Zeng, D.; Yang, CC.; Tseng, VS., et al., editors. *Smart health. Lecture notes in computer science*. Berlin, Heidelberg, Germany: Springer; 2013.
9. Paek H-J, Kim S, Hove T, Huh JY. Reduced harm or another gateway to smoking? Source, message, and information characteristics of e-cigarette videos on YouTube. *J Health Commun*. 2013 In press.
10. Etter J-F, Bullen C, Flouris AD, Laugesen M, Eissenberg T. Electronic nicotine delivery systems: a research agenda. *Tob Control*. 2011; 20(3):243–248. [PubMed: 21415064]
11. Avery, R.; Kenkel, D.; Lillard, DR.; Mathios, A. *Private profits and public health: does advertising smoking cessation products encourage smokers to quit?*. Cambridge MA: National Bureau of Economic Research; 2006.
12. Hammond D, McDonald PW, Fong GT, Borland R. Do smokers know how to quit? Knowledge and perceived effectiveness of cessation assistance as predictors of cessation behaviour. *Addiction*. 2004; 99(8):1042–1108. [PubMed: 15265101]
13. Rogers, EM. *Diffusion of innovations*. 5th ed.. New York: Free Press; 2003.
14. Weinstein ND. The precaution adoption process. *Health Psychol*. 1988; 7(4):355–386. [PubMed: 3049068]
15. Norman P, Conner M, Bell R. The theory of planned behavior and smoking cessation. *Health Psychol*. 1999; 18(1):89–94. [PubMed: 9925050]
16. DiClemente CC, Prochaska JO, Fairhurst SK, Velicer WF, Velasquez MM, Rossi JS. The process of smoking cessation: an analysis of precontemplation, contemplation, and preparation stages of change. *J Consult Clin Psychol*. 1991; 59(2):295–304. [PubMed: 2030191]
17. Rooke C, Amos A. News media representations of electronic cigarettes: an analysis of newspaper coverage in the UK and Scotland. *Tob Control*. 2013 In press.

18. Trumbo CW, Harper R. Use and perception of electronic cigarettes among college students. *J Am Coll Health*. 2013; 61(3):149–155.
19. Caponnetto P, Campagna D, Cibella F, et al. Efficiency and safety of an eElectronic cigAreTte (ECLAT) as tobacco cigarettes substitute: a prospective 12-month randomized control design study. *PLoS One*. 2013; 8(6):e66317. [PubMed: 23826093]
20. Polosa R, Caponnetto P, Morjaria JB, Papale G, Campagna D, Russo C. Effect of an electronic nicotine delivery device (e-Cigarette) on smoking reduction and cessation: a prospective 6-month pilot study. *BMC Public Health*. 2011; 11(1):786. [PubMed: 21989407]
21. Bullen C, Howe C, Laugesen M, et al. Electronic cigarettes for smoking cessation: a randomised controlled trial. *Lancet*. 2013; 382(9905):1629–1637. [PubMed: 24029165]
22. Zhu S-H, Gamst A, Lee M, Cummins S, Yin L, Zoref L. The use and perception of electronic cigarettes and snus among the U.S. population. *PLoS One*. 2013; 8(10):e79332. [PubMed: 24250756]
23. Regan AK, Promoff G, Dube SR, Arrazola R. Electronic nicotine delivery systems: adult use and awareness of the “e-cigarette” in the U.S. *Tob Control*. 2013; 22(1):19–23. [PubMed: 22034071]
24. Adkison SE, O'Connor RJ, Bansal-Travers M, et al. Electronic nicotine delivery systems: International Tobacco Control Four-Country Survey. *Am J Prev Med*. 2013; 44(3):207–215. [PubMed: 23415116]
25. Vickerman KA, Carpenter KM, Altman T, Nash CM, Zbikowski SM. Use of electronic cigarettes among state tobacco cessation quitline callers. *Nicotine Tob Res*. 2013; 15(10):1787–1791. [PubMed: 23658395]
26. Pearson JL, Richardson A, Niaura RS, Vallone DM, Abrams DB. e-Cigarette awareness, use, and harm perceptions in U.S. adults. *Am J Public Health*. 2012; 102(9):1758–1766. [PubMed: 22813087]
27. King BA, Alam S, Promoff G, Arrazola R, Dube SR. Awareness and ever use of electronic cigarettes among U.S. Adults, 2010–2011. *Nicotine Tob Res*. 2013; 15(9):1623–1627. [PubMed: 23449421]
28. Pepper JK, Brewer NT. Electronic nicotine delivery system (electronic cigarette) awareness, use, reactions and beliefs: a systematic review. *Tob Control*. 2013 In press.
29. Hesse BW, Moser RP, Rutten LJ, Kreps GL. The health information national trends survey: research from the baseline. *J Health Commun*. 2006; 11(1S):Svii–Sxvi.
30. Dillman, DA.; Smyth; Christian, LM.; Dillman, DA. Internet, mail, and mixed-mode surveys: the tailored design method. Hoboken NJ: Wiley & Sons; 2009.
31. The American Association for Public Opinion Research. Final disposition of case codes and outcome rates for surveys. 6th edition. Deerfield IL: American Association for Public Opinion Research; 2009. Standard definitions. www.aapor.org/Content/NavigationMenu/ResourcesforResearchers/StandardDefinitions/StandardDefinitions2009new.pdf.
32. Chou WS, Liu B, Post S, Hesse B. Health-related Internet use among cancer survivors: data from the Health Information National Trends Survey, 2003–2008. *J Cancer Surviv*. 2011; 5(3):263–270. [PubMed: 21505861]
33. Westat. Health Information National Trends Survey 4 (HINTS 4) Cycle 2 Methodology Report. Rockville MD: Westat; 2013. hints.cancer.gov/docs/HINTS_4_Cycle2_Methods_Report.pdf.
34. Elliott S. E-cigarette makers' ads echo tobacco's heyday. *The New York Times*. 2013 Aug 29. nytimes.com/2013/08/30/business/media/e-cigarette-makers-ads-echo-tobaccosheyday.html.
35. Koch W. Big Tobacco back in TV viewers' faces as FDA cuts in. *USA Today*. 2013 Sep 4. usatoday.com/story/news/nation/2013/09/03/big-tobacco-tv-ecigarette-ads/2608427/.
36. Kesmodel D. E-Cigarettes spark new smoking war. *The Wall Street Journal*. 2010 Aug 25. online.wsj.com/news/articles/SB10001424052748704557704575437710870116450.
37. Chen I. FDA summary of adverse events on electronic cigarettes. *Nicotine Tob Res*. 2013; 15(2): 615–616. [PubMed: 22855883]
38. Parker L, Green R. Small number of e-cigarette batteries explode. *NBC Chicago*. 2013 Oct 25. nbcchicago.com/investigations/Exploding-E-Cigarettes-229183561.html.
39. Goniewicz ML, Knysak J, Gawron M, et al. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tob Control*. 2013 In press.

40. Vardavas CI, Anagnostopoulos N, Kougias M, Evangelopoulou V, Connolly GN, Behrakis PK. Short-term pulmonary effects of using an electronic cigarette: impact on respiratory flow resistance, impedance, and exhaled nitric oxide. *Chest*. 2012; 141(6):1400–1406. [PubMed: 22194587]
41. Flouris AD, Chorti MS, Poulianiti KP, et al. Acute impact of active and passive electronic cigarette smoking on serum cotinine and lung function. *Inhal Toxicol*. 2013; 25(2):91–101. [PubMed: 23363041]
42. Choi K, Fabian L, Mottey N, Corbett A, Forster J. Young adults' favorable perceptions of snus, dissolvable tobacco products, and electronic cigarettes: findings from a focus group study. *Am J Public Health*. 2012; 102(11):2088–2093. [PubMed: 22813086]
43. Choi K, Forster J. Characteristics associated with awareness, perceptions, and use of electronic nicotine delivery systems among young U.S. Midwestern adults. *Am J Public Health*. 2013; 103(3):556–561. [PubMed: 23327246]
44. Pokhrel P, Fagan P, Little MA, Kawamoto CT, Herzog TA. Smokers who try e-cigarettes to quit smoking: findings from a multiethnic study in Hawaii. *Am J Public Health*. 2013; 103(9):e1–e6.

Figure 1.

Population awareness of e-cigarettes, 2009–2013

Note: Population awareness data from 2009 to 2011 were obtained from separate national surveys.^{23,26,27}

Figure 2.

Perceptions that e-cigarettes are less harmful than regular cigarettes among current smokers aware of e-cigarettes, 2010 and 2013

Note: Perceptions of e-cigarette harm for 2010 were obtained from separate national surveys.²⁶

Table 1

Characteristics of study sample: unweighted sample size and weighted percentages

	<i>n</i>	Weighted %
Age (years)		
18–34	529	29.8
35–49	845	25.8
50–64	1168	25.0
65–74	555	9.2
75	414	7.8
Missing	119	2.6
Gender		
Male	1390	47.7
Female	2172	50.5
Missing	68	1.8
Race/Ethnicity		
White	2043	61.6
African-American	496	10.0
Hispanic	511	13.7
Other ^a	208	6.5
Missing	372	8.2
Education		
High school or below	329	13.2
Completed high school	775	19.9
Some college	1057	36.8
College graduate or higher	1380	28.1
Missing	89	1.9
Annual household income		
<\$20,000	837	21.3
\$20,000–34,999	550	14.5
\$35,000–49,999	51	15.2
\$50,000–74,999	570	16.1
\$75,000	1043	30.2
Missing	119	2.6
Health status		
Excellent	382	11.2
Very good	1232	35.9
Good	1277	34.1
Fair	509	12.9
Poor	123	3.0
Missing	107	2.9
U.S. Census region		
Northeast	561	18.2

	<i>n</i>	Weighted %
Midwest	684	21.5
South	1542	37.1
West	843	23.1
Smoking status ^b		
Non-smoker	2052	58.1
Current	586	18.5
Former	939	22.4
Missing	53	1.0

^aParticipants who were categorized as “other” were those who identified as non-Hispanic American Indian or Alaskan, Asian, Native Hawaiian or other Pacific Islander, or multiple races. Smoking status was based on self-reported smoking consistent with other research.

^bRespondents who have never smoked up to 100 cigarettes in their lives were classified as non-smokers. Current smokers were those who smoked at least 100 cigarettes in their lives and were smoking daily or on some days. Former smokers were those who smoked at least 100 cigarettes in their lives but were currently not smoking at all.

Table 2

Point estimates and AORs for predicting e-cigarette awareness and perceived harmfulness

Predictor variables	Awareness of e-cigarettes (n=3487) ^a		Believe e-cigarettes are less harmful (n=2609) ^b	
	% (95% CI)	AOR (95% CI) ^c	% (95% CI)	AOR (95% CI) ^c
Overall	77.1 (74.5, 79.7)	—	50.7 (47.8, 53.7)	—
Age (years)				
18–34 (ref)	81.5 (74.7, 88.3)	1.00	60.1 (53.4, 66.8)	1.00
35–49	83.5 (80.3, 86.7)	1.00 (0.56, 1.79)	51.2 (45.5, 56.9)	0.62 * (0.41, 0.94)
50–64	78.6 (74.4, 82.8)	0.73 (0.41, 1.29)	47.3 (43.4, 51.1)	0.55 ** (0.38, 0.81)
65–74	67.8 (61.9, 73.7)	0.38 ** (0.21, 0.69)	42.5 (36.0, 49.0)	0.48 ** (0.30, 0.79)
75	46.6 (40.1, 53.2)	0.18 *** (0.10, 0.31)	27.5 (19.7, 35.3)	0.28 *** (0.16, 0.47)
Gender				
Male (ref)	79.6 (76.0, 83.2)	1.00	58.4 (53.4, 63.4)	1.00
Female	75.6 (72.3, 78.8)	0.82 (0.65, 1.02)	42.9 (39.2, 46.5)	0.84 (0.50, 1.40)
Race/ethnicity				
White (ref)	81.3 (78.7, 84.0)	1.00	53.8 (50.1, 57.5)	1.00
African American	78.1 (73.1, 83.0)	0.82 (0.52, 1.29)	42.2 (32.0, 52.4)	0.61 (0.35, 1.07)
Hispanic	64.3 (54.9, 73.8)	0.38 *** (0.24, 0.62)	53.5 (44.1, 63.0)	0.95 (0.64, 1.41)
Other	72.3 (53.2, 91.4)	0.60 (0.21, 1.73)	42.9 (30.8, 54.9)	0.57 (0.31, 1.08)
Education				
High school or below (ref)	65.0 (56.7, 73.2)	1.00	41.0 (28.2, 53.9)	1.00
Completed high school	72.8 (67.3, 78.3)	1.15 (0.69, 1.93)	45.9 (37.8, 54.0)	1.23 (0.65, 2.34)
Some college	80.3 (75.9, 84.8)	1.54 (0.90, 2.62)	51.4 (46.5, 56.3)	1.42 (0.77, 2.65)
College graduate or higher	81.8 (79.3, 84.2)	1.75 * (1.07, 2.87)	57.1 (52.2, 62.0)	2.06 * (1.06, 4.02)
Annual household income				
<\$20,000 (ref)	72.3 (65.4, 79.3)	47.1 (38.3, 55.9)	1.00	
\$20,000–34,999	76.5 (69.6, 83.3)	1.31 (0.83, 2.06)	49.7 (42.8, 56.7)	1.22 (0.77, 1.91)
\$35,000–49,999	72.4 (66.2, 78.7)	0.93 (0.54, 1.59)	48.0 (38.5, 57.4)	1.04 (0.63, 1.70)
\$50,000–74,999	82.5 (78.4, 86.5)	1.26 (0.73, 2.19)	52.6 (46.0, 59.3)	1.18 (0.74, 1.89)
\$75,000	81.2 (78.1, 84.3)	1.10 (0.64, 1.91)	54.2 (48.8, 59.6)	1.29 (0.83, 2.00)
Health status				
Excellent (ref)	75.2 (66.2, 84.2)	1.00	49.4 (37.3, 61.6)	1.00
Very good	81.2 (77.7, 84.7)	1.38 (0.79, 2.42)	52.4 (46.9, 58.0)	1.21 (0.67, 2.19)
Good	77.4 (73.7, 81.1)	1.24 (0.72, 2.11)	47.7 (42.6, 52.7)	1.07 (0.58, 1.98)
Fair	65.2 (53.7, 76.7)	0.76 (0.33, 1.76)	53.7 (43.5, 63.8)	1.75 (0.89, 3.44)
Poor	78.2 (66.8, 89.7)	1.28 (0.54, 3.03)	44.5 (20.5, 68.6)	0.96 (0.20, 4.56)
U.S. Census region				
Northeast (ref)	75.9 (70.1, 81.7)	1.00	51.2 (43.0, 59.3)	1.00
Midwest	79.6 (74.4, 84.7)	1.15 (0.72, 1.81)	48.7 (41.2, 56.3)	0.86 (0.55, 1.35)
South	78.1 (74.7, 81.6)	1.19 (0.77, 1.87)	52.4 (47.0, 57.9)	1.06 (0.73, 1.52)

Predictor variables	Awareness of e-cigarettes (<i>n</i> =3487) ^a		Believe e-cigarettes are less harmful (<i>n</i> =2609) ^b	
	% (95% CI)	AOR (95% CI) ^c	% (95% CI)	AOR (95% CI) ^c
West	74.2 (67.9, 80.4)	1.04 (0.67, 1.62)	49.6 (42.1, 57.0)	0.85 (0.54, 1.34)
Smoking status				
Non-smoker (ref)	73.0 (68.6, 77.3)	1.00	45.9 (42.1, 49.6)	1.00
Current	88.6 (84.4, 92.8)	3.37*** (2.02, 5.63)	65.0 (57.6, 72.4)	2.53*** (1.72, 3.72)
Former	78.3 (74.3, 82.3)	1.62** (1.16, 2.26)	49.5 (43.6, 55.3)	1.33 (0.99, 1.78)

Note: Boldface indicates statistical significance.

^a Missing cases for awareness of e-cigarettes variable (*n*=143) were excluded from analysis.

^b The model predicting belief that e-cigarettes are less harmful included only respondents who are aware of e-cigarettes.

^c ORs are adjusted for all other covariates in the model.

*
p<0.05;

**
p<0.01;

p<0.00005.

Table 3Predicting past-year quit attempt and intention to quit with e-cigarette awareness^a

Predictor variable	Past-year quit attempt ^b		Intention to quit ^c	
	% (95% CI)	AOR (95% CI) ^d	% (95% CI)	AOR (95% CI) ^d
Awareness of e-cigarettes				
Never heard of e-cigarettes (ref)	43.0 (33.9, 52.1)	1.00	68.0 (50.5, 85.6)	1.00
Aware of e-cigarettes	47.4 (42.7, 52.1)	1.25 (0.54, 2.91)	66.4 (58.9, 74.0)	0.96 (0.32, 2.92)

^aThese analyses included only current smokers.^bMissing cases for quit attempt in the past year ($n=217$) are excluded from these analyses.^cMissing cases for intention to quit ($n=8$) are excluded from these analyses.^dORs are adjusted for gender, race/ethnicity, education level, household income, health status, and Census region.

Table 4Predicting past-year quit attempt and intention to quit with perceived e-cigarette harmfulness^a

Predictor variable	Past-year quit attempt ^b		Intention to quit ^c	
	% (95% CI)	AOR (95% CI) ^d	% (95% CI)	AOR (95% CI) ^d
Perceived harmfulness				
Equally or more harmful than cigarettes (ref)	47.0 (39.8, 54.2)	1.00	69.4 (56.5, 82.3)	1.00
Less harmful than cigarettes	47.7 (41.0, 54.5)	0.48 (0.21, 1.09)	64.9 (54.9, 74.8)	0.95 (0.42, 2.12)

^aThese analyses included only current smokers who are aware of e-cigarettes.^bMissing cases for quit attempt in the past year ($n=131$) are excluded from these analyses.^cMissing cases for intention to quit ($n=3$) are excluded from these analyses.^dORs are adjusted for gender, race/ethnicity, education level, household income, health status, and Census region.