Published in final edited form as:

Mol Cancer Ther. 2012 December ; 11(12): 2549–2555. doi:10.1158/1535-7163.MCT-12-0473.

It's About Time: *Lessons For Solid Tumors From Chronic Myelogenous Leukemia Therapy*

Jason R. Westin¹ and Razelle Kurzrock²

¹Department of Lymphoma & Myeloma, The University of Texas MD Anderson Cancer Center, Houston, USA

²Department of Investigational Cancer, Therapeutics (Phase I Clinical Trials Program) The University of Texas MD Anderson Cancer Center, Houston, USA

Abstract

The use of imatinib in chronic myelogenous leukemia (CML) transformed the disease, rapidly changing the median survival from 4 years to at least 20 years. In this review, we outline the causes of this revolution, including the identification of a critical driving molecular aberration--*BCR-ABL*--and the development of a potent and specific inhibitor--imatinib. Equally important was the timing of the targeted therapy, specifically its administration to patients with newly diagnosed disease. In solid tumors, targeted therapies are often both developed and used in metastatic malignancies after conventional approaches have failed.

We postulate that this strategy is similar to using imatinib in blast crisis CML, where response rates are <15%, all patients relapse, and median survival remains only about one year. We hypothesize that the imatinib-led revolution in CML, including the critically important factor of timing, may be applicable to other cancers as well. Therefore, it will be important to utilize promising targeted therapies in the earliest phases of biomarker-defined solid tumors, prior to metastatic progression, to determine if outcomes can be significantly improved, and hence establish if the success of imatinib in CML is an anomaly or a paradigm.

Keywords

Cancer; chronic myelogenous leukemia; early stage cancer; imatinib; targeted therapy; time

INTRODUCTION

When torrential water tosses boulders, it is because of its momentum. When the strike of a hawk breaks the body of its prey, it is because of timing.

Sun Tzu, The Art of War

Life is really simple, but we insist on making it complicated.

Corresponding author: Dr. Razelle Kurzrock, M.D., Department of Investigational Cancer Therapeutics (Phase I Clinical Trials, Program) The University of Texas MD Anderson Cancer Center, 1515 Holcombe Blvd, Unit 0455, P.O. Box 301402, Houston, TX 77230-1402, USA, 1-713-794-1226 (telephone), 1-713-563-0566 (fax), rkurzroc@mdanderson.org.

Confucius

This thing all things devours:

Birds, beasts, trees, flowers;

Gnaws iron, bites steel;

Grinds hard stones to meal;

Slays king, ruins town,

And beats high mountain down

J.R.R Tolkien, *The Hobbit*, Gollum's Riddle About Time

The use of imatinib in chronic myelogenous leukemia (CML) transformed the disease, rapidly changing the median overall survival from four years to an estimated 19–25 years and counting. This revolution occurred due to several factors, including the identification of a critical driving molecular aberration--*BCRABL*--and the development of an effective and specific inhibitor--imatinib. However, these two factors alone were not transformative. Equally important was the timing of the use of the targeted therapy, specifically its administration to patients with newly diagnosed disease. Indeed, if therapy of CML with imatinib had been restricted to advanced disease, imatinib may have been regarded as having moderate efficacy, since response rates in blast crisis are <15%, all patients relapse, and median survival remains approximately one year.

Like blast crisis CML, metastatic solid malignancies are significantly more complex genetically and challenging to effectively treat with targeted therapies than newly diagnosed early stage disease. Responses can be achieved, but they are generally transient. We hypothesize that the imatinib-led revolution in CML, including the critically important factor of timing, may be applicable to other cancers as well. Therefore, it will be important to utilize promising targeted therapies in the earliest phases of biomarker-defined solid tumors, prior to metastatic progression, to determine if outcomes can be significantly improved, and hence establish if the success of imatinib in CML is an anomaly or a paradigm.

THE ERA OF TARGETED THERAPY

The era of molecularly targeted cancer therapy ushered in by imatinib began with incredible expectations. Magazine covers proclaimed “A New Hope For Cancer” and news reports questioned if the “magic bullet” could represent a long-awaited turning point in the war on cancer.(1) Imatinib specifically and effectively targets cells harboring the 9;22 translocation, the *sine qua non* of CML, and is considered a modern day version of Erhlich's “magic bullet” (Figure 1).(2–4) If left untreated, CML inevitably progresses from chronic to accelerated phase, and finally to end-stage blast crisis, exemplifying the malignant evolution believed to occur in most cancers.(2) Prior to imatinib, therapies for CML were mainly “cosmetic”. Hydroxyurea normalized blood counts but had no impact on survival. Interferon-based treatments yielded complete cytogenetic responses (absence of the 9;22 translocation in the bone marrow) in <15% of newly-diagnosed patients and a median

survival of ~ 4 years.(5,6) In contrast, in newly diagnosed CML, imatinib achieves complete cytogenetic responses in ~ 70% of individuals, and the median survival has increased to an estimated 19 – 25 years (Table 1).(7–20) Imatinib has unequivocally revolutionized the treatment of CML, transforming a uniformly fatal disease into one easily managed with non-toxic agents.

In the eleven years since the FDA approval of imatinib, dozens of targeted agents have been approved and many more are currently under investigation, confirming the dawn of a new era in oncology. However, few if any have lived up to the initial high expectations that a dramatic shift in solid tumor patient survival would occur, as it did for CML.

IMPACT OF TARGETED THERAPY IN SOLID TUMORS

The development of imatinib in CML has been described as a paradigm for targeted therapy. (21) First, a target critical to the survival of the cancer was identified. Second, a potent, specific, and relatively non-toxic inhibitor was developed. Yet, despite heroic efforts to emulate the imatinib in CML story in solid tumors, most targeted therapies have modest effects.(22) Targets critical to the survival of other cancers, such as *EGFR*, *KIT*, *BRAF*, and *ALK*, have been identified and targeted by relatively specific inhibitors (Table 2). (15,16,23–30) Gastrointestinal stromal tumor (GIST), the most common mesenchymal tumor of the gastrointestinal track, is notoriously chemotherapy-refractory, but responds to imatinib. GIST has driving mutations in *KIT* or *PDGFR-α* in the vast majority of cases, which predict for response to imatinib.(31) In newly diagnosed and resected high risk GIST, adjuvant imatinib results in a significant recurrence-free interval, but the disease often returns when imatinib is stopped.(32) The implication of this finding, similar to disease recurrence after cessation of imatinib in CML, is that a sub-clinically detectable clone has a therapeutically-dependent dormancy and thus may require continuous exposure to an inhibitor or novel strategies to eliminate the dormant clone.(33) Of note, CML that recurred after imatinib cessation was responsive to resumption of therapy in all patients treated on the STIM (Stop Imatinib) trail. (33)

Many solid tumors, like small cell lung and bladder cancers, do not yet have a known driving aberration(s) amenable to current targeted therapies despite significant genetic derangements found in tumor samples. As our understanding of the complex signaling networks that drive these cancers improves, it is possible that even cancers currently considered a “black box” will be amenable to therapies targeting the yet undefined aberration-associated critical survival pathways.

Unlike CML and resected GIST, targeted therapies for advanced solid tumors may occasionally demonstrate evidence of dramatic disease regression, even in extensively pre-treated patients, but often prove fleeting.(24,34–36) For example, nearly 90% of patients achieved tumor regression in the phase I trials of vemurafenib in *BRAF*-mutant melanoma and crizotinib in *ALK*-rearranged lung cancer.(24,26) These responses are generally achieved with minimal side effects, especially compared with cytotoxic chemotherapy. However, they are rarely durable. The “Lazarus” restaging images featured in publications are often from patients who still die of their disease (Figure 2). These remarkable but

transient responses confirm the continued relevance of the target to the survival of cancer cells in at least a portion of patients with advanced disease.

If the target is present and the drug active, why are these dramatic responses infrequent and impermanent? Numerous models of targeted therapy resistance demonstrate the activation of alternate survival pathways, mutations in the target of the inhibitor, or both.(37–40) These mechanisms of resistance are also found in blast crisis CML, and are thought to explain its persistently poor outcomes.(41) In hindsight, it is now clear that the timing of targeted therapy in CML, specifically its use prior to the development of complex resistance mechanisms, was critical in the transformation of disease outcomes.

THE SIGNIFICANCE OF TIMING

The imatinib in CML paradigm speaks to the critical nature of timing of targeted therapy: patients with accelerated phase and blast crisis CML do not receive the same benefit from Bcr-Abl inhibition as do newly-diagnosed chronic phase CML patients. Furthermore, there is a precipitous fall in the benefit even patients with previously treated chronic-phase CML receive compared to those with newly diagnosed disease (Table 1). In 2001, simultaneous landmark publications of imatinib in chronic phase and blast crisis CML (along with relapsed t(9;22)+ acute lymphoblastic leukemia) demonstrated striking differences in outcome.(7,14) In newly-diagnosed chronic phase CML, imatinib has shown an astounding 65% to 74% rate of complete cytogenetic remission. In previously treated chronic phase, this rate drops to 13 to 40%. With transformation to blast crisis CML, only 0 to 8% of patients achieve complete cytogenetic responses, and these responses are transient (Table 1). Although the efficacy of imatinib in blast crisis CML is disappointing, the fact that late-stage disease could respond provided proof-of principle for the utility of imatinib, and demonstrated the continued importance of Bcr-Abl activity in at least a portion of blast crisis CML.(7) However, even today, median survival of patients with blast crisis remains at less than one year. Transforming outcomes in CML required moving imatinib therapy to newly diagnosed patients.

As CML progresses, much of the initial resistance to imatinib is due to mutations in the kinase domain of *BCR-ABL*. The resulting resistance to imatinib led to the development of a second generation of tyrosine kinase inhibitors (TKIs), dasatinib and nilotinib. Both agents, now FDA-approved for newly diagnosed and relapsed CML, have demonstrated superior efficacy, at least molecularly, to imatinib in front-line phase III trials (Table 1).(18,19) However, in accelerated phase and blast crisis CML, the efficacy of second generation TKIs is essentially equivalent to imatinib.(7–11) This may be due to other genetic changes supplanting the role of Bcr-Abl in late-stage disease. Despite superior efficacy in newly diagnosed chronic phase CML, dasatinib and nilotinib are no match for CML once transformation has occurred.

MOVING FORWARD

IS ADVANCED CANCER ANALOGOUS TO ADVANCED CML?

In 1957, Foulds defined neoplastic progression as “the acquisition of permanent, irreversible changes” in a neoplasm.(42) Three years later, Nowell and Hungerford observed a minute chromosome in CML, the first genetic example of a “permanent, irreversible change” in cancer.(43)

Advanced cancers have multiple genetic derangements that accumulate as subsequent generations of tumor cells develop, selected for the advantages they confer on sub-populations.(44) Metastatic lesions are often significantly heterogeneous compared to their parent lesions, including in clinically utilized biomarkers.(45) Compared with CML, even newly diagnosed localized solid tumors are considered genetically complex, with multiple sub-populations with unique biological characteristics.(46,47) Indeed, a rapid autopsy study in pancreatic cancer found that mutations present in metastatic disease were present in subclones of the original primary tumor.(48) However, even the “genetically simple” chronic phase CML, if left untreated, will eventually develop therapeutic resistance and increasing genetic complexity, remarkably similar to metastatic solid tumors.(49,50) Newly diagnosed localized solid tumors may not be as genetically simple as chronic phase CML, but are certainly less complex than refractory metastatic solid tumors.

Based on both biologic and clinical observations, by the time a solid tumor metastasizes, it is likely analogous to accelerated phase or blast crisis CML. Therefore, based on the CML paradigm, even newly diagnosed metastatic disease is too late for targeted agents to achieve their maximal effect.

IS THIS PARADIGM APPLICABLE IN OTHER CANCERS?

The CML paradigm supports three conclusions that warrant exploration in the context of solid tumors.

1. The evaluation of a targeted therapy in advanced disease is valuable for proof-of-principle, but may dramatically underestimate the efficacy in early-stage disease—Patients with either advanced CML or metastatic solid tumors respond to matched targeted therapy, albeit usually without achieving complete or durable remissions (Table 2). Similar to the situation in CML, response rates decline with progression in solid tumors. For instance, lapatinib, a dual EGFR-erbB2 TKI, is FDA approved to treat advanced or metastatic HER2+ breast cancer. The response rates decline from 24% to 6.9% when lapatinib is used to treat patients with previously untreated vs. previously treated metastatic HER2+ breast cancer.(28,51) Although previously untreated and treated metastatic breast cancer are both likely analogous to advanced CML, they serve as an example of the further degradation of outcomes with increasingly advanced disease. It is unclear how much the efficacy of targeted agents would improve if they were utilized in newly diagnosed, localized disease. The innovative I-SPY2 breast cancer trial aims to evaluate neoadjuvant targeted agents along with standard chemotherapy in newly diagnosed localized disease, and develop predictive biomarkers to optimally match patients and therapy.

2. Compared to current standard therapies, drugs with superior efficacy in early-stage disease may be equivalent in late-stage disease—Dasatinib and nilotinib display superior molecular outcomes in chronic phase CML, but achieve similar dismal results as imatinib in blast crisis CML (Table 1). The lack of efficacy of the more potent TKIs in late stage CML speaks to the complexity of the disease after transformation occurs. The *BCR-ABL* translocation remains present, but inhibitors have little effect on survival. It is plausible that this phenomenon may also apply in solid tumors. Due to the significant financial cost of drug development, it is unfortunately possible that drugs which are superior pre-clinically but essentially equivalent in relapsed/refractory patients may have their development halted, despite the possibility they would prove superior in newly diagnosed patients.

3. The optimal time to employ the most potent targeted therapies is early in the course of the disease, prior to transformation—In CML, the increases in survival would be much less significant had investigators stayed focused on advanced disease, or even later chronic phase disease (Table 1). The lesson learned in CML is that, for optimal results, matched targeted therapy must be given to newly diagnosed patients in order to prevent progression, rather than to disease which has already progressed. Although there is evidence that remarkable disease regression may occasionally occur with targeted therapy in metastatic solid tumors, durable responses are the exception.(22,24,26) Metastatic solid tumors are generally considered incurable with cytotoxic chemotherapies: it is not unreasonable to extrapolate the same expectation to targeted therapies. It is therefore worth investigating whether or not both response rates and duration would improve if targeted agents were used in the earliest phases of solid tumors, before development of advanced disease.

CHALLENGES IN TREATING NEWLY DIAGNOSED CANCER

The predominant approach to early-stage solid tumors is surgical resection, perhaps with adjuvant cytotoxic chemotherapy or radiation. If the disease recurs, targeted agents previously evaluated in relapsed disease may be employed, but often patients are treated with cytotoxic chemotherapy alone (e.g., erlotinib, an EGFR inhibitor, is approved to treat non-small cell lung cancer that has failed one prior chemotherapy regimen).

A concern of potential harm emerges when considering changing this treatment paradigm to include targeted agents earlier in the disease course, especially if cure is possible with surgery, cytotoxic chemotherapy, and/or radiation therapy. These standard therapies have known efficacy, but also result in serious sequelae that impact quality of life.(52) Moving forward, oncologists should aim to eliminate cancer without permanently disrupting the lives of their patients. Early-stage solid tumors are more genetically complex than early-stage CML and may contain significant heterogeneity, and thus utilizing targeted therapies in lieu of surgery may pose a significant risk of harm to patients with potentially curable cancers. If a portion of a small primary tumor harbors *de novo* resistance to a targeted therapy, metastatic disease may result and prove difficult to control. However, it is conceivable that with improved knowledge of disease-critical aberrations and greater technology to account for tumor heterogeneity, it may eventually be possible to give

targeted therapy to biomarker-selected patients with newly diagnosed, early-stage solid tumors in lieu of current “curative” therapies. Initially, this approach would be most appropriate for patients who are unfit for standard therapies (e.g., high risk of peri-operative mortality). If these patients receive benefit greater than expected based on initial trials in relapsed or refractory patients, it would support the further investigation of targeted therapy in frontline cancer therapeutics. Other investigators have treated borderline resectable renal cell carcinomas with neoadjuvant targeted agents typically reserved for advanced disease, and demonstrated efficacy in a retrospective series.(53)

Less controversially, newly diagnosed patients with localized disease could receive standard treatments along with neoadjuvant targeted agents previously found efficacious and tolerable in relapsed patients. Examples of this approach include the innovative I-SPY2 breast cancer trial. Another approach would be to add tumor-specific targeted therapy as an adjuvant to patients rendered free of disease who remain at high-risk for early relapse, similar to imatinib in GIST or trastuzumab in HER2+ breast cancer.(32,54)

An additional essential step to long-term disease control or cure with targeted therapy is identification of biomarkers predictive of response to targeted therapy. The neoadjuvant breast cancer I-SPY 1 and 2 clinical trials aim to develop biomarkers correlating with pathologic response and adaptively randomize patients to targeted therapies with a higher likelihood of success.(55,56) These innovative and informative clinical trial designs can facilitate the rapid translation of new targeted agents and disease knowledge from the bench to the therapy of early stage cancers.

CONCLUSION

It is estimated that 12.7 million cancer diagnoses and 7.6 million cancer deaths occurred worldwide in 2008 alone.(57) In 2030, these numbers are projected to increase to 21.3 million new cases and 13.1 million deaths.(58) Clearly, new innovative strategies are needed to deal with this problem.

A critical question is whether the lessons learned from the dramatic shift in CML outcomes attained with imatinib when used in the earliest phases of the disease are applicable to any solid cancers. Administering potent, specific inhibitors to patients with advanced disease has yielded occasional impressive responses and served as a proof-of-principle in both CML and solid tumors (Tables 1 and 2). However, the full therapeutic potential of these inhibitors in CML, which have transformed median overall survival from 4 years to 19 – 25 years, was not realized until they were used in newly diagnosed patients. Leukemia investigators found the key to improved survival was preventing, rather than treating, progression. Late-stage CML has significant similarities, both in genetic complexity and therapeutic responses, to metastatic disease in solid tumors.

It is possible that after a localized solid tumor is established, an agent targeting a driver molecular aberration may not be sufficient to control the cancer, even if employed immediately. However, even if dramatic and durable effects are not achieved with targeted therapy in newly diagnosed solid tumors, their efficacy should exceed what is currently

observed in the relapsed or refractory setting. Furthermore, unraveling resistance is likely to be a far less herculean task early in the disease than when multiple events of molecular evolution have occurred in its later stages.

In conclusion, although scientific knowledge regarding cancer biology, genomics, and complex networks is advancing rapidly, successful translation to the clinic lags behind.⁽⁵⁹⁾ To lessen this gap, a third factor may need to be added to the original description of CML as a paradigm for targeted therapy: 1) identification of the crucial driving molecular aberration, 2) development of potent, specific inhibitors, and importantly 3) use of the inhibitors early enough in the disease course to allow for maximum efficacy. To date, it has been assumed that the biological properties of CML are unique, and thus amenable to therapy in a way unheard of for solid tumors. Yet, fundamentally, the response of patients with advanced CML and those with metastatic solid tumors to matched target therapy are similar: only a minority of patients achieve remission, all relapse, and survival increases are measured in months, not years. The treatment of CML was revolutionized, not by a targeted therapy alone, but by giving it to patients with newly diagnosed disease in order to prevent progression to blast crisis. This strategy has not been fully explored in solid tumors. In order to determine if CML is indeed a biological exception or whether the lessons learned from CML are a paradigm for success, it will be necessary to administer matched targeted therapy to patients with newly diagnosed, localized solid tumors.

Acknowledgments

None

Grant Support: No funding was used to support this paper

References

1. Lemonick M, Park A. New Hope For Cancer. TIME. 2001 May 28. 2001.
2. Kurzrock R, Gutterman JU, Talpaz M. The molecular genetics of Philadelphia chromosome-positive leukemias. N Engl J Med. 1988; 319:990–998. [PubMed: 3047582]
3. Druker BJ, Tamura S, Buchdunger E, Ohno S, Segal GM, Fanning S, et al. Effects of a selective inhibitor of the Abl tyrosine kinase on the growth of Bcr-Abl positive cells. Nat Med. 1996; 2:561–566. [PubMed: 8616716]
4. Strebhardt K, Ullrich A. Paul Ehrlich's magic bullet concept: 100 years of progress. Nat Rev Cancer. 2008; 8:473–480. [PubMed: 18469827]
5. Kantarjian H, Talpaz M, O'Brien S, Giles F, Faderl S, Verstovsek S, et al. Survival benefit with imatinib mesylate therapy in patients with accelerated-phase chronic myelogenous leukemia--comparison with historic experience. Cancer. 2005; 103:2099–2108. [PubMed: 15830345]
6. Björkholm M, Ohm L, Eloranta S, Derolf Å, Hultcrantz M, Sjöberg J, et al. Success Story of Targeted Therapy in Chronic Myeloid Leukemia: A Population-Based Study of Patients Diagnosed in Sweden From 1973 to 2008. J Clin Oncol. 2011; 29:2514–2520. [PubMed: 21576640]
7. Druker BJ, Sawyers CL, Kantarjian H, Resta DJ, Reese SF, Ford JM, et al. Activity of a Specific Inhibitor of the BCR-ABL Tyrosine Kinase in the Blast Crisis of Chronic Myeloid Leukemia and Acute Lymphoblastic Leukemia with the Philadelphia Chromosome. N Engl J Med. 2001; 344:1038–1042. [PubMed: 11287973]
8. Kantarjian HM, Cortes J, O'Brien S, Giles FJ, Albitar M, Rios MB, et al. Imatinib mesylate (STI571) therapy for Philadelphia chromosome-positive chronic myelogenous leukemia in blast phase. Blood. 2002; 99:3547–3553. [PubMed: 11986206]

9. Sawyers CL, Hochhaus A, Feldman E, Goldman JM, Miller CB, Ottmann OG, et al. Imatinib induces hematologic and cytogenetic responses in patients with chronic myelogenous leukemia in myeloid blast crisis: results of a phase II study. *Blood*. 2002; 99:3530–3539. [PubMed: 11986204]
10. Cortes J, Kim DW, Raffoux E, Martinelli G, Ritchie E, Roy L, et al. Efficacy and safety of dasatinib in imatinib-resistant or -intolerant patients with chronic myeloid leukemia in blast phase. *Leukemia*. 2008; 22:2176–2183. [PubMed: 18754032]
11. Kantarjian H, Giles F, Wunderle L, Bhalla K, O'Brien S, Wassmann B, et al. Nilotinib in Imatinib-Resistant CML and Philadelphia Chromosome–Positive ALL. *N Engl J Med*. 2006; 354:2542–2551. [PubMed: 16775235]
12. Talpaz M, Silver RT, Druker BJ, Goldman JM, Gambacorti-Passerini C, Guilhot F, et al. Imatinib induces durable hematologic and cytogenetic responses in patients with accelerated phase chronic myeloid leukemia: results of a phase 2 study. *Blood*. 2002; 99:1928–1937. [PubMed: 11877262]
13. Guilhot F, Apperley J, Kim D-W, Bullorsky EO, Baccarani M, Roboz GJ, et al. Dasatinib induces significant hematologic and cytogenetic responses in patients with imatinib-resistant or -intolerant chronic myeloid leukemia in accelerated phase. *Blood*. 2007; 109:4143–4150. [PubMed: 17264298]
14. Druker BJ, Talpaz M, Resta DJ, Peng B, Buchdunger E, Ford JM, et al. Efficacy and Safety of a Specific Inhibitor of the BCR-ABL Tyrosine Kinase in Chronic Myeloid Leukemia. *N Engl J Med*. 2001; 344:1031–1037. [PubMed: 11287972]
15. O'Brien SG, Guilhot F, Larson RA, Gathmann I, Baccarani M, Cervantes F, et al. Imatinib compared with interferon and low-dose cytarabine for newly diagnosed chronic-phase chronic myeloid leukemia. *N Engl J Med*. 2003; 348:994–1004. [PubMed: 12637609]
16. Deininger M, O'Brien SG, Guilhot F, Goldman JM, Hochhaus A, Hughes TP, et al. International Randomized Study of Interferon Vs STI571 (IRIS) 8-Year Follow up: Sustained Survival and Low Risk for Progression or Events in Patients with Newly Diagnosed Chronic Myeloid Leukemia in Chronic Phase (CML-CP) Treated with Imatinib. *ASH Annual Meeting Abstracts*. 2009; 114:1126.
17. Hochhaus A, Baccarani M, Deininger M, Apperley JF, Lipton JH, Goldberg SL, et al. Dasatinib induces durable cytogenetic responses in patients with chronic myelogenous leukemia in chronic phase with resistance or intolerance to imatinib. *Leukemia*. 2008; 22:1200–1206. [PubMed: 18401416]
18. Kantarjian H, Shah NP, Hochhaus A, Cortes J, Shah S, Ayala M, et al. Dasatinib versus Imatinib in Newly Diagnosed Chronic-Phase Chronic Myeloid Leukemia. *N Engl J Med*. 2010; 362:2260–2270. [PubMed: 20525995]
19. Saglio G, Kim D-W, Issaragrisil S, le Coutre P, Etienne G, Lobo C, et al. Nilotinib versus Imatinib for Newly Diagnosed Chronic Myeloid Leukemia. *N Engl J Med*. 2010; 362:2251–2259. [PubMed: 20525993]
20. Reed SD, Anstrom KJ, Li Y, Schulman KA. Updated Estimates of Survival and Cost Effectiveness for Imatinib versus Interferon-[alpha] Plus Low-Dose Cytarabine for Newly Diagnosed Chronic-Phase Chronic Myeloid Leukaemia. *Pharmacoeconomics*. 2008; 26:435–446. [PubMed: 18429659]
21. Druker BJ, David A. Karnofsky Award lecture. Imatinib as a paradigm of targeted therapies. *J Clin Oncol*. 2003; 21:239s–245s. [PubMed: 14645403]
22. Stewart DJ, Kurzrock R. Cancer: the road to Amiens. *J Clin Oncol*. 2009; 27:328–333. [PubMed: 19064964]
23. Kubota T. Gastrointestinal stromal tumor (GIST) and imatinib. *International J Clin Oncol*. 2006; 11:184–189.
24. Flaherty KT, Puzanov I, Kim KB, Ribas A, McArthur GA, Sosman JA, et al. Inhibition of Mutated, Activated BRAF in Metastatic Melanoma. *N Engl J Med*. 2010; 363:809–819. [PubMed: 20818844]
25. Mok TS, Wu Y-L, Thongprasert S, Yang C-H, Chu D-T, Saijo N, et al. Gefitinib or Carboplatin–Paclitaxel in Pulmonary Adenocarcinoma. *N Engl J Med*. 2009; 361:947–957. [PubMed: 19692680]

26. Kwak EL, Bang Y-J, Camidge DR, Shaw AT, Solomon B, Maki RG, et al. Anaplastic Lymphoma Kinase Inhibition in Non-Small-Cell Lung Cancer. *N Engl J Med*. 2010; 363:1693–1703. [PubMed: 20979469]
27. Von Hoff DD, LoRusso PM, Rudin CM, Reddy JC, Yauch RL, Tibes R, et al. Inhibition of the Hedgehog Pathway in Advanced Basal-Cell Carcinoma. *N Engl J Med*. 2009; 361:1164–1172. [PubMed: 19726763]
28. Gomez HL, Doval DC, Chavez MA, Ang PC-S, Aziz Z, Nag S, et al. Efficacy and Safety of Lapatinib As First-Line Therapy for ErbB2-Amplified Locally Advanced or Metastatic Breast Cancer. *J Clin Oncol*. 2008; 26:2999–3005. [PubMed: 18458039]
29. Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *The New England journal of medicine*. 2004; 350:2129–2139. [PubMed: 15118073]
30. Joensuu H, Roberts PJ, Sarlomo-Rikala M, Andersson LC, Tervahartiala P, Tuveson D, et al. Effect of the Tyrosine Kinase Inhibitor STI571 in a Patient with a Metastatic Gastrointestinal Stromal Tumor. *N Engl J Med*. 2001; 344:1052–1056. [PubMed: 11287975]
31. Heinrich MC, Corless CL, Demetri GD, Blanke CD, von Mehren M, Joensuu H, et al. Kinase Mutations and Imatinib Response in Patients With Metastatic Gastrointestinal Stromal Tumor. *J Clin Oncol*. 2003; 21:4342–4349. [PubMed: 14645423]
32. Joensuu H, Eriksson M, Sundby Hall K, et al. One vs three years of adjuvant imatinib for operable gastrointestinal stromal tumor: A randomized trial. *JAMA: The Journal of the American Medical Association*. 2012; 307:1265–1272. [PubMed: 22453568]
33. Mahon F-X, Rea D, Guilhot J, Guilhot F, Huguet F, Nicolini F, et al. Discontinuation of imatinib in patients with chronic myeloid leukaemia who have maintained complete molecular remission for at least 2 years: the prospective, multicentre Stop Imatinib (STIM) trial. *The lancet oncology*. 2010; 11:1029–1035. [PubMed: 20965785]
34. Woodman SE, Trent JC, Stemke-Hale K, Lazar AJ, Priol S, Pavan GM, et al. Activity of dasatinib against L576P KIT mutant melanoma: Molecular, cellular, and clinical correlates. *Molecular Cancer Therapeutics*. 2009; 8:2079–2085. [PubMed: 19671763]
35. Rudin CM, Hann CL, Laterra J, Yauch RL, Callahan CA, Fu L, et al. Treatment of Medulloblastoma with Hedgehog Pathway Inhibitor GDC-0449. *N Engl J Med*. 2009; 361:1173–1178. [PubMed: 19726761]
36. Tsimberidou AM, Iskander NG, Hong DS, Wheeler JJ, Fu S, Piha-Paul SA, et al. Personalized medicine in a phase I clinical trials program: The M. D. Anderson Cancer Center Initiative. *J Clin Oncol*. 2011; 29(suppl) (abstr CRA2500).
37. Katayama R, Khan TM, Benes C, Lifshits E, Ebi H, Rivera VM, et al. Therapeutic strategies to overcome crizotinib resistance in non-small cell lung cancers harboring the fusion oncogene EML4-ALK. *PNAS*. 2011; 108:7535–7540. [PubMed: 21502504]
38. Pao W, Miller VA, Politi KA, Riely GJ, Somwar R, Zakowski MF, et al. Acquired Resistance of Lung Adenocarcinomas to Gefitinib or Erlotinib Is Associated with a Second Mutation in the EGFR Kinase Domain. *PLoS Medicine*. 2005; 2:225–235.
39. Villanueva J, Vultur A, Herlyn M. Resistance to BRAF inhibitors: Unraveling mechanisms and future treatment options. *Cancer Research*. 2011; 71:7137–7140. [PubMed: 22131348]
40. Engelman JA, Zejnullahu K, Mitsudomi T, Song Y, Hyland C, Park JO, et al. MET Amplification Leads to Gefitinib Resistance in Lung Cancer by Activating ERBB3 Signaling. *Science*. 2007; 316:1039–1043. [PubMed: 17463250]
41. Shah NP, Nicoll JM, Nagar B, Gorre ME, Paquette RL, Kuriyan J, et al. Multiple BCR-ABL kinase domain mutations confer polyclonal resistance to the tyrosine kinase inhibitor imatinib (STI571) in chronic phase and blast crisis chronic myeloid leukemia. *Cancer Cell*. 2002; 2:117–125. [PubMed: 12204532]
42. Foulds L. Tumor Progression. *Cancer Research*. 1957; 17:355–356. [PubMed: 13437294]
43. Nowell P, Hungerford DA. A minute chromosome in human chronic granulocytic leukemia. *Science*. 1960; 132:1488–1501. [PubMed: 17739576]

44. Nowell P. The clonal evolution of tumor cell populations. *Science*. 1976; 194:23–28. [PubMed: 959840]
45. Curigliano G, Bagnardi V, Viale G, Fumagalli L, Rotmensz N, Aurilio G, et al. Should liver metastases of breast cancer be biopsied to improve treatment choice? *Annals of Oncology*. 2011; 22:2227–2233. [PubMed: 21343379]
46. Fidler I, Hart I. Biological diversity in metastatic neoplasms: origins and implications. *Science*. 1982; 217:998–1003. [PubMed: 7112116]
47. Gerlinger M, Rowan AJ, Horswell S, Larkin J, Endesfelder D, Gronroos E, et al. Intratumor Heterogeneity and Branched Evolution Revealed by Multiregion Sequencing. *N Engl J Med*. 2012; 366:883–892. [PubMed: 22397650]
48. Yachida S, Jones S, Bozic I, Antal T, Leary R, Fu B, et al. Distant metastasis occurs late during the genetic evolution of pancreatic cancer. *Nature*. 2010; 467:1114–1117. [PubMed: 20981102]
49. Mori N, Morosetti R, Lee S, Spira S, Ben-Yehuda D, Schiller G, et al. Allelotype Analysis in the Evolution of Chronic Myelocytic Leukemia. *Blood*. 1997; 90:2010–2014. [PubMed: 9292536]
50. Vogelstein B, Fearon ER, Kern SE, Hamilton SR, Preisinger AC, Nakamura Y, et al. Allelotype of Colorectal Carcinomas. *Science*. 1989; 244:207–211. [PubMed: 2565047]
51. Blackwell KL, Burstein HJ, Storniolo AM, Rugo H, Sledge G, Koehler M, et al. Randomized Study of Lapatinib Alone or in Combination With Trastuzumab in Women With ErbB2-Positive, Trastuzumab-Refractory Metastatic Breast Cancer. *J Clin Oncol*. 2010; 28:1124–1130. [PubMed: 20124187]
52. Kennedy ED, Schmocker S, Victor C, Baxter NN, Kim J, Brierley J, et al. Do patients consider preoperative chemoradiation for primary rectal cancer worthwhile? *Cancer*. 2011; 117:2853–2862. [PubMed: 21692046]
53. Thomas AA, Rini BI, Stephenson AJ, Garcia JA, Fergany A, Krishnamurthi V, et al. Surgical Resection of Renal Cell Carcinoma After Targeted Therapy. *Journal of Urology*. 2009; 182:881–886. [PubMed: 19616232]
54. Mackey J, McLeod D, Ragaz J, Gelmon K, Verma S, Pritchard K, et al. Adjuvant targeted therapy in early breast cancer. *Cancer*. 2009; 115:1154–1168. [PubMed: 19165856]
55. Esserman LJ, Berry DA, DeMichele A, Carey L, Davis SE, Buxton M, et al. Pathologic Complete Response Predicts Recurrence-Free Survival More Effectively by Cancer Subset: Results From the I-SPY 1 TRIAL—CALGB 150007/150012, ACRIN 6657. *J Clin Oncol*. 2012 May 29. [Epub ahead of print].
56. Berry DA. Adaptive clinical trials in oncology. *Nat Rev Clin Oncol*. 2012; 9:199–207. [PubMed: 22064459]
57. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA: A Cancer Journal for Clinicians*. 2011; 61:69–90. [PubMed: 21296855]
58. Ferlay, JSH.; Bray, F.; Forman, D.; Mathers, C.; Parkin, DM. GLOBOCAN 2008 v1.2, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10. Lyon, France: International Agency for Research on Cancer; <http://globocan.iarc.fr> [accessed on 8/16/12]
59. Chin L, Andersen JN, Futreal PA. Cancer genomics: from discovery science to personalized medicine. *Nat Med*. 2011; 17:297–303. [PubMed: 21383744]

Figure 1.
Chemical structure of imatinib

Figure 2.

Transient response to targeted therapy in a heavily pretreated patient. FDG PET images of a patient with metastatic medulloblastoma pre-therapy, after 2 months, and after 3 months of vismodegib. Adapted with permission from reference 35.

Table 1

Efficacy of BCR-ABL Tyrosine Kinase Inhibitors by Disease Stage

Phase	CHR	MCyR	CCyR	MMR	Median OS (months)	OS at 12 months	Ref
Blast Crisis							
Imatinib	I	11%	5%	0%	--	--	7
Imatinib	I/II	23%	3%	8%	--	22%	8
Imatinib	II	15%	16%	7%	--	32%	9
Dasatinib ¹	II	27%	36%	26%	--	50%	10
Nilotinib ¹	I	6%	18%	6%	--	--	11
Accelerated Phase Relapse							
Imatinib	II	53%	24%	17%	--	NR	12
Dasatinib ¹	II	39%	33%	24%	--	NR	13
Previously Treated Chronic Phase							
Imatinib	I	77%	31%	13%	--	--	14
Imatinib	III	82%	55%	39.6% ³	--	NR	15, 16
Dasatinib ¹	II	91%	59%	49%	--	NR	17
Newly diagnosed Chronic Phase							
Imatinib	III	95%	85%	73.8% ³	--	NR	15, 16
Imatinib	III	--	--	66% ⁴	28% ⁴	NR	18
Imatinib	III			65% ⁴	22% ⁴	NR	19
Dasatinib	III	--	--	77% ⁴	46% ⁴	NR	18
Nilotinib	III	--	--	79% ⁴	44% ⁴	--	19

CHR = Complete Hematologic Response, MCyR = Major cytogenetic response, CCyR = Complete cytogenetic response, MMR = Major molecular response, OS = Overall Survival.

¹ Imatinib-resistant or intolerant,² Progression Free Survival at 12 months,³ Best observed response,⁴ response at 12 months,

-- Not reported.

NIH-PA Author Manuscript

NIH-PA Author Manuscript

NIH-PA Author Manuscript

Table 2

Activity of Targeted Agents in Molecularly Defined Populations

Cancer	Common Genetic Driver	Targeted Therapy	Primary	Relapse	Clinical Benefit Rate	Ref
CML	BCR-ABL	Imatinib/Dasatinib/Nilotinib	X		>95%	15, 16
GIST	c-KIT	Imatinib	X		84%	23
Melanoma	V600E BRAF	PLX4032		X	>90%	24
Lung Cancer	EGFR	Erlotinib/Gefitinib		X	92%	25
Lung Cancer	EML4-ALK	Crizotinib		X	>90%	26
Basal Cell Carcinoma	PTCH1/SMO	GDC-0449		X	87%	27
Breast Cancer	ERBB2	Lapatinib		X	31%	28

Clinical benefit: complete response, partial response, and stable disease