Published in final edited form as:

Angew Chem Int Ed Engl. 2014 June 2; 53(23): 5912–5916. doi:10.1002/anie.201402834.

Catalytic Asymmetric Synthesis of 8-Oxabicyclooctanes by Intermolecular [5+2] Perylium Cycloadditions**

Michael R. Witten and Prof. Dr. Eric N. Jacobsen

Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford St., Cambridge, MA 02138 (USA), Fax: (+1) 617-496-1880

Eric N. Jacobsen: jacobsen@chemistry.harvard.edu

Abstract

Highly enantioselective intermolecular [5+2] cycloadditions of perylium ion intermediates with electron-rich alkenes are promoted by a dual catalyst system composed of an achiral thiourea and a chiral primary aminothiurea. The observed enantioselectivity is highly dependent on the substitution pattern of the 5 π component, and the basis for this effect is analyzed using experimental and computational evidence. The resultant 8-oxabicyclo[3.2.1]octane derivatives possess a scaffold common in natural products and medicinally active compounds and are also versatile chiral building blocks for further manipulations. Several stereoselective complexity-generating transformations of the 8-oxabicyclooctane products are presented.

Keywords

asymmetric catalysis; conformational analysis; cycloaddition; hydrogen bonds; organocatalysis

Chiral 8-oxabicyclo[3.2.1]octane architectures reside at the core of numerous natural products and biologically significant compounds (Figure 1).^[1,2] Heterocycles of this class have also proven to be valuable intermediates in the stereoselective synthesis of oxygenated seven-membered carbocycles^[3] and tetrahydrofuran derivatives.^[4] Successful stereoselective approaches to the 8-oxabicyclo[3.2.1]octane framework have relied on transition metal-catalyzed [3+2] cycloadditions,^[5] [4+3] cycloadditions,^[6] diastereoselective [5+2] cycloadditions,^[7] and diastereoselective cascade cyclizations.^[8]

The [5+2] cycloaddition of a perylium ylide precursor with a two-carbon dipolarophile (e.g. **6** to **7**, Scheme 1) provides a concise approach to the 8-oxabicyclo[3.2.1]octane framework, while also embedding a reactive α,β -unsaturated ketone in the bicyclic core.^[9,10] The products can thus be subjected to stereoselective elaboration to afford a diverse array of chiral structures in just one or two steps (*vide infra*). We reported recently that a dual

**This work was supported by the NIH (GM-43214) and an NDSEG Predoctoral Fellowship to M.R.W. (32CFR168a). We thank Dr. Shao-Liang Zheng for crystal structure determination and Dr. Noah Z. Burns and Mr. Andreas R. Rötheli for helpful discussions. Panels b and c of Figure 2 were generated using CYLview.^[25]

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Correspondence to: Eric N. Jacobsen, jacobsen@chemistry.harvard.edu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201402834>.

catalyst system consisting of the chiral primary amine **2** and the achiral thiourea **3** catalyzes enantioselective intramolecular [5+2] cycloadditions of alkenylpyranones of structure **1** through the proposed intermediacy of thiourea-complexed aminopyrylium salts (**4**, Scheme 1).^[11,12,13] While this methodology enables the preparation of a range of complex tricyclic products, the corresponding intermolecular reaction would provide more general access to 8-oxabicyclo[3.2.1]octane derivatives from simpler and more accessible starting materials.^[14] However, initial efforts to effect intermolecular variants of these reactions proved unsuccessful (<40% *ee*).^[11] Here, we disclose that closer examination of this reaction manifold has revealed striking substrate structure-enantioselectivity effects and has led to the successful development of highly enantioselective intermolecular [5+2] cycloadditions.

The racemic pyranone **8a** (Scheme 2), which is superficially analogous to pyranone substrates (**1**) identified as optimal for the intramolecular reaction,^[11] was examined initially as a model substrate for the study of the intermolecular cycloaddition. Treatment of **8a** with a series of terminal alkenes under the dual catalyst conditions developed in the original study revealed that the electronic properties of the 2 π component had a strong effect on reactivity.^[15] Electron-deficient olefins such as acrylonitrile and methyl acrylate were completely inert; styrene displayed measurable but low reactivity to afford racemic product; electron-rich olefins such as benzyl vinyl ether and ethyl vinyl ether underwent cycloaddition with higher conversions and encouraging enantioselectivity. The observation that a nucleophilic 2 π reactant is required in the catalytic reaction is consistent with a mechanism involving a cationic, electron-poor aminopyrylium intermediate analogous to **4** (Scheme 1) rather than an ambiphilic, zwitterionic oxidopyrylium intermediate.^[9f,g]

It proved possible to affect the enantioselectivity of the intermolecular cycloaddition reaction to a remarkable degree through relatively subtle modifications to the structure of the pyrylium precursor. All substrates shown to undergo intramolecular [5+2] cycloadditions with high enantioselectivity in the original study possessed a linkage to the 2 π component through the 6-position of the pyranone (**1**, Scheme 1).^[11] The possibility that the presence of a 6-substituent might affect enantioselectivity in the intermolecular reaction was evaluated by comparing the reaction of **8a** with that of methylated pyranone **9a** (Table 1, entries 1–2).^[15] The dramatic improvement in enantioselectivity observed with methylated derivative **9a** suggests a critical role of that substituent in controlling the transition structure geometry in the catalytic reaction (*vide infra*).

Further improvement in both the enantioselectivity and the yield of the cycloaddition reaction was achieved through electronic tuning of the leaving group on the pyranone. While the *p*-thiomethylbenzoate leaving group had been identified as optimal in the intramolecular reaction, electron-deficient analogs **9b–9d** were found to be advantageous in the intermolecular manifold (entries 3–5).^[16] In particular, 3,4,5-trifluorobenzoate derivative **9d** provided the best balance of reactivity and enantioselectivity, and further improvement in *ee* could be achieved by carrying out the reaction at ambient temperature and decreasing the amount of ethyl vinyl ether from 10 to 5 equivalents relative to pyranone (entry 6).

Having thus established viable parameters for effecting highly enantioselective intermolecular [5+2] cycloadditions of a pyrylium ion precursor, an investigation of the

reaction substrate scope was carried out (Table 2).^[17] Pyranones bearing longer alkyl chains at the 6-position also undergo cycloaddition with ethyl vinyl ether with high enantioselectivity (entry 2); however β -branching in the side chain has a deleterious effect on reactivity (entries 3–4). Cycloadducts bearing siloxymethylene groups could be accessed in moderate-to-good enantioselectivity under the catalytic conditions (entries 5–7). Products of this type have found extensive synthetic applications.^[3a,d,7a,e,f] Benzyl vinyl ether (entry 8) and (methoxymethoxy)ethene (entry 9) were also used successfully as 2π components in the cycloaddition reaction, affording products with readily cleavable ethers. However, styrene derivatives remain poorly reactive in reactions with **9d**, with the resultant cycloadducts formed in very low (<15%) *ee*'s. Enamines and polysubstituted vinyl ethers also proved to be poor substrates, the former undergoing hydrolysis under the reaction conditions and the latter exhibiting very low reactivity.^[15]

We sought to elucidate the basis for the significant difference in enantioselectivity between the reactions of pyranones **8** and **9** (e.g., Table 1, entries 1–2) as a possible path toward gleaning insight into the mechanism of stereoinduction in these reactions. The formation of the adduct **2·9** (Figure 2a) was confirmed by mass spectrometric analysis of a reaction mixture sampled prior to addition of the vinyl ether component.^[15] The lowest energy structures of the two aminopyrylium ions **2·8** and **2·9** were located computationally (Figure 2b,c left).^[18] Rotation about the C(sp²)-N bond in each structure and reoptimization then yielded ground state structures in which the opposite face of the aminopyrylium ion is exposed (Figure 2b,c right).^[19] In the minor conformer of **2·9**, a steric interaction between the methyl substituent and the catalyst's cyclohexyl backbone causes a significant distortion and energetic penalty (>6 kcal/mol relative to the major conformer). That steric interaction is absent if the methyl group is omitted (Figure 2c), resulting in a much smaller energy difference between the conformations of **2·8**. We propose that the methyl substituent in **9** thus enforces a dominant reactive conformation of the aminopyrylium ion adduct, resulting in a more highly enantioselective cycloaddition pathway.

The 8-oxabicyclooctane cycloadducts possess several functional group handles for potential elaboration (Scheme 3). The synthesis of **10** was carried out successfully on 0.70 g batches of starting material under the conditions described in Table 2 above to give cycloadduct in 69% yield and 96% *ee*. Products resulting from conjugate addition (**26**), epoxidation (**27**),^[20] Diels–Alder reaction (**28**),^[21] or Luche reduction (**29**) and subsequent acylation (**30**) were all generated as single diastereomers within detection limits. The cycloadduct **10** can also be subjected to hydrogenation to another key intermediate, saturated ketone **31**.^[22] This heterocycle undergoes Grignard addition stereoselectively to afford tertiary alcohol **32**. Tosylhydrazone **33**, which is primed to undergo Shapiro or Bamford–Stevens reactions, can also be readily synthesized from intermediate **31**. Finally, σ -bond insertion of cyclohexyne yields an unusual 9,6-fused ring system (**34**) in a single step.^[23]

The oxygen bridge of the 8-oxabicyclo[3.2.1]octane framework can also be cleaved reductively. In the case of iodide **36**, this results in formation of an interesting 7-membered product possessing an exocyclic enone (Scheme 4).^[24] As in all of the previous examples, the product was generated as a single diastereomer and without compromise of the optical purity of the initial cycloadduct.

In summary, the catalytic, asymmetric, intermolecular [5+2] pyrylium cycloadditions developed in this study provide multifunctional chiral building blocks that are immediate precursors to a wide range of compound classes. Studies into the mechanism of catalyst cooperativity, as well as extension of the principle into other transformations, are the focus of continuing efforts in this laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. For example, Englerin A: Ratnayake R, Covell D, Ransom TT, Gustafson KR, Beutler JA. *Org Lett.* 2009; 11:57–60. [PubMed: 19061394] ; Descurainin: Sun K, Li X, Li W, Wang J, Liu J, Sha Y. *Chem Pharm Bull.* 2004; 52:1483–1486. [PubMed: 15577251] Intricarene: Marrero J, Rodríguez AD, Barnes CL. *Org Lett.* 2005; 7:1877–1880. [PubMed: 15844929] Anthecularin: Karioti A, Skaltsa H, Linden A, Perozzo R, Brun R, Tasdemir D. *J Org Chem.* 2007; 72:8103–8106. [PubMed: 17880246] Cortistatin A: Aoki S, Watanabe Y, Sanagawa M, Setiawan A, Kotoku N, Kobayashi M. *J Am Chem Soc.* 2006; 128:3148–3149. [PubMed: 16522087] Hedyosumin C: Su Z-S, Yin S, Zhou Z-W, Wu Y, Ding J, Yue J-M. *J Nat Prod.* 2008; 71:1410–1413. [PubMed: 18651771]
2. Compounds containing the chiral 8-oxabicyclo[3.2.1]octane core have been identified recently as psychoactive analogs of the tropane alkaloids with promising pharmaceutical potential: Meltzer PC, Liang AY, Blundell P, Gonzalez MD, Chen Z, George C, Madras BK. *J Med Chem.* 1997; 40:2661–2673. [PubMed: 9276012] Meltzer PC, Liu S, Blanchette HS, Blundell P, Madras BK. *Bioorg Med Chem.* 2002; 10:3583–3591. [PubMed: 12213473] Meltzer PC, Kryatova O, Pham-Huu DP, Donovan P, Janowsky A. *Bioorg Med Chem.* 2008; 16:1832–1841. [PubMed: 18053732]
3. a) Wender PA, Lee HY, Wilhelm RS, Williams PD. *J Am Chem Soc.* 1989; 111:8954–8957.b) Wender PA, Kogen H, Lee HY, Munger JD, Wilhelm RS, Williams PD. *J Am Chem Soc.* 1989; 111:8957–8958.c) Bromidge SM, Sammes PG, Street LJ. *J Chem Soc, Perkin Trans 1.* 1985:1725–1730.d) Wender PA, Jesudason CD, Nakahira H, Tamura N, Tebbe AL, Ueno Y. *J Am Chem Soc.* 1997; 119:12976–12977.e) Roethle PA, Hernandez PT, Trauner D. *Org Lett.* 2006; 8:5901–5904. [PubMed: 17134301] f) Li Y, Nawrat CC, Pattenden G, Winne JM. *Org Biomol Chem.* 2009; 7:639–640. [PubMed: 19194574] g) Nicolaou KC, Kang Q, Ng SY, Chen DYK. *J Am Chem Soc.* 2010; 132:8219–8222. [PubMed: 20496885]
4. a) Fishwick CWG, Mitchell G, Pang PFW. *Synlett.* 2005:285–286.b) Krishna UM. *Tetrahedron Lett.* 2010; 51:2148–2150.c) Yadav AA, Sarang PS, Trivedi GK, Salunkhe MM. *Synlett.* 2007:989–991.d) Ali MA, Bhogal N, Findlay JBC, Fishwick CWG. *J Med Chem.* 2005; 48:5655–5658. [PubMed: 16134933]
5. a) Kitagaki S, Anada M, Kataoka O, Matsuno K, Umeda C, Watanabe N, Hashimoto S. *J Am Chem Soc.* 1999; 121:1417–1418.b) Hodgson DM, Labande AH, Pierard FYTM, Expósito Castro MÁ. *J Org Chem.* 2003; 68:6153–6159. [PubMed: 12895044] c) Hodgson DM, Brückl T, Glen R, Labande AH, Selden DA, Dossetter AG, Redgrave AJ. *Proc Natl Acad Sci USA.* 2004; 101:5450–5454. [PubMed: 15037752] d) Shimada N, Anada M, Nakamura S, Nimbu H, Tsutsui H, Hashimoto S. *Org Lett.* 2008; 10:3603–3606. [PubMed: 18616257] e) Ishida K, Kusama H, Iwasawa N. *J Am Chem Soc.* 2010; 132:8842–8843. [PubMed: 20540576]
6. a) Harmata M, Ghosh SK, Hong X, Wacharasindhu S, Kirchhoefer P. *J Am Chem Soc.* 2003; 125:2058–2059. [PubMed: 12590528] b) Huang J, Hsung RP. *J Am Chem Soc.* 2005; 127:50–51. [PubMed: 15631443]
7. a) Wender PA, Rice KD, Schnute ME. *J Am Chem Soc.* 1997; 119:7897–7898.b) López F, Castedo L, Mascareñas JL. *Org Lett.* 2000; 2:1005–1007. [PubMed: 10768208] c) López F, Castedo L, Mascareñas JL. *Org Lett.* 2002; 4:3683–3685. [PubMed: 12375918] d) Krishna UM, Deodhar KD, Trivedi GK. *Tetrahedron.* 2004; 60:4829–4836.e) Wender PA, Bi FC, Buschmann N, Gosselin F, Kan C, Kee JM, Ohmura H. *Org Lett.* 2006; 8:5373–5376. [PubMed: 17078721] f) Wender PA,

- Buschmann N, Cardin NB, Jones LR, Kan C, Kee JM, Kowalski JA, Longcore KE. Nat Chem. 2011; 3:615–619. [PubMed: 21778981]
8. Li B, Zhao YJ, Lai YC, Loh TP. Angew Chem. 2012; 124:8165–8169. Angew Chem Int Ed. 2012; 51:8041–8045.
9. For recent reviews, see: Singh V, Krishna UM, Vikrant, Trivedi GK. Tetrahedron. 2008; 64:3405–3428. Pellissier H. Adv Synth Catal. 2011; 353:189–218. Ylijoki KEO, Stryker JM. Chem Rev. 2013; 113:2244–2266. [PubMed: 23153111] ; for pioneering reports, see: Hendrickson JB, Farina JS. J Org Chem. 1980; 45:3359–3361. Sammes PG, Street LJ. J Chem Soc, Chem Commun. 1982:1056–1057. Sammes PG, Street LJ. J Chem Soc, Perkin Trans 1. 1983:1261–1265. Sammes PG, Street LJ. J Chem Res, Synop. 1984:196–197.
10. Other than our earlier work (ref. [11]), there is a single report of an enantioselective [5+2] pyrylium cycloaddition (<20% product ee): Hodgson DM, Stuppel PA, Johnstone C. ARKIVOC. 2003; vii:49–58.
11. Burns NZ, Witten MR, Jacobsen EN. J Am Chem Soc. 2011; 133:14578–14581. [PubMed: 21848300]
12. For a review of anion-binding asymmetric catalysis, see: Brak K, Jacobsen EN. Angew Chem. 2013; 125:558–588. Angew Chem Int Ed. 2013; 52:534–561.
13. For preparation and use of catalyst **3**, see: Schreiner PR, Wittkopp A. Org Lett. 2002; 4:217–220. [PubMed: 11796054] Wittkopp A, Schreiner PR. Chem—Eur J. 2003; 9:407–414. [PubMed: 12532289]
14. For diastereoselective intermolecular [5+2] cycloadditions, see: Tchabanenko K, Sloan C, Bunetel YM, Mullen P. Org Biomol Chem. 2012; 10:4215–4219. [PubMed: 22517727] Garnier EC, Liebeskind LS. J Am Chem Soc. 2008; 130:7449–7458. [PubMed: 18479131]
15. See the Supporting Information for details.
16. In ref. [11], we proposed that the low solubility of the *p*-thiomethylbenzoic acid byproduct was a crucial driving force in the intramolecular reaction. In contrast, 3,4,5-trifluorobenzoic acid is fully soluble under the reaction conditions employed in this study. The improved reactivity imparted by the 3,4,5-trifluorobenzoate group in **9d** relative to less electron-deficient analogs may be ascribed simply to its superior properties as a leaving group. However, the 3,4,5-trifluorobenzoate leaving group is inferior in the intramolecular reaction. See the Supporting Information for details.
17. For certain indicated substrates, catalyst loading could be reduced to 15 mol% **2** and 15 mol% **3** without detrimental effect on yield or enantioselectivity. In all other cases, this reduction in catalyst resulted in significantly diminished yields.
18. B3LYP/6–31G(d) has been established as an appropriate level of theory for studying [5+2] pyrylium cycloadditions. See: López F, Castedo L, Mascareñas JL. J Org Chem. 2003; 68:9780–9786. [PubMed: 14656107] Wang SC, Tantillo DJ. J Org Chem. 2008; 73:1516–1523. [PubMed: 18205383]
19. These calculations apply to the conformations of the aminopyrylium ground states prior to the cycloaddition step, rather than to the actual transition structures for cycloaddition. They are further simplified in that they omit the thiourea **3**-benzoate counteranion. Nonetheless, we propose that the pronounced energy difference calculated for the conformations of **2**·**9d** is expected to translate to the relevant cycloaddition transition structures that lead to the two enantiomers **10**.
20. Marshall KA, Mapp AK, Heathcock CH. J Org Chem. 1996; 61:9135–9145.
21. Nair V, Anilkumar G, Sujatha TS, Nair JS. Synth Commun. 1998; 28:2549–2557.
22. Archer DA, Bromidge SM, Sammes PG. J Chem Soc, Perkin Trans 1. 1988:3223–3228.
23. Gampe CM, Boulos S, Carreira EM. Angew Chem. 2010; 122:4186–4189. Angew Chem Int Ed. 2010; 49:4092–4095.
24. Ohmori N. J Chem Soc, Perkin Trans 1. 2002:755–767.
25. Legault, CY. CYLview, version 1.0b. Université de Sherbrooke; 2009. <http://www.cylview.org>

Figure 1.
The 8-oxabicyclo[3.2.1]octane core in selected natural products.

Figure 2.

a) Formation of intermediate aminopyrylium. b) and c) Lowest energy calculated structures (B3LYP/6–31G(d)) of the aminopyrylium ions **2•8** and **2•9**. In both cases, the major conformer has the face of the aminopyrylium ion that leads to the major observed enantiomer of cycloadduct exposed, while the other face is shielded. The difference in energy between the major and minor conformers is substantially higher for **2•9**, however, and that is a potential explanation for the higher enantioselectivities obtained in cycloadditions of **9**.

Previous work:

Ref. [11]

This work: **Scheme 1.**

Enantioselective [5+2] cycloadditions.

Scheme 2.
Screen of 2π components.

Scheme 3.

Derivatization reactions. Reaction conditions: a) **2** (15 mol%), **3** (15 mol%), AcOH (15 mol %), ethyl vinyl ether, toluene; b) Me_2CuLi , Et_2O ; c) $t\text{BuOOH}$, NaHCO_3 , H_2O , MeOH ; d) cyclopentadiene, ZnCl_2 , C_6H_6 ; e) $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$, NaBH_4 , MeOH ; f) $p\text{-BrBzOH}$, DMAP, pyr., CH_2Cl_2 ; g) SiH_2Ph_2 , ZnCl_2 (10 mol%), $\text{Pd}(\text{PPh}_3)_4$ (2 mol%), CHCl_3 ; h) MeMgI , Et_2O ; i) H_2NNHTs , THF, H_2O ; j) cyclohexenylphenyl iodonium tetrafluoroborate, KOEt_3 , THF. All products are isolated in >20:1 *dr*.

Scheme 4.

Stereocontrolled preparation of cycloheptanone **37**. Reaction conditions: a) SiH_2Ph_2 , ZnCl_2 (10 mol%), $\text{Pd}(\text{PPh}_3)_4$ (2 mol%), CHCl_3 ; b) AcOH , HCl , THF , H_2O ; c) I_2 , PPh_3 , imid., C_6H_6 ; d) Zn , MeOH .

Table 1

Reaction Optimization.^[a]

Entry	Substrate	<i>t</i> [h]	Yield ^[b] [%]	<i>ee</i> ^[c] [%]
1	8a	48	53	47
2	9a	48	55	90
3	9b	48	63	94
4	9c	48	65	93
5	9d	48	73	92
6 ^[d]	9d	72	75	96

^[a] Reactions were performed on a 0.07 mmol scale.

^[b] Determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard.

^[c] Determined by HPLC using commercial chiral columns.

^[d] 5 equiv ethyl vinyl ether, room temperature.

Table 2

Substrate Scope.^[a]

Yield [%]		ee[b] [%]	
Entry	Substrate	R ³	Product		
1[c]	9d	Et	10	69	96
2	12	Et	13	59	90
3[c]	14	Et	15	22	88
4	16	Et	17	26	67
5	18	Et	19	64	86

Entry	Substrate	R ³	Product	Yield [%]	<i>ee</i> [<i>b</i>] [%]
6 ^[c,d]		Et		95	64
7		Et		75	89
8 ^[c]		Bn		88	89
9 ^[e]		MOM		54	91

^[a] Reactions were performed on a 0.2 mmol scale. Isolated yields after chromatography on silica gel. For details on substrate synthesis, see Supporting Information.

^[b] Determined by HPLC using commercial chiral columns.

^[c] 15 mol% **2** + 15 mol% **3**.

^[d] 96 h at 0 °C.

^[e] 10 equiv vinyl ether.

^[f] The absolute stereochemistry of a derivative of **10** was determined by X-ray crystallography [see Supporting Information], and that of all other products was assigned by analogy.

^[g] methoxymethyl.