Published in final edited form as:

Am J Alzheimers Dis Other Dement. 2012 November ; 27(7): 520–529. doi:10.1177/1533317512459798.

NEUROPSYCHOLOGICAL AND NEUROIMAGING MARKERS IN EARLY VS. LATE-ONSET ALZHEIMER'S DISEASE

Natalie C. Kaiser^{*,a}, Rebecca J. Melrose^b, Collin Liu^a, David L. Sultzer^{a,b}, Elvira Jimenez^a, Michael Su^b, Lorena Monserratt^a, and Mario F. Mendez^a

^aVA Greater Los Angeles Healthcare System, 11301 Wilshire Blvd., Los Angeles, CA 90073 USA

^bDavid Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA 90095 USA

Abstract

Background—Early-onset Alzheimer's disease (EOAD) has been overshadowed by the more common late-onset AD (LOAD). Yet, the literature indicates EOAD may have less hippocampal-memory presentations and more focal neocortical localization early in the disease.

Objective—To evaluate these proposed differences between these two forms of AD and to explore what they inform about differences in AD-pathophysiology.

Methods—21 EOAD and 24 LOAD patients matched for disease progression and severity were compared on neurocognitive measures and resting state fluorodeoxy-glucose positron emission tomography (FDG-PET).

Results—EOAD patients had worse executive functions with greater hypometabolism in the parietal regions; whereas LOAD patients had worse confrontation naming and verbal recognition memory with greater hypometabolism in inferior frontotemporal regions.

Conclusions—In addition to highlighting significant differences between EOAD and LOAD, these results reveal dissociation between executive deficits in AD and frontal hypometabolism, suggesting early disturbances of the parietal-frontal network in EOAD.

Keywords

Early-onset; Late-onset; Neuropsychology; Alzheimer's disease; Dementia; Neuroimaging; FDG-PET

INTRODUCTION

In 1901, Alois Alzheimer observed a patient at the Frankfurt Asylum named Auguste Deter; a 51-year-old patient with strange behavioral symptoms and a memory disturbance. Five years later, when this patient passed away, neuropathology revealed neuritic plaques and neurofibrillary tangles consistent with what became known as "Alzheimer's disease" (AD). Over the years, the much more common late-onset AD (LOAD) overshadowed this early-

*Address correspondence to: Natalie Kaiser, PhD, VA Greater Los Angeles Healthcare System, 11301 Wilshire Blvd., Bldg 401 Room A235, Los Angeles, CA 90073, Phone: 310-478-3711, x42393 Natalie.Kaiser@va.gov.

onset AD (EOAD) characterized by an onset before age 65. The study of EOAD is important, not only because it affects up to 2.4 out of 100,000 persons,¹ but also because it tends to strike people when they are in the prime of their career and family responsibilities.

The literature suggests that EOAD differs substantially from LOAD²⁻⁴. There is general agreement that patients with EOAD have greater cortical atrophy and hypometabolism compared to patients with LOAD at a similar stage of disease.⁵ Functional neuroimaging studies have demonstrated significant cortical atrophy, hypoperfusion, and hypometabolism in EOAD, particularly in parietal and lateral temporal cortices, and significant lesions or hypometabolism in medial temporal and hippocampal regions in LOAD⁵⁻¹⁴. There are conflicting studies, however, on the cognitive aspects of EOAD. Some studies find that EOAD patients show greater impairments in attention, language, visuospatial and executive functions; and LOAD patients show comparatively greater deficits in episodic memory^{2-5,15-22}. In contrast, other studies find no differences in cognition according to age of onset^{7,23}. Given the differences on functional neuroimaging, it would be important to know whether the cognitive aspects correspond with the differences in functional neuroimaging results in EOAD vs. LOAD.

This study compares EOAD with LOAD on both cognitive features and functional neuroimaging with positron emission tomography (PET). There have been few direct comparisons between EOAD and LOAD patients, and much of the literature has been restricted to examination of solely neuropsychological differences between the groups^{2,3,15,24-27}; or solely neuroimaging differences^{6,8-11,13,28-30}. We hypothesize that such comparison would confirm the presence of greater non-memory deficits and more prominent neocortical hypometabolism, particularly in the parietal lobes, in EOAD as compared to LOAD. Finally, this study has implications for understanding differences in the underlying pathophysiology of AD.

METHODS

Participants

Participants were recruited from the Departments of Neurology and Geriatric Psychiatry at the Veterans Affairs (VA) Greater Los Angeles Healthcare Center and the University of California, Los Angeles (UCLA) Geffen School of Medicine. All participants were native English speakers. Diagnosis of AD was determined by the National Institute of Communicable Diseases and Stroke-Alzheimer's Disease and Related Disorders Association (NINCDS-ARDA) criteria for clinically probable AD³¹. Whether one was classified as "early-onset" vs. "late-onset" was based on age of onset; those with a disease onset prior to age 65 were "early-onset" and those with a disease onset subsequent to age 65 were "late-onset." Individuals with major medical illnesses (except hypertension or diabetes) or psychiatric illnesses not due to the dementia process were excluded from the study. The study was reviewed and approved by the local Institutional Review Board (IRB) and study participants were enrolled according to IRB guidelines. The participants in this study were specifically selected in order to match the EOAD and LOAD groups on five variables which are well-established confounds in previous studies.

The participants were group matched on education, time since diagnosis, gender distribution, functional status (e.g., Activities of daily living (ADLs)/Instrumental Activities of daily living (IADLs)) and global cognitive severity (as measured by the Mini Mental State Exam, MMSE).

Procedures

The patients underwent neuropsychological testing, and neuroimaging with FDG-PET of the brain. Caregivers (study partners) who accompanied patients to appointment also filled out questionnaires to provide information regarding their experiences/ impressions of the patient.

Measures

Demographics—The study partners (caregivers) completed a demographic questionnaire on behalf of the subjects, which included information such as age, education level, marital and living situation status, and occupation. Disease-related information including diagnosis and time of onset was also provided by caregivers.

Global Cognitive Severity and Functional Status (ADLs/IADLs)—

1. All participants completed the MMSE as a measure of global cognitive severity. Data was normed, adjusting for age and education level ³².
2. Alzheimer's Disease Cooperative Study-Activities of Daily Living Inventory (ADCS-ADL). The ADCS-ADL is a 23-item inventory wherein the caregiver is asked to focus on the patient's observed performance over the past month..

Language—

1. Boston Naming Test. This is a 60-item confrontation naming task measuring the word retrieval process of patients when presented with a picture of an item. Data was normed, adjusting for age, education, and gender ³³.
2. Controlled Oral Word Association Test (COWAT), wherein participants are given 60 seconds to produce as many words as possible from a semantic and phonemic category. Data was normed, adjusting for age, education, and gender ³⁴.

Visuospatial Skills—A subindex from the intersecting pentagon task on the MMSE ³⁵³⁶ was utilized to evaluate the types of visuospatial errors made in this task. The eight errors in this index were relating to size of the figures, number of figures drawn, intersection of the pentagons, tremor or segmentation errors, angle errors, rotation, motor perseveration, and pull-to-stimulus tendencies.

Attention and Executive Control—

1. Digit Span subtest of the Wechsler Adult Intelligence Scale, 3rd edition (WAIS-III). Participants are presented with a series of numbers presented auditorily, and asked to repeat these numbers back to examiner in the exact order read to them (digits

forward) or in the reverse order the digits were read to them (digits backward). Normative data adjusting for age was obtained from the WAIS-III manual.

2. Stroop Task, which provides participants with 3 different conditions (word reading, color naming, and interference task). In each condition, participants are timed to see how many words they can successfully read/name within a given amount of time. Normative data adjusting for age was obtained from the Golden manual.
3. Trail Making Test (TMT, Part A only). This is a test of visual attention and psychomotor speed, where the participant is provided with a sheet of paper with numbers from 1–25 within circles and instructed to connect the numbers in consecutive order using straight lines. Data was normed, adjusting for age and education was provided ³⁷.
4. COWAT (“FAS”, see above).

Verbal Learning and Memory—Neuropsychological battery of the Consortium to Establish a Registry for Alzheimer’s Disease (CERAD); 10-item word list learning and recognition only). The list learning/recognition task on the CERAD includes 10 common nouns presented consecutively and read aloud by the participant, with a different order used on each of three successive occasions. Following this task, there is a free recall condition and recognition paradigm presented. Data was normed, adjusting for age, gender, and education level ³⁸.

(FDG) PET Imaging: Data Acquisition and Processing—All patients underwent resting state functional neuroimaging with PET scans as part of this study in accordance with the technique of Hamacher Coenen & Stocklin (1986)³⁹. They received an intravenous infusion of 5–10 mCi of [18F]-FDG and then waited forty minutes quietly in a dimly lighted room with eyes open (uptake phase). Participants were next placed in the scanner with the imaging plane parallel to the canthomeatal plane and metabolic data were acquired for 40 minutes. LOAD patients were scanned on different tomographs over the course of the study. A total of 3 patients were scanned on a Siemens 953/31 tomographic scanner (voxel size 1.96×1.96×3.38 mm, Siemens Medical Solutions, Hoffman Estates, Illinois), 11 on a mobile GE Advance (voxel size 2.34×2.34×4.25 mm, 2.73×2.73×4.25, or 2.34×2.34×4.25m, or 2.34×2.34×3.27GE Medical Systems, Milwaukee, WI), and 7 on a Philips Gemini TF-PET-CT (2×2×2). All early AD patients (n=20) were scanned on a Philips Gemini TF-PET-CT (2×2×2). Most participants completed PET imaging on the same day as the cognitive assessment; however, all of the PET scans were done within 2 months of the respective cognitive testing. Raw data in DICOM format were converted to ANALYZE format. In all images, origins were manually set to the anterior cingulate cortex, before the spatial normalization step. Images were normalized to Montreal Neurological Institute (MNI) space using tri-linear interpolation and resampled to 2 × 2 × 2 mm voxels, and smoothed using a 6-mm FWHM smoothing kernel, using SPM8 (Wellcome Trust Centre for Neuroimaging, London, United Kingdom) in MATLAB R2010b (MathWorks, Inc.).

We investigated the data to determine if there were any systematic differences between groups with respect to global metabolism. We obtained mean values of the entire brain using

custom MatLab software, and cerebellum gray matter using Automatic Anatomic Labeling (AAL) toolkit in SPM8. The ratio of whole-brain/cerebellar gray matter was calculated for each subject.

Analyses

Demographic and neuropsychological variables were compared in SPSS 17.0 using Chi Square analyses as well as independent samples t-tests to evaluate between-group differences. Both raw scores as well as normed, standard scores (adjusting for age, education, gender in order to directly compare levels of performance by expressing each patient's scores in standard deviations from the mean of normal control performance) were compared. Group differences were also evaluated with analysis of covariance (ANCOVA), using global cognitive severity (MMSE scores) as a covariate. The significance level was set at the $p < 0.05$ level. Due to the small sample size, we did not correct for multiple comparisons.

Voxel-Based Statistical Analysis—Statistical maps of the FDG-PET images were generated using SPM8. Two-sample t-test was performed between EOAD and LOAD groups. Subject images were normalized to the global mean. Threshold masking was used to remove signal from structures outside of the brain (set to 0.8). Between-group difference maps were set at a voxel threshold of $p < 0.01$ (uncorrected for multiple comparisons) and considered significant at the cluster level at $p < 0.05$, corrected for multiple comparisons using the family-wise error procedure (FWE).

RESULTS

Demographics

Table 1 depicts the demographic makeup of our sample. The only demographic variables that significantly differed between the EOAD and LOAD groups was age (as anticipated), and ethnicity, wherein the EOAD group had more non-white individuals.

Neuropsychological Comparisons of EOAD and LOAD Patients

Global Cognitive Severity and Functioning—EOAD and LOAD patients were matched in regards to global cognitive functioning as measured by the MMSE (*See* Table 2). When normed (age, education adjusted) data was compared, the non-significant between-group differences remained. There were no significant group differences on the Alzheimer's Disease Cooperative Study Scale (measuring ADL/IADL functionality, $t(34) = -0.15$, n.s.). Similarly, when broken down and analyzed by subscales indicating degrees of daily functioning, no significant group differences were documented (Basic: $t(27) = -1.18$, n.s.; Autonomy: $t(32) = 0.04$, n.s.; Higher Level Function, $t(20) = 0.26$, n.s.).

Language—LOAD patients performed significantly worse than EOAD patients on the Boston Naming Test (number correct with or without stimulus cue), $t(38) = 3.15$, $p < 0.01$; This finding remained significant when comparing normed data (adjusting for age, gender, and education). On verbal fluency raw scores, there were no significant differences with respect to either phonemic or semantic fluency. When normed data was compared, the

EOAD patients had poorer performances when compared to LOAD patients on the phonemic fluency measure.

Visuospatial Skills—Although there were no statistically significant differences in gross visuoconstruction skills as assessed by the overlapping pentagons portion on the MMSE, there were differences between the EOAD and LOAD groups in type of errors made on visuoconstruction task^{35,36}. More of the EOAD patients (26.3%) made pull-to-stimulus errors (e.g., patient drew, wrote, or copied over the figure model), compared to the LOAD patients (0%) ($\chi^2(1) = 6.59, p = 0.01$).

Attention and Executive Control—On a measure of simple attention, there were no significant between-group differences; WAIS Digit Span Forward, $t(41) = -0.83$, n.s.; overall digit span total score, $t(41) = -1.74$, n.s.. In contrast, on measures of complex attention or working memory, significant between-group differences were found on the Wechsler Adult Intelligence Scale Digit Span (backwards, raw) subtest, $t(41) = -2.36$, $p < 0.05$. The EOAD patients had lower performances ($M = 3.0$, $sd = 1.75$) on this subtest when compared to their LOAD counterparts ($M = 4.42$, $sd = 2.20$). Measures of visual attention and psychomotor speed did not yield significant differences when raw scores were compared; however, a significant difference emerged on Trails A performance when normed data (adjusting for age and education levels) was compared, $t(16.10) = -2.16$, $p < 0.05$. The LOAD patients tended to perform comparatively better than EOAD patients.

On executive measures, the EOAD group consistently performed worse than the LOAD group on a number of tasks. Although raw score differences were null, normed data comparisons (controlling for age) revealed significant between-group differences on the Stroop interference or response inhibition task ($t(35) = -4.48$, $p < 0.001$), with LOAD patients performing comparatively better to EOAD patients. Additionally, although semantic fluency scores were comparable, a phonemic fluency task indicated significant differences with poorer performances for the EOAD group compared to the LOAD group when analyses were adjusted for demographics (FAS, $t(42) = -2.15$, $p < 0.05$). The other “executive” deficits in EOAD, compared to LOAD, have been noted and include problems with backward digits and “pull-to-stimulus” visuospatial functioning.

Verbal Learning and Memory—There was not significant group differences on Trials 1–3, CERAD total recall, $t(32) = 0.13$, n.s.; however, the delayed recall nearly reached significance with the LOAD patients performance the worst, $t(32) = 1.92$, $p = 0.06$ scores. In addition, there was a significant difference between true-positive hits on a recognition paradigm; LOAD patients performed worse on this task than EOAD patients did.

Normed-data comparisons (controlling for age, gender, and education level) yielded similar findings, with only the age, gender, and education corrected recognition scores yielding significant differences (CERAD Recognition, $t(31.84) = 4.17$, $p < 0.001$).

Covariate-adjusted analyses: Controlling for Global Cognitive Functioning—When MMSE scores were entered as a covariate in analyses examining between-group differences, all previously significant findings persisted with the sole exception of the Trail

Making Test, Part A. This test was no longer significant when controlling for MMSE scores, $F(2,31) = 2.67$, n.s.

FDG-PET Imaging

Statistical parametric mapping (SPM) was employed to compare FDG- PET images of the EOAD and LOAD groups. Results are illustrated in Figure 1. See Table 3 for coordinates and statistics. There were no differences between groups with respect to the whole brain/cerebellum ratio. Between-group two-sample t-test ($p < .05$, corrected at the cluster level using the FWE procedure) revealed relative hypometabolism in the EOAD group compared to the LOAD group in a large cluster in the posterior cortex, including bilateral precuneus, bilateral superior parietal lobule (L>R), left supramarginal gyrus, left angular gyrus, posterior regions of the left lateral temporal cortex (superior, middle, and inferior temporal gyri), and left occipital lobe ($p < .05$, FWE corrected at the cluster level). There was a trend towards relative hypometabolism in left lateral frontal cortex, primarily encompassing the left precentral gyrus and extending anteriorly to include posterior middle frontal and inferior frontal gyri ($p = .07$ FWE corrected at the cluster level). In contrast, relative hypometabolism in the LOAD group compared to the EOAD group was observed in inferior portions of the brain, including inferior frontal lobe and temporal cortex. Specifically, there was a large cluster encompassing bilateral orbitofrontal gyri and medial frontal lobe (including the cingulate, gyrus rectus, and superior frontal gyrus). Moving posteriorly it included parts of bilateral inferior frontal gyrus. This also included the right superior, middle, and inferior temporal gyri, fusiform, parahippocampal gyrus, and hippocampus. A second cluster revealed relative hypometabolism in LOAD vs. EOAD in left parahippocampal gyrus and fusiform. Additional between group differences were observed in bilateral cortex surrounding the Sylvian fissure and cerebellum. Figure 2 provides view of one representative patient from the EOAD and LOAD group, respectively, to demonstrate the pattern of parietal hypometabolism in the EOAD group, while showing the pattern of inferior frontal hypometabolism in the LOAD group. The frontal pattern within the LOAD group, however, is more subtle, as the true differences are more apparent in the between-group comparisons.

DISCUSSION

This study found neuropsychological and functional imaging differences between patients with AD of early onset compared to those with AD of late onset. Compared to the LOAD patients, the EOAD patients performed worse on a several measures of executive functioning (working memory on digit span backwards; phonemic vs. semantic fluency; a pull-to-stimulus copy performance, and the Stroop interference task). This finding of greater executive dysfunction in the EOAD cohort has been inconsistent in previous studies; while some studies have confirmed this finding,²¹ others have not found group differences on executive tasks⁷. In contrast, the LOAD patients, compared to the EOAD patients, had worse confrontation naming and recognition memory scores, findings more typical of AD^{2-4,15,22,25}. These changes correspond with relative parietal hypometabolism among the EOAD patients and relative frontotemporal hypometabolism among the LOAD patients.

These results are consistent with the literature in some findings but not others. Similar to other studies,²⁶ this study finds that EOAD patients perform worse than LOAD patients on executive tasks, including working memory as reflected in digit span backwards. The finding that EOAD are significantly worse on a measure of phonemic fluency (but not semantic fluency) further supports the pattern of executive dysfunction, as research has demonstrated that disproportionate letter fluency correlates with frontal-executive dysfunction^{40,41}. In addition, a pull-to-stimulus (e.g., patient wrote, drew, or copied over figure model) error on the visuoconstruction task is much more common in EOAD rather than LOAD patients. This type of visuospatial error is consistent with an executive control problem in that it indicates environmental-dependency⁴². In contrast, the LOAD patients, as compared to the EOAD patients, have worse language functioning as evidenced by poorer performance on a confrontation naming task. This finding differs from some previous studies which report more severe language impairment as a feature of EOAD^{4,17,43}. Similarly, there have been different findings regarding worse memory disturbances in LOAD vs. EOAD^{7,16,17}. This study reports worse recognition memory for the LOAD group compared to the EOAD group.

On FDG-PET, the EOAD patients, compared to the LOAD patients, have lower metabolic activity in the parietal (left worse than right) lobes. Other studies of EOAD indicate the presence of significant parietal lobe involvement along with additional changes in other neocortical regions early in the disease^{6–14}, whereas prior neuroimaging studies of LOAD show early changes in hippocampal and inferior temporal regions⁵. In terms of laterality, although the current study suggests that EOAD patients have greater lateralization of Alzheimer changes to the left hemisphere, other studies are inconsistent on whether AD affects the left or right hemisphere disproportionately^{5,14}. Finally, the degree of underlying atrophy can impact regional hypometabolism, but it would still reflect selective, disproportionate disease of the parietal lobes in EOAD compared to the frontotemporal lobes in LOAD.

A significant finding of this study is the presence of worse executive functions without worse frontal metabolism in EOAD as compared to LOAD. This dissociation has been previously noted but not specifically studied⁴¹. Such a dissociation supports recent views of regional dysfunction in that executive skills are not exclusively frontally mediated, and are rather the product of an interconnected network. Namely, the dorsolateral frontal cortex has dense connections with the parietal lobe; a network that has been implicated in tasks of executive functioning^{40,41}, especially in early-onset dementia⁴¹. There is an increasing understanding that cognition is a product of these networks and tracts, and investigations of white matter pathways indicate disruption of important parietal lobe-dorsolateral frontal tracts in AD⁴⁰. Early involvement of the parietal lobe in EOAD would affect gray matter as well as its main dorsolateral frontal networks. In addition, in this study there was a trending pattern of lower metabolic activity in the dorsolateral frontal regions in EOAD.

This study has potential limitations. First, sample size is relatively small, which has implications for generalizability of these findings as well as statistical limitations. Nevertheless, this report revealed several important differences between EOAD and LOAD, which were generally adequately powered (Post-hoc power analyses ranged from .66–.99 for

significant group differences reported herein). Thus, the results herein, which may be characterized as exploratory, may require confirmatory analyses with a larger sample size in future research. Second, some cognitive domains, in particular visuospatial functioning, had limited assessment. Future research may include a more comprehensive cognitive comparison of both early and late-onset groups. Third, the presence of greater hypometabolism in the medial/inferior frontal lobe in the LOAD group may reflect age-related changes in this older cohort. Previous work has found that medial prefrontal regions are sensitive to normal aging^{44,45}. This could not be addressed in the current study given the lack of normal controls. However, prior investigations comparing EOAD or LOAD subjects with normal controls report the same pattern of metabolic changes (e.g. ^{5,11}). Thus, although not directly compared in the present study, the patterns of worse parietal metabolism in EOAD and frontal metabolism in LOAD are consistent with qualitatively different patterns than would be observed just from normal aging profile. Furthermore, given the lack of overlap in age between the two cohorts, controlling for age would remove innate variance. Finally, the LOAD group included FDG-PET data from subjects scanned on multiple PET tomographs. The between-group differences were analyzed using a normalization procedure in SPM, which yields a ratio scale for each subject (voxel divided by global mean), and ratio values appear to vary by less than 5% across scanners⁴⁶. We assessed the global ratio values in each group and observed a nearly identical mean and range of values in each group. This suggests that while the use of multiple scanners may have impacted our ability to detect differences between groups, it is unlikely a large effect.

In conclusion, these findings suggest differences in AD-pathophysiology between EOAD and LOAD; a notion that has been debated since Auguste Deter was first encountered by Alois Alzheimer. This study confirms some previous findings and reports new findings with respect to differences between EOAD and LOAD. Specifically, the results implicate the differential early involvement in parietal lobes with probable dysfunction of, not only cognition from parietal gray matter disturbances, but also executive dysfunction presumably from the parietal-frontal network. Much more research is needed in order to clarify this finding and the differences in AD-pathophysiology in EOAD as compared to the much more common, and typical, LOAD. Future studies could profitably evaluate other important areas, such as white matter integrity in order to assess the fronto-parietal network and gene status, as this has a large impact on the development of early-onset dementias⁴⁷.

Acknowledgments

Research supported by NIH Grant #R01AG034499-3 (M.Mendez PI); Department of Veterans Affairs VA Merit Review (2 separate awards; 1 to M. Mendez and another to D. Sultzer); Career Development Award to R. Melrose; NIMH #R01-MH56031 (D. Sultzer,) PI

References

1. Renvoize E, Hanson M, Dale M. Prevalence and causes of young onset dementia in an English health district. *Int J Geriatr Psychiatry*. 2011 Jan; 26(1):106–107. [PubMed: 21157856]
2. Loring DW, Lorgen JW. Neuropsychological patterns of presenile and senile dementia of the Alzheimer type. *Neuropsychologia*. 1985; 23(3):351–357. [PubMed: 4022302]
3. Filley CM, Kelly J, Heaton RK. Neuropsychologic features of early- and late-onset Alzheimer's disease. *Arch Neurol*. 1986 Jun; 43(6):574–576. [PubMed: 3718284]

4. Seltzer BSI. A comparison of clinical features in early-and-late-onset primary degenerative dementia. One entity or two? *Archives of Neurology*. 1983; 40(3):143–146. [PubMed: 6830452]
5. Kim EJ, Cho SS, Jeong Y, et al. Glucose metabolism in early onset versus late onset Alzheimer's disease: an SPM analysis of 120 patients. *Brain*. 2005 Aug; 128(Pt 8):1790–1801. [PubMed: 15888536]
6. Frisoni GB, Testa C, Sabattoli F, Beltramello A, Soininen H, Laakso MP. Structural correlates of early and late onset Alzheimer's disease: voxel based morphometric study. *J Neurol Neurosurg Psychiatry*. 2005 Jan; 76(1):112–114. [PubMed: 15608008]
7. Grady CL, Haxby JV, Horwitz B, Berg G, Rapoport SI. Neuropsychological and cerebral metabolic function in early vs late onset dementia of the Alzheimer type. *Neuropsychologia*. 1987; 25(5):807–816. [PubMed: 3501553]
8. Jagust WJ, Reed BR, Seab JP, Budinger TF. Alzheimer's disease. Age at onset and single-photon emission computed tomographic patterns of regional cerebral blood flow. *Arch Neurol*. 1990 Jun; 47(6):628–633. [PubMed: 2346388]
9. Yasuno F, Imamura T, Hirano N, et al. Age at onset and regional cerebral glucose metabolism in Alzheimer's disease. *Dement Geriatr Cogn Disord*. 1998 Mar-Apr; 9(2):63–67. [PubMed: 9524795]
10. Salmon E, Collette F, Degueldre C, Lemaire C, Franck G. Voxel-based analysis of confounding effects of age and dementia severity on cerebral metabolism in Alzheimer's disease. *Hum Brain Mapp*. 2000 May; 10(1):39–48. [PubMed: 10843517]
11. Sakamoto S, Ishii K, Sasaki M, et al. Differences in cerebral metabolic impairment between early and late onset types of Alzheimer's disease. *J Neurol Sci*. 2002 Aug 15; 200(1–2):27–32. [PubMed: 12127672]
12. Kemp PM, Holmes C, Hoffmann SM, et al. Alzheimer's disease: differences in technetium-99m HMPAO SPECT scan findings between early onset and late onset dementia. *J Neurol Neurosurg Psychiatry*. 2003 Jun; 74(6):715–719. [PubMed: 12754337]
13. Ishii K, Kawachi T, Sasaki H, et al. Voxel-based morphometric comparison between early- and late-onset mild Alzheimer's disease and assessment of diagnostic performance of z score images. *AJNR Am J Neuroradiol*. 2005 Feb; 26(2):333–340. [PubMed: 15709131]
14. Karas G, Scheltens P, Rombouts S, et al. Precuneus atrophy in early-onset Alzheimer's disease: a morphometric structural MRI study. *Neuroradiology*. 2007 Dec; 49(12):967–976. [PubMed: 17955233]
15. Binetti G, Magni E, Padovani A, Cappa SF, Bianchetti A, Trabucchi M. Neuropsychological heterogeneity in mild Alzheimer's disease. *Dementia*. 1993 Nov-Dec; 4(6):321–326. [PubMed: 8136895]
16. Jacobs D, Sano M, Marder K, et al. Age at onset of Alzheimer's disease: relation to pattern of cognitive dysfunction and rate of decline. *Neurology*. 1994 Jul; 44(7):1215–1220. [PubMed: 8035918]
17. Koss E, Edland S, Fillenbaum G, et al. Clinical and neuropsychological differences between patients with earlier and later onset of Alzheimer's disease: A CERAD analysis, Part XII. *Neurology*. 1996 Jan; 46(1):136–141. [PubMed: 8559362]
18. Fujimori M, Imamura T, Yamashita H, et al. Age at onset and visuocognitive disturbances in Alzheimer disease. *Alzheimer Dis Assoc Disord*. 1998 Sep; 12(3):163–166. [PubMed: 9772018]
19. Imamura T, Takatsuki Y, Fujimori M, et al. Age at onset and language disturbances in Alzheimer's disease. *Neuropsychologia*. 1998 Sep; 36(9):945–949. [PubMed: 9740367]
20. Snowden JS, Stopford CL, Julien CL, et al. Cognitive phenotypes in Alzheimer's disease and genetic risk. *Cortex*. 2007 Oct; 43(7):835–845. [PubMed: 17941342]
21. Stopford CL, Snowden JS, Thompson JC, Neary D. Variability in cognitive presentation of Alzheimer's disease. *Cortex*. 2008 Feb; 44(2):185–195. [PubMed: 18387548]
22. Skantzakis E, Tzallas P, Kruse J, Kalpouzos C, Charalambidis D. Coherent continuum extreme ultraviolet radiation in the sub-100-nJ range generated by a high-power many-cycle laser field. *Opt Lett*. 2009 Jun 1; 34(11):1732–1734. [PubMed: 19488164]
23. Toyota Y, Ikeda M, Shinagawa S, et al. Comparison of behavioral and psychological symptoms in early-onset and late-onset Alzheimer's disease. *Int J Geriatr Psychiatry*. 2007 Sep; 22(9):896–901. [PubMed: 17343292]

24. Licht EA, McMurtray AM, Saul RE, Mendez MF. Cognitive differences between early- and late-onset Alzheimer's disease. *Am J Alzheimers Dis Other Dement*. 2007 Jun-Jul;22(3):218–222. [PubMed: 17606531]
25. Suribhatla S, Baillon S, Dennis M, et al. Neuropsychological performance in early and late onset Alzheimer's disease: comparisons in a memory clinic population. *Int J Geriatr Psychiatry*. 2004 Dec; 19(12):1140–1147. [PubMed: 15526308]
26. Reid W, Broe G, Creasey H, et al. Age at onset and pattern of neuropsychological impairment in mild early-stage Alzheimer disease. A study of a community-based population. *Arch Neurol*. 1996 Oct; 53(10):1056–1061. [PubMed: 8859068]
27. Smits LL, Pijnenburg YA, Koedam EL, et al. Early Onset Alzheimer's Disease is Associated with a Distinct Neuropsychological Profile. *J Alzheimers Dis*. 2012 Jan 1; 30(1):101–108. [PubMed: 22366769]
28. Small GW, Kuhl DE, Riege WH, et al. Cerebral glucose metabolic patterns in Alzheimer's disease. Effect of gender and age at dementia onset. *Arch Gen Psychiatry*. 1989 Jun; 46(6):527–532. [PubMed: 2730277]
29. Ichimiya A, Herholz K, Mielke R, Kessler J, Slansky I, Heiss WD. Difference of regional cerebral metabolic pattern between presenile and senile dementia of the Alzheimer type: a factor analytic study. *J Neurol Sci*. 1994 May; 123(1–2):11–17. [PubMed: 8064302]
30. Hanyu H, Nakano S, Abe S, Arai H, Iwamoto T, Takasaki M. [Differences of neuroimaging between early-onset and late-onset Alzheimer-type dementia]. *Rinsho Shinkeigaku*. 1995 Oct; 35(10):1104–1109. [PubMed: 8821493]
31. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*. 1984 Jul; 34(7):939–944. [PubMed: 6610841]
32. Crum RM, Anthony JC, Bassett SS, Folstein MF. Population-based norms for the Mini-Mental State Examination by age and educational level. *JAMA*. 1993 May 12; 269(18):2386–2391. [PubMed: 8479064]
33. Heaton, RKM.; Walden, S.; Taylor, Micheal J.; Grant, Igor. Revised Comprehensive Norms for an Expanded Halstead-Reitan Battery: Demographically Adjusted Neuropsychological Norms for African American and Caucasian Adults. Psychological Assessment Resources, Inc.; 2004.
34. Gladsjo JA, Schuman CC, Evans JD, Peavy GM, Miller SW, Heaton RK. Norms for letter and category fluency: demographic corrections for age, education, and ethnicity. *Assessment*. 1999 Jun; 6(2):147–178. [PubMed: 10335019]
35. Lamar M, Podell K, Carew TG, et al. Perseverative behavior in Alzheimer's disease and subcortical ischemic vascular dementia. *Neuropsychology*. 1997 Oct; 11(4):523–534. [PubMed: 9345696]
36. Jefferson AL, Cosentino SA, Ball SK, et al. Errors produced on the mini-mental state examination and neuropsychological test performance in Alzheimer's disease, ischemic vascular dementia, and Parkinson's disease. *J Neuropsychiatry Clin Neurosci*. 2002 Summer;14(3):311–320. [PubMed: 12154156]
37. Tombaugh TN. Trail Making Test A and B: normative data stratified by age and education. *Arch Clin Neuropsychol*. 2004 Mar; 19(2):203–214. [PubMed: 15010086]
38. Welsh KA, Butters N, Mohs RC, et al. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part V. A normative study of the neuropsychological battery. *Neurology*. 1994 Apr; 44(4):609–614. [PubMed: 8164812]
39. Hamacher K, Coenen HH, Stocklin G. Efficient stereospecific synthesis of no-carrier-added 2-[18F]-fluoro-2-deoxy-D-glucose using aminopolyether supported nucleophilic substitution. *J Nucl Med*. 1986 Feb; 27(2):235–238. [PubMed: 3712040]
40. Woo BK, Harwood DG, Melrose RJ, et al. Executive deficits and regional brain metabolism in Alzheimer's disease. *Int J Geriatr Psychiatry*. 2010 Nov; 25(11):1150–1158. [PubMed: 20069587]
41. Schroeter ML, Vogt B, Frisch S, et al. Executive deficits are related to the inferior frontal junction in early dementia. *Brain*. 2012 Jan; 135(Pt 1):201–215. [PubMed: 22184615]

42. Cosentino S, Jefferson A, Chute DL, Kaplan E, Libon DJ. Clock drawing errors in dementia: neuropsychological and neuroanatomical considerations. *Cogn Behav Neurol*. 2004 Jun; 17(2):74–84. [PubMed: 15453515]
43. Chui HC, Teng EL, Henderson VW, Moy AC. Clinical subtypes of dementia of the Alzheimer type. *Neurology*. 1985 Nov; 35(11):1544–1550. [PubMed: 4058744]
44. Kalpouzos G, Chetelat G, Baron JC, et al. Voxel-based mapping of brain gray matter volume and glucose metabolism profiles in normal aging. *Neurobiol Aging*. 2009 Jan; 30(1):112–124. [PubMed: 17630048]
45. Mielke R, Kessler J, Szekely B, Herholz K, Wienhard K, Heiss WD. Normal and pathological aging--findings of positron-emission-tomography. *J Neural Transm*. 1998; 105(8–9):821–837. [PubMed: 9869321]
46. Grady CL, Berg G, Carson RE, Daube-Witherspoon ME, Friedland RP, Rapoport SI. Quantitative comparison of cerebral glucose metabolic rates from two positron emission tomographs. *J Nucl Med*. 1989 Aug; 30(8):1386–1392. [PubMed: 2787850]
47. Rogers BS, Lippa CF. A clinical approach to early-onset inheritable dementia. *Am J Alzheimers Dis Other Dement*. 2012 May; 27(3):154–161. [PubMed: 22573281]

Figure 1. Base Comparison of EOAD relative to LOAD

Two-sample t-test comparison shows relative hypometabolism in the EAD group (blue) and relative hypometabolism in the LAD group (red) (uncorrected $p=0.01$, $T=2.43$, $k_c=10$ voxels). Picture is in neurological convention; brain's left is your left.

Figure 2. Axial Image of Regional Hypometabolism in Representative EOAD and LOAD patient
Arrows highlight areas of hypometabolism for each respective patient; for EOAD patient, hypometabolism seen in the parietal region whereas in LOAD, hypometabolism seen in the inferior frontal region. Picture is in neurological convention; brain's left is your left.

Table 1

Demographics of early and late-onset Alzheimer’s disease participants

Demographic Variable	Early-onset Alzheimer’s disease N = 21	Late-onset Alzheimer’s Disease N = 24
Age*** (years)	$\overline{M} = 57.78$ ($sd = 4.35$)	$\overline{M} = 80.32$ ($sd = 5.89$)
Gender		
Male	N = 15 (65%)	N = 15 (68%)
Female	N = 8 (35%)	N = 7 (32%)
Education (years)	$\overline{M} = 15.95$ ($sd = 2.63$)	$\overline{M} = 16.05$ ($sd = 2.38$)
Handedness		
Right	N = 20 (91%)	N = 21 (95%)
Left	N = 1 (4%)	N = 1 (4.5%)
Ambidextrous	N = 1 (5%)	N = 0
Time since onset (years)	$\overline{M} = 3.12$ ($sd = 1.61$)	$\overline{M} = 3.91$ ($sd = 2.76$)
Family history of dementia		
Yes	N = 8 (35%)	N = 11 (50%)
No	N = 12 (52%)	N = 11 (50%)
Unsure	N = 3 (13%)	N = 0
Ethnicity*		
White	N = 21 (91%)	N = 15 (68%)
Non-white	N = 2 (9%)	N = 7 (32%)

Note:

* indicates significant between-group difference at the p = 0.05 level;

** indicates significant between-group difference at the p < 0.01 level;

*** indicates significant between-group difference at the p < 0.001 level

Table 2

Neuropsychological comparisons of EAD and LAD patients

Neuropsychological Measure		<i>t</i>	Df	Sig.	EAD <i>M, sd</i>		LAD <i>M, sd</i>	
<i>Global Cognitive Severity</i>								
Mini Mental State Examination (MMSE)		-1.13	41	NS	19.19	6.35	21.09	4.03
<i>Language</i>								
Boston Naming Test (with or without stimulus cue)		3.15	38	A, B, C	49.85	15.58	35.30	13.57
FAS (Phonemic Fluency)		-1.23	42	B, C	21.55	14.64	26.68	13.08
Animals (Semantic Fluency)		0.58	35	NS	8.55	6.51	7.59	3.99
<i>Visuospatial Skills</i>		χ^2	Df	Sig.	% Positive EAD		% Positive LAD	
MMSE Pentagons			1	p = .01	26.3%		0%	
Pull-to-Stimulus Copy		6.59	1					
Size of Figure		0.48	1	NS	10.5%		18.2%	
Number of Figures		1.92	1	NS	31.6%		13.6%	
Pentagon Intersection		2.18	1	NS	68.4%		45.5%	
Trenor or Segmentation		0.98	1	NS	36.8%		22.7%	
Five Angles		0.11	1	NS	36.8%		31.8%	
Rotation		0.02	1	NS	10.5%		9.1%	
Motor Perseveration		0.01	1	NS	26.3%		27.3%	
<i>Attention/ Executive Control</i>		<i>t</i>	Df	Sig.	EAD <i>M, sd</i>		LAD <i>M, sd</i>	
Stroop Task								
Word Reading		.306	35	NS	55.56	21.76	58.58	19.49
Color Naming		.294	35	NS	32.78	16.06	32.63	18.03
Interference (Color and Word)		-1.79	35	B, C	7.83	5.89	13.32	11.59
Trail Making Test-Part A		1.16	13.82	B	165.07	240.66	99.05	52.87
Wechsler Adult Intelligence Scale Digit Span								
Forward		-.831	41	NS	6.77	2.33	7.33	2.08
Backward		-2.36	41	A, B, C	3.00	1.75	4.42	2.20
Total		-1.74	41	NS	9.77	3.83	11.76	3.65
<i>Verbal Learning/Memory</i>								
CERAD								
Word List Memory (trials 1-3)		.131	32	NS	9.00	5.49	8.77	4.44
Word List Recall		1.92	32	NS	1.33	1.78	0.32	1.29

Neuropsychological Measure	<i>t</i>	Df	Sig.	EAD $\overline{M}$, <i>sd</i>	LAD $\overline{M}$, <i>sd</i>
Word List Recognition	2.94	33	A, B, C	2.55	2.84
Savings [(Delayed Recall/Trial 3) × 100]	1.67	33	NS	34.49	32.81

Note: "A", "B", & "C" denote statistically significant findings; "A" denotes significance in raw data analyses (t-tests); "B" indicates significance when comparing normed-data; "NS" denotes not significant in raw data or normed data analyses; "C" indicates maintained significance when adjusting for global cognitive severity scores; CERAD stands for Consortium to Establish a Registry for Alzheimer's Disease

Table 3

Between-group differences in metabolism

	Cluster P (FWE corrected)	k_c	Peak Voxel MNI (x y z)	Peak T
EAD < LAD				
Bilateral Precuneus & Superior parietal lobule, left Inferior parietal lobule, left Posterior temporal cortex, left Occipital lobe	<0.0005	12311	-8, -64, 54	6.6
LAD < EAD				
Bilateral inferior & medial frontal cortex, right temporal cortex	<0.001	29872	52, 24, -2	8.24
Left parahippocampal/fusiform gyri, cerebellum	0.016		-44, -54, -46	5.69