Published in final edited form as:

Psychol Med. 2012 October ; 42(10): 2167–2179. doi:10.1017/S0033291712000426.

Visual information processing dysfunction across the developmental course of early psychosis

V. B. Perez¹, K. M. Shafer², and K. S. Cadenhead^{2,3,*}

¹University of California, San Francisco (UCSF), CA, USA

²University of California, San Diego (UCSD), CA, USA

³Veteran's Affairs San Diego Health Care, La Jolla, CA, USA

Abstract

Background—Patients with schizophrenia consistently demonstrate information processing abnormalities assessed with visual masking (VM) tasks, and these deficits have been linked to clinical and functional severity. It has been suggested that VM impairments may be a vulnerability marker in individuals at risk for developing psychosis.

Method—Forward and backward VM performance was assessed in 72 first-episode (FE) psychosis patients, 98 subjects at risk (AR) for psychosis and 98 healthy controls (HC) using two identification tasks (with either a high- or low-energy mask) and a location task. VM was examined for stability in a subgroup (FE, $n=15$; AR, $n=35$; HC, $n=21$) and assessed relative to clinical and functional measures.

Results—In the identification tasks, backward VM deficits were observed in both FE and AR relative to HC whereas forward VM deficits were only present in FE patients compared to HC. In the location task, AR subjects demonstrated superior performance in forward VM relative to HC. VM performance was stable over time, and VM deficits were associated with baseline functional measures and predicted future negative symptom severity in AR subjects.

Conclusions—Visual information processing deficits, as indexed by backward VM, are present before and after the onset of frank psychosis, and probably represent a stable vulnerability marker that is associated with negative symptoms and functional decline. Additionally, the paradoxically better performance of AR subjects in select forward tasks suggests that early compensatory changes may characterize an emerging psychotic state.

Keywords

At-risk; information processing; prodrome; psychosis; schizophrenia

Introduction

Impairments of visual information processing have been assessed with visual masking (VM) tasks in patients with schizophrenia (Saccuzzo & Braff, 1981; Green *et al.* 1994a,b, 2003; Cadenhead *et al.* 1998; Rassovsky *et al.* 2005). In VM paradigms, a target stimulus is flashed briefly and the detectability of the target is reduced by a masking stimulus that either precedes the target in forward masking or follows the target in backward masking. Although patients with schizophrenia display deficits in both forward (Slaghuis & Bakker, 1995; Slaghuis & Curran, 1999; Green *et al.* 2003; Rassovsky *et al.* 2004) and backward (Braff & Saccuzzo, 1981; Saccuzzo & Braff, 1981; Schwartz *et al.* 1983; Green & Walker, 1986; Knight, 1992; Green *et al.* 2003) VM in comparison to controls, evidence has consistently shown that visual backward masking (VBM) deficits are robust across task manipulations and subtypes of schizophrenia (Saccuzzo *et al.* 1996). Furthermore, masking deficits in both directions have been reported in unmedicated remitted schizophrenia patients (Miller *et al.* 1979; Green *et al.* 1999), unaffected first-degree relatives of schizophrenia patients, and schizotypal personality disordered subjects (Saccuzzo & Schubert, 1981; Saccuzzo *et al.* 1996; Green *et al.* 1997, 2006; Keri *et al.* 2000), suggesting that masking deficits may represent an endophenotypic marker (Gottesman & Gould, 2003).

The prodromal phase of schizophrenia reflects a vulnerable state to the full disorder with evidence of significant clinical and functional disability ; yet these declines in psychosocial functioning may also be present in individuals who do not go on to develop a psychotic illness (Ballon *et al.* 2007; Addington *et al.* 2011). Conversion rates from the prodrome, as defined by empirical clinical criteria (Miller *et al.* 2002, 2003), to psychosis have been reported to be approximately 25–35% over 1 to 2.5 years (Yung *et al.* 2004; Olsen & Rosenbaum, 2006; Cannon *et al.* 2008). As a means of improving the positive predictive power of these criteria, biobehavioral markers (Cornblatt & Erlenmeyer-Kimling, 1985; Gur *et al.* 1998; Cadenhead *et al.* 2002, 2005; Niendam *et al.* 2006b; Turetsky *et al.* 2007; Perez *et al.* 2011) and neurocognitive assessment (Eastvold *et al.* 2007; Jahshan *et al.* 2010) have been implemented as additional strategies to increase the specificity of prodromal criteria, but VM has yet to be assessed in individuals at risk for psychosis meeting these criteria.

To improve the sensitivity of VM as a putative biomarker for psychosis, and to better understand the mechanism by which psychosis develops, the stimuli used in the VM task can be modified to stimulate specific subcortical neuroanatomical pathways (a low spatial frequency mask stimulates the magnocellular and a high spatial frequency mask the parvocellular pathway) that originate in the eye and project to primary visual cortex and the corresponding dorsal and ventral processing streams (Breitmeyer & Ganz, 1976; Breitmeyer & Ogmen, 2000; Van Essen *et al.* 1992). The masking paradigms can require participants to locate (dorsal stream) *versus* identify (ventral stream) target stimuli, activating the cortical component (Balogh & Merritt, 1987; Green *et al.* 1994b; Cadenhead *et al.* 1998; Slaghuis & Curran, 1999). Patients with psychosis exhibit VM deficits when they are required to locate the target stimuli (Green *et al.* 1994b, 2006; Cadenhead *et al.* 1998), suggesting dysfunction in the magnocellular channel and dorsal stream. Other findings of deficits in identification tasks (Purushothaman *et al.* 2000) implicated dysfunction in the parvocellular channel and ventral stream.

The severity of VM abnormalities has been linked to negative symptoms (Green & Walker, 1986; Slaghuis & Bakker, 1995; Slaghuis & Curran, 1999), poorer prognosis for recovery (Saccuzzo & Braff, 1981; Rund *et al.* 1993), and impaired social perception (Sergi & Green, 2003). Furthermore, Green & Braff (2001) discussed the importance of determining how these information processing deficits are related to specific functional outcomes such as social skills acquisition, problem solving, and the ability to function within the community. Therefore, early identification of processing deficits that are predictive of functional outcome may provide treatment targets linked to specific dysfunction in patients in the early stages of psychosis.

The key to using VM as a measure of future clinical severity is its stability over time, and thus far, two studies have assessed VM stability in schizophrenia (Rund *et al.* 1993; Lee *et al.* 2008). Neither of these studies found a significant effect of time on VM in patients tested over a 2-year period. However, in both studies, patient age spanned a large range and was not accounted for in stability measures, confounding the purported stability of VM with normative age-related changes.

The present study is the first to characterize visual information processing in subjects at risk for psychosis relative to first-episode (FE) psychosis patients and healthy controls (HC). We expected to find VM deficits in FE patients, with less pronounced impairments in at-risk (AR) subjects, of whom only a portion are predicted to go on to develop a psychotic disorder. We hypothesized that VM performance would show high stability over time and predict clinical and functional outcome. In exploratory analyses we investigated whether VM deficits are predictive of later psychosis in subjects at risk for psychosis, representing a vulnerability marker for future psychotic illness.

Method

Participants

Baseline assessment included 98 AR subjects, 72 FE patients and 98 HC. Clinical ratings and demographic data are presented in Table 1. Subjects were part of the Cognitive Assessment and Risk Evaluation (CARE) Program at the University of California, San Diego (UCSD). AR subjects were identified as at risk for psychosis based on criteria from the Structured Interview for Prodromal Syndromes (SIPS; Miller *et al.* 2002, 2003), which includes three prodromal syndromes: attenuated psychotic symptoms (APS), brief intermittent psychotic states (BIPS) and/or genetic risk with deterioration in psychosocial functioning (GRD). FE subjects met DSM-IV criteria for a psychotic disorder [52 schizophrenia, 16 psychosis not otherwise specified (NOS), four psychotic mood disorder] according to the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID), with psychosis onset within 24 months. HC were recruited through advertisements. Exclusion criteria for HC consisted of current psychiatric medications, current or past diagnosis of an Axis I disorder, Cluster A personality disorder, or family history of psychotic illness. Exclusion criteria for all participants consisted of past head injury, current drug abuse/dependence, neurological disorder, visual acuity of <20/50, or IQ <70. This study was approved by the Institutional Review Board (IRB) of UCSD, and all participants provided written assent/ consent.

Follow-up data are included for three time points over 2 years. At Time 2 (mean 9.07 ± 5.7 months), participants included 60 AR, 38 FE and 49 HC. At Time 3 (mean 16.4 ± 8.3 months), participants included 35 AR, 15 FE and 21 HC.

Only AR subjects who had follow-up clinical data were included in psychotic conversion *versus* non-conversion analyses. Our conversion rates were: 8/66 (12.1%) at 12 months after baseline, 9/45 (20.0%) at 24 months, and 10/31 (32.3%) at 36 months.

Clinical ratings

Clinical ratings were collected by a clinician researcher within 1 month of VM data collection for Time 1, Time 2 and Time 3. Assessment measures included: the Scale for the Assessment of Positive Symptoms (SAPS; Andreasen, 1984), the Scale for the Assessment of Negative Symptoms (SANS; Andreasen, 1983), the Brief Psychiatric Ratings Scale (BPRS; Overall & Gorham, 1962), the Global Assessment of Functioning (GAF) scale (Hall, 1995), and additionally for AR, the Scale of Prodromal Symptoms (SOPS) from the SIPS (Miller *et al.* 2002, 2003). At Time 1 only, AR subjects and a subset of FE subjects ($n=32$) were scored on the Global Functioning: Role (GF: Role; Niendam *et al.* 2006a) and the Global Functioning: Social (GF: Social; Auther *et al.* 2006) scales.

Masking task

Stimuli (Green *et al.* 2002) were driven at 150 Hz/ 6.67 ms per screen sweep. The target was a square with a gap on one of three sides that appeared in one computer screen quadrant 1° from fixation. The mask comprised four clustered squares that spatially overlapped each possible target location. The three conditions included a high-energy target location task (LOC), a high-energy target identification task (HI), and a low-energy target identification task (LO). Energy is defined as duration $\times$ intensity; a high-energy mask was twice the target (four screen sweeps per mask and two per target) and a low-energy mask was half the target (one screen sweep per mask and two per target). For forward and backward directions, six interstimulus intervals (ISIs; 26, 52, 78, 104, 130, 156 ms) were administered for LOC, and seven ISIs (26, 52, 78, 104, 130, 156, 234 ms) were administered for HI and LO. Trials with simultaneous presentation of target and mask, and unmasked target trials, are included in each condition. Twelve trials were administered for each run (each of three targets presented at each of four locations). Conditions were in a block design with random forward and backward trial presentation. In LOC, participants indicated in which quadrant the target appeared. In HI and LO, participants indicated which side of the target stimulus contained the notch. Staircasing methods (Wetherill & Levitt, 1965) were used to arrive at an individualized perceptual threshold to equate unmasked performance. The gray scale value (i.e. critical stimulus intensity) of the target was systematically varied upward and downward, based on whether the subject's response was correct or not, until performance yielded 84% accuracy. During thresholding, target duration was held constant for 13.3 ms. Subjects who did not perform above chance on unmasked target trials were excluded (three AR and three FE subjects). Analyses were conducted with the remaining participants.

Statistical analyses

Group effects in baseline VM—Separate repeated-measures ANCOVAs were conducted at baseline (Time 1) for each of the three masking conditions (LOC, HI and LO) with group as the between-subjects factor, direction (forward, backward) and ISI as within-subject factors, and age as a covariate. Significant interactions involving group and direction were parsed with follow-up one-way ANCOVAs. Greenhouse–Geisser and Bonferroni corrections were used where appropriate.

Conversion effects in baseline VM—AR subjects were parsed into individuals who later converted to psychosis and those who did not develop a psychotic disorder within 12 months of clinical follow-up. Because of the small sample sizes (converters, $n=8$; non-converters, $n=66$), we developed *a priori* hypotheses to examine VM performance, with emphasis at those ISIs that best differentiated AR and FE from HC. As mentioned earlier, patients with schizophrenia display both forward and backward deficits, although some studies suggest more severe impairment in backward masking (Saccuzzo *et al.* 1996). Based on these previous findings, converter/ non-converter analyses were limited to VBM performance.

Stability of VM—Data were collected at three time points over 2 years. To examine the stability of masking effects over time, a repeated-measures ANCOVA was conducted separately for forward and backward conditions for each task (LOC, HI, LO) with group as the between-subjects factor, age as a covariate, and time (Time 1, Time 2, Time 3), direction (forward, backward) and ISI (six intervals for LOC, seven intervals for HI and LO conditions) as within-subject factors. Only those subjects (35 AR, 15 FE, 21 HC) who had data for each time point were included. To assess potential contributions of attrition in the characteristics of the longitudinal sample, subjects who were followed for each of the three time points (completers) were compared to those who did not remain in the study (non-completers) on demographic, clinical and functional measures.

Correlates of VM to clinical and functional outcome—To examine the relationship between baseline VM and clinical and functional outcome, each of the six VM conditions (forward and backward LOC, HI, LO) at Time 1 was averaged independently across ISI and entered into a Pearson correlation with clinical and functional measures at Time 1 (98 AR, 72 FE, 98 HC), Time 2 (60 AR, 38 FE, 49 HC) and Time 3 (35 AR, 15 FE, 21 HC) for each patient group separately. Significant correlations ($p<0.05$) informed follow-up analyses, where correlated baseline VM conditions were entered as predictors of future clinical/ functional outcome into backward multiple regression analyses for Time 2 and Time 3 separately. The assumptions of normality, linearity and homoscedasticity of residuals were met.

Results

Group characteristics

As seen in Table 1, gender and age differed statistically between groups and were included as a between-subjects factor and a covariate, respectively, in VM analyses. *Post-hoc* tests

revealed that AR tended to be younger than FE ($p<0.001$) and HC ($p<0.01$), but FE and HC did not differ from each other. AR presented with fewer overall symptoms and higher functioning than FE patients.

Group effects for critical stimulus intensity and unmasked targets

VM performance is shown in Fig. 1. Values for critical stimulus intensity were comparable across groups ($p>0.5$) and accuracy for the unmasked targets did not differ between groups in any condition (all $p>0.1$), suggesting that non-specific factors such as task difficulty or baseline performance differences did not affect group differences in VM.

Group masking effects

Analyses for group, age, direction and ISI are shown in Table 2.

Gender—There was a significant gender effect in overall VM performance in LOC ($p<0.01$; females<males) but not in HI or LO. Gender did not significantly interact with group, direction, ISI, time or age in any condition in any analysis. Subsequent report of results, therefore, does not include gender in the model.

Age—We failed to find a significant group $\times$ age interaction in any of the ANCOVA models, indicating that each group's VM performance was related to age equivalently, and the interaction term was dropped from the model. Subsequent analyses revealed significant improvement in HI and LO performance with increased age, but not in LOC.

Direction—A significant direction effect was present in LOC, but not in the HI or LO masking conditions. Significant group $\times$ direction interactions in each condition justified the decision to parse each task with follow-up ANCOVAs for each direction.

Location mask—We observed main effects of group and ISI, but the group $\times$ ISI interaction did not reach significance in either direction. Of note, planned contrasts suggested that AR performed better than HC ($p<0.05$) in forward masking, yet, in backward masking, AR performed worse than HC ($p<0.05$). FE did not differ from HC in either direction.

Identification: high-energy mask—Forward masking analyses showed main effects of group and ISI, but no group $\times$ ISI interaction was found. Across ISIs, FE performance was reduced significantly compared to HC ($p<0.05$), although AR did not differ statistically from HC. In backward masking, there were main effects of group and ISI. Planned contrasts showed that FE patients performed worse than HC ($p<0.05$), and AR performed intermediate to HC and FE, but were not significantly different from either group. We also found a group $\times$ ISI interaction, justifying a parsing of ISI. *Post-hoc* tests revealed that FE performed worse than HC in the backward 104 ms ($p<0.05$), 130 ms ($p<0.005$) and 234 ms ($p<0.005$) ISI conditions, and AR performed worse than HC in the 130 ms ($p=0.052$) and 156 ms ($p<0.05$) ISI conditions. There were no differences in VM between FE and AR when each ISI was examined individually.

Identification: low-energy mask—Forward masking analyses revealed a significant main effect of group, a marginal effect of ISI, but no interaction effect was observed. FE performance was significantly reduced compared to HC ($p<0.001$), but HC and AR groups did not differ from each other. In backward masking, we found main effects of group and ISI. We also found a marginal group $\times$ ISI interaction. *Post-hoc* tests revealed that FE performed worse than HC in the backward 78 ms ($p<0.001$), 130 ms ($p<0.05$), 156 ms ($p<0.01$) and 234 ms ($p<0.05$) ISI conditions, and AR performed worse than HC in the 52 ms ($p<0.05$), 78 ms ($p<0.001$), 104 ms ($p<0.01$) and 130 ms ($p<0.005$) ISI conditions. There were no differences in backward VM between FE and AR when each ISI was examined individually.

Masking performance among converters to psychosis

Because of the small sample sizes, we conducted exploratory analyses of converter ($n=8$) and non-converter ($n=66$) differences at Time 1 based on *a priori* hypotheses that pairwise differences would be greatest in conditions that best differentiated AR and FE from HC in VBM. Therefore, planned contrasts were performed at each ISI between 52 and 234 ms in the backward conditions. Converters showed a significant reduction in baseline VBM performance relative to non-converters at the 78 ms ISI in the backward HI condition ($p<0.05$, Cohen's $d=0.91$), and at the 130 ms ISI in the backward LO condition ($p<0.05$, Cohen's $d=0.66$), as shown in Fig. 2.

Effect of time on masking across groups

VM performance by ISI for each condition is presented for each group for Time 1, Time 2 and Time 3 in the sample that received all three test sessions in Fig. 3. In the backward LOC condition exclusively, a main effect of time was observed (LOC: $F_{2,142}=3.4$, $p<0.05$), where all groups improved over time. There were no significant interactions involving time with group, age, or ISI in any condition, indicating stable performance over time for all groups and across ISI. Furthermore, correlational analyses examining performance at Time 1 with performance at Time 2 indicated that good performers remained good, and poor performers remained poor (r values range from 0.3 to 0.7 across all VM tasks, all $p<0.001$). Similar associations occurred between performance at Time 1 and Time 3 (r values range from 0.3 to 0.6 across all VM tasks, all $p<0.005$).

Attrition—To address any potential contributions of attrition in the baseline sample relative to the longitudinal sample included in the stability analyses, we examined baseline demographic, clinical and functional measures between subjects who remained in the study compared to those who did not complete all three assessments (Table 1). Subjects who completed the study were somewhat younger than those who did not complete each of the three time points ($F_{1,267}=4.2$, $p<0.05$) and, as such, age remained a covariate in stability analyses. Demographics did not differ on any other dimension (parental education, gender, handedness all $p<0.2$). Furthermore, non-completers showed no evidence of increased symptom severity or psychosocial decline: FE and AR completers did not differ from non-completers on symptoms (all $p>0.35$ and $p>0.2$, respectively) or functional measures (all $p>0.7$ and $p>0.3$, respectively).

Potential medication effects

To address effects of dopamine D₂-receptor blocking antipsychotic medication, exploratory analyses comparing antipsychotic medicated ($n=53$) to unmedicated ($n=19$) FE patients, and antipsychotic medicated ($n=19$) to unmedicated ($n=79$) AR subjects, were conducted. No VM differences were observed in either group.

Relationship between masking performance and functional ability

Baseline VM was not correlated with GAF scores for either patient group at any time point. In AR, poor role functioning at Time 1 was associated with worse performance on baseline forward identification tasks (HI: $r=0.23$, $p<0.05$; LO: $r=0.28$, $p<0.01$). In a subset of FE, poor performance on forward identification tasks (HI: $r=0.44$, $p<0.01$; LO: $r=0.36$, $p<0.05$) was correlated with social functioning, but not role functioning at Time 1. Masking performance was not predictive of future functional outcome in role or social domains in either patient group. Because of the lack of association between baseline masking and future functional measures, predictive regression analyses were not conducted.

Prediction of clinical symptom profile

In FE, no significant correlations were observed between clinical symptoms and VM in any condition at baseline or follow-up. In AR, baseline VM was not correlated with clinical symptoms at Time 1. However, baseline VM was associated with total negative symptoms on the SANS in AR at Time 2 (forward HI: $r=-0.27$, $p=0.036$; backward HI: $r=-0.31$, $p=0.015$; forward LO: $r=-0.37$, $p<0.01$; backward LO: $r=-0.31$, $p=0.015$) and Time 3 (forward HI: $r=-0.33$, $p=0.038$; backward HI: $r=-0.39$, $p=0.014$; forward LO: $r=-0.51$, $p<0.001$; backward LO: $r=-0.28$, $p=0.055$). VM was not correlated with positive symptoms in either patient group.

VM and clinical symptom profile did not reveal any association in the FE group. As such, follow-up regression analyses were only performed in the AR group. Two separate backward multiple regression analyses examined future negative symptoms at Time 2 and Time 3. Baseline (Time 1) forward HI and LO and backward HI and LO conditions were entered as predictors. The results of the Time 2 initial model accounted for 17.1% of the variance in negative symptoms, and showed that forward LO masking was negatively predictive, indicating that AR with worse performance were expected to have greater negative symptoms at Time 2 ($\beta=-0.146$, $p<0.05$). All other variables did not contribute significantly to the model, and secondary stepwise models were not predictive above and beyond the initial model ($F_{\text{change } 1,59}=0.01$, $p=0.97$, $R^2=0.00$). In the Time 3 model, we observed a significant regression coefficient that accounted for 33.5% of the variance in negative symptoms ($F_{4,34}=5.1$, $p<0.02$, $R^2=0.335$, at step 1). Forward LO ($\beta=-0.25$, $p<0.01$) and backward HI ($\beta=-0.21$, $p<0.05$) VM significantly predicted negative symptoms at Time 3. Additional stepwise models were not significantly predictive above and beyond the initial model ($F_{\text{change } 1,34}=0.6$, $p=0.46$, $R^2=-0.01$).

Discussion

Using a VM paradigm, we examined information processing across putative developmental phases of schizophrenia. Previously, it has been shown that performance in VM tasks was diminished across the schizophrenia spectrum (Green *et al.* 1994a,b, 1997, 1999; Rassovsky *et al.* 2004; Lee *et al.* 2008). The current study replicates these findings by showing abnormal masking performance in schizophrenia patients early in their illness course, and further extends the current literature to show that these deficits are present in individuals at risk for psychosis, although to a lesser degree.

Findings of less pronounced neurobiological abnormalities in AR relative to FE may reflect the heterogeneity of putatively prodromal patients and the fact that only a small percentage will convert to schizophrenia (Cannon *et al.* 2008). However, the low-energy backward masking condition showed that AR performance resembled the performance of FE, and that both of these groups showed impairment relative to HC. In addition, AR subjects with the most severe impairments in the identification tasks were more likely to convert to psychosis (although the small samples necessitate cautious interpretation). Notably, our observation of impairment in AR subjects across backward location and identification tasks implicates both the dorsal and ventral processing streams, as observed in FE patients.

An unexpected finding was that, although the AR sample had deficits in the backward location VM task relative to the HC sample, they had superior performance in the forward location task. Patients with schizophrenia spectrum disorders are known to have difficulty integrating perceptual information, leading to superior performance on tasks that require interpretation of individual features of a stimulus or visual illusion (Parnas *et al.* 2001; Silverstein *et al.* 2006). Although we can speculate that AR patients, who are already showing signs of clinical symptomatology and/or psychosocial deterioration, are showing evidence of early difficulties with perceptual integration on this task, clarification of the mechanism underlying superior performance warrants further study.

Theoretically, age-related differences in VM may be reflective of the typical developmental course of speed and accuracy improvements in information processing. Some (Haith, 1971; Miller, 1972; Welsandt *et al.* 1973) but not all (Buss *et al.* 1999; Green *et al.* 2003) studies have found an inverse relationship between age and susceptibility to the disruptive effects of the mask. Importantly, VM performance was analyzed in parallel with aging, and over time. We confirmed that VM performance improved with age equally across groups. Thus, the magnitude of the deficits observed in FE relative to those observed in AR is not due to age-related decline. Furthermore, age did not interact significantly with time for any group.

In the sample who received repeated assessment, VM performance was stable, with significant correlations over time on each task. These results are consistent with previous studies examining the stability of the VM paradigm (Rund *et al.* 1993; Lee *et al.* 2008). Therefore, stability analyses suggest that VM remains a useful biomarker as an endophenotype and as a vulnerability marker for psychosis.

Disruptions in information processing in schizophrenia have been linked to specific clinical and functional outcomes such as negative symptoms, social skills acquisition, problem

solving and the ability to function within the community (Cadenhead *et al.* 1997; Slaghuys & Curran, 1999; Green & Braff, 2001; Sergi & Green, 2003). The present evidence converges with other studies (Ballon *et al.* 2007; Cornblatt *et al.* 2007) indicating that VM deficits observed in patient groups are associated with a poorer level of functioning, and that these declines begin even before the onset of psychosis. Regarding symptom profile, we did not find an association between symptoms and VM in FE, consistent with other studies (Green *et al.* 2003), although many studies do not report these relationships (Saccuzzo *et al.* 1996; Rassovsky *et al.* 2004; Lee *et al.* 2008; Green *et al.* 2009). Additionally, we did not find a relationship between performance and AR symptom profile at baseline. However, poor VM in each identification task was associated with future negative symptoms in AR. Such findings suggest that masking performance may identify not only those individuals who go on to develop a psychotic illness but also those who develop deficit symptoms and associated psychosocial decline (Cannon *et al.* 2008). The relationship between VM impairment and negative symptomatology that emerges prior to the onset of psychosis may be helpful in identifying those at highest risk for severe illness.

We have shown that VM tasks are reliably able to detect information processing impairments in FE patients, and in subjects at risk for developing psychosis, and that these deficits are not due to the influence of aging. By comparing masking performance across different tasks, the current findings show that neural mechanisms underlying both forward and backward masking are dysfunctional (Perkins *et al.* 2005; Barnes *et al.* 2008), and that information processing deficits are evident in tasks that favor both the dorsal and ventral processing streams, even before psychosis onset. As target and mask parameters can be modified to emphasize different information processing streams (i.e. dorsal *versus* ventral pathways), using specific target–mask combinations that are highly sensitive to information processing deficits across the phases of schizophrenia may be most effective for detecting those AR patients who may go on to develop psychosis. Impaired VM performance in putatively prodromal patients may help to identify traits that represent surrogate end-points, and lead to targeted early intervention of specific deficits.

Acknowledgments

We thank M. Green and K. Neuchterlein for allowing us to use the computerized visual masking paradigm they developed at University of California, Los Angeles (UCLA). In addition, we thank the clinicians who provided careful assessment of all at-risk and first-episode subjects (N. Haroun, K. Kristensen, A. Hurria, T. Kaur, A. Eslami, C. Jashan, T. Alderman and I. Domingues) and the dedicated research assistants who spent many hours with our subjects and their data (A. Eastvold, I. Marks, J. Nunag and D. Roman). This study was supported by National Institutes of Health (NIH)-supported grants R01 MH60720 and K24 MH76191.

References

- Addington J, Cornblatt BA, Cadenhead KS, Cannon TD, McGlashan TH, Perkins DO, Seidman LJ, Tsuang MT, Walker EF, Woods SW, Heinssen R. At clinical high risk for psychosis : outcome for nonconverters. *American Journal of Psychiatry*. 2011; 168:800–805. [PubMed: 21498462]
- Andreasen, NC. The Scale for the Assessment of Negative Symptoms (SANS). Iowa City, IA: University of Iowa; 1983.
- Andreasen, NC. The Scale for the Assessment of Positive Symptoms (SAPS). Iowa City, IA: University of Iowa; 1984.

- Annett, M. Left, Right, Hand, and Brain: The Right Shift Theory. Hove, UK: Lawrence Erlbaum Associates Ltd; 1985.
- Auther, A.; Smith, CW.; Cornblatt, BA. Global Functioning : Social Scale (GF: Social). Glen Oaks, NY: Zucker-Hillside Hospital; 2006.
- Ballon JS, Kaur T, Marks II, Cadenhead KS. Social functioning in young people at risk for schizophrenia. *Psychiatry Research*. 2007; 151:29–35. [PubMed: 17383739]
- Balogh DW, Merritt RD. Visual masking and the schizophrenia spectrum: interfacing clinical and experimental methods. *Schizophrenia Bulletin*. 1987; 13:679–698. [PubMed: 3326163]
- Barnes TR, Leeson VC, Mutsatsa SH, Watt HC, Hutton SB, Joyce EM. Duration of untreated psychosis and social function : 1-year follow-up study of first-episode schizophrenia. *British Journal of Psychiatry*. 2008; 193:203–209. [PubMed: 18757977]
- Braff DL, Saccuzzo DP. Information processing dysfunction in paranoid schizophrenia : a two-factor deficit. *American Journal of Psychiatry*. 1981; 138:1051–1056. [PubMed: 7258380]
- Breitmeyer BG, Ganz L. Implications of sustained and transient channels for theories of visual pattern masking, saccadic suppression, and information processing. *Psychological Review*. 1976; 83:1–36. [PubMed: 766038]
- Breitmeyer BG, Ogmen H. Recent models and findings in visual backward masking: a comparison, review, and update. *Perception & Psychophysics*. 2000; 62:1572–1595. [PubMed: 11140180]
- Buss E, Hall JW 3rd, Grose JH, Dev MB. Development of adult-like performance in backward, simultaneous, and forward masking. *Journal of Speech, Language, and Hearing Research*. 1999; 42:844–849.
- Cadenhead KS, Geyer MA, Butler RW, Perry W, Sprock J, Braff DL. Information processing deficits of schizophrenia patients : relationship to clinical ratings, gender and medication status. *Schizophrenia Research*. 1997; 28:51–62. [PubMed: 9428064]
- Cadenhead KS, Light GA, Geyer MA, McDowell JE, Braff DL. Neurobiological measures of schizotypal personality disorder : defining an inhibitory endophenotype? *American Journal of Psychiatry*. 2002; 159:869–871. [PubMed: 11986147]
- Cadenhead KS, Light GA, Shafer KM, Braff DL. P50 suppression in individuals at risk for schizophrenia: the convergence of clinical, familial, and vulnerability marker risk assessment. *Biological Psychiatry*. 2005; 57:1504–1509. [PubMed: 15953486]
- Cadenhead KS, Serper Y, Braff DL. Transient versus sustained visual channels in the visual backward masking deficits of schizophrenia patients. *Biological Psychiatry*. 1998; 43:132–138. [PubMed: 9474445]
- Cannon TD, Cadenhead K, Cornblatt B, Woods SW, Addington J, Walker E, Seidman LJ, Perkins D, Tsuang M, McGlashan T, Heinssen R. Prediction of psychosis in youth at high clinical risk : a multisite longitudinal study in North America. *Archives of General Psychiatry*. 2008; 65:28–37. [PubMed: 18180426]
- Cornblatt BA, Auther AM, Niendam T, Smith CW, Zinberg J, Bearden CE, Cannon TD. Preliminary findings for two new measures of social and role functioning in the prodromal phase of schizophrenia. *Schizophrenia Bulletin*. 2007; 33:688–702. [PubMed: 17440198]
- Cornblatt BA, Erlenmeyer-Kimling L. Global attentional deviance as a marker of risk for schizophrenia : specificity and predictive validity. *Journal of Abnormal Psychology*. 1985; 94:470–486. [PubMed: 3865953]
- Eastvold AD, Heaton RK, Cadenhead KS. Neurocognitive deficits in the (putative) prodrome and first episode of psychosis. *Schizophrenia Research*. 2007; 93:266–277. [PubMed: 17467955]
- Gottesman II, Gould TD. The endophenotype concept in psychiatry : etymology and strategic intentions. *American Journal of Psychiatry*. 2003; 160:636–645. [PubMed: 12668349]
- Green MF, Braff DL. Translating the basic and clinical cognitive neuroscience of schizophrenia to drug development and clinical trials of antipsychotic medications. *Biological Psychiatry*. 2001; 49:374–384. [PubMed: 11239909]
- Green MF, Lee J, Cohen MS, Engel SA, Korb AS, Nuechterlein KH, Wynn JK, Glahn DC. Functional neuroanatomy of visual masking deficits in schizophrenia. *Archives of General Psychiatry*. 2009; 66:1295–1303. [PubMed: 19996034]

- Green MF, Nuechterlein KH, Breitmeyer B. Backward masking performance in unaffected siblings of schizophrenic patients. Evidence for a vulnerability indicator. *Archives of General Psychiatry*. 1997; 54:465–472. [PubMed: 9152100]
- Green MF, Nuechterlein KH, Breitmeyer B. Development of a computerized assessment for visual masking. *International Journal of Methods in Psychiatric Research*. 2002; 11:83–89. [PubMed: 12459798]
- Green MF, Nuechterlein KH, Breitmeyer B, Mintz J. Backward masking in unmedicated schizophrenic patients in psychotic remission : possible reflection of aberrant cortical oscillation. *American Journal of Psychiatry*. 1999; 156:1367–1373. [PubMed: 10484946]
- Green MF, Nuechterlein KH, Breitmeyer B, Mintz J. Forward and backward visual masking in unaffected siblings of schizophrenic patients. *Biological Psychiatry*. 2006; 59:446–451. [PubMed: 16139818]
- Green MF, Nuechterlein KH, Breitmeyer B, Tsuang J, Mintz J. Forward and backward visual masking in schizophrenia : influence of age. *Psychological Medicine*. 2003; 33:887–895. [PubMed: 12877403]
- Green MF, Nuechterlein KH, Mintz J. Backward masking in schizophrenia and mania. I. Specifying a mechanism. *Archives of General Psychiatry*. 1994a; 51:939–944. [PubMed: 7979881]
- Green MF, Nuechterlein KH, Mintz J. Backward masking in schizophrenia and mania. II. Specifying the visual channels. *Archives of General Psychiatry*. 1994b; 51:945–951. [PubMed: 7979882]
- Green MF, Walker E. Symptom correlates of vulnerability to backward masking in schizophrenia. *American Journal of Psychiatry*. 1986; 143:181–186. [PubMed: 3946651]
- Gur RE, Cowell P, Turetsky BI, Gallacher F, Cannon T, Bilker W, Gur RC. A follow-up magnetic resonance imaging study of schizophrenia. Relationship of neuroanatomical changes to clinical and neurobehavioral measures. *Archives of General Psychiatry*. 1998; 55:145–152. [PubMed: 9477928]
- Haith MM. Developmental changes in visual information processing and short-term visual memory. *Human Development*. 1971; 14:249–261. [PubMed: 5126591]
- Hall RC. Global assessment of functioning. A modified scale. *Psychosomatics*. 1995; 36:267–275. [PubMed: 7638314]
- Jahshan C, Heaton RK, Golshan S, Cadenhead KS. Course of neurocognitive deficits in the prodrome and first episode of schizophrenia. *Neuropsychology*. 2010; 24:109–120. [PubMed: 20063952]
- Keri S, Antal A, Szekeres G, Benedek G, Janka Z. Visual information processing in patients with schizophrenia : evidence for the impairment of central mechanisms. *Neuroscience Letters*. 2000; 293:69–71. [PubMed: 11065140]
- Knight RA. Specifying cognitive deficiencies in premorbid schizophrenics. *Progress in Experimental Personality and Psychopathology Research*. 1992; 15:252–289. [PubMed: 1343108]
- Lee J, Nuechterlein KH, Subotnik KL, Sugar CA, Ventura J, Gretchen-Doorly D, Kelly K, Green MF. Stability of visual masking performance in recent-onset schizophrenia : an 18-month longitudinal study. *Schizophrenia Research*. 2008; 103:266–274. [PubMed: 18450427]
- Miller LK. Visual masking and developmental differences in information processing. *Child Development*. 1972; 43:704–709. [PubMed: 5034742]
- Miller S, Saccuzzo D, Braff D. Information processing deficits in remitted schizophrenics. *Journal of Abnormal Psychology*. 1979; 88:446–449. [PubMed: 479468]
- Miller TJ, McGlashan TH, Rosen JL, Cadenhead K, Cannon T, Ventura J, McFarlane W, Perkins DO, Pearlson GD, Woods SW. Prospective diagnosis of the initial prodrome for schizophrenia based on the Structured Interview for Prodromal Syndromes: preliminary evidence of interrater reliability and predictive validity. *American Journal of Psychiatry*. 2002; 159:863–865. [PubMed: 11986145]
- Miller TJ, McGlashan TH, Rosen JL, Cadenhead K, Cannon T, Ventura J, McFarlane W, Perkins DO, Pearlson GD, Woods SW. Prodromal assessment with the structured interview for prodromal syndromes and the scale of prodromal symptoms: predictive validity, interrater reliability, and training to reliability. *Schizophrenia Bulletin*. 2003; 29:703–715. [PubMed: 14989408]
- Niendam, TA.; Bearden, CE.; Johnson, JK.; Cannon, TD. Global Functioning : Role Scale (GF: Role). Los Angeles, CA: L. A. University of California; 2006a.

- Niendam TA, Bearden CE, Johnson JK, McKinley M, Loewy R, O'Brien M, Nuechterlein KH, Green MF, Cannon TD. Neurocognitive performance and functional disability in the psychosis prodrome. *Schizophrenia Research*. 2006b; 84:100–111. [PubMed: 16563699]
- Olsen KA, Rosenbaum B. Prospective investigations of the prodromal state of schizophrenia : assessment instruments. *Acta Psychiatrica Scandinavica*. 2006; 113:273–282. [PubMed: 16638071]
- Overall JE, Gorham DR. The Brief Psychiatric Rating Scale. *Psychological Reports*. 1962; 10:790–812.
- Parnas J, Vianin P, Saebye D, Jansson L, Volmer-Larsen A, Bovet P. Visual binding abilities in the initial and advanced stages of schizophrenia. *Acta Psychiatrica Scandinavica*. 2001; 103:171–180. [PubMed: 11240573]
- Perez VB, Ford JM, Roach BJ, Woods SW, McGlashan TH, Srihari VH, Loewy RL, Vinogradov S, Mathalon DH. Error monitoring dysfunction across the illness course of schizophrenia. *Journal of Abnormal Psychology*. 2011 Published online: 7 November 2011.
- Perkins DO, Gu H, Boteva K, Lieberman JA. Relationship between duration of untreated psychosis and outcome in first-episode schizophrenia : a critical review and meta-analysis. *American Journal of Psychiatry*. 2005; 162:1785–1804. [PubMed: 16199825]
- Purushothaman G, Ogmen H, Bedell HE. Gamma-range oscillations in backward-masking functions and their putative neural correlates. *Psychological Review*. 2000; 107:556–577. [PubMed: 10941280]
- Rassovsky Y, Green MF, Nuechterlein KH, Breitmeyer B, Mintz J. Paracontrast and metacontrast in schizophrenia : clarifying the mechanism for visual masking deficits. *Schizophrenia Research*. 2004; 71:485–492. [PubMed: 15474919]
- Rassovsky Y, Green MF, Nuechterlein KH, Breitmeyer B, Mintz J. Modulation of attention during visual masking in schizophrenia. *American Journal of Psychiatry*. 2005; 162:1533–1535. [PubMed: 16055778]
- Rund BR, Landrø NI, Orbeck AL. Stability in backward masking performance in schizophrenics, affectively disturbed patients, and normal subjects. *Journal of Nervous and Mental Disease*. 1993; 181:233–237. [PubMed: 8473875]
- Saccuzzo DP, Braff DL. Early information processing deficit in schizophrenia. New findings using schizophrenic subgroups and manic control subjects. *Archives of General Psychiatry*. 1981; 38:175–179. [PubMed: 7212944]
- Saccuzzo DP, Schubert DL. Backward masking as a measure of slow processing in schizophrenia spectrum disorders. *Journal of Abnormal Psychology*. 1981; 90:305–312. [PubMed: 7264060]
- Saccuzzo DS, Cadenhead KS, Braff DL. Backward versus forward visual masking deficits in schizophrenic patients : centrally, not peripherally, mediated? *American Journal of Psychiatry*. 1996; 153:1564–1570. [PubMed: 8942452]
- Schwartz BD, Winstead DK, Adinoff B. Temporal integration deficit in visual information processing by chronic schizophrenics. *Biological Psychiatry*. 1983; 18:1311–1320. [PubMed: 6652164]
- Sergi MJ, Green MF. Social perception and early visual processing in schizophrenia. *Schizophrenia Research*. 2003; 59:233–241. [PubMed: 12414080]
- Silverstein S, Uhlhaas PJ, Essex B, Halpin S, Schall U, Carr V. Perceptual organization in first episode schizophrenia and ultra-high-risk states. *Schizophrenia Research*. 2006; 83:41–52. [PubMed: 16497484]
- Slaghuis WL, Bakker VJ. Forward and backward visual masking of contour by light in positive- and negative-symptom schizophrenia. *Journal of Abnormal Psychology*. 1995; 104:41–54. [PubMed: 7897052]
- Slaghuis WL, Curran CE. Spatial frequency masking in positive- and negative-symptom schizophrenia. *Journal of Abnormal Psychology*. 1999; 108:42–50. [PubMed: 10066991]
- Turetsky BI, Calkins ME, Light GA, Olincy A, Radant AD, Swerdlow NR. Neurophysiological endophenotypes of schizophrenia : the viability of selected candidate measures. *Schizophrenia Bulletin*. 2007; 33:69–94. [PubMed: 17135482]
- Van Essen DC, Anderson CH, Felleman DJ. Information processing in the primate visual system: an integrated systems perspective. *Science*. 1992; 255:419–423. [PubMed: 1734518]

- Welsandt RF Jr, Zupnick JJ, Meyer PA. Age effects in backward visual masking (Crawford paradigm). *Journal of Experimental Child Psychology*. 1973; 15:454–461. [PubMed: 4714976]
- Wetherill GB, Levitt H. Sequential estimation of points on a psychometric function. *British Journal of Mathematical and Statistical Psychology*. 1965; 18:1–10. [PubMed: 14324842]
- Yung AR, Phillips LJ, Yuen HP, McGorry PD. Risk factors for psychosis in an ultra high-risk group : psychopathology and clinical features. *Schizophrenia Research*. 2004; 67:131–142. [PubMed: 14984872]

Fig. 1.

Visual masking (VM) performance for first-episode (FE) patients (green), at-risk (AR) subjects (red) and healthy controls (HC; blue) is shown for all three masking conditions (Location, Identification : high-energy masking, Identification : low-energy masking). Interstimulus intervals between the target and the mask are shown on the x axis. Forward masking trials present the mask and then the target, and are depicted to the left of the 0 ms time point. Backward masking trials present the target and then the mask, and are depicted to the right of the 0 ms time point. At time 0 ms, the target and the mask are presented

concurrently. The target presented alone without the mask occurs at the ‘unmasked’ time point. Performance accuracy (percent correct) is shown on the y axis.

Fig. 2.

Visual backward masking (VBM) accuracy (group means and standard error bars) is displayed for the interstimulus intervals in each masking condition predicted to show the greatest deficits in patient groups. *Top panels:* VBM performance is shown across first-episode patients (green), at-risk subjects (red) and healthy controls (blue). *Bottom panels:* From the at-risk group, follow-up clinical data 12 months after baseline parses VBM performance for converters (crimson) and non-converters (gray) to psychosis. Data show poorer VBM performance among converters to psychosis. * $p < 0.05$.

Fig. 3.

Visual masking (VM) performance is displayed at each interstimulus interval for each group at Time 1 (black), Time 2 (magenta) and Time 3 (blue). In the location condition, chance performance has an accuracy of 25%. For both identification tasks, chance performance has an accuracy of 33%. The sample size at Time 1 included 98 healthy controls (HC), 72 at-risk (AR) subjects, and 98 first-episode (FE) patients. At Time 2, samples comprised 49 HC, 60 AR subjects and 38 FE patients. At Time 3, samples comprised 21 HC, 35 AR subjects and 15 FE patients.

Table 1

Group demographics^a

	Baseline sample ^b				Longitudinal sample ^c			
	Healthy controls (n=98)	FE patients (n=72)	At-risk subjects (n=98)	<i>F</i> / χ^2	<i>p</i> value	Healthy controls (n=21)	FE patients (n=15)	At-risk subjects (n=35)
Age (years)	20.7 (4.5)	21.1 (3.8)	18.8 (4.2)	7.9	0.001	19.0 (5.2)	20.7 (5.1)	18.8 (4.0)
Parental education (years)	14.9 (2.6)	14.5 (2.8)	14.0 (2.7)	2.2	0.113	14.8 (1.6)	14.6 (3.3)	14.7 (2.3)
Gender				14.9	0.001			
Female	52 (53.1)	17 (23.6)	40 (40.8)			12 (57.1)	4 (26.7)	10 (28.6)
Male	46 (46.9)	55 (76.4)	58 (59.2)			9 (42.9)	11 (73.3)	25 (71.4)
Handedness ^d				3.6	0.465			
Right	88 (89.8)	53 (73.6)	84 (85.7)			19 (90.5)	11 (73.3)	29 (82.9)
Left	7 (7.1)	9 (12.5)	7 (7.1)			1 (4.8)	3 (20.0)	3 (8.6)
Both	3 (3.1)	3 (4.2)	4 (4.1)			1 (4.8)	1 (6.7)	3 (8.6)
Prodromal criteria ^e								
APS			87 (88.8)					24 (68.6)
BIPS			9 (9.2)					3 (8.6)
GRD			34 (34.7)					5 (14.3)
No longer met criteria								5 (14.3)
Antipsychotic type								
Atypical		50 (69.4)	19 (19.4)				14 (73.7)	7 (20.0)
Typical		1 (1.4)	–				1 (5.3)	–
Both		2 (2.8)	–				–	–
None		19 (26.4)	79 (80.6)				4 (21.1)	28 (80.0)
SAPS total ^f								
Time 1		6.8 (4.3)	5.2 (3.2)				6.5 (4.3)	5.4 (2.8)
Time 2							5.3 (4.4)	2.8 (2.1)
Time 3							5.9 (3.2)	2.8 (2.7)
SANS total ^f								
Time 1		9.8 (5.6)	6.7 (4.4)				8.4 (5.6)	5.9 (3.7)

	Baseline sample ^b			Longitudinal sample ^c		
	Healthy controls (n=98)	FE patients (n=72)	At-risk subjects (n=98)	Healthy controls (n=21)	FE patients (n=15)	At-risk subjects (n=35)
					F/χ^2	p value
Time 2					8.6 (5.6)	4.3 (4.1)
Time 3					6.1 (3.8)	4.1 (3.8)
BPRS total ^{*†}						
Time 1		17.4 (7.8)	15.8 (6.6)		16.7 (8.2)	15.6 (5.6)
Time 2					12.2 (4.4)	9.7 (5.4)
Time 3					12.7 (6.9)	9.8 (6.5)
SOPS total [†]						
Time 1			33.8 (13.9)			33.6 (14.4)
Time 2						15.3 (10.2)
Time 3						15.5 (11.9)
GAF score ^{*†}						
Time 1		42.6 (12.7)	53.7 (9.7)		43.3 (11.9)	54.1 (9.5)
Time 2					47.7 (7.9)	60.9 (10.1)
Time 3					51.7 (8.9)	59.6 (15.2)
Role functioning ^f						
Time 1		5.0 (1.9)	6.3 (1.7)		5.0 (1.2)	6.8 (1.6)
Social functioning ^g						
Time 1		5.9 (1.5)	6.4 (1.5)		6.1 (1.4)	6.5 (1.3)

FE, First-episode psychosis; APS, attenuated positive symptoms; BIPS, brief intermittent psychotic symptoms; GRD, genetic risk and deterioration; SAPS, Scale for the Assessment of Positive Symptoms; SANS, Scale for the Assessment of Negative Symptoms; BPRS, Brief Psychiatric Rating Scale; SOPS, Scale of Prodromal Symptoms; GAF, Global Assessment of Functioning.

^aValues are given as number and percentage of subjects for gender, handedness, prodromal criteria and antipsychotic type. Group means with the standard deviation for age, parental education, SAPS, SANS, BPRS, SOPS, GAF, Global Role, and Global Social scales are reported. Age and education were analyzed with one-way ANOVAs. Gender and handedness were analyzed with Pearson χ^2 tests.

^bDemographics for all participants included in the baseline analysis at Time 1: healthy controls (HC) $n=98$, FE $n=72$, at-risk (AR) subjects $n=98$.

^cDemographics exclusively for subjects participating across Time 1, Time 2 and Time 3: HC $n=21$, FE $n=15$, AR $n=35$.

^dThe Annett Handedness (1985) questionnaire was used to measure handedness.

^eProdromal criteria APS, BIPS and GRD are not mutually exclusive.

^fThe Global Functioning : Role (GF : Role) scale (Niendam *et al.* 2006a) was used to measure role functioning.

^gThe Global Functioning : Social (GF : Social) scale (Auther *et al.* 2006) was used to measure social functioning.

Baseline FE subjects included for each measure are as follows : Handedness $n=65$; SAPS, SANS, BPRS, GAF $n=68$; GF : Role/Social : T1 $n=32$.

Baseline AR subjects included for each measure are as follows : Handedness, SAPS, SANS, BPRS, SOPS, GAF $n=95$; GF : Role/Social : T1 $n=85$.

* Significant improvement within the FE group over time. * $p<0.05$.

• Significant improvement within the AR group over time. † $p<0.05$.

Table 2

ANCOVA results showing group differences in visual masking (VM) performance

	Partial χ^2	df	<i>F</i>	<i>p</i>
Location				
Group	0.01	2, 264	0.92	0.40
Age	0.00	1, 264	0.01	0.91
Direction	0.02	1, 264	4.67	0.03
Group $\times$ direction	0.06	2, 264	8.38	<0.001
Forward location				
Group	0.03	2, 264	3.44	0.03
Age	0.00	1, 264	0.11	0.74
ISI	0.10	5, 1320	29.39	<0.001
Group $\times$ ISI	0.01	10, 1320	1.41	0.19
Backward location				
Group	0.03	2, 264	3.74	0.03
Age	0.00	1, 264	0.03	0.88
ISI	0.02	5, 1320	6.60	<0.001
Group $\times$ ISI	0.01	10, 1320	1.72	0.10
Identification: high-energy mask				
Group	0.04	2, 264	5.14	0.01
Age	0.05	1, 264	14.32	<0.001
Direction	0.01	1, 264	1.66	0.20
Group $\times$ direction	0.02	2, 264	3.20	0.04
Forward: high identification				
Group	0.04	2, 264	5.46	0.01
Age	0.02	1, 264	5.33	0.02
ISI	0.02	6, 1584	5.80	<0.001
Group $\times$ ISI	0.01	12, 1584	1.41	0.16
Backward: high identification				
Group	0.03	2, 264	3.73	0.03
Age	0.07	1, 264	18.26	<0.001
ISI	0.03	6, 1584	8.44	<0.001
Group $\times$ ISI	0.02	12, 1584	2.55	0.00
Identification: low-energy mask				
Group	0.08	2, 264	10.85	<0.001
Age	0.04	1, 264	10.97	0.00
Direction	0.01	1, 264	1.51	0.22
Group $\times$ direction	0.05	2, 264	7.40	<0.001
Forward: low identification				
Group	0.08	2, 264	11.47	<0.001
Age	0.02	1, 264	4.71	0.03
ISI	0.01	6, 1584	1.98	0.07

	Partial χ^2	df	F	p
Group $\times$ ISI	0.01	12, 1584	1.39	0.17
Backward: low identification				
Group	0.06	2, 264	13.27	0.00
Age	0.05	1, 264	7.80	<0.001
ISI	0.01	6, 1584	2.82	0.02
Group $\times$ ISI	0.01	12, 1584	1.45	0.08

ISI, Interstimulus interval; df, degrees of freedom.