Published in final edited form as:

Commun Stat Theory Methods. 2014 July 29; 43(16): 3536–3551. doi:10.1080/03610926.2012.700379.

An Empirical Bayes Method for Multivariate Meta-analysis with an Application in Clinical Trials

Yong Chen, Sheng Luo, Haitao Chu, Xiao Su, and Lei Nie

Abstract

We propose an empirical Bayes method for evaluating overall and study-specific treatment effects in multivariate meta-analysis with binary outcome. Instead of modeling transformed proportions or risks via commonly used multivariate general or generalized linear models, we directly model the risks without any transformation. The exact posterior distribution of the study-specific relative risk is derived. The hyperparameters in the posterior distribution can be inferred through an empirical Bayes procedure. As our method does not rely on the choice of transformation, it provides a flexible alternative to the existing methods and in addition, the correlation parameter can be intuitively interpreted as the correlation coefficient between risks.

Keywords

Bivariate beta-binomial model; Exact method; Hypergeometric function; Meta-analysis; Relative risk; Sarmanov family

1 Introduction

Meta-analysis is a statistical procedure of combining the evidence from several independent studies concerned with the same scientific question. Recently, the rapid growth of evidence-based medicine has led to a dramatic increasing attention to statistical methods for meta-analysis¹. One of the most important settings considered in meta-analysis is the study of the association between a binary outcome and a binary exposure. Such examples include meta-analysis in clinical trials where subjects are randomized to treatment and placebo groups and the outcome of interest is a certain event. When both outcome and exposure are binary variables, the relative risk of an event comparing subjects who received treatment to those who received placebo provides an important measure of treatment effects².

Two general statistical approaches can be used to make inference on the relative risk. The first is the univariate analysis which estimates the overall relative risk by the weighted sum of the study-specific relative risks. Under homogeneous treatment effect assumption, the common relative risk can be estimated by a fixed effect model. When homogeneity is in doubts, a random effect model can be adopted³. The second approach, which draws lots of attention recently^{1,4,5,6,7,8}, is the multivariate meta-analysis. This approach accounts for the between-group correlation which is ignored by the univariate analysis. By borrowing

*Corresponding author: Yong Chen is Assistant Professor, Division of Biostatistics, The University of Texas Health Science Center at Houston, 1200 Herman Pressler, Houston, TX 77030, USA (yong.chen@uth.tmc.edu; Phone: 713-500-9569).

strength between multiple groups, the multivariate approach can be more efficient than the univariate analysis in estimating the overall treatment effect^{6,9,8}. For studies involving binary outcome and exposure, two modeling strategies have been commonly used in the second approach: a bivariate general linear mixed effect model on the transformed proportions⁴ and a bivariate generalized linear mixed effect model on the transformed risks⁵ (e.g., logit or probit transformations). Both models allow for between-group correlation and perform well in most of cases. It is worth pointing out that when the sample sizes for studies are moderate and the event is rare, the latter model can outperform the former⁸.

In this paper, we propose an alternative method in the framework of multivariate meta-analysis. Instead of modeling the transformed proportions or risks, we model the risks directly. One advantage of this proposed method is that the inference does not rely on the choice of transformation. As a consequence, the correlation parameter can be intuitively interpreted as the correlation coefficient between risks. Another benefit is that the study-specific relative risk has a posterior distribution with close form expression, which offers computational advantageous when posterior distributions of study-specific treatment effects are also of interest.

This paper is organized as follows. In section 2, we state the main results where we describe the proposed model and its inference for multiple 2×2 tables in meta-analysis and its extension for meta-regression analysis. In Section 3, we conduct simulation studies to evaluate the performance of the proposed method in finite samples. We illustrate our method in Section 4 with an example of meta-analysis of type 2 diabetes mellitus followed by a brief summary in Section 5.

2 Main Results

2.1 Notations and Models

For the i -th study, let n_{ji} , y_{ji} and p_{ji} be the number of subjects, the number of subjects experienced a certain event, and the risk of experiencing the event in the j -th group, respectively ($j = 1, \dots, J$). For simplicity, we consider the cases with two groups under comparison (i.e., $J = 2$) and the extension to cases with more than 2 groups is straightforward. We consider the following random effects model

$$\begin{aligned} (p_{1i}, p_{2i}) | (a_1, b_1, a_2, b_2, \rho) &\sim g(p_1, p_2; a_1, b_1, a_2, b_2, \rho), \\ (y_{1i}, y_{2i}) | (n_{1i}, n_{2i}, p_{1i}, p_{2i}) &\sim \text{Binomial}(y_{1i} | n_{1i}, p_{1i}) \times \text{Binomial}(y_{2i} | n_{2i}, p_{2i}), \end{aligned} \quad (1)$$

where $g(p_1, p_2; a_1, b_1, a_2, b_2, \rho)$ is the joint distribution of study-specific random effects p_{1i} and p_{2i} such that the marginal distribution of random effects in j th group p_{ji} is the beta distribution with shape parameters (a_j, b_j) and the correlation coefficient between p_{1i} and p_{2i} is ρ . Similar to a standard random effect model, the model (1) implies that for given study-specific random effects p_{1i} and p_{2i} , y_{1i} and y_{2i} follow independent binomial distributions.

Such a joint distribution of p_{1i} and p_{2i} can be constructed using the procedure proposed by Sarmanov¹⁰. This procedure has later been studied extensively by Cole et al.¹¹, Lee¹², Shubina and Lee¹³, and Danaher and Hardie¹⁴. Specifically, let $f_1(p_1)$ and $f_2(p_2)$ be two

univariate probability density functions, and $\psi_1(p_1)$ and $\psi_2(p_2)$ be bounded integrable non-constant functions that satisfy $\int \psi_j(p)f_j(p)dt = 0$ for $j = 1, 2$. The function defined by

$$h(p_1, p_2) = f_1(p_1)f_2(p_2)\{1 + \omega\psi_1(p_1)\psi_2(p_2)\}$$

is a bivariate joint density with specified marginals $f_1(p_1)$ and $f_2(p_2)$, provided ω is a number which satisfies the condition that $1 + \omega\psi_1(p_1)\psi_2(p_2) \geq 0$ for all p_1 and p_2 .

There are a variety of choices for the functions $\psi_1(p_1)$ and $\psi_2(p_2)$, which lead to different interpretations of the parameter ω . Here we let

$f_i(p_j) = \text{beta}(p_j; a_j, b_j) = \{B(a_j, b_j)\}^{-1} p_j^{a_j-1} (1-p_j)^{b_j-1}$ with $B(a, b)$ denotes the beta function, and $\psi_j(p_j) = (p_j - \mu_j)/\delta_j$ with $\mu_j = a_j/(a_j + b_j)$ and

$\delta_j^2 = \mu_j(1 - \mu_j)/(a_j + b_j + 1)$ for $j = 1, 2$. One advantage of this choice is that the parameter ω has an intuitive interpretation as the correlation coefficient between p_1 and p_2 , i.e., $\omega = \rho$. The joint distribution of p_1 and p_2 , referred to as Sarmanov bivariate beta distribution in Danaher and Hardie¹⁴, is

$$g(p_1, p_2; a_1, b_1, a_2, b_2, \rho) = \text{beta}(p_1; a_1, b_1) \text{beta}(p_2; a_2, b_2) \{1 + \rho\psi_1(p_1)\psi_2(p_2)\}. \quad (2)$$

With the distribution of random effects specified in equation (2), the log likelihood function for the unknown parameters $(a_1, b_1, a_2, b_2, \rho)$ is

$$\begin{aligned} \log L(a_1, b_1, a_2, b_2, \rho) &= \sum_{i=1}^n \log \iint \Pr(y_{1i}, y_{2i} | p_{1i}, p_{2i}) g(p_{1i}, p_{2i}; a_1, b_1, a_2, b_2, \rho) dp_{1i} dp_{2i} \\ &= \sum_{i=1}^n \log [P_{BB}(y_{1i}; n_{1i}, a_1, b_1) P_{BB}(y_{2i}; n_{2i}, a_2, b_2) \\ &\quad \left\{ 1 + \frac{\rho}{\sqrt{\frac{a_1 b_1}{(a_1 + b_1)^2 (a_1 + b_1 + 1)}} \sqrt{\frac{a_2 b_2}{(a_2 + b_2)^2 (a_2 + b_2 + 1)}} \left(\frac{y_{1i} + a_1}{n_{1i} + a_1 + b_1} - \frac{a_1}{a_1 + b_1} \right) \left(\frac{y_{2i} + a_2}{n_{2i} + a_2 + b_2} - \frac{a_2}{a_2 + b_2} \right) \right\} \right], \end{aligned} \quad (3)$$

where $P_{BB}(y; n, a, b)$ is the probability mass function of a beta-binomial distribution, i.e.

$$P_{BB}(y; n, a, b) = \binom{n}{y} \frac{B(y+a, n-y+b)}{B(a, b)}$$

The last expression in equation (3) has been derived by Danaher and Hardie¹⁴ and an outline of derivation is provided in the Appendix Section A. We refer the model (3) as Sarmanov beta-binomial model. Notice that when $\rho = 0$, the Sarmanov beta-binomial model reduces to the independent beta-binomial model, i.e., product of two beta-binomial distributions.

In meta-analysis, the primary quantity of interest is the overall treatment effect, which can be measured by the overall relative risk, defined by $\theta = \mu_2/\mu_1 = \{a_2/(a_2 + b_2)\}/\{a_1/(a_1 + b_1)\}$.

$b_1)\}$. In practice, biomedical researchers may be also interested in the statistical evidence on the treatment effects contributed by individual studies. Such quantity can be quantified by the posterior density function of the study-specific relative risk $\theta_i = p_{2i}/p_{1i}$ if we treat $g(p_1, p_2, a_1, b_1, a_2, b_2, \rho)$ as a prior distribution on (p_{1i}, p_{2i}) . We next discuss the inference procedures for the overall and the study-specific relative risks.

2.2 Estimation of Overall Relative Risk

At least two methods can be used to make inference on the overall relative risk θ . The first is to introduce another level of hierarchy by imposing prior distributions on the hyperparameters $(a_1, b_1, a_2, b_2, \rho)$. The posterior distribution of θ can be obtained through Monte Carlo simulations. Here we use the second method, which estimates the hyperparameters $(a_1, b_1, a_2, b_2, \rho)$ by maximizing the log likelihood $\log L(a_1, b_1, a_2, b_2, \rho)$. This can be done by using commonly used statistical software such as *SAS* and *SPLUS/R*. We implement it through *R* (*R Development Core Team, Version 2.14.1*) with the *optim* function, which uses a quasi-Newton method with box constraints on the ranges of parameters. A *SPLUS/R* program to fit this model with a working example is attached in Appendix Section B.

Denote $(\hat{a}_1, \hat{b}_1, \hat{a}_2, \hat{\rho})$ the maximum likelihood estimates based on the log likelihood function (3). The variance of the overall relative risk estimate $\hat{\theta}$ can be estimated by the delta method. Specifically, to avoid computational problems, we reparametrize the parameters (a_1, b_1, a_2, b_2) in their log scale and denote H the corresponding hessian matrix of the log likelihood function. The variance of the overall relative risk estimate $\hat{\theta}$ is estimated by $\hat{\theta}^2 D^T (-H)^{-1} D$ where $D = (-b_1/\hat{a}_1 + b_1, b_1/(\hat{a}_1 + b_1), b_2/(\hat{a}_2 + b_2), -b_2/(\hat{a}_2 + b_2), 0)^T$.

2.3 Inference on Study-specific Relative Risk

The statistical evidence from individual studies can be quantified by the posterior distributions of study-specific relative risks θ_i , namely $\Pr(\theta_i | \text{data}_i, a_1, b_1, a_2, b_2, \rho)$ where $\text{data}_i = (y_{1i}, n_{1i}, y_{2i}, n_{2i})$. Notice that the hyperparameters are unknown. One way to circumvent this problem is to simply replace the hyperparameters by their estimates. Such an approach is called the empirical Bayes method^{15,16,17,18}. The inference based on $\Pr(\theta_i | \text{data}_i, \hat{a}_1, \hat{b}_1, \hat{a}_2, \hat{b}_2, \hat{\rho})$ ignores the uncertainty on the hyperparameter estimates, hence may lead to liberal credible intervals. To obtain credible intervals with coverage probability close to the nominal level, one can use the bias correction method or the bootstrap method^{19,20}. In Section 3, we will investigate the coverage probability of the credible intervals using the empirical Bayes method ignoring the uncertainty on the hyperparameter estimates.

We next derive the posterior distribution of the study-specific relative risk. Note that an important property of Sarmanov beta prior is that it is pseudo-conjugate for binomial distributions¹². Specifically, Sarmanov beta prior can be expressed as a linear combination of products of independent beta distributions. Therefore, the posterior distribution of the study-specific relative risk can be written as a linear combination of posterior distributions under independent beta priors. In fact, the posterior distribution of relative risk under independent beta priors for a single 2×2 table has been considered by several researchers. Nurminen and Mutanen²¹ and Gupta et al.²² derived the posterior distribution under

independent beta priors with integer hyperparameters. Hora and Kelley²³ and Hashemi et al.²⁴ extended the previous results to beta priors with any positive hyperparameters. Now we restate their results using expression with hypergeometric functions.

Assume p_{1i} and p_{2i} are independent and follow beta distributions with hyperparameters (a_1, b_1) and (a_2, b_2) , respectively. The posterior distributions of p_{1i} and p_{2i} are beta distributions with parameters (α_1, β_1) and (α_2, β_2) respectively, where $\alpha_j = y_j + a_j$ and $\beta_j = n_j - y_j + b_j$ ($j = 1, 2$). The exact posterior distributions of θ_i can be expressed as

$$f(\theta_i; \alpha_1, \beta_1, \alpha_2, \beta_2) = \begin{cases} \theta_i^{\alpha_2-1} \{B(\alpha_1, \beta_1)B(\alpha_2, \beta_2)\}^{-1} B(\alpha_1 + \alpha_2, \beta_1) \\ \times F(1 - \beta_2, \alpha_1 + \alpha_2; \alpha_1 + \alpha_2 + \beta_1; \theta_i) & \text{for } \theta_i \in [0, 1), \\ \theta_i^{-\alpha_1-1} \{B(\alpha_1, \beta_1)B(\alpha_2, \beta_2)\}^{-1} B(\alpha_1 + \alpha_2, \beta_2) \\ \times F(1 - \beta_1, \alpha_1 + \alpha_2; \alpha_1 + \alpha_2 + \beta_2; 1/\theta_i) & \text{for } \theta_i \in [1, \infty), \end{cases} \quad (4)$$

where $F(\cdot, \cdot, \cdot, \cdot)$ denotes the hypergeometric function²⁵ defined by

$$F(\alpha, \beta; \gamma, z) = \frac{1}{B(\beta, \gamma - \beta)} \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-tz)^{-\alpha} dt, \quad \text{for } \gamma > \beta > 0.$$

With the results in (4), the exact posterior distribution of relative risk θ_i under Sarmanov beta prior can be calculated as

$$\begin{aligned} f^*(\theta_i; \alpha_1, \beta_1, \alpha_2, \beta_2, \rho) \\ = \omega_1 f(\theta_i; \alpha_1, \beta_1, \alpha_2, \beta_2) \\ + \omega_2 f(\theta_i; \alpha_1 + 1, \beta_1, \alpha_2, \beta_2) \\ + \omega_3 f(\theta_i; \alpha_1, \beta_1, \alpha_2 \\ + 1, \beta_2) \\ + \omega_4 f(\theta_i; \alpha_1 \\ + 1, \beta_1, \alpha_2 + 1, \beta_2), \end{aligned} \quad (5)$$

where $f(\theta_i, \alpha_1, \beta_1, \alpha_2, \beta_2)$ is the posterior density function of relative risk under independent beta priors and is defined in equation (4), the weights ω_k ($k = 1, \dots, 4$) are functions of a_1, b_1, a_2, b_2, ρ given in Appendix Section A. When the correlation is zero, i.e., $\rho = 0$, in which the weights are $\omega_1 = 1$ and $\omega_2 = \omega_3 = \omega_4 = 0$, equation (5) reduces to equation (4).

2.4 Extension to Meta-regression Analysis

When a meta-analysis contains a relatively large number of studies and the study level covariates are available, multivariate meta-regression analysis can assess the impact of study size, trial quality and other trial characteristics on effect sizes. To account for the heterogeneity between trials and the correlation between treatment groups, models have been proposed in the framework of multivariate generalized linear mixed effect models, such as Van Houwelingen *et al.* (1993)²⁶, Smith *et al.* (1995)²⁷, Skrandal and Rabe-Hesketh (2004)²⁸ and Chu and Cole (2006)⁵. Now we extend the model (1) to regression setting. Specifically, we assume that the study-specific risk p_{ji} for $j = 1, 2$ have beta distributions with mean parameters μ_{ji} and dispersion parameters ϕ_j , respectively,

$$p_{ji} | (\phi_j, \mu_{ji}) \sim \text{beta}\{p_{ji}; \mu_{ji}/(1/\phi_j - 1), (1 - \mu_{ji})/(1/\phi_j - 1) \text{ for } j=1, 2.$$

Then $E[p_{ji} | \phi_j, \mu_{ji}] = \mu_{ji}$ and $\text{var}(p_{ji} | \phi_j, \mu_{ji}) = \delta_{ji}^2 = \phi_j \mu_{ji} (1 - \mu_{ji})$. The mean of each beta distribution is a function of covariates

$$\mu_{ji} = h^{-1}(X_i \eta_j) \text{ for } j=1, 2,$$

where $h(\cdot)$ is a link function and X_i are the study-specific covariates such as the trial quality and race of the study population. To allow for correlation between groups, we assume the paired study-specific risks (p_{1i}, p_{2i}) follow the Sarmanov beta prior distribution, i.e.,

$$(p_{1i}, p_{2i}) | (\phi_1, \mu_{1i}, \phi_2, \mu_{2i}) \sim \text{beta}\{p_{1i}; \mu_{1i}/(1/\phi_1 - 1), (1 - \mu_{1i})/(1/\phi_1 - 1)\} \\ \times \text{beta}\{p_{2i}; \mu_{2i}/(1/\phi_2 - 1), (1 - \mu_{2i})/(1/\phi_2 - 1)\} \left\{ 1 + \rho \frac{(p_{1i} - \mu_{1i})}{\delta_{1i}} \frac{(p_{2i} - \mu_{2i})}{\delta_{2i}} \right\}.$$

This model allows different dispersion parameter ϕ_j across different groups. Similar to the estimation procedure for model (1), this bivariate beta-binomial regression model can be fitted by maximizing the corresponding log likelihood function.

3 Simulation Studies

In this section, we evaluate the performance of the model (1) through simulation studies. In the first simulation study, we examine the results in equation (4) and equation (5) empirically via Markov chain Monte Carlo (MCMC) methods. Specifically, the histograms of MCMC samples are overlaid with the corresponding theoretical density functions. The empirical results confirms that the derived density functions are correct. For the interest of space, the details of this simulation are summarized in the Web Supplement.

In the second simulation study, we evaluate the finite sample performance of the maximum likelihood estimator for the overall relative risk. We set the true values as $a_1 = b_1 = a_2 = b_2 = 0.5$, $\rho = 0, 0.2$ and 0.4 , and the number of studies $n = 20, 40$ and 60 with the study sample size 50 in each group. Table 1 compares the bias, SE (the standard deviation of the overall log relative risk estimates), SEM (the sampling mean of the standard error estimates), and CP (the coverage probability of Wald-type confidence intervals of the overall log relative risk) estimated from the Sarmanov and independent beta-binomial model. When $\rho = 0$, the independent model gives unbiased estimates, SEM close to SE, and coverage probabilities close to nominal level while the Sarmanov model performed equally well in terms of bias, SE, SEM, only with coverage probabilities negligibly worse. When ρ increases to 0.2 or 0.4 , both models provide unbiased estimates, the SEs of both models decrease due to “borrowing strength” across groups within the same study⁹. The Sarmanov model provides SEM close to SE and coverage probabilities close to 0.95 , while the independent model produces unchanged SEMs and over-conservative confidence intervals because it does not account for the between-group correlation. In addition, by “borrowing strength” between multiple groups, the required number of studies to achieve certain efficiency can be significantly

reduced, e.g., the SE of log relative risk of the Sarmanov model when $n = 40$ and $\rho = 0.4$ is even smaller than the SE from the independent model when $n = 60$ and $\rho = 0$. These simulation results indicate that Sarmanov model provides valid and efficient inference and is robust to the between-group correlation with moderate number of studies, while the independent model only give valid inference when the between-group correlation is zero.

In Section 2.3, we propose to estimate the study-specific relative risk based on an empirical Bayes method with hyperparameters fixed at their estimates. The corresponding credible interval could be liberal due to the ignored uncertainty on the hyperparameters. In the third simulation study, we investigate the performance of the credible interval based on $\Pr(\theta_i | \text{data}_i, \hat{a}_1, \hat{b}_1, \hat{a}_2, \hat{b}_2, \hat{\rho})$ and compare with the conventional Wald confidence interval which is based on data of the i th study only. The selected parameters are the same as the second simulation study. For each simulated n studies, we obtain the estimates of the hyperparameters by maximizing the log likelihood in equation (3) and then estimate the 95% credible interval from the exact posterior distribution of the first study's relative risk using equation (5) (refer to as "Sarmanov CI"). In contrast, we obtain the 95% Wald confidence interval of the first study's relative risk based only on the first study's data (refer to as "Wald CI"). Table 2 compares the coverage probability (CP) and the expected length of Sarmanov CI with those of Wald CI. The simulation results suggest that under all settings considered, the coverage probability of Sarmanov CI is close to the nominal level, and so is Wald CI. This indicates that when the number of studies is moderate, Sarmanov CI still performs reasonably well, even without accounting for the uncertainty on the hyperparameters.

4 An application to a meta-analysis of type 2 diabetes mellitus after gestational diabetes

Gestational diabetes mellitus (GDM), affecting around 18% of pregnancies, is a form of diabetes that is diagnosed during pregnancy and increases risk of type 2 diabetes mellitus (T2DM)²⁹. Recently, Bellamy et al.³⁰ conducted a comprehensive systematic review and meta-analysis to assess the strength of association between GDM and T2DM. Twenty cohort studies conducted from 1991 to 2008 in multiple countries are selected. The data (summarized in Table 1 of the Web Supplement) has 6,862 and 3,997 T2DM incidences among 643,588 and 31,867 individuals without and with GDM, respectively. We define relative risk as the ratio of risks of T2DM comparing those with GDM to those without. Based on the Sarmanov beta-binomial model, the overall relative risk is estimated as 9.20 (95% CI: 5.17, 16.35), indicating significantly increased risk of developing T2DM for the women with GDM. We note that the study by Feig³¹ has extremely large sample size ($N = 659, 164$) and account for 97.6% of total sample size. To assess the sensitivity of the results to this study, we conduct a sensitivity analysis without this study and the overall relative risk is estimated as 9.11 (95% CI: 4.96, 16.73). These results suggest that our results are not sensitive to this study.

By applying the Sarmanov beta prior distribution with the aforementioned hyperparameters estimates, we obtain each individual study's posterior density distribution. We apply bisection root-finding method to compute the 95% credible intervals. Figure 1 displays the forest plot of the estimates and the 95% CIs of each study-specific relative risk and the

overall relative risk. To visually display how each study contributes to the overall relative risk, Figure 2 plots the posterior density distributions of four randomly selected studies. It shows that while in Gunderson et al³², most of the density of relative risk are between 2 and 5 (mean: 3.91, 95% CI: 2.84, 5.17), the density shifts away from 1 in Lee et al³³ with the majority of the density lying between 2 and 7 (mean: 4.81, 95% CI: 3.02, 7.52). In the studies of Benjamin et al³⁴ and O'Sullivan et al³⁵, most of the relative risk density lies in the intervals of [2, 10] (mean: 8.81, 95% CI: 2.70, 24.92) and [5, 10] (mean: 7.53, 95% CI: 4.74, 11.77), respectively. Note that the curve of Benjamin et al³⁴ is much more spread out than the other studies because of its relatively small study sample size.

5 Summary

In this paper, we propose an alternative model for multivariate meta-analysis of binary outcomes. In contrast to the commonly used methods which either model the transformed proportions through a general linear model, or model the transformed risks through a generalized linear model, this alternative method directly models the risks without any transformation. One advantage of this method is that the inference does not rely on the choice of transformation. The correlation parameter can be intuitively interpreted as the correlation coefficient between risks.

Our contributions in this paper are to propose the use of the Sarmanov beta-binomial model in meta-analysis of clinical trials and to derive the exact posterior distribution function of study-specific treatment effects which enables a simple empirical Bayes inference. With simulation studies, we investigate the efficiency gained by accounting for the between-group correlation in the Sarmanov beta-binomial models and the frequentist performance of the credible intervals based on the empirical Bayes method.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Yong Chen's research was partially supported by a start-up fund from the University of Texas School of Public Health. Sheng Luo's research was partially supported by two NIH/NINDS grants: U01 NS043127 and U01 NS43128. Haitao Chu was supported in part by the U.S. Department of Health and Human Services Agency for Healthcare Research and Quality Grant R03HS020666 and P01CA142538 from the U.S. National Cancer Institute. Views expressed in this paper are the authors' professional opinions and do not necessarily represent the official positions of the U.S. Food and Drug Administration. We gratefully acknowledge the editor, Dr. Narayanaswamy Balakrishnan, and an anonymous reviewer for constructive comments that greatly improved the manuscript.

References

1. Jackson D, Riley R, White IR. Multivariate meta-analysis: Potential and promise. *Statistics in Medicine*. 2011
2. Piantadosi, S. *Clinical trials: a methodologic perspective*. Vol. 593. Wiley-Blackwell; 2005.
3. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled clinical trials*. 1986; 7(3):177–188. [PubMed: 3802833]

4. Reitsma J, Glas A, Rutjes A, Scholten R, Bossuyt P, Zwinderman A. Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *Journal of clinical epidemiology*. 2005; 58(10):982–990. [PubMed: 16168343]
5. Chu H, Cole SR. Bivariate meta-analysis of sensitivity and specificity with sparse data: a generalized linear mixed model approach. *Journal of clinical epidemiology*. 2006; 59(12):1331. [PubMed: 17098577]
6. Riley RD, Abrams KR, Sutton AJ, Lambert PC, Thompson JR. Bivariate random-effects meta-analysis and the estimation of between-study correlation. *BMC Medical Research Methodology*. 2007; 7(1):3. [PubMed: 17222330]
7. Riley RD, Thompson JR, Abrams KR. An alternative model for bivariate random-effects meta-analysis when the within-study correlations are unknown. *Biostatistics*. 2008; 9(1):172. [PubMed: 17626226]
8. Hamza TH, Reitsma JB, Stijnen T. Meta-analysis of diagnostic studies: A comparison of random intercept, normal-normal, and binomial-normal bivariate summary ROC approaches. *Medical Decision Making*. 2008; 28(5):639–649. [PubMed: 18753684]
9. Riley R, Abrams K, Lambert P, Sutton A, Thompson J. An evaluation of bivariate random-effects meta-analysis for the joint synthesis of two correlated outcomes. *Statistics in medicine*. 2007; 26(1): 78–97. [PubMed: 16526010]
10. Sarmanov O. Generalized normal correlation and two-dimensional Fréchet classes. *Soviet Mathematics Doklady*. 1966; 7:596–599.
11. Cole BF, Lee MLT, Whitmore GA, Zaslavsky AM. An empirical Bayes model for Markov-dependent binary sequences with randomly missing observations. *Journal of the American Statistical Association*. 1995; 90(432):1364–1372.
12. Lee MLT. Properties and applications of the Sarmanov family of bivariate distributions. *Communications in Statistics-Theory and Methods*. 1996; 25(6):1207–1222.
13. Shubina M, Lee MLT. On maximum attainable correlation and other measures of dependence for the Sarmanov family of bivariate distributions. *Communications in Statistics-Theory and Methods*. 2004; 33(5):1031–1052.
14. Danaher PJ, Hardie BGS. Bacon with your eggs? Applications of a new bivariate beta-binomial distribution. *The American Statistician*. 2005; 59(4):282–286.
15. Efron B, Morris C. Stein's estimation rule and its competitors-an empirical Bayes approach. *Journal of the American Statistical Association*. 1973; 68(341):117–130.
16. Efron B, Morris C. Data analysis using Stein's estimator and its generalizations. *Journal of the American Statistical Association*. 1975; 70(350):311–319.
17. Gelman, A.; Carlin, J.; Stern, H.; Rubin, D. *Bayesian data analysis*. CRC press; 2004.
18. Carlin, BP.; Louis, TA. *Bayesian methods for data analysis*. Chapman & Hall/CRC; 2009.
19. Deely J, Lindley D. Bayes empirical Bayes. *Journal of the American Statistical Association*. 1981; 76(376):833–841.
20. Carlin BP, Gelfand AE. A sample reuse method for accurate parametric empirical Bayes confidence intervals. *Journal of the Royal Statistical Society Series B (Methodological)*. 1991; 53(1):189–200.
21. Nurminen M, Mutanen P. Exact Bayesian analysis of two proportions. *Scandinavian Journal of Statistics*. 1987; 14(1):67–77.
22. Gupta R, Albanese R, Penn J, White T. Bayesian estimation of relative risk in biomedical research. *Environmetrics*. 1997; 8(2):133–143.
23. Hora SC, Kelley GD. Bayesian inference on the odds and risk ratios. *Communications in Statistics-Theory and Methods*. 1983; 12(6):681–692.
24. Hashemi L, Nandram B, Goldberg R. Bayesian analysis for a single 2x2 table. *Statistics in medicine*. 1997; 16(1):1311–1328. [PubMed: 9232755]
25. Gauss CF. *Disquisitiones generales circa seriem infinitam*. *Comm Gott*. 1813; 2:123–161.
26. Van Houwelingen HC, Zwinderman KH, Stijnen T. A bivariate approach to meta-analysis. *Statistics in Medicine*. 1993; 12(24):2273–2284. [PubMed: 7907813]

27. Smith TC, Spiegelhalter DJ, Thomas A. Bayesian approaches to random-effects meta-analysis: A comparative study. *Statistics in Medicine*. 1995; 14(24):2685–2699. [PubMed: 8619108]
28. Skrondal, A.; Rabe-Hesketh, S. Generalized latent variable modeling: multilevel, longitudinal, and structural equation models. CRC Press; 2004.
29. Coustan DR, Lowe LP, Metzger BE, Dyer AR. The Hyperglycemia and Adverse Pregnancy Outcome (HAPO) study: paving the way for new diagnostic criteria for gestational diabetes mellitus. *American journal of obstetrics and gynecology*. 2010; 202(6):654-e1. [PubMed: 20510967]
30. Bellamy L, Casas JP, Hingorani AD, Williams D. Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis. *The Lancet*. 2009; 373(9677):1773–1779.
31. Feig DS, Zinman B, Wang X, Hux JE. Risk of development of diabetes mellitus after diagnosis of gestational diabetes. *Canadian Medical Association Journal*. 2008; 179(3):229. [PubMed: 18663202]
32. Gunderson EP, Lewis CE, Tsai AL, Chiang V, Carnethon M, Quesenberry CP, et al. A 20-Year Prospective Study of Childbearing and Incidence of Diabetes in Young Women, Controlling for Glycemia Before Conception. *Diabetes*. 2007; 56(12):2990. [PubMed: 17898128]
33. Lee H, Jang HC, Park HK, Metzger BE, Cho NH. Prevalence of type 2 diabetes among women with a previous history of gestational diabetes mellitus. *Diabetes research and clinical practice*. 2008; 81(1):124–129. [PubMed: 18456364]
34. Benjamin E, Winters D, Mayfield J, Gohdes D. Diabetes in pregnancy in Zuni Indian women. Prevalence and subsequent development of clinical diabetes after gestational diabetes. *Diabetes Care*. 1993; 16(9):1231. [PubMed: 8404425]
35. O'Sullivan, JB. Carbohydrate metabolism in pregnancy and the newborn London. Springer-Verlag; 1989. The Boston gestational diabetes studies: review and perspectives; p. 287-294.

Appendix

Section A: Some derivations

Derivation of equation (3)

For simplicity of notation, we suppress the index “i”. With some algebra, we can show

$$\int \text{Binomial}(y|n;p)\text{beta}(p;a,b)\{1+c(p-\mu)\}dp = P_{BB}(y|n;a,b)\left\{1+C\left(\frac{y+a}{n+a+b}-\mu\right)\right\}.$$

Denote $\mu_j = a_j/(a_j+b_j)$ and $\sigma_j^2 = \mu_j(1-\mu_j)/(a_j+b_j+1)$ ($j=1,2$). We have

$$\begin{aligned} & \int \int \text{Pr}(y_1|n_1;p_1)\text{Pr}(y_2|n_2;p_2)\text{beta}(p_1;a_1,b_1)\text{beta}(p_2;a_2,b_2)\left\{1+\frac{\rho}{\sigma_1\sigma_2}(p_1-\mu_1)(p_2-\mu_2)\right\}dp_1dp_2 \\ &= \int \text{Pr}(y_2|n_2;p_2)\text{beta}(p_2;a_2,b_2) \int \text{Pr}(y_1|n_1;p_1)\text{beta}(p_1;a_1,b_1)\left\{1+\frac{\rho}{\sigma_1\sigma_2}(p_1-\mu_1)(p_2-\mu_2)\right\}dp_1dp_2 \\ &= \int \text{Pr}(y_2|n_2;p_2)\text{beta}(p_2;a_2,b_2)P_{BB}(y_1|n_1;a_1,b_1)\left\{1+\frac{\rho}{\sigma_1\sigma_2}\left(\frac{y_1+a_1}{n_1+a_1+b_1}-\mu_1\right)(p_2-\mu_2)\right\}dp_2 \end{aligned}$$

$$= \left[P_{BB}(y_{1i}; n_{1i}, a_1, b_1) P_{BB}(y_{2i}; n_{2i}, a_2, b_2) \left\{ 1 + \frac{\rho}{\sigma_1 \sigma_2} \left(\frac{y_{1i} + a_1}{n_{1i} + a_1 + b_1} - \mu_1 \right) \left(\frac{y_{2i} + a_2}{n_{2i} + a_2 + b_2} - \mu_2 \right) \right\} \right].$$

Derivation of equation (5)

With beta marginals and mixing functions $\phi_i = (p_i - \mu_i)/\delta_i$, the Sarmanov prior distribution of p_1 and p_2 can be written as linear combination of products of independent beta distributions as follows,

$$\begin{aligned} g(p_1, p_2; a_1, b_1, a_2, b_2, \rho) \\ = v_1 \text{beta}(p_1; a_1, b_1) \text{beta}(p_2; a_2, b_2) + v_2 \text{beta}(p_1; a_1 + 1, b_1) \text{beta}(p_2; a_2, b_2) \\ + v_3 \text{beta}(p_1; a_1, b_1) \text{beta}(p_2; a_2 + 1, b_2) + v_4 \text{beta}(p_1; a_1 + 1, b_1) \text{beta}(p_2; a_2 + 1, b_2), \end{aligned}$$

where $\text{beta}(\cdot; a_j, b_j)$ is the beta distribution, v_k ($k = 1, \dots, 4$) are weights, defined by, $v_1 = 1 + \rho d$, $v_2 = v_3 = -\rho d$, $v_4 = \rho d$, $d = (\mu_1 \mu_2)/(\delta_1 \delta_2)$. After some algebra, the posterior distribution of p_1 and p_2 given data is also a linear combination of products of independent beta distributions,

$$\begin{aligned} \Pr(p_1, p_2 | y_1, y_2, a_1, b_1, a_2, b_2, \rho) \\ = \omega_1 \text{beta}(p_1; \alpha_1, \beta_1) \text{beta}(p_2; \alpha_2, \beta_2) + \omega_2 \text{beta}(p_1; \alpha_1 + 1, \beta_1) \text{beta}(p_2; \alpha_2, \beta_2) \\ + \omega_3 \text{beta}(p_1; \alpha_1, \beta_1) \text{beta}(p_2; \alpha_2 + 1, \beta_2) + \omega_4 \text{beta}(p_1; \alpha_1 + 1, \beta_1) \text{beta}(p_2; \alpha_2 + 1, \beta_2), \end{aligned}$$

where the weights ω_k ($k = 1, \dots, 4$) are defined by,

$$\omega_1 = \frac{v_1 B(\alpha_1, \beta_1) B(\alpha_2, \beta_2)}{CB(a_1, b_1) B(a_2, b_2)}, \quad \omega_2 = \frac{v_2 B(\alpha_1 + 1, \beta_1) B(\alpha_2, \beta_2)}{CB(a_1 + 1, b_1) B(a_2, b_2)},$$

$$\omega_3 = \frac{v_3 B(\alpha_1, \beta_1) B(\alpha_2 + 1, \beta_2)}{CB(a_1, b_1) B(a_2 + 1, b_2)}, \quad \text{and} \quad \omega_4 = \frac{v_4 B(\alpha_1 + 1, \beta_1) B(\alpha_2 + 1, \beta_2)}{CB(a_1 + 1, b_1) B(a_2 + 1, b_2)},$$

and C , the normalizing constant, is calculated as

$$C = \frac{v_1 B(\alpha_1, \beta_1) B(\alpha_2, \beta_2)}{B(a_1, b_1) B(a_2, b_2)} + \frac{v_2 B(\alpha_1 + 1, \beta_1) B(\alpha_2, \beta_2)}{B(a_1 + 1, b_1) B(a_2, b_2)} + \frac{v_3 B(\alpha_1, \beta_1) B(\alpha_2 + 1, \beta_2)}{B(a_1, b_1) B(a_2 + 1, b_2)} + \frac{v_4 B(\alpha_1 + 1, \beta_1) B(\alpha_2 + 1, \beta_2)}{B(a_1 + 1, b_1) B(a_2 + 1, b_2)}.$$

The proof is completed following the derivation of equation (4).

Some results on the moments of relative risk

Under independent beta priors, the k -th posterior moment of relative risk exists for $k < \min(\alpha_1, \beta_2)$ and is given by

$$E[\theta^k; \alpha_1, \beta_1, \alpha_2, \beta_2] = \frac{B(\alpha_1 - k, \beta_1 + k) B(\alpha_2 + k, \beta_2 - k)}{B(\alpha_1, \beta_1) B(\alpha_2, \beta_2)}. \quad (6)$$

Specifically, the mean and variance are given by $E[\theta; \alpha_1, \beta_1, \alpha_2, \beta_2] = \beta_1 \alpha_2 / \{(\alpha_1 - 1)(\beta_2 - 1)\}$ and

$$\text{var}(\theta; \alpha_1, \beta_1, \alpha_2, \beta_2) = \frac{\beta_1(\beta_1 + 1)\alpha_2(\alpha_2 + 1)}{(\alpha_1 - 1)(\alpha_1 - 2)(\beta_2 - 1)(\beta_2 - 2)} - \left\{ \frac{\beta_1 \alpha_2}{(\alpha_1 - 1)(\beta_2 - 1)} \right\}^2.$$

Under correlated beta priors, the k -th posterior moment of relative risk exists for $k < \min(\alpha_1, \beta_2)$ and is given by

$$\begin{aligned} E[\theta^k; \alpha_1, \beta_1, \alpha_2, \beta_2, \rho] &= \omega_1 E[\theta^k; \alpha_1, \beta_1, \alpha_2, \beta_2] \\ &+ \omega_2 E[\theta^k; \alpha_1 + 1, \beta_1, \alpha_2, \beta_2] \\ &+ \omega_3 E[\theta^k; \alpha_1, \beta_1, \alpha_2 + 1, \beta_2] \\ &+ \omega_4 E[\theta^k; \alpha_1 + 1, \beta_1, \alpha_2 + 1, \beta_2], \end{aligned} \quad (7)$$

where $E[\theta^k; \alpha_1, \beta_1, \alpha_2, \beta_2]$ is the k -th posterior moment of relative risk under independent beta priors, defined in equation (6).

Some results on correlation coefficients

Denote $y_{ji} = \sum_{k=1}^{n_{ji}} z_{jim}$ where Z_{jim} is the indicator of whether the k th subject in the j th group of i th study experienced an event. Denote the dispersion parameter $\phi_j = 1/(a_j + b_j + 1)$. Note that Y_{ji} follows a beta-binomial distribution with parameters (a_j, b_j) . Therefore, $\text{corr}(Z_{jik}, Z_{jik'}) = \phi_j$, i.e., the correlation between the outcomes for two subjects from the same study and the same j th group is ϕ_j . Using double expectation technique, we can show that $(Z_{1ik}, Z_{2ik'}) = \rho \sqrt{\phi_1 \phi_2}$, i.e., the correlation between the outcomes for two subjects from the same study but different groups is $\rho \sqrt{\phi_1 \phi_2}$.

Section B: SPLUS/R program to fit model (1) and a working example

```
# function to compute the log-likelihood in equation (5)
myLik <- function(mypar, mydat) {
  par <- par.cal(mypar); a1 <- par[1]; b1 <- par[2]; a2 <- par[3]; b2 <- par[4]
  temp1 <- (lgamma(a1+mydat$y1) + lgamma(b1+mydat$n1-mydat$y1) + lgamma(a2+mydat$y2) + lgamma(b2+mydat$n2-mydat$y2)
  + lgamma(a1+b1) + lgamma(a2+b2))
  temp2 <- (lgamma(a1) + lgamma(b1) + lgamma(a2) + lgamma(b2)) +
```

```

lgamma(a1+b1+mydat$n1) + lgamma(a2+b2+mydat$n2))
if (flag == 0) myLogLik <- sum(temp1 - temp2) # if independent beta-
binomial model
if (flag == 1) { # if Sarmanov beta-binomial model
rho <- par[5]
mul <- a1/(a1+b1); mu2 <- a2/(a2+b2)
delta1 <- sqrt(mul*(1-mul)/(a1+b1+1)); delta2 <- sqrt(mu2*(1-mu2)/(a2+b2+1))
temp3 <- (log(1+rho/delta1/delta2*(mydat$y1-mydat$n1*mul)*(mydat$y2-mydat
$n2*mu2)/(a1+b1+mydat$n1)/(a2+b2+mydat$n2)))
myLogLik <- sum(temp1 - temp2 + temp3)}
return(myLogLik)
}
# Back-transform the parameters (a1,b1,a2,b2,rho) to original scale
par.cal <- function(mypar) {
a1 <- exp(mypar[1]); b1 <- exp(mypar[2]); a2 <- exp(mypar[3]); b2 <-
exp(mypar[4])
if (flag == 0) return(c(a1,b1,a2,b2))
if (flag == 1) {
eta <- mypar[5]; cc <- sqrt(a1*a2*b1*b2)/sqrt((a1+b1+1)*(a2+b2+1))
upper.bound <- cc/max(a1*b2, a2*b1); lower.bound <- -cc/max(a1*a2, b1*b2)
rho <- (upper.bound-lower.bound)*exp(eta)/(1+exp(eta)) + lower.bound
return(c(a1,b1,a2,b2,rho))}
}
# note: we use Delta method to get the variance of log(RR) and use Wald
interval on log(RR)
RR.comp.log <- function(par, hessian) {
a1 <- par[1]; b1 <- par[2]; a2 <- par[3]; b2 <- par[4]
myRR.overall <- log(a2/(a2+b2)/(a1/(a1+b1)))
myVar <- solve(-hessian)
if (flag == 0) myD <- matrix(c(-b1/(a1+b1), b1/(a1+b1), b2/(a2+b2), -b2/
(a2+b2)), nrow=1)
if (flag == 1) myD <- matrix(c(-b1/(a1+b1), b1/(a1+b1), b2/(a2+b2), -b2/
(a2+b2), 0), nrow=1)
myRR.overall.Var <- as.numeric(myD %*% myVar %*% t(myD))
myRR.overall.sd <- sqrt(myRR.overall.Var)
myRR.left.bound <- myRR.overall-1.96*sqrt(myRR.overall.Var)
myRR.right.bound <- myRR.overall+1.96*sqrt(myRR.overall.Var)
return(list(RR=exp(myRR.overall), RR.left=exp(myRR.left.bound),
RR.right=exp(myRR.right.bound)))
}
# Dataset from Bellamy (2009) Lancet
y1<-c(6628,22,0,150,1,16,7,8,0,0,0,1,0,1,7,0,0,3,18,0)
n1<-
c(637341,868,39,2242,111,783,108,489,11,435,70,61,52,39,431,35,57,47,328,41)

```

```

y2<-c(2874,71,21,43,53,405,6,13,7,23,44,21,10,15,105,10,33,14,224,5)
n2<-
c(21823,620,68,166,295,5470,70,35,23,435,696,229,28,45,801,15,241,47,615,145)
# remove the first study with extremely large sample size
#y1 <- y1[-1]; n1 <- n1[-1]; y2 <- y2[-1]; n2 <- n2[-1]
init.val <- rep(0, 5)
# maximization of the likelihood of independent beta-binomial model
flag <- 0 # flag = 0: independent beta-binomial model
results.indep <- optim(init.val[1:4], myLik, method = "L-BFGS-B",
lower=rep(-20,4), upper=rep(20,4), control = list(fnscale=-1,maxit=1000),
hessian = T, mydat=list(y1=y1,n1=n1,y2=y2,n2=n2))
RR.comp.log(par.cal(results.indep$par), results.indep$hessian)
# maximization of the likelihood of Sarmanov beta-binomial model
flag <- 1 # flag = 1: Sarmanov beta-binomial model
results <- optim(init.val, myLik, method = "L-BFGS-B",
lower=rep(-20,5), upper=rep(20,5), control = list(fnscale=-1,maxit=1000),
hessian=T, mydat=list(y1=y1,n1=n1,y2=y2,n2=n2))
RR.comp.log(par.cal(results$par), results$hessian)

```

Figure 1.

Forest plot of 20 study-specific and the overall relative risk with 95% credible intervals (numbers listed on the right). Relative risk is defined as the ratio of risks of T2DM comparing those with GDM to those without. The numbers on the y-axis indicate the study years. X-axis is log scale and is truncated at 300.

Figure 2.

Posterior distributions of study-specific relative risks for four studies: Lee et al.³³, Gunderson et al.³², Benjamin et al.³⁴, and O'Sullivan et al.³⁵. Relative risk is defined as the ratio of risks of T2DM comparing those with GDM to those without. The numbers in the legend are the study years.

Table 1

Estimates of the bias, SE (the standard deviation of the estimates), SEM (the sampling mean of the standard error estimate), CP (coverage probability), and RE (relative efficiency) of log relative risk in 5,000 simulations based on Sarmanov beta-binomial model and independent beta-binomial model, with various numbers of studies n and various between-group correlations ρ . The study sample sizes are 50 in each group. $(a_1, b_1, a_2, b_2) = (0.5, 0.5, 0.5, 0.5)$.

n	ρ	Sarmanov model				Independent model			
		Bias	SE	SEM	CP (%)	Bias	SE	SEM	CP (%)
20	0	-0.007	0.229	0.215	93.4	0.001	0.223	0.214	94.4
	0.2	0.001	0.198	0.195	95.2	0.006	0.201	0.214	96.5
	0.4	0.003	0.165	0.172	97.3	0.002	0.183	0.214	97.9
40	0	-0.003	0.156	0.151	93.8	0.003	0.155	0.151	94.7
	0.2	-0.001	0.141	0.137	93.8	-0.002	0.140	0.151	96.7
	0.4	0.001	0.115	0.118	96.5	0.000	0.125	0.151	98.3
60	0	-0.001	0.125	0.123	94.5	0.003	0.123	0.123	95.3
	0.2	-0.001	0.114	0.111	94.3	0.003	0.112	0.123	96.8
	0.4	0.001	0.095	0.096	95.8	0.001	0.101	0.123	98.3

Table 2

Estimates of the expected length and coverage probability (CP) of 95% Sarmanov CI and Wald CI for the first study's relative risk in 5,000 simulations with various number of studies n , for various between-group correlations ρ . The study sample sizes are 400 in both groups. $(a_1, b_1, a_2, b_2) = (0.5, 0.5, 0.5, 0.5)$.

n	ρ	Sarmanov CI		Wald CI	
		Length	CP (%)	Length	CP (%)
20	0.00	0.885	94.8	0.934	95.3
	0.20	0.868	95.4	0.927	94.9
	0.40	0.983	94.6	1.067	94.6
40	0.00	0.882	94.7	0.904	95.3
	0.20	1.023	94.6	1.097	94.6
	0.40	0.964	94.2	1.022	94.1
60	0.00	0.852	95.0	0.898	95.0
	0.20	0.937	94.5	0.991	95.0
	0.40	0.951	95.0	1.019	94.7