Published in final edited form as:

J Psychosom Res. 2014 August ; 77(2): 151–154. doi:10.1016/j.jpsychores.2014.05.006.

Comparing Weight Gain in the Year Prior to Treatment for Overweight and Obese Patients with and without Binge Eating Disorder in Primary Care

Valentina Ivezaj, Ph.D.¹, Roushig Kalebjian, M.A., R.N.², Carlos M. Grilo, Ph.D.¹, and Rachel D. Barnes, Ph.D.¹

¹Department of Psychiatry, Yale University School of Medicine

²Yale School of Nursing

Abstract

Objective—To examine weight change trajectories among overweight and obese patients with binge eating disorder (BED) versus without (NBO) during the year prior to seeking treatment.

Methods—Participants were 97 (75 women, 22 men) overweight and obese patients recruited for the same weight-loss treatment in primary care; 26 (27%) met *DSM-5* BED criteria. Participants were assessed with the Eating Disorder Examination and completed self-report questionnaires about their weight histories and the Beck Depression Inventory-II.

Results—Participants' self-reported current weight and measured current weight were significantly correlated and did not statistically differ. Reported weight changes during the year prior to seeking treatment differed significantly by group: BED patients gained an average of 18.3 lb (8.2 kg) whereas NBO patients gained an average of 1.5 lb (0.7 kg). Among BED patients, but not NBO, weight change during the prior year was positively correlated with greater eating-disorder psychopathology, binge-eating frequency, frequency of overeating at lunch and dinner, and depression scores. For the overall group, BED status and binge-eating frequency each made independent significant contributions to predicting weight change in the past year.

Conclusion—Findings suggest BED patients are gaining considerably more weight during the year prior to treatment than NBO patients. BED treatment may interrupt a steep weight gain trajectory and prevent further weight gain for BED patients suggesting need for early intervention. Primary care physicians should screen for BED when overweight and obese patients present with rapid weight gain.

© 2014 Elsevier Inc. All rights reserved.

Correspondence: Valentina Ivezaj, Ph.D., Yale University School of Medicine, P.O. Box 208098, New Haven, CT 06520-8098; Phone: (203) 785-7807; Fax: (203) 785-7855; valentina.ivezaj@yale.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure of Conflicts

The authors have no competing interests to report.

Keywords

Binge Eating; Eating Disorder; Obesity; Primary Care; Weight Change; Weight Gain

Introduction

Binge eating disorder (BED), defined as recurrent binge-eating accompanied by marked distress but without extreme weight-compensatory behaviors, is a prevalent eating disorder associated strongly with obesity [1]. BED is linked with heightened medical and psychosocial problems [1, 2] and greater health-care utilization [3] relative to obese individuals without BED (NBO). Although certain psychological/behavioral [4] and pharmacological [5] treatments are effective for reducing binge-eating and associated psychopathology, most of these interventions generally result in minimal weight loss with obese BED patients [5, 6]. It has been suggested, however, that treatments that reduce binge eating may interrupt *further weight gain* [7]. Indeed, in a community-based study of non-treated young adults with BED, the rate of obesity increased from 22% to 39% during a five-year naturalistic follow-up [8].

Three recent studies with treatment-seeking BED patients [9–11] reported steep weight gain trajectories with self-reported weights averaging 9.5 lb (4.3 kg) [9] to 16.6 lb (7.5 kg) [11] during the year prior to seeking treatment in both primary care [9] and specialty treatment clinics [10,11]. These studies converged in suggesting that many patients with BED gain substantial amounts of weight prior to seeking treatment but differed somewhat in findings regarding the magnitudes of the weight fluctuations and their associations with binge-eating. Prior studies have also reported weight fluctuations during the weeks between patients' intake and first weight-loss treatment session among obese patients [12] and obese patients with BED [13].

Importantly, no studies have compared weight trajectories between overweight/obese patients with versus without BED prior to treatment. Thus, it remains unclear whether the reported weight gains [9–11] are specific to treatment-seeking BED patients. The present study compared weight changes during the past year by BED and NBO groups recruited to participate in the same weight-loss treatment study and examined the relationships between reported weight changes to eating behaviors and mood.

Methods

Participants were a series of 97 (75 women, 22 men) overweight/obese (body mass index [BMI] ≥ 25) patients recruited through primary care provider referrals and flyers placed in waiting/patient rooms for a behavioral weight loss treatment study being performed in primary care in a large university-based medical health-care center in an urban setting. Participants were informed that the behavioral treatments were provided as part of a research study and would involve no cost. Twenty-six (27%) participants met *DSM-5* BED criteria [14]. The majority of participants were White ($n=70$, 72%) and had at least some college education ($n=87$, 90%). Recruitment for the treatment study was intended to enhance generalizability by utilizing relatively few exclusionary criteria (BMI ≥ 55 , age over 65,

select severe psychiatric problems (e.g., schizophrenia, bipolar disorder), severe medical problems (e.g., cardiac disease), current or planned pregnancy/breastfeeding, and uncontrolled liver disease, hypertension, thyroid disease, or diabetes. Regular access to the internet and telephone was required.

Procedures

Participants provided informed consent (study had IRB approval) and completed questionnaires. Participants were assessed by trained research-clinicians. During the screening phase of the study, the following two questions were added as part of the exclusion criteria: 1) How much has your weight changed in the last 6 months? This was only an exclusion if weight change was greater than 9 pounds, due to a medical condition such as pregnancy or hormonal therapy. 2) Are you currently receiving any treatment for your eating or weight (weight loss programs)?

Measures

Participants were asked to provide self-reported lowest and highest adult weights, ages for those weights, height, and weight currently, 3, 6, and 12 months ago. Actual weight was measured using a high-capacity digital scale and height was measured.

The Eating Disorder Examination (EDE [15]) is a well-established semi-structured interview that assesses eating disorders. The EDE has demonstrated good reliability in BED patients [16]. Analyses examined the EDE Global score (reflecting eating-disorder psychopathology), patients' perceptions of overeating at specific meals, and average number of days with *objective bulimic episodes* (OBEs; eating unusually large quantities of food with a subjective sense of loss of control, which corresponds to binge-eating) during the past six months. These OBEs were considered binge eating episodes.

The Beck Depression Inventory-II (BDI-II [17]) assesses current depression level. It is a well-established measure with excellent reliability and validity [18].

Data Analyses

Weight Change during the past 12-months was calculated by subtracting Self-Reported Weight 12 Months Ago from Self-Reported Current Weight. Three groups also were created: Maintained Weight (gained or lost ≤ 5 lb or 2.25 kg), Lost Weight (lost ≥ 5 lb or 2.25 kg), and Gained Weight (gained ≥ 5 lb or 2.25 kg) during the past 12 months [9]. Percent Weight Change also was calculated to help address different starting weights between the NBO and BED groups. Correlations were examined to test the relationships between Weight Change and the clinical measures for each of the BED and NBO groups. Additionally, *t*-tests were used to compare differences in clinical measures between the BED and NBO groups. A stepwise multiple regression analysis was conducted to examine joint and independent contributions of variables predicting weight change during the past 12 months. Variables that differed between groups and variables that were associated significantly with Weight Change at the bivariate level were entered into the regression analyses.

Results

The BED and NBO groups did not differ on sex, race, or education. BED patients (age $M=43.6$, $SD=9.4$) were significantly younger than NBO patients (age $M=48.9$, $SD=10.6$), $t=2.22$, $p=.029$. Age was not significantly correlated with Weight Change ($p=.195$). BED patients were significantly heavier than NBO patients and therefore Percent Weight Change also was examined (Table 1).

Accuracy of Self-Reported Weight

Overall, participants' self-reported current weight ($M=217.9$, $SD=50.1$ lb or $M=98.1$, $SD=22.5$ kg) and measured current weight ($M=218.4$, $SD=50.2$ lb or $M=98.3$, $SD=22.6$ kg) were significantly correlated (Pearson's $r=.999$, $p<.0005$) and did not statistically differ ($p=.069$); mean difference was less than one pound. Similar results emerged for the BED group; self-reported current weight and measured current weight were significantly correlated (Pearson's $r=.999$, $p<.0005$) and did not statistically differ ($p=.243$). For the NBO group, self-reported weight and actual measured weight were significantly correlated (Pearson's $r=.998$, $p<.0005$), but were statistically different ($p=.001$), albeit a difference of less than 1 pound.

Weight Trajectories during the Year Prior to Treatment

Overall, patients gained an average of 6.0 lb or 2.7 kg ($SD=20.5$ lb or $SD=9.2$ kg), which corresponds to 2.36% ($SD=8.8$) of their body weight in the 12 months before weight loss treatment. Weight Change differed significantly; BED patients gained $M=18.3$ lb or 8.2 kg ($SD=24.8$ lb or 11.2 kg; range=-4.0 lb to +99.3 lb or -1.8 kg to 44.7 kg) and NBO patients gained $M=1.5$ lb or 0.7 kg ($SD=16.6$ lb or 7.5 kg; range=-77.6 lb to +50.0 lb or -34.9 kg to +22.5 kg) (Table 1, Figure 1a). Lowest adult weight and age at lowest adult weight did not differ by group (Table 1).

Among BED patients, 34.6% ($n=9$) maintained weight (gained or lost < 5 lb or 2.25 kg), 0% ($n=0$) lost weight (lost ≥ 5 lb or 2.25 kg), and 65.4% ($n=17$) gained weight (gained ≥ 5 lb or 2.25 kg). Among NBO patients, 36.8% ($n=25$) maintained weight, 19.1% ($n=13$) lost weight, and 44.1% ($n=30$) gained weight during the year before treatment. Fisher's Exact Test indicated that there were more NBO patients in the Lost Weight category than BED patients, $\chi^2(N=94)=5.77$, $p=.017$, but not the Weight Gained, $p=.065$, or Weight Maintained, $p=.846$, categories (see Figure 1b for Weight Change categories).

Eating behaviors and depression

Table 1 summarizes the correlates of 12-month Weight Change with eating-disorder psychopathology, eating behaviors, and depression. Among BED patients, Weight Change was positively correlated with EDE Global, binge-eating frequency, frequency of overeating at lunch and dinner, and BDI-II scores. Among NBO patients, Weight Change was not significantly associated with any of the clinical measures. A stepwise multiple regression analysis was conducted to examine the joint and independent contributions of the variables (EDE Global, binge-eating frequency, frequency of overeating at lunch and dinner, and BDI-II scores), when controlling for age, current BMI, highest adult weight, and BED

status, to predict Weight Change. Analyses revealed that BED status and binge-eating frequency accounted for 41.1% of the variance of Weight Change, $F(2, 90) = 31.43$, $p < .0005$, and both made significant independent contributions: BED status ($\beta = 0.384$, $t = -2.73$, $p = .008$) and binge-eating frequency ($\beta = 0.916$, $t = 6.52$, $p < .0005$).

Discussion

This is the first study, to our knowledge, to compare weight change for overweight/obese individuals with and without BED in primary care during the year prior to seeking weight-loss treatment. Several notable findings emerged. First, there was a striking difference in weight change trajectories; BED patients gained, on average, 18.3 lb (8.2 kg), while NBO patients gained, on average, only 1.5 lb (0.7 kg), during the year prior to seeking treatment. Second, the magnitude of weight fluctuations varied for both groups; weight change of BED patients ranged from 4 lb lost to 99.3 lb gained (1.8 kg lost to 44.7 kg gained), while weight change of NBO patients ranged from 77.6 lb lost to 50.0 lb gained (34.9 kg lost to 22.5 kg gained). When categorizing patients as Maintained Weight (gained or lost < 5 lb or 2.25 kg), Lost Weight (lost ≥ 5 lb or 2.25 kg), or Gained Weight (gained ≥ 5 lb or 2.25 kg), nearly two-thirds of BED patients gained weight and none lost, while less than half of NBO patients gained and nearly 20% lost weight during that year. Finally, Weight Change during the past year was predicted significantly (jointly and independently) by BED status and binge-eating frequency.

Both groups experienced their lowest adult weight at the same age (early twenties), which corresponds with the average age of BED onset in epidemiologic studies [1]. Despite starting at a similar age for lowest adult weight, the BED group currently weighed more even though they were *younger* than the NBO group. Our findings replicate the previously documented sharp weight gain trajectory of BED patients [9–11] but provide specificity to the findings by showing that such weight gains are considerably greater than treatment-seeking NBO patients, who, in turn, appear to show what is generally expected in the general population [19, 20]. Importantly, among the BED group, weight change was strongly related to binge-eating frequency and moderately with eating-disorder psychopathology and overeating. When examining predictors of weight change over the past 12 months for the overall study group, BED status and frequency of binge eating emerged as the only significant predictors of weight change, jointly accounting for approximately 41% of the variance. This finding highlights the relationship between binge eating and weight gain and calls for early screening for BED to help prevent significant weight gains, particularly among BED patients. Of note, this BED group was similar to, and perhaps less severe than, BED groups in previous studies [9–11], indicating that differences between the BED and NBO groups did not emerge due to a particularly or unusually severe BED group.

Collectively, these findings suggest the importance of early identification and treatment of BED within primary care settings. For example, in addition to screening for medical conditions, primary care providers should screen for BED when overweight and obese patients present with rapid weight gain. These findings support previous suggestions that treating disordered eating may prevent excess weight gain and/or the worsening of obesity

among those with BED [7]. Future research should also investigate pre-treatment weight trajectories in relation to BED and weight loss treatment outcomes.

We note potential limitations. Our findings pertain to treatment-seeking overweight and obese persons in primary care settings who choose to participate in research. Findings may not generalize to other treatment settings or to overweight/obese groups with different characteristics. However, our recruitment and ascertainment of the overweight/obese patients with and without BED within primary care for the same obesity treatment reduces some potential confounds and clinic biases and enhances generalizability to the extent that primary care tends to serve as a gate-keeper in our health-care system. Another potential limitation is reliance on self-reported weight history, although consistent with previous reports [9–11, 21], the group was extremely accurate in reporting current weight. While the accuracy of retrospective weights for previous time periods is uncertain, epidemiologic research has found that although there tends to be some biases, overall, adults are relatively accurate in reporting weight for periods as long as ten years prior [22]. In addition, participants were not asked about their weight loss treatment during the previous year, which would aid in our understanding of the vast weight fluctuations observed in both groups. Lastly, the study was cross-sectional in nature and included a small sample size. Future studies with larger sample sizes are needed to establish these findings prospectively and to test the prognostic significance of pre-treatment weight trajectories.

In conclusion, BED patients gained considerably more weight during the year prior to treatment than NBO patients. Results provide further evidence [9–11] to support the notion that BED treatments may serve an important role in weight stabilization and prevention of future weight gain for BED patients. Early detection, prevention, and interventions for BED, particularly among weight-loss treatment seeking individuals in primary care, may be critical in preventing more severe obesity.

Acknowledgments

This study was supported, in part, by grants from the National Institutes of Health (K23 DK092279, K24 DK070052). The National Institutes of Health were not involved in study design, collection, analysis, interpretation of data, the writing of the report, or in the decision to submit this article for publication. The article does not represent the views of the NIH. No additional funding was obtained for submitting this article.

Conflicts of Interest Statements

Drs. Grilo, Barnes, Ivezaj, and Roushig Kalebjian report grants from NIH during the conduct of the study. Dr. Grilo also reports personal fees from Shire, other from American Psychological Association, and from Guilford Press and Taylor and Francis Publishers, outside the submitted work.

References

1. Hudson JL, Hiripi E, Pope HG, Kessler RC. The prevalence and correlates of eating disorder in the NCS Replication. *Biol Psychiatry*. 2007; 61:348–358. [PubMed: 16815322]
2. Grilo CM, Hrabosky HI, White MA, Allison KC, Stunkard AJ, Masheb RM. Overvaluation of shape and weight in binge eating disorder and overweight controls: refinement of a diagnostic construct. *J Abnorm Psychol*. 2008; 117:414–419. [PubMed: 18489217]
3. Johnson JG, Spitzer RL, Williams JB. Health problems, impairment and illnesses associated with bulimia nervosa and binge eating disorder among primary care and obstetric gynecology patients. *Psychol Med*. 2001; 31:1455–1466. [PubMed: 11722160]

4. Wilson GT, Grilo CM, Vitousek KM. Psychological treatment of eating disorders. *Am Psychol*. 2007; 62:199–216. [PubMed: 17469898]
5. Reas DL, Grilo CM. Review and meta-analysis of pharmacotherapy for binge-eating disorder. *Obesity*. 2008; 16:2024–2038. [PubMed: 19186327]
6. Grilo CM, Masheb RM, Wilson GT. Efficacy of cognitive behavioral therapy and fluoxetine for the treatment of binge eating disorder: a randomized double-blind placebo-controlled comparison. *Biol Psychiatry*. 2005; 57:301–309. [PubMed: 15691532]
7. Yanovski SZ. Binge eating disorder and obesity in 2003: could treating an eating disorder have a positive effect on the obesity epidemic? *Int J Eat Disord*. 2003; 34/1(Supp):117–120.
8. Fairburn CG, Cooper Z, Doll HA, Norman P, O'Conner M. The natural course of bulimia nervosa and binge eating disorder in young women. *Arch Gen Psychiatry*. 2000; 57:659–665. [PubMed: 10891036]
9. Barnes RD, Blomquist KK, Grilo CM. Exploring pretreatment weight trajectories in obese patients with binge eating disorder. *Compr Psychiatry*. 2011; 52:312–318. [PubMed: 21497226]
10. Blomquist K, Barnes RD, White MA, Masheb RM, Morgan PT, Grilo CM. Exploring weight gain in year before treatment for binge eating disorder: a different context for interpreting limited weight losses in treatment studies. *Int J Eat Disord*. 2011; 44:435–439. [PubMed: 20635382]
11. Masheb RM, White MA, Grilo CM. Substantial weight gains are common prior to treatment-seeking in obese patients with binge eating disorder. *Compr Psychiatry*. 2013; 54:880–884. [PubMed: 23639407]
12. West DS, Harvey-Berino J, Krukowski RA, Skelly JM. Pretreatment weight change is associated with obesity treatment outcomes. *Obesity*. 2011; 19:1791–1795. [PubMed: 21331064]
13. Eldredge KL, Agras WS, Arnow B. The last supper: Emotional determinants of pretreatment weight fluctuation in obese binge eaters. *Int J Eat Disord*. 1994; 16:83–88. [PubMed: 7920585]
14. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (5th ed). Arlington, VA: American Psychiatric Publishing. In text: American Psychiatric Association; 2013.
15. Fairburn, CG.; Cooper, Z. The eating disorder examination (12th ed.). In: Fairburn, CG.; Wilson, GT., editors. *Binge Eating: Nature, Assessment, and Treatment*. New York: Guilford Press; 1993. p. 317-360.
16. Grilo CM, Masheb RM, Lozano-Blanco C, Barry DT. Reliability of the Eating Disorder Examination in patients with binge eating disorder. *Int J Eat Disord*. 2004; 35:80–85. [PubMed: 14705160]
17. Beck, AT.; Steer, R. *Manual for revised Beck Depression Inventory*. New York: Psychological Corporation; 1987.
18. Beck AT, Steer R, Garbin MG. Psychometric properties of the beck depression inventory 25 years of evaluation. *Clin Psychol Rev*. 1988; 8:77–100.
19. He XZ, Baker DW. Changes in weight among a nationally representative cohort of adults aged 51 to 61, 1992 to 2000. *Am J Prev Med*. 2004; 27:8–15. [PubMed: 15212769]
20. Williamson DF, Kahn HS, Byers T. The 10-year incidence of obesity and major weight gain in black and white US women aged 30–55 years. *Am J Clin Nutr*. 1991; 53/6(Suppl):1515–1518.
21. Barnes RD, White MA, Masheb RM, Grilo CM. Accuracy of self-reported weight and height and resulting body mass index among obese binge eaters in primary care: Relationship with eating disorder and associated psychopathology. *Prim Care Companion J Clin Psychiatry*. 2010; 12:e1–e6.
22. Perry GS, Byers TE, Mokdad AH, Serdula MK, Williamson DF. The validity of self-reports of past body weights by U.S. adults. *Epidemiol*. 1995; 6:61–66.

Highlights

- Binge eating disorder (BED) is a common eating disorder associated with obesity.
- Obese patients with BED report steep weight gains during the year before treatment.
- Compared pretreatment weight trajectories of patients with BED vs. without (NBO).
- Pretreatment year weight gain was higher for obese/overweight BED vs. NBO patients.
- BED status and binge-eating frequency each made independent significant contributions to predicting weight change in the past year.

a. Weight trajectories during the year prior to seeking treatment shown for the overall participant group and separately for the binge eating disorder (BED) and non-BED overweight/obese (NBO) groups.

b. Weight changes during the year prior to seeking treatment reported by percentage of the binge eating disorder (BED) and non-BED overweight/obese (NBO) groups.

Figure 1.

a. Weight trajectories during the year prior to seeking treatment shown for the overall participant group and separately for the binge eating disorder (BED) and non-BED overweight/obese (NBO) groups.

b. Weight changes during the year prior to seeking treatment reported by percentage of the binge eating disorder (BED) and non-BED overweight/obese (NBO) groups.

Table 1

Descriptive statistics, bivariate correlations of clinical variables with weight change^a, and t-tests comparing study groups.

	BED n=26	BED n=26	NBO n=70	NBO n=70	Test Statistic
	M (SD)	Weight Change 12 Months (r)	M (SD)	Weight Change 12 Months (r)	t-test
Self-reported weight	247.4 (61.6)	-	207.0 (40.8)	-	-3.11**
Actual weight	246.7 (62.4)	-	207.9 (41.0)	-	-2.94**
Current BMI	38.6 (7.8)	-	33.6 (6.0)	-	-3.34***
WC % 12 months	6.9 (9.0)	-	0.7 (8.2)	-	-3.22**
WC 12 months	18.3 (24.8)	-	1.5 (16.6)	-	-3.20**
WC 6 months	7.8 (17.4)	-	3.2 (9.0)	-	-1.25
WC 3 months	4.0 (8.3)	-	2.1 (5.8)	-	-1.26
Lowest weight	159.2 (39.9)	-	146.1 (31.2)	-	-1.69
Adult age	21.2 (4.7)	-	23.5 (7.3)	-	1.50
Highest weight	254.2 (72.9)	-	216.5 (45.6)	-	-2.47*
Adult age	40.2 (9.4)		45.9 (11.1)		2.36*
BDI-II	11.1 (7.6)	.382*	6.7 (5.7)	-.003	-3.05*
EDE global	2.2 (0.8)	.473*	1.6 (0.9)	.093	-2.98**
Overeating Frequency ^b					
Breakfast	7.0 (9.0)	.236	3.8 (6.7)	.105	-1.68
Lunch	8.8 (9.4)	.500**	7.2 (8.6)	.137	-0.78
Dinner	13.3 (10.2)	.417*	11.6 (9.0)	.181	-0.79
OBE Frequency	2.9 (1.6)	.845***	0.1 (0.2)	.049	-8.52***

Note. BED = binge eating disorder group. NBO = non-BED obese group. BMI = body mass index. WC = weight change. BDI-II = Beck Depression Inventory-II. EDE = Eating Disorder Examination. OBE = objective bulimic episodes; frequency represents the average during the last 6 months.

*
p .05,
**
p .01,

p .001.

^a Pearson's *r* correlations between clinical variables and participants' Weight Change during the past 12 months (Current Weight – Weight 12 Months Ago).

^b Multiply by 0.45 to convert to kg.

^c Overeating represents frequency of overeating at meals during the past 28 days.