Published in final edited form as:

Compr Psychiatry. 2008 ; 49(1): 6–12. doi:10.1016/j.comppsy.2007.07.006.

Time to Emergence of Severe Suicidal Ideation among Psychiatric Patients as a Function of Suicide Attempt History

Margaret S. Andover, Ph.D.¹, Brandon E. Gibb, Ph.D.², and Ivan W. Miller, Ph.D.¹

¹Psychosocial Research Program, Butler Hospital and Brown Medical School, Providence, RI

²Department of Psychology, Binghamton University (SUNY), Binghamton, NY

Abstract

Background—Little is known about the emergence of suicidal ideation among psychiatric inpatients with histories of no, single, or multiple suicide attempts. We investigated differences in time to reemergence of severe suicidal ideation among psychiatric patients as a function of their suicide attempt histories.

Method—One hundred seventeen individuals meeting criteria for a major depressive disorder who were recently discharged from a psychiatric hospital and participating in a larger study of treatments for depression were included in the current study. Suicidal ideation, depressive symptoms, hopelessness, and depressogenic cognitions were assessed at baseline, and suicidal ideation was assessed at 3, 6, 12, and 18 month follow up, as well as inpatient readmission if applicable. Time to the reemergence of severe suicidal ideation was analyzed using survival analysis.

Results—Twenty-two percent of our sample reported the occurrence of severe suicidal ideation over an 18-month period. Severe suicidal ideation emerged earlier among patients who had a history of prior suicide attempts than those who did not, but single and multiple suicide attempters did not differ significantly in time to severe suicidal ideation. Suicide attempt history remained a significant predictor of time to severe suicidal ideation when statistically controlling for hopelessness, depressive symptoms, depressogenic cognitions, and suicidal ideation at admission and initial treatment group assignment, especially between single and non-attempters.

Conclusions—Although nearly a quarter of participants endorsed severe, clinically significant suicidal ideation within 18 months of discharge, those with suicide attempt histories reported the occurrence of severe suicidal ideation significantly earlier than those without suicide attempt histories.

Suicidal ideation, attempted suicide, and death by suicide represent a range of suicidal behaviors. Although death by suicide is a low base rate behavior, occurring in approximately 0.01% of the population in 2003 [1], the prevalence of suicidal ideation is significantly higher. Between 11 and 17% of individuals living in the United States report some suicidal ideation during their lifetime [2, 3]. Suicidal ideation represents an important public health

Correspondence should be addressed to: Margaret S. Andover, Butler Hospital, Psychosocial Research Program, 345 Blackstone Blvd., Providence, RI 02906 (Tel: 401-455-6456; Fax: 401-455-6235; Margaret_Andover@brown.edu).

The authors report no conflicts of interest.

problem because of increased utilization of mental health services [4], the distress it causes the individual and others, its implications for the mental health of the individual, and its role as a risk factor for death by suicide [5]. Individuals reporting suicidal ideation or attempted suicide are likely to experience enduring suicidal ideation [6], and those reporting suicidal ideation are more likely to attempt suicide in the future [5, 7].

The emergence of suicidal ideation is of both empirical and clinical interest. For instance, the Food and Drug Administration's public health advisories of increased suicidal behavior in children and adolescents [8], and later adults [9], being treated with antidepressants has generated recent interest in the emergence of suicidal thoughts and behaviors among individuals receiving pharmacotherapy [10, 11]. However, the length of time to emergence of severe suicidal ideation following inpatient treatment has been little studied. In one study among adolescents, emergence was most common within the first two years after the initial assessment [12]. Further research on the emergence of suicidal ideation is necessary to better understand suicidality and to develop intervention and prevention efforts better timed to periods of heightened risk. Research on the topic generally has focused on the presence or absence of suicidal ideation without accounting for ideation severity. However, the severity of suicidal ideation is of clinical importance and may vary in its clinical significance (i.e., experiencing thoughts of death versus having a suicide plan). To the best of our knowledge, there have been no studies investigating time to the emergence of clinically significant suicidal ideation either in adult psychiatric patients or as a function of suicide attempt status.

Although traditionally classified together, recent research has suggested that individuals with a history of multiple suicide attempts differ from those with a history of a single attempt. Specifically, multiple attempters report more symptoms of hopelessness, depression, and anger than single attempters following a suicide attempt [13, 14]. Individuals with a history of multiple attempts report more suicidal intent [15] and are more likely to attempt suicide in the five years following inpatient discharge than those with a history of a single attempt [16, 17]. Further, individuals with a history of multiple suicide attempts report higher levels of suicidal ideation than those with a history of a single attempt [14] and are more likely than single and non-attempters to report increased suicidal ideation within 12 months following a suicidal crisis [18].

The purpose of this study was to investigate differences in time to emergence of clinically significant suicidal ideation, reflecting high levels of ideation, desire for death, preparation, and resolve [19], among adult psychiatric patients with histories of no, single, or multiple suicide attempts. Using survival analysis, we explored length of time from hospital discharge to emergence of clinically significant, severe, suicidal ideation as a function of suicide attempt status. In addition, we investigated whether suicide attempt status would be a stronger predictor of time to severe suicidal ideation than other factors associated with suicidal ideation, such as hopelessness and depression [20], negative cognitive styles [21, 22], and recent suicidal ideation [23].

Method

Participants

This study consisted of 121 patients recently admitted to a psychiatric inpatient or partial hospital unit who were participating in a larger longitudinal study of the efficacy of combined psychosocial and pharmacological treatments for depression [24], which was approved by the hospital's institutional review board. Additional details regarding the protocol and outcomes of the treatment study can be found in Miller et al [24]. Individuals included in the study were between 18 and 65 years of age, met criteria for a diagnosis of Major Depressive Disorder obtained through the Structured Clinical Interview for the DSM-III (SCID-P) [25], reported scores greater than 17 on both the Beck Depression Inventory [26] and the Modified Hamilton Rating Scale for Depression [27], had sufficient reading skills to complete questionnaires, were currently living with at least one family member, and informed consent was obtained from both the patient and family. Individuals were excluded from participating if they met criteria for a current diagnosis of borderline personality disorder, bipolar disorder, schizophrenia, substance dependence, or somatization disorder, had a primary diagnosis of post-traumatic stress disorder, had a medical illness that contraindicated antidepressant medication, or if significant cognitive impairment prevented them from completing questionnaires.

Assessments

Assessments took place at hospital admission, hospital discharge, and 3 months, 6 months, 12 months, and 18 months following discharge. In addition, assessment measures were administered at the time of hospital readmission, if applicable. Assessments were conducted in person whenever possible and by telephone when necessary. The majority of patients (76%; 92/121) completed assessments through the 18-month follow up.

Measures

Patients were interviewed for specific information regarding suicide attempts, such as number of attempts and age at first attempt. In addition, demographic variables, including patient age and ethnicity, were obtained during this interview. The Modified Scale for Suicidal Ideation (MSSI) [28], an 18-item interview modified from the Scale for Suicidal Ideation [29], was used to assess patients' levels of suicidal ideation. The scale consists of two factors: Suicidal Desire and Ideation and Resolved Plans and Preparation [19]. Excellent internal consistency and concurrent validity have been reported in clinical samples [30], and the scale has been used in a number of research studies. A score of 21 on the MSSI was defined as a cutoff for severe levels of suicidal ideation [28], reflecting high levels and frequency of suicidal thoughts and desires, resolve, and a developed plan [19]. Depressive symptoms were assessed using the Beck Depression Inventory (BDI) [26], a self-report measure, and the Modified Hamilton Rating Scale for Depression (MHRSD) [27], a clinician-administered interview. The BDI and MHRSD are commonly used in research, and research has demonstrated strong reliability and validity in clinical samples for both measures [27, 31]. The Beck Hopelessness Scale (BHS) [32] was used to assess the individual's negative expectations about the future. This commonly used measure has demonstrated strong reliability validity across several populations [33]. Depressogenic

cognitions were measured with the Dysfunctional Attitude Scale (DAS) [34] and the Cognitive Bias Questionnaire (CBQ) [35]. The DAS consists of 40-items designed to measure beliefs that may represent a cognitive vulnerability to depression. Studies have reported excellent internal consistency and moderate validity against a measure of depressive symptoms [36]. The CBQ was designed to measure negatively biased cognitive styles by assessing an individual's typical response for a given scenario. The depressed-distorted subscale, which assesses unrealistically negative cognitive style, was included in this study's analyses. The CBQ correlates significantly with measures of depression and negative self-statements [37].

Treatment

Upon discharge from the hospital, participants were randomized to receive one of four treatment conditions that was either "matched" or "mismatched" to their pretreatment level of deficits for a six-month period. The treatment conditions included: pharmacotherapy alone; pharmacotherapy and cognitive therapy; pharmacotherapy and family therapy; or pharmacotherapy, cognitive therapy, and family therapy. See Miller et al [24] for greater detail regarding the randomization procedures and treatment conditions.

During this six month period, a subset of patients (19%; 23/121) who manifested significant deterioration or suicide risk were removed from their randomized treatment and, if clinically appropriate, were provided with another or additional study treatment or referred outside the study for treatment. Another 20% (24/121) of patients dropped out of randomized treatment during the first six months. After 6 months, individuals who completed randomized treatment and had responded to treatment continued with that intervention for an additional 12 months (55.4%; 67/121). Those who had not responded were referred outside the study to another treatment. During the entire study, patient assessment continued regardless of treatment status. That is, we attempted to complete assessments on all patients initially entered into the study. Thus, it is important to note that data included in this report is largely naturalistic, as treatment was randomly assigned for only for the first 6 months of the 18-month study, and only a subset of patients who did not drop out actually received randomized treatment.

Statistical Analysis

Group differences were analyzed using analyses of variance (ANOVAs) and chi-square analyses when appropriate. Time to emergence of severe suicidal ideation was investigated using survival analysis. This set of statistical procedures is used to model time until an event occurs, or survival time, and uses survival time as the dependent variable in analyses. Survival analysis also incorporates incomplete survival time by analyzing censored data. In the current study, the Kaplan-Meier method was used to determine differences in survival time among the groups, and the Cox proportional hazard regression was used to investigate predictors to survival time [38, 39]. All data analyses were performed using SPSS version 13.

Results

Although 121 psychiatric patients participated in the original treatment outcome study [24], 4 of these participants (1 single attempter and 3 multiple attempters) were excluded from the current study because they endorsed severe suicidal ideation (MSSI score of 21 or greater) at discharge. Of the 117 participants included in this study, 70 (57.9%) reported no history of a suicide attempt, while 20 (17.1%) reported a single attempt and 27 (23.1%) reported multiple attempts. Descriptive statistics for our sample are presented in Table 1. The groups did not differ significantly in terms of sex, age, ethnicity, marital status, or treatment group assignment at randomization. However, significant group differences were found for mean suicidal ideation at admission ($F(2, 117) = 17.01, p < .001$) and discharge ($F(2, 117) = 6.73, p = .002$), with single and multiple attempters reporting more suicidal ideation at admission and discharge than non-attempters, and multiple attempters reporting more suicidal ideation at admission than single attempters. It is important to note that group means at discharge from the hospital reflected low suicidal ideation according to the measure's cutoff scores [28]. Groups also differed significantly in depressive symptoms at admission as reported on the BDI ($F(2, 116) = 3.06, p = .05$) and MHRSD ($F(2, 116) = 3.05, p = .05$), with multiple attempters reporting more depressive symptoms than non-attempters. Groups did not differ in the occurrence of rehospitalization after hospital discharge ($\chi^2(2, N = 117) = 2.32, p = .31$). Initial treatment group assignment was not significantly associated with suicide attempt status ($\chi^2(6, N = 117) = 9.81, p = .13$) or the MSSI at admission ($F(3, 115) = 0.67, p = .57$).

A review of clinical team notes revealed five suicide attempts during the assessment period. Two individuals attempting suicide during the study reported no previous attempts at baseline, one reported one previous attempt, and two reported multiple previous attempts. There were no deaths by suicide during the assessment period. Twenty-six (22.2%) of the 117 participants reported the occurrence of severe levels of suicidal ideation during follow up, and approximately 7% ($n = 8$) of participants reported recurrent severe suicidal ideation (defined as severe suicidal ideation at more than one time point during the follow up period). Although patients with a history of suicide attempts were more likely to report the emergence of severe suicidal ideation during follow up than non-attempters ($\chi^2(1, N = 117) = 15.06, p = .001$), the difference between single and multiple attempters was not significant ($\chi^2(1, N = 47) = 0.003, p = .96$). As shown on Table 2, participants reporting the emergence of severe suicidal ideation during follow up were more likely to report greater suicidal ideation at admission ($t(114) = 3.94, p < .001$) and discharge ($t(112) = 2.53, p = .01$), and greater dysfunctional attitudes on the DAS ($t(115) = 2.16, p = .03$). They did not differ with regard to initial treatment group assignment ($\chi^2(3, N = 117) = 5.28, p = .15$), depressive symptoms at admission (BDI: $t(115) = 1.17, p = .24$; MHRSD: $t(115) = 1.78, p = .08$), age ($t(115) = .49, p = .63$), gender ($\chi^2(1, N = 117) = 0.003, p = .96$), ethnicity ($\chi^2(1, N = 117) = 0.27, p = .60$), or marital status ($\chi^2(1, N = 117) = 0.34, p = .56$).

Next, we investigated differences in time to emergence of severe suicidal ideation using survival analysis. Cumulative survival time is presented in Table 3. We performed a Kaplan-Meier analysis to examine whether individuals differed in the likelihood of eventually reporting severe levels of suicidal ideation based on suicide attempt status. Including individuals who did not report severe suicidal ideation during the follow up period,

estimated mean time to severe suicidal ideation differed significantly among non-attempters, single attempters, and multiple attempters (17.06 months, 12.79 months, and 11.98 months, respectively; log-rank = 19.51, $df = 2$, $p < .001$; Breslow = 20.30, $df = 2$, $p < .001$; Tarone-Ware = 20.02, $df = 2$, $p < .001$; see Table 4). Pairwise comparisons revealed significant differences between non-attempters and single attempters (Wilcoxon = 12.07, $df = 1$, $p = .001$) and non-attempters and multiple attempters (Wilcoxon = 15.42, $df = 1$, $p < .001$), with non-attempters reporting a longer duration than single or multiple attempters. The difference between single and multiple attempters was not significant (Wilcoxon = 0.03, $df = 1$, $p = .87$).

A Cox proportional hazard regression was used to investigate whether suicide attempt status predicted time to severe suicidal ideation above other known predictors of suicidal ideation. We found that history one or more suicide attempts was predictive of time to severe suicidal ideation (Wald = 5.01, $df = 1$, $p = .03$, $\beta = 1.09$), even when controlling for initial treatment group assignment and hopelessness, depressive symptoms, depressogenic cognitions, and suicidal ideation at admission. Compared with non-attempters, single attempt status predicted time to suicidal ideation (Wald = 5.94, $df = 1$, $p = .02$, $\beta = 1.35$), but multiple attempt status did not (Wald = 2.20, $df = 1$, $p = .14$, $\beta = .95$). Multiple attempters did not differ significantly from single attempters in time to severe suicidal ideation (Wald = .29, $df = 1$, $p = .59$, $\beta = .30$). Time to severe suicidal ideation was also predicted by suicidal ideation (Wald = 4.37, $df = 1$, $p = .02$, $\beta = .04$), but not depressive symptoms (BDI: Wald = 0.04, $df = 1$, $p = .84$, $\beta = .01$; MHRSD: Wald = 0.14, $df = 1$, $p = .71$, $\beta = .01$), hopelessness[§] (Wald = 0.29, $df = 1$, $p = .59$, $\beta = .03$), depressogenic cognitions (DAS: Wald = 0.83, $df = 1$, $p = .36$, $\beta = .01$; CBQ: Wald = 0.02, $df = 1$, $p = .88$, $\beta = .02$), or initial treatment group assignment (Wald = 0.93, $df = 1$, $p = .33$, $\beta = .20$). The overall model fit was significant ($\chi^2(11) = 30.33$, $p = .001$), and the survival curve as a function of suicide attempt status is shown in Figure 1.

Discussion

The purpose of this study was to investigate differences in time to emergence of severe suicidal ideation among psychiatric patients as a function of their suicide attempt histories. Overall, we found that the attempt history groups differed in their time to severe suicidal ideation. Specifically, severe suicidal ideation emerged earlier for single and multiple attempters than for patients with no prior history of suicide attempts. Single and multiple attempters, however, did not differ significantly from each other. Further, although suicidal ideation at admission was a significant predictor of time to emergence of severe suicidal ideation during follow up, history of a suicide attempt continued to predict time to emergence when statistically controlling for treatment group assignment at randomization and the effects of suicidal ideation, hopelessness, depressive symptoms, and depressogenic cognitions at admission.

[§]Although hopelessness was predictive of time to emergence at lower severities of suicidal ideation (Cutoff of 8: Wald = 5.27, $df = 1$, $p = .02$, $\beta = .09$; Cutoff of 15: Wald = 4.76, $df = 1$, $p = .03$, $\beta = .11$), hopelessness did not predict above suicide attempt status, suicidal ideation at admission, depressive symptoms, depressogenic cognitions, and treatment group assignment (Cutoff of 8: Wald = .54, $df = 1$, $p = .46$, $\beta = .03$; Cutoff of 15: Wald = 1.58, $df = 1$, $p = .21$, $\beta = .07$).

Time to the emergence of severe suicidal ideation has implications for our understanding of and clinical work with psychiatric patients. These findings suggest that psychiatric patients who have attempted suicide will likely experience severe suicidal ideation earlier following hospital discharge than those who have not attempted suicide. Despite intensive treatment, nearly a quarter of our sample (22.2%) reported the occurrence of severe suicidal ideation within the 18-month follow up period, and 7% reported recurrent severe suicidal ideation. The prevalence of emerging severe suicidal ideation, especially among those who have attempted suicide, emphasizes the need for continued clinical care and monitoring.

Results of this study suggested no difference in time to severe, clinically significant suicidal ideation based on frequency of suicide attempts (single versus multiple) and indicated that individuals with a history of suicide attempts are similar in their time to emergence of clinically significant suicidal ideation, regardless of frequency. Data also suggested that history of multiple suicide attempts did not predict time to severe suicidal ideation above other predictors of suicidal ideation, including hopelessness, depressive symptoms, and depressogenic cognitions, while history of a single attempt did. However, these findings contradict previous research suggesting that multiple attempters may be at higher risk for suicidal behaviors than single attempters [14]. In a naturalistic study investigating time to suicide attempt following hospital discharge among adolescents, Goldston et al [17] reported that those with histories of multiple attempts were twice as likely as single or non-attempters to attempt suicide in five years following discharge. However, posthospitalization treatment was not reported in the study, and differences in sample characteristics and outcome variable limit generalizability from Goldston et al [17] to the current study. It is possible that although single and multiple attempters do not differ in time to the emergence of severe suicidal ideation, multiple attempters may simply be more likely to act on this ideation and make a subsequent attempt. Future studies focused on the emergence of both ideation and attempts are needed to test this hypothesis.

Patients' participation in treatment may have impacted reported suicidal ideation in the current study and may account for inconsistencies between this study's findings and previous research. It is important to note that all participants received mental health treatment while on the inpatient unit and outpatient treatment for at least the first six months after discharge from the hospital. In addition, participants either continued treatment for another twelve months or were referred to another type of treatment based on their response to the initial six months of treatment. Due to the large number of potential treatments and the uncontrolled nature of much of the treatment received by participants, we were unable to provide substantial data on the relationship of treatment to the course of suicidal ideation. However, Kessler et al [40] found that although treatment utilization increased over a 10-year period from 1990–1992 to 2001–2003, there was no corresponding decrease in suicidal behaviors. Future studies designed to investigate the effect of specific treatments on the course of suicidal ideation and behavior and needed to address this important issue.

Nearly a quarter of participants did not complete assessments to the 18-month follow up point. While survival analysis considers censored data of those who do not experience the event prior to their termination from the study, this may limit the generalizability of the findings. In addition, although the current authors chose to investigate time to severe

suicidal ideation because of its clinical utility and importance, future studies may investigate time to mild or moderate suicidal ideation. Future studies may also investigate the effects of changes in predictors of suicidal ideation over time, such as changes in depressive symptoms, on the emergence of suicidal ideation.

This study adds to the limited body of literature on the emergence of severe suicidal ideation in psychiatric patients and single, multiple, and non-attempters. To the best of our knowledge, this is the first study to investigate time to emergence of clinically significant suicidal ideation in adult psychiatric patients, or as a function of previous suicide attempts. Suicidal ideation affects a significant percentage of the population and is an important area of research of its own right and as a risk factor for death by suicide [5]. Understanding the natural course of its emergence, as well as risk and protective factors to time of emergence, will enable us to better understand suicidal ideation and its clinical implications, such as optimal timing of prevention measures.

Acknowledgments

This article was supported by National Institute of Mental Health grants MH44779 and MH58866 to Ivan W. Miller.

References

1. Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. [Accessed Jul 13, 2006] Web-based Injury Statistics Query and Reporting System (WISQARS) [online]. 2005. Available at: www.cdc.gov/ncipc/wisqars
2. Kessler RC, Borges G, Walters EE. Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1999; 56:617–26. [PubMed: 10401507]
3. Weissman MM, Bland RC, Canino GJ, Greenwald S, Hwu HG, Joyce PR, et al. Prevalence of suicide ideation and suicide attempts in nine countries. *Psychol Med*. 1999; 29:9–17. [PubMed: 10077289]
4. Brook R, Klap R, Liao D, Wells KB. Mental health care for adults with suicide ideation. *Gen Hosp Psychiatry*. 2006; 28:271–7. [PubMed: 16814625]
5. Brown GK, Beck AT, Steer RA, Grisham JR. Risk factors for suicide in psychiatric outpatients: a 20-year prospective study. *J Consult Clin Psychol*. 2000; 68:371–7. [PubMed: 10883553]
6. Norlev J, Davidsen M, Kjoller M. Six year follow-up of health status changes in Danish adults with suicide tendency in 1994. *Suicide Life Threat Behav*. 2006; 36:103–12. [PubMed: 16676631]
7. Pfeffer CR, Klerman GL, Hurt SW, Kakuma T, Peskin JR, Siefker CA. Suicidal children grow up: Rates and psychosocial risk factors for suicide attempts during follow-up. *J Am Acad Child Adolesc Psychiatry*. 1993; 31:106–13. [PubMed: 8428862]
8. Food and Drug Administration. [Accessed Oct 16, 2006] Public Health Advisory: Suicidality in children and adolescents being treated with antidepressant medications. 2004. Available at: <http://www.fda.gov/cder/drug/advisory/SSRI200410.htm>
9. Food and Drug Administration. [Accessed Oct 16, 2006] Public Health Advisory: Suicidality in adults being treated with antidepressant medications. 2005. Available at: <http://www.fda.gov/cder/drug/advisory/SSRI200507.htm>
10. Rihmer Z, Akiskal H. Do antidepressants t(h)reat(en) depressives? Toward a clinically judicious formulation of the antidepressant-suicidality FDA advisory in light of declining national suicide statistics from many countries. *J Affect Disord*. 2006; 94:3–13. [PubMed: 16712945]
11. Vasa RA, Carlino AR, Pine DS. Pharmacotherapy of depressed children and adolescents: current issues and potential directions. *Biol Psychiatry*. 2006; 59:1021–8. [PubMed: 16406250]

12. Pfeffer CR, Hurt SW, Kakuma T, Peskin JR, Siefker CA, Nagabhairava S. Suicidal children grow up: Suicidal episodes and effects of treatment during follow-up. *J Am Acad Child Adolesc Psychiatry*. 1994; 33:225–30. [PubMed: 8150794]
13. Esposito C, Spirito A, Boergers J, Donaldson D. Affective, behavioral, and cognitive functioning in adolescents with multiple suicide attempts. *Suicide Life Threat Behav*. 2003; 33:389–99. [PubMed: 14695054]
14. Forman EM, Berk MS, Henriques GR, Brown GK, Beck AT. History of multiple suicide attempts as a behavioral marker of severe psychopathology. *Am J Psychiatry*. 2004; 161:437–43. [PubMed: 14992968]
15. Gispert M, Davis MS, Marsh L, Wheeler K. Predictive factors in repeated suicide attempts by adolescents. *Hosp Community Psychiatry*. 1987; 38:390–3. [PubMed: 3570186]
16. Goldston DB, Daniel S, Reboussin DM, Kelley A, Ievers C, Brunstetter R. First-time suicide attempters, repeat attempters, and previous attempters on an adolescent inpatient psychiatry unit. *J Am Acad Child Adolesc Psychiatry*. 1996; 35:631–9. [PubMed: 8935210]
17. Goldston DB, Daniel SS, Reboussin DM, Reboussin BA, Frazier PH, Kelley AE. Suicide attempts among formerly hospitalized adolescents: A prospective naturalistic study of risk during the first 5 years after discharge. *J Am Acad Child Adolesc Psychiatry*. 1999; 38:660–71. [PubMed: 10361783]
18. Walker RL, Joiner TE Jr, Rudd MD. The course of post-crisis suicidal symptoms: how and for whom is suicide “cathartic”? *Suicide Life Threat Behav*. 2001; 31:144–52. [PubMed: 11459247]
19. Joiner TE Jr, Rudd MD, Rajab MH. The Modified Scale for Suicidal Ideation: Factors of suicidality and their relation to clinical and diagnostic variables. *J Abnorm Psychol*. 1997; 106:260–5. [PubMed: 9131846]
20. Konick LC, Gutierrez PM. Testing a model of suicide ideation in college students. *Suicide Life Threat Behav*. 2005; 35:181–92. [PubMed: 15843335]
21. Pinto A, Whisman MA. Negative affect and cognitive biases in suicidal and nonsuicidal hospitalized adolescents. *J Am Acad Child Adolesc Psychiatry*. 1996; 35:158–65. [PubMed: 8720625]
22. Smith JM, Alloy LB, Abramson LY. Cognitive vulnerability to depression, rumination, hopelessness, and suicidal ideation: multiple pathways to self-injurious thinking. *Suicide Life Threat Behav*. 2006; 36:443–54. [PubMed: 16978098]
23. Garrison CZ, Addy CL, Jackson KL, McKeown RE, Waller JL. A longitudinal study of suicidal ideation in young adolescents. *J Am Acad Child Adolesc Psychiatry*. 1991; 30:597–603. [PubMed: 1890093]
24. Miller IW, Keitner GI, Ryan CE, Solomon DA, Cardemil EV, Beevers CG. Treatment matching in the posthospital care of depressed patients. *Am J Psychiatry*. 2005; 162:2131–8. [PubMed: 16263854]
25. First, M.; Spitzer, R.; Williams, J.; Gibbons, M. Structured Clinical Interview for DSM-IV--Patient version. New York: Biometrics Research Department, New York State Psychiatric Institute; 1995.
26. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry*. 1961; 4:561–71. [PubMed: 13688369]
27. Miller IW, Bishop S, Norman WH, Maddever H. The Modified Hamilton Rating Scale for Depression: reliability and validity. *Psychiatry Res*. 1985; 14:131–42. [PubMed: 3857653]
28. Miller IW, Norman WH, Bishop SB, Dow MG. The Modified Scale for Suicidal Ideation: reliability and validity. *J Consult Clin Psychol*. 1986; 54:724–5. [PubMed: 3771893]
29. Beck AT, Kovacs M, Weissman A. Assessment of suicidal intention: the Scale for Suicide Ideation. *J Consult Clin Psychol*. 1979; 47:343–52. [PubMed: 469082]
30. Rudd MD, Rajab MH. Use of the modified scale for suicidal ideation with suicide ideators and attempters. *J Clin Psychol*. 1995; 51:632–5. [PubMed: 8801239]
31. Beck AT, Steer RA, Garbin MG. Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clin Psychol Rev*. 1988; 8:77–100.
32. Beck AT, Weissman A, Lester D, Trexler L. The measurement of pessimism: the hopelessness scale. *J Consult Clin Psychol*. 1974; 42:861–5. [PubMed: 4436473]

33. Glanz LM, Haas GL, Sweeney JA. Assessment of hopelessness in suicidal patients. *Clin Psychol Rev.* 1995; 15:49–64.
34. Weissman, A. Unpublished doctoral dissertation. University of Pennsylvania; 1979. Dysfunctional Attitudes Scale: A validation study.
35. Krantz S, Hammen C. Assessment of cognitive bias in depression. *J Abnorm Psychol.* 1979; 88:611–9. [PubMed: 512187]
36. Dobson KS, Breiter HJ. Cognitive assessment of depression: Reliability and validity of three measures. *J Abnorm Psychol.* 1983; 92:107–9. [PubMed: 6833629]
37. Frost RO, MacInnis DJ. The Cognitive Bias Questionnaire: Further evidence. *J Pers Assess.* 1983; 47:173–7. [PubMed: 16367602]
38. Luke DA, Homan SM. Time and change: Using survival analysis in clinical assessment and treatment evaluation. *Psychol Assessment.* 1998; 10:360–78.
39. Kleinbaum, DG. Survival analysis: A self-learning text. New York: Springer; 1996.
40. Kessler RC, Berglund P, Borges G, Nock M, Wang PS. Trends in suicide ideation, plans, gestures, and attempts in the United States, 1990–1992 to 2001–2003. *JAMA.* 2005; 293:2487–95. [PubMed: 15914749]

Figure 1.
Cox Proportional Hazard Regression Survival Curve

Table 1
Group Demographics (No Attempt, Single Attempt, Multiple Attempt) on Relevant Variables

	No Attempt (<i>n</i> = 70)	Single Attempt (<i>n</i> = 20)	Multiple Attempt (<i>n</i> = 27)	<i>df</i>	χ^2/F
Sex (% women)	68.6	85.0	77.8	2	2.49
Ethnicity (% Caucasian)	94.3	95.0	92.6	2	0.14
Age (years)	39.43 (12.24)	35.40 (10.60)	36.89 (11.58)	2, 116	1.10
Marital Status (% Married)	68.6	76.2	60.0	2	1.53
Admission MSSI	14.19 (12.19) ^a	20.90 (11.90) ^b	29.56 (13.38) ^c	2, 115	15.15 ^{***}
Discharge MSSI	0.99 (3.03) ^a	3.10 (4.11) ^b	2.92 (5.04) ^b	2, 113	3.83 [*]
Admission BDI	32.24 (7.78) ^a	32.85 (9.32) ^{a,b}	36.78 (8.23) ^b	2, 116	3.06 [*]
Admission MHRSD	36.56 (6.10) ^a	36.50 (6.48) ^{a,b}	39.89 (6.15) ^b	2, 116	3.05 [*]
Admission BHS	14.49 (5.01)	15.50 (5.11)	16.15 (3.84)	2, 116	1.29
Admission DAS	154.50 (42.37)	162.15 (38.74)	164.30 (45.22)	2, 116	0.63
Admission CBQ	2.94 (2.98)	3.75 (3.13)	4.19 (3.63)	2, 116	1.68
Treatment Group				6	7.37

Notes. Unless otherwise specified, values in cells represent means, and values in parentheses represent standard deviations. Chi-square tests were used for tests involving sex and ethnicity; all other analyses were conducted using analyses of variance.

MSSI = Modified Scale for Suicidal Ideation. BDI = Beck Depression Inventory. MHRSD = Modified Hamilton Rating Scale for Depression. BHS = Beck Hopelessness Scale. DAS = Dysfunctional Attitudes Scale. CBQ = Cognitive Bias Questionnaire.

Means with different superscripts differ significantly at $p < .05$.

* $p < .05$,

*** $p < .001$.

Table 2

Group Demographics (Reemergence of Severe Suicidal Ideation) on Relevant Variables

	No Reemergence (<i>n</i> = 91)	Reemergence (<i>n</i> = 26)	<i>df</i>	χ^2/F
Sex (% women)	73.6	73.1	1	0.003
Ethnicity (% Caucasian)	93.4	96.2	1	0.27
Age (years)	38.44 (11.71)	37.15 (12.52)	115	0.49
Marital Status (% Married)	67.0	73.1	1	0.34
Admission MSSI	16.36 (13.33)	27.81 (12.11)	114	3.94****
Discharge MSSI	1.31 (3.08)	3.46 (5.71)	112	2.53**
Admission BDI	32.91 (8.15)	35.08 (8.77)	115	1.17
Admission MHRSD	36.77 (6.16)	39.23 (6.45)	115	1.78
Admission BHS	14.73 (4.93)	16.15 (4.20)	115	1.34
Admission DAS	153.57 (40.44)	173.81 (45.75)	115	2.19*
Admission CBQ	3.07 (2.97)	4.42 (3.71)	115	1.94
Treatment Group			3	5.28

Notes. Unless otherwise specified, values in cells represent means, and values in parentheses represent standard deviations. Chi-square tests were used for tests involving sex and ethnicity; all other analyses were conducted using analyses of variance.

MSSI = Modified Scale for Suicidal Ideation. BDI = Beck Depression Inventory. MHRSD = Modified Hamilton Rating Scale for Depression. BHS = Beck Hopelessness Scale. DAS = Dysfunctional Attitudes Scale. CBQ = Cognitive Bias Questionnaire.

*
 $p < .05$,

**
 $p < .01$,

 $p < .005$,

 $p < .001$.

Table 3

Cumulative Percent Reporting the Occurrence of Severe Suicidal Ideation

	Non-Attempters	Single Attempters	Multiple Attempters
Discharge	0%	0%	0%
3 months	1%	0%	8%
6 months	3%	22%	24%
9 months	6%	34%	41%
12 months	6%	40%	41%
15 months	9%	40%	46%
18 months	9%	40%	46%

Table 4

Comparison of Survival Times by Attempt Status

Group	No. of Events	Mean survival time (months)	95% CI
Non-Attempter	7	17.06	16.23, 17.90
Single Attempter	8	12.79	9.51, 16.07
Multiple Attempter	11	11.98	9.25, 14.72
All Participants	26	15.26	14.23, 16.29

Note. CI = confidence interval. For the log rank significance test, the value was 19.51 ($df = 2$), $p < .001$; for Breslow, the value was 20.30 ($df = 2$), $p < .001$; for Tarone-Ware, the value was 20.02 ($df = 2$), $p < .001$.