Published in final edited form as:

Nat Rev Rheumatol. 2010 January ; 6(1): 13–20. doi:10.1038/nrrheum.2009.240.

SLE nephritis – learning from murine models

Anne Davidson and Cynthia Aranow

Center for Autoimmune and Musculoskeletal Diseases, Feinstein Institute for Medical Research,
Manhasset NY 11030

Abstract

SLE nephritis is a challenging clinical condition for which current therapies are unsatisfactory with respect to both remission induction and unwanted toxicities. Despite intervention the rates of end stage renal disease appear to be increasing in the US. New discoveries over the last decade have greatly improved our understanding of immune activation and effector inflammatory pathways in SLE nephritis but these have not yet translated into an effective new approved therapeutic. An analysis of the mechanisms of new immunomodulatory drugs in multiple models of murine SLE shows clearly that interacting networks of immune and effector pathways are recruited as disease progresses. It is therefore difficult to reverse established disease by targeting a single cell population or inflammatory pathway once long-lived autoreactive lymphocyte populations are present and peripheral organs are inflamed. These data suggest that we need to consider new paradigms for the management of SLE that include earlier immune intervention, long-term maintenance therapies and protection of target organs.

Introduction

Nephritis is a challenging problem affecting between 30–60% of SLE patients^{1, 2}. Outcomes after broad spectrum immunosuppression remain unsatisfactory with complete remission rates of 50% at best³, relapse rates of up to 30% over a two year period⁴ and unacceptable toxicities. Despite the implementation of maintenance regimens^{5–7} the incidence of end stage renal disease (ESRD) in SLE patients is increasing^{8, 9} especially in non-Caucasians. Barriers to performing informative clinical trials of new drugs for SLE nephritis remain formidable. Disease heterogeneity, the confounding effects of other immunosuppressive medications and concurrent medical problems contribute to these difficulties.

The last decade has witnessed an exciting increase in our understanding of the immunopathogenesis of autoimmunity. Despite many new discoveries, the path to a new approved therapy for SLE is strewn with a growing list of failures, even of drugs with efficacy in other diseases. Because some disease mechanisms are shared between humans and mice, the study of diverse mouse models has enabled acquisition of pre-clinical data that support clinical trials of novel therapies (Table 1)¹⁰. The major disadvantages of relying on murine models are differences in physiology between mouse and man and the extensive genetic heterogeneity in an outbred human population that complicates predictions of

treatment efficacy. Nevertheless, mechanistic studies of therapeutic interventions in murine SLE may help explain the difficulties in identifying new drugs for human SLE.

Pathogenesis of SLE nephritis

SLE nephritis is initiated by the glomerular deposition of immune complexes (IC) that trigger a cascade of inflammatory events including activation of Fc receptors¹¹ and complement¹² recruitment of inflammatory cells and eventual fibrosis. IC also activate resident renal cells through Toll-like receptors (TLRs) to produce inflammatory mediators¹³. Secondary lymphoid tissue has been reported in the renal parenchyma in some patients^{14, 15}. Interstitial nephritis occurs in one murine model in the complete absence of circulating immunoglobulins¹⁶; similarly, pauci-immune nephritis has been observed in humans¹⁷. Renal microvascular damage and thromboses also occur, especially in patients with anti-phospholipid antibodies¹⁸.

Several renal cell types including endothelial cells, podocytes, interstitial cells and renal dendritic cells are the focus of new studies in nephritis (Figure 1)¹⁹. Renal migration of inflammatory cells requires endothelial cell activation²⁰. Hypoxia, due to loss of glomerular and peritubular capillaries contributes to cell stress and cell death and induces molecules that further activate innate immune receptors²¹. There is currently an increased interest in molecules that protect the endothelium from hypoxia, such as oxygen sensors that regulate endothelial leak and microthrombus formation²² and receptors that regulate endothelial adhesiveness. Patients with active lupus have increased levels of angiopoietin-2, an antagonist of Tie2, a receptor that maintains endothelial integrity and prevents leukocyte recruitment to the kidney; this may be a more specific marker of renal involvement than soluble VCAM-1²³. Signals delivered via the vagus nerve to nicotinic acetylcholine receptors (nAChRs) may exert a renoprotective effect by decreasing inflammatory cytokines and preserving endothelial integrity²⁴. All these pathways are potential therapeutic targets for SLE nephritis. In addition, circulating endothelial cells²⁵ and endothelial protein C receptor^{26, 27} may be biomarkers of endothelial dysfunction and damage.

The role of podocytes in renal inflammation is increasingly recognized²⁸ and their loss in chronic disease causes glomerulosclerosis. Podocytes form a critical barrier to proteinuria and protect the endothelium by producing both glomerular basement membrane and VEGF²⁹. Activation of angiotensin receptors on podocytes induces the release of inflammatory mediators and cell death³⁰; blockade of angiotensin receptors protects podocytes and may be beneficial for SLE nephritis³¹.

Interstitial cell infiltration, tubular atrophy that follows vascular compromise, activation of smooth muscle myofibroblast cells and interstitial fibrosis are predictors of poorer prognosis and/or progression to ESRD in SLE nephritis^{32, 33}. Activation of intrinsic renal dendritic cells occurs in both murine and human SLE^{20, 34} and is associated with expression of inflammatory cytokines, proteolytic activity, increased caspase 1 activity^{20, 35} and upregulation of cell surface molecules including CD11b²⁰. Genetic differences involving both immune and non-immune genes may influence the long term outcome of renal inflammation although there are as yet no rigorous studies in SLE nephritis; these studies are

challenging to perform even in the setting of renal transplant in which many more patients are available³⁹. Recent identification of polymorphisms of genes that enhance or protect from SLE nephritis such as ITGAM (the α chain of CD11b)³⁶, IL-18³⁷ and kallikreins³⁸ suggest new avenues for investigation. Further studies of the mechanisms involved in intrinsic renal cell activation, interstitial renal inflammation and recruitment of fibrosis pathways should lead to new strategies for renal protection from the consequences of antibody deposition.

New immunologic therapies for SLE

Although the trigger initiating SLE in a genetically susceptible host is not known, initiation of murine lupus can occur in a variety of ways. As disease progresses interactions between innate and acquired immune activation pathways create amplification loops and interacting networks that become more difficult to regulate (Figure 2).

B cell directed therapies

B cells are culprits in autoimmunity not only as producers of pathogenic autoantibodies but because they have diverse effector functions including antigen uptake and transport⁴⁰, antigen presentation to T cells^{41, 42}, production of cytokines⁴³ and chemokines⁴⁴, and migration to sites of inflammation where they participate in further recruitment of inflammatory cells^{45, 46}. In mice, B cell deficiency prevents SLE completely, whereas the complete absence of soluble antibodies only results in attenuated disease¹⁶. In humans, active SLE is associated with high numbers of circulating autoreactive plasmablasts/plasma cells, memory B cell expansion and a dearth of naïve B cells. The plasma cells seen in active SLE are most likely newly generated as they are responsive to cyclophosphamide, (indicating that they are still dividing)⁴⁷ or anti-CD40L (that dissolves germinal centers)⁴⁸.

Although B-cell depletion seems an intuitive means to induce lupus remission, two recent trials of rituximab, an anti-CD20 antibody that depletes all B cells except plasma cells, have failed. Although anti-CD20 depletes memory B cells and expands the transitional ones in some patients, others relapse with expanding memory cells⁴⁹. Anecdotal reports have suggested that combination cyclophosphamide-corticosteroids-anti-CD20, induces remission of SLE, including nephritis. However, in a trial in which nephritis patients received background mycophenolic acid and steroids with or without two courses of rituximab, no benefit of rituxan was observed at 52 weeks. Moreover, an increase in BAFF occurring during the reconstitution phase⁵⁰ might expand naïve autoreactive or memory B-cells. Anti-CD20 is effective for both rheumatoid arthritis and multiple sclerosis each of which is associated with effector B cells in target organs but with less systemic inflammation than SLE. Murine studies have indicated that secondary lymphoid tissues in SLE-prone mice are resistant to B cell depletion even with depletion in the peripheral blood⁵¹. It is still debatable whether more aggressive therapy with anti-CD20, concomitant cyclophosphamide, or BAFF blockade during reconstitution will be beneficial in human SLE.

Modulation of B cell selection and survival can also be achieved by inhibition of BAFF and its homolog APRIL. BAFF blockade does not prevent formation of germinal centers⁵² but results in partial B cell depletion and more stringent selection of the naïve autoreactive

repertoire⁵³. Although memory B-cells are BAFF-independent in normal mice⁵⁴, BAFF boosts reactivation of memory B cells when inflammatory cytokines are present⁵⁵. Depletion of plasma cells requires blockade of both BAFF and APRIL; IgM-producing plasma cells are more sensitive to this blockade than IgG-producing ones⁵⁶, probably because plasma cells in bone marrow niches receive alternative survival signals, (especially during inflammation). Intriguingly, activated macrophages and DCs express BAFF receptors; in mice BAFF blockade prevents activation of DCs in inflammatory sites resulting in local decreases of IL-6 release, maturation of T_H17 cells and plasma cell survival⁵⁷.

BAFF blockade attenuates murine lupus especially when administered before disease onset but with considerable heterogeneity between models⁵⁸; furthermore, in most models, blockade of BAFF alone is as effective as blockade of both BAFF and APRIL, indicating that plasma cell depletion is not required for this beneficial outcome. In some strains the combination of BAFF blockade and CTLA4Ig induces remission of nephritis more effectively than either drug alone. Complete absence of BAFF does not prevent eventual glomerular damage and proteinuria in a spontaneous SLE model but does prevent renal failure and death⁵⁹, perhaps because of a decreased inflammatory load. This suggests that careful analysis of short and long term outcomes needs to be performed in clinical trials using B-cell depletion for SLE nephritis.

Belimumab, a selective inhibitor of BAFF has been evaluated in a Phase II study of generalized SLE. While the trial failed to meet its primary endpoint, post-hoc analyses using a composite disease activity measure showed clinical efficacy in a subset of patients; preliminary results from a phase III study in non-renal lupus show that patients receiving belimumab plus standard of care have an improved clinical response, including prevention of flare, compared with patients receiving placebo over 52 weeks. Patients receiving belimumab depleted their peripheral naïve and transitional B cells within 3–6 months with minimal effects on memory B cells and plasmablasts; autoantibody levels were only modestly affected but a small subset of patients became seronegative over a long time period⁶⁰. Whether there is an effect on B cell selection over time or whether chronic B cell depletion will induce homeostatic expansion of autoreactive memory B cells remain to be determined. These findings confirm the mechanisms identified in murine studies; explanations for the salutary effects of BAFF blockade include the secondary effect of B cell depletion on the expansion of other inflammatory cell populations and/or the inhibitory effect of BAFF blockade on immature DCs. While administration of TACI-Ig, an agent blocking both BAFF and APRIL might lead to a more consistent depletion of plasma cells, preferential depletion of IgM-secreting plasma cells that produce protective antibodies^{61, 62} may result in unacceptable toxicity.

T cell directed therapies

CTLA4Ig, an antagonist of B7-CD28 costimulation is beneficial in murine SLE, especially when given before class switched and somatically mutated autoantibodies arise^{63, 64}. The follicular T helper cells (T_{FH}) that drive B cell help in germinal centers are completely dependent on CD28 costimulation⁶⁵. However TLR mediated signals together with

combinations of cytokines including BAFF, IL-21 and IL-17 can substitute for B7-CD28 costimulation and drive reactivation of memory B cells^{55, 66}. Thus administration of CTLA4Ig may be futile in active lupus or chronic disease with persistent memory T cells and long-lived B cells. This might account for the failure of monthly abatacept (CTLA4Ig), given with tapering doses of steroids, to alter autoantibody titers or rates of flare at one year in patients with non-renal flares. These studies confirm the challenges of using CTLA4Ig alone in established disease. A study of abatacept in lupus nephritis is in progress. Remission of active SLE nephritis can be achieved in mice using a combination of cyclophosphamide and CTLA4Ig. Activated T and B cells are depleted by this regimen, with reversal of the activation phenotype of intrinsic renal cells^{20, 64}. A clinical trial of this combination in humans is also currently in progress.

Follicular T helper cells (T_{FH}) might be targeted with IL-17, IL-21, CD40 and ICOS antagonists, some of which are in early development in human clinical trials. While IL-21 is required for disease onset in some murine models⁶⁷ it is dispensable for others, even when T_{FH} are involved⁶⁸ and its inhibition has only modest effects on late disease. Anti-CD40L is effective at preventing murine SLE but has less value for established nephritis⁶⁹. Thus, while T_{FH} are required for disease initiation either they are not sufficient for disease perpetuation or their inhibition in late disease requires blocking more than one molecule. It remains to be determined whether targeting of T_{FH} is a viable therapeutic option.

The role of regulatory T cells (T_{reg}) in SLE nephritis is still controversial. Initial studies in spontaneous murine models have shown expansion of regulatory T cell subsets in the periphery and in the kidney over time⁷⁰ and adoptive transfer of large numbers of CD4+/CD25+ T_{reg} in the NZB/W model had only a very modest effect on disease onset and survival⁷¹. This may be because T_{reg} can be inactivated or convert to TH_{17} cells in an inflammatory environment⁷². In contrast, induction of T_{reg} by non-depleting anti-CD3 antibodies had very robust effects in the SNF_1 model; in this case suppression was mediated by inducible CD4+/CD25-/LAP+/Foxp3- T_{reg} ⁷³. Intriguing new data suggests that partial inhibition of the mTOR pathway may induce T_{reg} ⁷⁴ pointing to an additional mechanism for the efficacy of mTOR inhibitors in murine SLE models. Clearly much remains to be learned about the potential therapeutic role of different types of T_{reg} in SLE.

Therapies directed at the innate immune system

Many SLE patients manifest a Type I interferon (IFN) signature in the peripheral blood, induced by exposure of PBMCs to IFN α and/or immune complexes⁷⁵. Type I IFNs can induce either anti-DNA or anti-Sm/RNP antibodies and accelerate disease in murine models by activating antigen presenting cells, enhancing expression of TLRs and BAFF receptors, enhancing germinal center formation and inducing release of BAFF and IL-6^{75, 76}. IFN α inhibitors can inhibit IFN α accelerated disease⁷⁷ but it is still unknown if IFNs drive disease after the onset of inflammation. Clinical trials of anti-IFN α antibodies in SLE are underway. In mice, Type I IFNs confer resistance to both CTLA4Ig and BAFF inhibitors (unpublished data). This has two important implications. First high serum levels of BAFF or of T cell derived cytokines do not necessarily indicate sensitivity to inhibition of those cytokines.

Second it may be necessary to randomize patients based on their interferon signature in order to analyze the effects of other immune modulators.

While SLE has classically been considered a T cell mediated disease, initiation of SLE in BAFF transgenic mice depends only on signaling through the TLR adaptor molecule MyD88⁷⁸. Intracellular TLRs can be targeted using short synthetic oligodeoxynucleotides (ODN). ODN are more effective for disease prevention than for inducing remission in mice and inhibitors of TLR7 are as effective as inhibitors of both TLR7 and TLR9⁷⁹. The precise role for specific TLR inhibitors will need to be addressed in human clinical trials. Selective TLR9 inhibition may not be an appropriate strategy because it may increase IFN production by immature dendritic cells and thereby induce B cells to secrete pathogenic anti-RNA antibodies⁸⁰. Moreover, humans express a functional TLR8 in monocytes and myeloid DCs⁸¹ and it is unclear whether this is blocked by currently available ODN. Nevertheless, the disease modifying effects of hydroxychloroquine, an inhibitor of autophagosome formation (in which TLRs encounter nucleic acid antigens)⁸², suggest that such reagents may be beneficial.

Conclusions

Despite advances in our understanding of lupus pathogenesis we have not yet identified an effective new therapy for SLE nephritis. A number of clear themes are emerging from the murine studies that bear upon this difficulty. First, mouse models are heterogeneous in their responses to therapies. In particular, mice with excessive activation of the innate immune system are resistant to T cell directed therapies. Second, depletion of circulating autoantibodies does not always translate into an immediate clinical effect. Conversely, if antibody effector functions are blunted it is not essential to remove autoantibodies in order to improve nephritis outcomes. Third, therapies that deplete naïve B cells do not have robust efficacy for active disease and could lead to expansion of autoreactive B cells in the face of generalized immunosuppression. Fourth, while many therapies prevent disease onset, most have significantly compromised efficacy when used at later stages. For established disease, combinations of therapies are more effective than single therapy alone but are considerably more immunosuppressive. Because the inflammatory milieu of chronic SLE profoundly alters T and B cell function and sets up dysregulated amplification loops, restoration of normal tolerance is difficult to achieve once disease has begun. Some changes may be persistent since they are associated with the generation of populations of long-lived effector cells and with alterations in DNA methylation⁸³. Finally, genetic polymorphisms that alter the susceptibility of the kidneys to inflammation and fibrosis will affect prognosis. The data from the murine models suggest that we need to reconsider our current paradigms for the management of SLE. At present we reserve immunologic interventions for patients with active disease and withdraw medications in inactive disease. The most logical strategy is to prevent activation of autoreactive cells before class switching and somatic mutation of immunoglobulin genes, formation of long-lived populations and establishment of multiple dysregulated inflammatory pathways have occurred. Approaches for the treatment of newly diagnosed SLE after remission of the first flare could include relatively safe forms of long-term immune modulation (Table 1) as well as some of the reagents discussed above. Concurrent medical problems such as hypertension and obesity need to be aggressively

treated at early stages. In addition, genetic and phenotypic profiling of patients may allow identification of subgroups more likely to respond to particular therapies. Once inflammation is present, intensive immunotherapy can deplete activated B and T cells but strategies need to be developed to avoid reconstitution of an autoreactive repertoire and/or homeostatic proliferation of long-lived populations. Therapies directed at effector pathways need to be developed and used early to protect target organs from damage. Both short and long term outcomes need to be examined in clinical trials. The recent success of belimumab, an agent directed at both autoreactive B cells and target organs, exemplifies the principle that measuring the results of moderate immunomodulation for the maintenance of long term remission requires large and prolonged clinical trials of reagents that target more than one immune pathway.

References

- References were selected using Medline search and the terms lupus and/or nephritis with therapies, inflammation, endothelial cells, podocytes, biomarkers, macrophages, dendritic cells, T cells, B cells, Rituximab, CTLA4Ig, interferon, TLR, BAFF, B cell depletion, cytokines. Articles published between 2006 and 2009 were given preference for inclusion. In addition I used a personal collection that includes approximately 3000 references related to lupus and novel therapies.
1. Sidiropoulos PI, Kritikos HD, Boumpas DT. Lupus nephritis flares. *Lupus*. 2005; 14:49–52. [PubMed: 15732288]
 2. Mok CC, et al. Predictors and outcome of renal flares after successful cyclophosphamide treatment for diffuse proliferative lupus glomerulonephritis. *Arthritis Rheum*. 2004; 50:2559–68. [PubMed: 15334470]
 3. Appel GB, et al. Mycophenolate mofetil versus cyclophosphamide for induction treatment of lupus nephritis. *J Am Soc Nephrol*. 2009; 20:1103–12. [PubMed: 19369404]
 4. Mok CC. Therapeutic options for resistant lupus nephritis. *Semin Arthritis Rheum*. 2006; 36:71–81. [PubMed: 16884971]
 5. Chan TM. Preventing renal failure in patients with severe lupus nephritis. *Kidney Int Suppl*. 2005:S116–9. [PubMed: 15752227]
 6. Contreras G, et al. Sequential therapies for proliferative lupus nephritis. *N Engl J Med*. 2004; 350:971–80. [PubMed: 14999109]
 7. Contreras G, Tozman E, Nahar N, Metz D. Maintenance therapies for proliferative lupus nephritis: mycophenolate mofetil, azathioprine and intravenous cyclophosphamide. *Lupus*. 2005; 14 (Suppl 1):s33–8. [PubMed: 15803929]
 8. Dooley MA, Hogan S, Jennette C, Falk R. Cyclophosphamide therapy for lupus nephritis: poor renal survival in black Americans. Glomerular Disease Collaborative Network. *Kidney Int*. 1997; 51:1188–95. [PubMed: 9083285]
 9. Costenbader KH, Solomon DH, Winkelmayer W, Brookhart MA. Incidence of End-Stage Renal Disease Due to Lupus Nephritis in the U.S., 1995–2004. *Arthritis and Rheumatism*. 2008:Abstract 1927.
 10. Wiesendanger M, Stanevsky A, Kovsky S, Diamond B. Novel therapeutics for systemic lupus erythematosus. *Curr Opin Rheumatol*. 2006; 18:227–35. [PubMed: 16582684]
 11. Clynes R, Dumitru C, Ravetch JV. Uncoupling of immune complex formation and kidney damage in autoimmune glomerulonephritis. *Science*. 1998; 279:1052–4. [PubMed: 9461440]
 12. Turnberg D, Cook HT. Complement and glomerulonephritis: new insights. *Curr Opin Nephrol Hypertens*. 2005; 14:223–8. [PubMed: 15821414]
 13. Anders HJ, Schlondorff D. Toll-like receptors: emerging concepts in kidney disease. *Curr Opin Nephrol Hypertens*. 2007; 16:177–83. [PubMed: 17420659]

14. Segerer S, Schlondorff D. B cells and tertiary lymphoid organs in renal inflammation. *Kidney Int.* 2008; 73:533–7. [PubMed: 18094677]
15. Jeruc J, et al. Tubulo-interstitial involvement in lupus nephritis with emphasis on pathogenesis. *Wien Klin Wochenschr.* 2000; 112:702–6. [PubMed: 11020960]
16. Chan OT, Hannum LG, Haberman AM, Madaio MP, Shlomchik MJ. A novel mouse with B cells but lacking serum antibody reveals an antibody-independent role for B cells in murine lupus. *J Exp Med.* 1999; 189:1639–48. [PubMed: 10330443]
17. Schwartz MM. The pathology of lupus nephritis. *Semin Nephrol.* 2007; 27:22–34. [PubMed: 17336686]
18. Lewis EJ, Schwartz MM. Pathology of lupus nephritis. *Lupus.* 2005; 14:31–8. [PubMed: 15732285]
19. Schlondorff DO. Overview of factors contributing to the pathophysiology of progressive renal disease. *Kidney Int.* 2008; 74:860–6. [PubMed: 18650795]
20. Schiffer L, et al. Activated renal macrophages are markers of disease onset and disease remission in lupus nephritis. *J Immunol.* 2008; 180:1938–47. [PubMed: 18209092]
21. Ferraccioli G, Romano G. Renal interstitial cells, proteinuria and progression of lupus nephritis: new frontiers for old factors. *Lupus.* 2008; 17:533–40. [PubMed: 18539706]
22. Mazzone M, et al. Heterozygous deficiency of PHD2 restores tumor oxygenation and inhibits metastasis via endothelial normalization. *Cell.* 2009; 136:839–51. [PubMed: 19217150]
23. Kumpers P, et al. The Tie2 receptor antagonist Angiopoietin-2 facilitates vascular inflammation in Systemic Lupus Erythematosus. *Ann Rheum Dis.* 2008
24. Yeboah MM, et al. Cholinergic agonists attenuate renal ischemia-reperfusion injury in rats. *Kidney Int.* 2008; 74:62–9. [PubMed: 18401335]
25. Yao G, et al. Evaluation of renal vascular lesions using circulating endothelial cells in patients with lupus nephritis. *Rheumatology (Oxford).* 2008; 47:432–6. [PubMed: 18292121]
26. Izmirly PM, et al. Expression of endothelial protein C receptor in cortical peritubular capillaries associates with a poor clinical response in lupus nephritis. *Rheumatology (Oxford).* 2009; 48:513–9. [PubMed: 19286697]
27. Sesin CA, Yin X, Esmon CT, Buyon JP, Clancy RM. Shedding of endothelial protein C receptor contributes to vasculopathy and renal injury in lupus: in vivo and in vitro evidence. *Kidney Int.* 2005; 68:110–20. [PubMed: 15954900]
28. Shankland SJ. The podocyte's response to injury: role in proteinuria and glomerulosclerosis. *Kidney Int.* 2006; 69:2131–47. [PubMed: 16688120]
29. Michaud JL, Kennedy CR. The podocyte in health and disease: insights from the mouse. *Clin Sci (Lond).* 2007; 112:325–35. [PubMed: 17291194]
30. Crowley SD, et al. Glomerular type 1 angiotensin receptors augment kidney injury and inflammation in murine autoimmune nephritis. *J Clin Invest.* 2009; 119:943–53. [PubMed: 19287096]
31. Duran-Barragan S, McGwin G Jr, Vila LM, Reveille JD, Alarcon GS. Angiotensin-converting enzyme inhibitors delay the occurrence of renal involvement and are associated with a decreased risk of disease activity in patients with systemic lupus erythematosus--results from LUMINA (LIX): a multiethnic US cohort. *Rheumatology (Oxford).* 2008; 47:1093–6. [PubMed: 18511474]
32. Faurschou M, Starklint H, Halberg P, Jacobsen S. Prognostic factors in lupus nephritis: diagnostic and therapeutic delay increases the risk of terminal renal failure. *J Rheumatol.* 2006; 33:1563–9. [PubMed: 16881113]
33. Mok CC. Prognostic factors in lupus nephritis. *Lupus.* 2005; 14:39–44. [PubMed: 15732286]
34. Segerer S, et al. Compartment specific expression of dendritic cell markers in human glomerulonephritis. *Kidney Int.* 2008; 74:37–46. [PubMed: 18368027]
35. Tucci M, et al. Glomerular accumulation of plasmacytoid dendritic cells in active lupus nephritis: role of interleukin-18. *Arthritis Rheum.* 2008; 58:251–62. [PubMed: 18163476]
36. Hom G, et al. Association of systemic lupus erythematosus with C8orf13-BLK and ITGAM-ITGAX. *N Engl J Med.* 2008; 358:900–9. [PubMed: 18204098]

37. Warchol T, Lianeri M, Wudarski M, Lacki JK, Jagodzinski PP. IL-18 105 A>C polymorphism contributes to renal manifestations in patients with SLE. *Rheumatol Int.* 2009
38. Liu K, et al. Kallikrein genes are associated with lupus and glomerular basement membrane-specific antibody-induced nephritis in mice and humans. *J Clin Invest.* 2009; 119:911–23. [PubMed: 19307730]
39. Kruger B, Schroppel B, Murphy BT. Genetic polymorphisms and the fate of the transplanted organ. *Transplant Rev (Orlando).* 2008; 22:131–40. [PubMed: 18631866]
40. Phan TG, Green JA, Gray EE, Xu Y, Cyster JG. Immune complex relay by subcapsular sinus macrophages and noncognate B cells drives antibody affinity maturation. *Nat Immunol.* 2009; 10:786–93. [PubMed: 19503106]
41. Mamula MJ. Epitope spreading: the role of self peptides and autoantigen processing by B lymphocytes. *Immunol Rev.* 1998; 164:231–9. [PubMed: 9795779]
42. Roosnek E, Lanzavecchia A. Efficient and selective presentation of antigen-antibody complexes by rheumatoid factor B cells. *J Exp Med.* 1991; 173:487–9. [PubMed: 1703209]
43. Lund FE. Cytokine-producing B lymphocytes-key regulators of immunity. *Curr Opin Immunol.* 2008; 20:332–8. [PubMed: 18417336]
44. Ngo VN, Cornall RJ, Cyster JG. Splenic T zone development is B cell dependent. *J Exp Med.* 2001; 194:1649–60. [PubMed: 11733579]
45. Cassese G, et al. Inflamed kidneys of NZB/W mice are a major site for the homeostasis of plasma cells. *Eur J Immunol.* 2001; 31:2726–32. [PubMed: 11536171]
46. Takemura S, et al. Lymphoid neogenesis in rheumatoid synovitis. *J Immunol.* 2001; 167:1072–80. [PubMed: 11441118]
47. Odendahl M, et al. Disturbed peripheral B lymphocyte homeostasis in systemic lupus erythematosus. *J Immunol.* 2000; 165:5970–9. [PubMed: 11067960]
48. Huang W, et al. The effect of anti-CD40 ligand antibody on B cells in human SLE. *Arthritis and Rheumatism.* 2002; 46:1554–62. [PubMed: 12115186]
49. Anolik JH, et al. Rituximab improves peripheral B cell abnormalities in human systemic lupus erythematosus. *Arthritis Rheum.* 2004; 50:3580–90. [PubMed: 15529346]
50. Vallerskog T, et al. Differential effects on BAFF and APRIL levels in rituximab-treated patients with systemic lupus erythematosus and rheumatoid arthritis. *Arthritis Res Ther.* 2006; 8:R167. [PubMed: 17092341]
51. Ahuja A, et al. Depletion of B cells in murine lupus: efficacy and resistance. *J Immunol.* 2007; 179:3351–61. [PubMed: 17709552]
52. Ramanujam M, et al. Mechanism of action of transmembrane activator and calcium modulator ligand interactor-Ig in murine systemic lupus erythematosus. *J Immunol.* 2004; 173:3524–34. [PubMed: 15322217]
53. Thien M, et al. Excess BAFF rescues self-reactive B cells from peripheral deletion and allows them to enter forbidden follicular and marginal zone niches. *Immunity.* 2004; 20:785–98. [PubMed: 15189742]
54. Scholz JL, et al. BLyS inhibition eliminates primary B cells but leaves natural and acquired humoral immunity intact. *Proc Natl Acad Sci U S A.* 2008; 105:15517–22. [PubMed: 18832171]
55. Doreau A, et al. Interleukin 17 acts in synergy with B cell-activating factor to influence B cell biology and the pathophysiology of systemic lupus erythematosus. *Nat Immunol.* 2009
56. Ramanujam M, et al. Similarities and differences between selective and nonselective BAFF blockade in murine SLE. *J Clin Invest.* 2006; 116:724–34. [PubMed: 16485042]
57. Lai Kwan Lam Q, King Hung Ko O, Zheng BJ, Lu L. Local BAFF gene silencing suppresses Th17-cell generation and ameliorates autoimmune arthritis. *Proc Natl Acad Sci U S A.* 2008; 105:14993–8. [PubMed: 18820032]
58. Ramanujam M, Davidson A. BAFF blockade for Systemic Lupus Erythematosus - will the promise be fulfilled? *Immunol Rev.* 2008; 223:156–174. [PubMed: 18613835]
59. Jacob CO, et al. Paucity of clinical disease despite serological autoimmunity and kidney pathology in lupus-prone New Zealand mixed 2328 mice deficient in BAFF. *J Immunol.* 2006; 177:2671–80. [PubMed: 16888029]

60. Wallace DJ, et al. A phase II, randomized, double-blind, placebo-controlled, dose-ranging study of belimumab in patients with active systemic lupus erythematosus. *Arthritis Care and Research*. 2009 In press.
61. Chen Y, et al. Regulation of dendritic cells and macrophages by an anti-apoptotic cell natural antibody that suppresses TLR responses and inhibits inflammatory arthritis. *J Immunol*. 2009; 183:1346–59. [PubMed: 19564341]
62. Manson JJ, Mauri C, Ehrenstein MR. Natural serum IgM maintains immunological homeostasis and prevents autoimmunity. *Springer Semin Immunopathol*. 2005; 26:425–32. [PubMed: 15611856]
63. Mihara M, et al. CTLA4Ig inhibits T cell-dependent B-cell maturation in murine systemic lupus erythematosus. *J Clin Invest*. 2000; 106:91–101. [PubMed: 10880052]
64. Schiffer L, et al. Short term administration of costimulatory blockade and cyclophosphamide induces remission of systemic lupus erythematosus nephritis in NZB/W F1 mice by a mechanism downstream of renal immune complex deposition. *J Immunol*. 2003; 171:489–97. [PubMed: 12817034]
65. Linterman MA, et al. Roquin differentiates the specialized functions of duplicated T cell costimulatory receptor genes CD28 and ICOS. *Immunity*. 2009; 30:228–41. [PubMed: 19217324]
66. Ettinger R, et al. IL-21 and BAFF/BLyS synergize in stimulating plasma cell differentiation from a unique population of human splenic memory B cells. *J Immunol*. 2007; 178:2872–82. [PubMed: 17312131]
67. Bubier JA, et al. A critical role for IL-21 receptor signaling in the pathogenesis of systemic lupus erythematosus in BXSB-Yaa mice. *Proc Natl Acad Sci U S A*. 2009; 106:1518–23. [PubMed: 19164519]
68. Linterman MA, et al. Follicular helper T cells are required for systemic autoimmunity. *J Exp Med*. 2009; 206:561–76. [PubMed: 19221396]
69. Kallal SL, Cutler AH, Datta SK, Thomas DW. Anti-CD40 ligand antibody treatment of SNF1 mice with established nephritis: preservation of kidney function. *J Immunol*. 1998; 160:2158–65. [PubMed: 9498753]
70. Forestier C, et al. Expansion and hyperactivity of CD1d-restricted NKT cells during the progression of systemic lupus erythematosus in (New Zealand Black x New Zealand White)F1 mice. *J Immunol*. 2005; 175:763–70. [PubMed: 16002672]
71. Scalapino KJ, Daikh DI. Suppression of glomerulonephritis in NZB/NZW lupus prone mice by adoptive transfer of ex vivo expanded regulatory T cells. *PLoS One*. 2009; 4:e6031. [PubMed: 19551149]
72. Beriou G, et al. IL-17-producing human peripheral regulatory T cells retain suppressive function. *Blood*. 2009; 113:4240–9. [PubMed: 19171879]
73. Wu HY, Center EM, Tsokos GC, Weiner HL. Suppression of murine SLE by oral anti-CD3: inducible CD4+CD25-LAP+ regulatory T cells control the expansion of IL-17+ follicular helper T cells. *Lupus*. 2009; 18:586–96. [PubMed: 19433458]
74. Perl A, et al. T-cell and B-cell signaling biomarkers and treatment targets in lupus. *Curr Opin Rheumatol*. 2009; 21:454–64. [PubMed: 19550330]
75. Pascual V, Farkas L, Banchereau J. Systemic lupus erythematosus: all roads lead to type I interferons. *Curr Opin Immunol*. 2006; 18:676–82. [PubMed: 17011763]
76. Mathian A, Weinberg A, Gallegos M, Banchereau J, Koutouzov S. IFN- α induces early lethal lupus in preautoimmune (New Zealand Black x New Zealand White) F1 but not in BALB/c mice. *J Immunol*. 2005; 174:2499–506. [PubMed: 15728455]
77. Zagury D, et al. IFN α kinoid vaccine-induced neutralizing antibodies prevent clinical manifestations in a lupus flare murine model. *Proc Natl Acad Sci U S A*. 2009; 106:5294–9. [PubMed: 19279210]
78. Groom JR, et al. BAFF and MyD88 signals promote a lupuslike disease independent of T cells. *J Exp Med*. 2007; 204:1959–71. [PubMed: 17664289]
79. Pawar RD, et al. Inhibition of Toll-like receptor-7 (TLR-7) or TLR-7 plus TLR-9 attenuates glomerulonephritis and lung injury in experimental lupus. *J Am Soc Nephrol*. 2007; 18:1721–31. [PubMed: 17460144]

80. Ehlers M, Ravetch JV. Opposing effects of Toll-like receptor stimulation induce autoimmunity or tolerance. *Trends Immunol.* 2007; 28:74–9. [PubMed: 17197239]
81. Iwasaki A, Medzhitov R. Toll-like receptor control of the adaptive immune responses. *Nat Immunol.* 2004; 5:987–95. [PubMed: 15454922]
82. Lafyatis R, York M, Marshak-Rothstein A. Antimalarial agents: closing the gate on Toll-like receptors? *Arthritis Rheum.* 2006; 54:3068–70. [PubMed: 17009223]
83. Crispin JC, Kyttaris VC, Juang YT, Tsokos GC. How signaling and gene transcription aberrations dictate the systemic lupus erythematosus T cell phenotype. *Trends Immunol.* 2008; 29:110–5. [PubMed: 18249583]
84. Gilliet M, Cao W, Liu YJ. Plasmacytoid dendritic cells: sensing nucleic acids in viral infection and autoimmune diseases. *Nat Rev Immunol.* 2008; 8:594–606. [PubMed: 18641647]
85. Chaturvedi A, Dorward D, Pierce SK. The B cell receptor governs the subcellular location of Toll-like receptor 9 leading to hyperresponses to DNA-containing antigens. *Immunity.* 2008; 28:799–809. [PubMed: 18513998]
86. Christensen SR, Shlomchik MJ. Regulation of lupus-related autoantibody production and clinical disease by Toll-like receptors. *Semin Immunol.* 2007; 19:11–23. [PubMed: 17276080]
87. Kulkarni O, et al. Anti-Ccl2 Spiegelmer permits 75% dose reduction of cyclophosphamide to control diffuse proliferative lupus nephritis and pneumonitis in MRL-Fas(lpr) mice. *J Pharmacol Exp Ther.* 2009; 328:371–7. [PubMed: 18997060]
88. Suwannaroj S, Lagoo A, Keisler D, McMurray RW. Antioxidants suppress mortality in the female NZB x NZW F1 mouse model of systemic lupus erythematosus (SLE). *Lupus.* 2001; 10:258–65. [PubMed: 11341102]

Key Points

- The incidence of end stage renal disease due to SLE nephritis is static or increasing in the US and is responsible for high expenditure of health care dollars in SLE patients
- Several high profile clinical trials of novel biologic therapies targeting the immune system have failed in SLE nephritis despite convincing evidence that SLE nephritis is immune mediated
- While inhibition of single cell types or inflammatory mediators can prevent disease in murine models there is significant loss of efficacy in established disease
- Multiple interactions between innate and adaptive immune pathways in established disease amplify inflammation and make it difficult to restore normal tolerance mechanisms
- An understanding of the pathways that mediate intrinsic renal cell activation, interstitial renal inflammation, renal hypoxia and fibrosis should yield new strategies for protection of the kidneys
- The role of genetic polymorphisms in determining susceptibility to SLE and to renal damage may allow targeting of therapies to particular patient subgroups

Figure 1.

Pathogenesis of SLE nephritis. Renal deposition of autoantibodies, exposure to circulating inflammatory mediators and activation of complement initiate an inflammatory program that involves upregulation of adhesion molecules on endothelial cells, activation of intrinsic renal cells, induction of chemokines and recruitment of inflammatory cells that release cytokines and other molecules. Podocyte injury leads to loss of the slit diaphragm resulting in proteinuria, as well as decreased glomerular basement membrane production resulting in vascular compromise. Microvascular thromboses and endothelial cell death contribute to renal hypoxia that in turn causes tubular atrophy. Progressive inflammation recruits an increasingly diverse set of inflammatory mediators that amplify injury. Endothelial cell and renal dendritic cell activation are reversible at early stages, but activation of fibrotic pathways and of smooth muscle myofibroblast cells and progressive podocyte loss eventually result in irreversible renal damage. These pathways may all be amenable to therapeutic intervention.

Figure 2.

SLE activity induces multiple amplification loops that result in dysregulation of immune responses. After disease is triggered in a genetically susceptible individual, T_{FH} cells initiate disease by facilitating class switching and somatic mutation of autoantibodies that bind DNA or RNA. $IFN\alpha$, released during responses to exogenous antigens, contributes to disease initiation by priming the adaptive immune system thereby providing T cell help for autoreactive B cells and inducing release of IL-6 that enhances plasma cell differentiation. Alternatively, excessive BAFF is sufficient to initiate disease without T cell help (Loop 1). Abnormal naïve B cell repertoires and dysregulated germinal center regulatory mechanisms allow expansion and escape of autoreactive B cells. IgG autoantibodies recognizing nucleic acids are internalized via Fc or B cell receptors (BCR) into TLR expressing dendritic cells or B-cells respectively. TLR engagement in plasmacytoid dendritic cells induces type I IFN production⁸⁴ whereas TLR engagement in B cells increases BCR signaling and antibody production⁸⁵. In mice TLR7 is required for the production of anti-RNA antibodies whereas TLR9 is required for the anti-DNA response⁸⁶. Once long-lived plasma cells that secrete autoreactive IgG are present, perpetuation of the autoreactive response may continue in a T cell independent fashion (Loop 2). After the onset of clinical disease activated effector cells in the spleen and inflamed organs (red stars) provide cytokines that synergize with BAFF to enhance naïve B cell activation, class switching and memory B cell reactivation^{55, 66}, and directly cause tissue damage. $IFN\alpha$ cooperates with BAFF and enhances class switching and plasma cell differentiation via induction of cytokines such as IL-6. (Loop 3). Memory B cells activate naïve T cells, resulting in epitope spreading; memory T and B cells may then interact in a costimulation independent manner (Loop 4). The multiple interacting and self-

amplifying pathways that arise as disease progresses make it difficult to induce remission of disease with a single immune modulating reagent.

Table 1

List of reagents successful in mice that can be translated to trials in generalized and renal SLE

Target	Status of human trial (see www.clinicaltrials.gov)
<i>Innate immunity</i>	
IFN α	In progress and planned
TLR (ODN)	Planned
BAFF (Belimumab)	Phase II failed (generalized - efficacy). Phase III (generalized) primary endpoint met, a second Phase III study is in progress
BAFF/APRIL (Atacicept)	Halted (nephritis - toxicity). Generalized SLE trial in progress
<i>Acquired immunity</i>	
Cytoxan/Cellcept comparison	Equivalent
mTOR antagonists	Effective. Further studies in progress
B7-CD28 (CTLA4Ig)	Failed (generalized - efficacy) Nephritis trial in progress Nephritis trial with cyclophosphamide in progress
ICOS ligand	In progress
CD40/CD40L	Failed (generalized - toxicity)
B cells (anti-CD20)	Failed (generalized and nephritis - efficacy) Humanized antibody (nephritis in progress)
Modulation of B cell signaling (anti-CD22)	In progress
Cytokines (IL-6, IL-17, IL-21, IL-10)	Some planned
<i>Therapies directed at target organs</i>	
Complement	Phase I completed, others planned
IFN γ	In progress
TNF α	In progress for nephritis
Chemokines (CCL2) ⁸⁷	In progress for non-SLE renal disease
<i>Immune ablation</i>	
Stem cell transplant	In progress
<i>Interventions that may be protective without causing immune suppression</i>	
Vitamin D	In progress
“Natural” IgM ⁶¹	
ACE inhibitors	
Cholinergic agonists	
Statins	In progress
N-Acetylcysteine ⁸⁸	In progress
PPAR agonists	In progress