Published in final edited form as:

Chem Asian J. 2011 August 1; 6(8): 2101–2106. doi:10.1002/asia.201100190.

Phosphine-Catalyzed [4+2] Annulations of 2-Alkylallenoates and Olefins: Synthesis of Multisubstituted Cyclohexenes

Yang S. Tran^a, Tioga J. Martin^a, and Ohyun Kwon^a [Prof.]

Ohyun Kwon: ohyun@chem.ucla.edu

^aDepartment of Chemistry and Biochemistry, University of California, Los Angeles, 607 Charles E. Young Drive East, Los Angeles, CA 90095-1569 (USA), Fax: (+1) 310-206-3722

Abstract

From our investigations on phosphine-catalyzed [4+2] annulations between α -alkyl allenoates and activated olefins for the synthesis of cyclohexenes, we discovered a hexamethylphosphorous triamide (HMPT)-catalyzed [4+2] reaction between α -alkyl allenoates **1** and arylidene malonates or arylidene cyanoacetates **2** that provides highly functionalized cyclohexenes **3** and **4** in synthetically useful yields (30–89%), with moderate to exclusive regioselectivity, and reasonable diastereoselectivity. Interestingly, the [4+2] annulations between the α -alkyl allenoates **1** and the olefins **2** manifested a polarity inversion of the 1,4-dipole synthon **1**, depending on the structure of the olefin, thus providing cyclohexenes **3** exclusively when using arylidene cyanoacetates. The polarity inversion of α -alkyl allenoates from a 1,4-dipole **A** to **B** under phosphine catalysis can be explained by an equilibrium between the phosphonium dienolate **C** and the phosphorous ylide **D**.

Keywords

alkenes; allenoates; annulation; phosphine catalysis; regioselectivity

Introduction

The prevalence of six-membered carbocycles in pharmaceutical agents and natural products renders this ring system among the most important chemical structural motifs.^[1] Not surprisingly, therefore, a considerable amount of effort has been directed toward the development of new technologies to access these systems. The Diels–Alder reaction is probably the most efficient and widely used method for the rapid construction of these carbocycles.^[2] Remarkably, relatively few other chemical methods are available for the preparation of cyclohexenes through intramolecular processes^[3] and, even less commonly, intermolecular annulations.^[4]

Nucleophilic phosphine catalysis has recently played an important role in the development of new reactions, thus providing access to a variety of molecular frameworks.^[5] In particular, allenates, in the presence of a suitable tertiary phosphine catalyst, react with a long list of activated electrophiles to form various carbocycles and heterocycles.^[6] In this area, we have disclosed our findings on the use of ethyl 2-alkyl allenates as a 1,4-dipole synthon and their reactions with activated imines to generate tetrahydropyridines under phosphine catalysis.^[7] Subsequent investigations have led to the construction of cyclohexenes when ethyl 2-alkyl allenates are treated with arylidene malononitriles.^[8] Intriguingly, we observed the highly regioselective formation of ethyl 5,5-dicyano-4-arylcyclohex-1-enecarboxylate and ethyl 4,4-dicyano-5-arylcyclohex-1-enecarboxylate, depending on the catalyst employed. In this process, the ethyl 2-alkyl allenates **1** serve as a 1,4-dipole synthon **A** in the presence of catalytic hexamethylphosphorous triamide (HMPT), but as a polarity-reversed 1,4-dipole **B** in the presence of electron-deficient triarylphosphines (Scheme 1). This polarity inversion from the 1,4-dipole **A** to the 1,4-dipole **B** appears to arise from a change in the equilibrium between the phosphonium dienolate **C** and the vinylogous ylide **D**. The resulting [4+2] adducts are valuable intermediates for further transformations, both in natural product synthesis^[9] and diversity-oriented synthesis.^[10] Herein, we provide a full account of our investigation into this [4+2] reaction using various classes of olefinic electrophiles.

Results and Discussion

Based on our previous investigations into tetrahydropyridine syntheses,^[7] we initially hypothesized that ethyl 2-methyl allenate could potentially react with alkenes to form cyclohexenes under the influence of an appropriate phosphine catalyst. Mechanistically, the first step involves the generation of a phosphonium dienolate **C** through conjugate addition of the phosphine to ethyl 2-methylallenate **1a** (Scheme 2). Michael addition of the dienolate **C** to an activated olefin **2** facilitates the formation of the zwitterionic intermediate **E**. Subsequent proton-transfer steps facilitate tautomerization of the vinylphosphonium zwitterion **E** to an allylphosphonium zwitterion **F**. The intermediate **F** undergoes an intramolecular 6-*endo* cyclization and β -elimination of the phosphine to furnish the cyclohexene **3**.

With this mechanism in mind, we set out to examine the effects of a range of activated alkenes, catalysts with varying degrees of nucleophilicity,^[11] and solvents of various polarity^[12] on the formation of cyclohexenes through the [4+2] annulations of 2-alkyl allenates. Our initial attempts were made using ethyl 2-methyl allenate and commercially available electrophiles, including ethyl and methyl acrylates, acrylonitrile, cyclohexenone, dimethyl maleate, dimethyl fumarate, ethyl maleimide, and ethyl cinnamate. Our preliminary results were disappointing: we could not detect any cyclohexene products under phosphine catalysis conditions when using triphenylphosphine or tributylphosphine as the catalyst in benzene or dichloromethane. We hypothesized that the incompatibility of these electrophiles for [4+2] annulation was due, in part, to inefficient isomerization of the zwitterion **E** to the intermediate **F** (Scheme 2), thus inhibiting the final cyclization step.^[13] To assist this isomerization process, we envisioned that a longer-living zwitterion **E** might

be required to facilitate the formation of the intermediate **F**. Therefore, we chose to employ benzyldiene malonate and benzyldiene cyanoacetate as the coupling electrophiles.

Promisingly, the expected cyclohexene **3a** was formed when diethyl benzyldiene malonate (**2a**, 1.0 mmol) was treated with ethyl 2-methyl allenoate (**1a**, 2.0 mmol) and triphenylphosphine (20 mol%) in benzene under reflux (Table 1, entry 1). Although product formation was apparent through ¹H NMR spectroscopic analysis, the reaction proceeded at a slow rate and reached completion only after 96 hours, thereby providing a moderate isolated yield of **3a** (32%). Triarylphosphines featuring benzene rings substituted with electron-rich or electron-poor groups did not yield any product after performing the reaction for one day (Table 1, entries 2–4). Electron-rich tertiary phosphines containing alkyl, aryl, alkoxy, or diethylamino substituents were ineffective at mediating cyclohexene formation (Table 1, entries 5–11). These catalysts led to oligomerization of the allenoate **1a** instead of incorporation of the benzyldiene malonate (Table 1, entries 2–11). Further examination of catalysts revealed that bis(diethylamino)phenylphosphine could induce cyclohexene formation with a moderate isolated yield (10%, Table 1, entry 12). This observation led us to explore HMPT as the catalyst. To our delight, we achieved an appreciable increased reaction efficiency, thus leading to higher mass recovery of the [4+2] adduct, with an isolated yield of 63%, albeit as a 28:72 mixture of **3a** and **4a** (Table 1, entry 13). Our assignments of the structures of compounds **3a** and **4a** are based on comparison with those of their dicyano counterparts derived from the benzyldiene malononitrile.^[8]

Having found the optimal catalyst, we further optimized the reaction parameters to improve the reaction yields and regioselectivities (Table 2). There appeared to be a trend toward better mass recovery and regioselectivity when less-polar solvents were used. For instance, aromatic solvents were optimal, with very little difference in yield or regioselectivity when using benzene, toluene, or xylene, with the notable exception of the poor performance in high boiling mesitylene (Table 2, entries 1–4).^[14] While hexanes provided regioselectivity similar to those of the aromatic solvents (Table 2, entry 5), pentane exhibited excellent selectivity for **4a** in good yield (Table 2, entry 6). Although pentane would have been an ideal solvent for the formation of cyclohexenes **4**, the arylidene malonates other than diethyl benzyldiene malonate did not dissolve in pentane and their reactions did not proceed. Speculating that the high regioselectivity might be due to the low reaction temperature, we employed dichloromethane and a series of dichloromethane/pentane mixtures, but without success (Table 2, entries 7 and 8). Polar aprotic solvents provided modest results (Table 2, entries 9–12). The use of freshly distilled diethyl benzyldien emalonate (**2a**) and HMPT, with slow addition (over 4 h) of ethyl 2-methyl allenoate (**1a**) to minimize oligomerization of the allenoate, improved the reaction yield dramatically (89%) with retention of regioselectivity (Table 2, entry 13). Use of a stoichiometric amount of HMPT provided a similar result (Table 2, entry 14). Although there are reports of nucleophilic phosphine catalyzed reactions of allenes being facilitated by water,^[15–17] we found that the presence of water had a detrimental effect on cyclohexene formation.

With the optimized conditions for cyclohexene formation in hand, we performed experiments to probe the scope of this [4+2] annulation, and found out that a wide range of arylidene malonates was tolerated (Table 3). For example, arylidene malonates containing

benzene rings with electron-withdrawing groups (e.g., nitro, cyano, halogen) at the *para*-position produced cyclohexenes in good yields with reasonable regioselectivity (favoring the cyclohexene **4**; Table 3, entries 2–6). The formation of the cyclohexenes was much slower when the electron-rich *para*-methyl benzylidene malonate **2g** was used, thus leading to lower mass recovery as a result of the competing oligomerization of the allenolate (Table 3, entry 7). Allene oligomerization was circumvented when the allenolate **1a** was added over 12 hours to a mixture of dimethyl 4-methylbenzylidene malonate (**2g**) and HMPT (20 mol%) in benzene under reflux; here, the reaction yield improved to 84% (Table 3, entry 8). The reaction worked well with *meta*-methyl, bromo, and methoxy-substituted benzylidene malonates, albeit with poorer regioselectivities (Table 3, entries 9–11). Furthermore, the reaction was compatible with heteroarylidenes malonates: dimethyl 2-furylidene malonate (**2k**), dimethyl 2-thienylidene malonate (**2l**), and dimethyl *N*-Boc-2-indolylidene malonate (**2m**) were viable substrates that furnished cyclohexene products with good yields and regioselectivities (Table 3, entries 12–14).

Next, we examined (Table 4) the use of arylidene cyanoacetates as activated olefin partners for the [4+2] annulations of ethyl 2-methyl allenolate (**1a**). Unlike the situation with arylidene malonates, here we isolated only one cyclohexene regioisomer **3** when arylidene cyanoacetates were employed under otherwise identical conditions. Slightly higher yields resulted from electron-deficient arylidenes (Table 4, entries 1–3 and 8) compared with those from electron-rich arylidenes (Table 4, entries 4, 5, and 7). Product yields of the desired cyclohexenes **3** diminished when 2-substituted arylidenes were used, presumably because of steric hindrance and subsequent decreased reactivity at the β -carbon atom (Table 4, entries 7 and 8).^[18] For a given substituent (e.g., chloro, bromo), the *para*-substituted arylidene provided a better reaction yield than its *ortho* or *meta* counterpart (Table 4, entries 2/8 and 3/9).

The overall mass recoveries for the [4+2] annulations with the arylidene cyanoacetates were lower than those of the corresponding arylidene malonates. ¹H NMR spectroscopic analysis of the crude reaction mixtures revealed the presence of trace amounts (typically <15%) of the other isomer **4**. All attempts to isolate and characterize these minor products **4** were unsuccessful. Further examination of the crude reaction mixture obtained from *ortho*-methoxybenzylidene cyanoacetate (**2t**) allowed the isolation of a new minor product in 15% yield; its structure was assigned as that of the [2+2] adduct **5**.

To further explore the substrate scope for the phosphine-catalyzed [4+2] cyclohexene formation, we investigated the effects of various α -alkyl allenolates (Table 5). This reaction tolerated a range of allenylic β' -substituents on the allenolate **1**, including aryl, substituted aryl, and carboxylic ester moieties, thus providing cyclohexenes **3** in good isolated yields with reasonable diastereoselectivities (Table 5, entries 1–3). For these β' -substituted

allenoates **1**, only the cyclohexenes **3** were formed, in accordance with previous observations.^[8] Notably, dimethyl benzylidene malonate (**2a**) and ethyl 2-benzylallenoate (**1b**) did not furnish any of their corresponding cyclohexenes **3**, presumably because of steric hindrance and the diminished reactivities of these olefins.

Mechanistic Considerations

Based on these observations, we rationalize that the formation of **3** and **4** arises through the γ - or β' -addition of the phosphonium intermediate (**C** $\rightleftharpoons$ **D**) to the activated olefin **2**. With more reactive arylidene cyanoacetates and/or sterically congested 2-alkylallenoates with α -alkyl groups larger than methyl, the addition occurs at the γ -carbon atom of the phosphonium enolate **C**; the formation of regioisomer **3** ensues (Scheme 2). With less-reactive arylidene malonates, however, the phosphonium dienolate **C** isomerizes into the vinylogous ylide **D**, which adds to the olefin **2** and produces intermediate **G**. Consecutive proton transfers provide the deconjugated enoate **H**, which undergoes 6-*endo* cyclization to generate the cyclic ylide **I**. Finally, 1,2-proton transfer and β -elimination of the phosphine catalyst furnish the cyclohexene **4** (Scheme 3).

Synthetic Application of Functionalized Cyclohexenes

Scheme 4 illustrates the synthetic utility of the *gem*-cyanoacetate adduct. Alkaline hydrolysis of the nitrile **3o** using aqueous potassium carbonate and hydrogen peroxide furnished the carboxamide **6** in 84% yield without saponification of the ethyl carboxylic ester groups.

Scheme 5 further demonstrates the potential utility of this [4+2] annulation for the synthesis of biologically active natural products. We achieved rapid entry into the tetracyclic framework **7**^[19] of isoborreverine^[20] through *N*-Boc deprotection of the cyclohexene triester **4m** by using trifluoroacetic acid followed by the lactamization with the *tert*-butyl-magnesium chloride.

Conclusions

We have further established that the novel phosphine-catalyzed [4+2] annulation between allenoates and activated alkenes is a valuable strategy for cyclohexene synthesis. Ethyl 2-alkyl allenoates can serve as either a 1,4-dipole **A** or an inverted dipole **B** by virtue of the steric environment around the allenoate or the reactivity of the activated alkene. Further manipulation of the cyclohexene adducts demonstrates the utility of the phosphine-catalyzed [4+2] annulations of allenoates with olefins.

Experimental Section

Materials and Methods

All reactions were performed under an atmosphere of argon with dry solvents and anhydrous conditions, unless otherwise noted. Benzene, toluene, and CH₂Cl₂ were freshly distilled from CaH₂. All other reagents were used as received from commercial sources. All ethyl 2-substituted methylallenoates **1** were synthesized according to procedures reported

previously. Arylidenene malonate and arylidene cyanoacetate **2** were synthesized through phosphine-catalyzed Knoevenagel condensation of the pertinent aldehyde and cyanoacetate, according to the procedure reported by Yadav,^[21] or through piperidine-catalyzed condensation of the corresponding aldehyde and malonate. HMPT was distilled under vacuum in three fractions; only the middle fraction was used. The distilled HMPT was then stored under an atmosphere of argon in a dry glove box. Reactions were monitored through thin layer chromatography (TLC) on 0.25 mm Silicycle silica gel plates (TLG-R10011B-323), visualizing with UV light or staining with permanganate or anisaldehyde. Flash column chromatography was performed using Silicycle Silia-P gel (50 μ m particle size, R12030B) and compressed air. Preparative HPLC was performed by using (*R,R*)-1,2-diaminocyclohexane(DACH)-3,5-dinitrobenzoyl(DNB)-modified silica gel. IR spectra were recorded using a Perkin–Elmer Paragon 1000 FTIR spectrometer. NMR spectra were obtained using Bruker Avance-500, ARX-500, or Avance-300 instruments and calibrated using $\text{CHCl}_3/\text{CDCl}_3$ as the internal reference (7.26 and 77.0 ppm for ^1H and ^{13}C NMR spectra, respectively). Data for ^1H NMR spectra are reported as follows: chemical shift (δ ppm), multiplicity, coupling constant (Hz), and integration. Data for ^{13}C NMR spectra are reported in terms of chemical shift and multiplicities, with coupling constants (Hz) in the case of J_{CF} coupling. The following abbreviations are used to explain the multiplicities: s, singlet; d, doublet; t, triplet; q, quartet; sept, septet; m, multiplet; br, broad; app, apparent. High-resolution electron ionization (EI) mass spectra were recorded after rapid thermal vaporization of samples deposited on a desorption ionization filament inserted directly into an EI (70 eV, 200°C) source of a triple-sector high-resolution instrument (VG/Micromass Autospec) tuned to 8000 static resolution (M/DM, 10% valley) using perfluorinated kerosene (formula weight 705, Lancaster Synthesis, Windham, NH) as the internal calibrant. High-resolution electrospray ionization (HRESI) mass spectra were recorded after flow injection of CHCl_3 solutions into an ESI source attached to a 7.5-T FTMS (Ion Spec Ultima, Irvine, CA) instrument. Data were analyzed using the instrument-supplied software. Melting points (m.p.) are uncorrected; they were recorded using an electrothermal capillary melting point apparatus.

General Procedure for the Synthesis of Cyclohexenes **3** and **4**

A solution of arylidene malonate or arylidene cyanoacetate (1.0 mmol) and HMPT (0.2 or 1.0 mmol) in benzene (5 mL) was heated under reflux in a 50 mL flame-dried round-bottom flask under an atmosphere of argon. The allenolate (3.0 mmol) in benzene (10 mL) was added dropwise over a period of at least 4 h. An orange-red solution was obtained; the mixture was stirred under reflux until TLC analysis revealed consumption of the olefin **2**. The reaction mixture was concentrated and the residue purified through flash column chromatography (gradient eluent: 50% EtOAc in hexanes), followed by preparative HPLC (gradient eluent: 50% CH_2Cl_2 in hexanes), to provide the corresponding cyclohexene **3** and **4**.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the U.S. National Institutes of Health (O.K.: R01M071779, P41M081282) for financial support.

References

1. a) Myers, RL. The 100 Most Important Chemical Compounds. Greenwood Press; Westport, Connecticut: 2007. b) Buss, AD.; Butler, MS., editors. Natural Product Chemistry for Drug Discovery. RSC Publishing; Cambridge, UK: 2010. c) Gringauz, A. Introduction to Medicinal Chemistry—How Drugs Act and Why. Wiley-VCH; New York: 1997. d) Fischer, J.; Ganellin, RC. Analogue-Based Drug Discovery. Wiley-VCH; Weinheim: 2006. e) Ho, TL. Carbocycle Construction in Terpene Synthesis. Wiley-VCH; Weinheim, Germany: 1998.
2. a) Fringuelli, F.; Taticchi, A., editors. The Diels–Alder Reaction: Selected Practical Methods. Wiley; West Sussex, England: 2002. b) Kobayashi, S.; Jorgensen, KA., editors. Cyclo-addition Reactions in Organic Synthesis. Wiley-VCH; Weinheim, Germany: 2002. c) Nicolaou KC, Snyder SA, Montagnon T, Vassilikogiannakis GE. Angew Chem. 2002; 114:1742. Angew Chem Int Ed. 2002; 41:1668. d) Takao KI, Munakata R, Tadano KI. Chem Rev. 2005; 105:4779. [PubMed: 16351062]
3. a) Jung ME, Ho DG. Org Lett. 2007; 9:375. [PubMed: 17217308] b) Wang LC, Luis AL, Agapiou K, Jang HY, Krische MJ. J Am Chem Soc. 2002; 124:2402. [PubMed: 11890765] c) Frank SA, Mergott DJ, Roush WR. J Am Chem Soc. 2002; 124:2404. [PubMed: 11890766] d) Krafft ME, Haxell TFN. J Am Chem Soc. 2005; 127:10168. [PubMed: 16028918] e) Grubbs RH, Miller SJ, Fu GC. Acc Chem Res. 1995; 28:446. f) Kim H, Goble SD, Lee C. J Am Chem Soc. 2007; 129:1030. [PubMed: 17263374]
4. Enders D, Huttli MRM, Grondal C, Raabe G. Nature. 2006; 441:861. [PubMed: 16778886]
5. For reviews, see: Lu X, Zhang C, Xu Z. Acc Chem Res. 2001; 34:535. [PubMed: 11456471] ; Valentine DH, Hillhouse JH. Synthesis. 2003; 3:317.; Methot JL, Roush WR. Adv Synth Catal. 2004; 346:1035.; Lu X, Du Y, Lu C. Pure Appl Chem. 2005; 77:1985.; Nair V, Menon RS, Sreekanth AR, Abhilash N, Biji AT. Acc Chem Res. 2006; 39:520. [PubMed: 16906748] ; Denmark SE, Beutner GL. Angew Chem. 2008; 120:1584.; Angew Chem Int Ed. 2008; 47:1560.; Ye L, Zhou J, Tang Y. Chem Soc Rev. 2008; 37:1140. [PubMed: 18497927] ; Kwong CKW, Fu MY, Lam CSL, Toy PH. Synthesis. 2008:2307.; Aroyan CE, Dermenci A, Miller SJ. Tetrahedron. 2009; 65:4069.; Cowen BJ, Miller SJ. Chem Soc Rev. 2009; 38:3102. [PubMed: 19847345] ; Marinetti A, Voituriez A. Synlett. 2010:174.; Beata K. Cent Eur J Chem. 2010:1147.
6. For selected examples of phosphine catalysis of allenes with electrophiles, see: Zhang C, Lu X. J Org Chem. 1995; 60:2906.; Xu Z, Lu X. Tetrahedron Lett. 1997; 38:3461.; Kumar K, Kapur A, Ishar MPS. Org Lett. 2000; 2:787. [PubMed: 10814429] ; Kumar K, Kapoor R, Kapur A, Ishar MPS. Org Lett. 2000; 2:2023. [PubMed: 10891220] ; Ishar MPS. Org Lett. 2000; 2:2023. [PubMed: 10891220] ; Zhu XF, Henry CE, Wang J, Dudding T, Kwon O. Org Lett. 2005; 7:1387. [PubMed: 15787513] ; Zhu XF, Schaffner AP, Li RC, Kwon O. Org Lett. 2005; 7:2977. [PubMed: 15987184] ; Zhu XF, Henry CE, Kwon O. Tetrahedron. 2005; 61:6276.; Wallace DJ, Sidda RL, Reamer RA. J Org Chem. 2007; 72:1051. [PubMed: 17253835] ; Henry CE, Kwon O. Org Lett. 2007; 9:3069. [PubMed: 17629288] ; Creech GS, Kwon O. Org Lett. 2008; 10:429. [PubMed: 18173275] ; Xu S, Zhou L, Ma R, Song H, He Z. Chem Eur J. 2009; 15:8698. [PubMed: 19693756] ; Guo H, Xu Q, Kwon O. J Am Chem Soc. 2009; 131:6318. [PubMed: 19374356] ; Wang T, Ye S. Org Lett. 2010; 12:4168. [PubMed: 20712333]
7. a) Zhu XF, Lan J, Kwon O. J Am Chem Soc. 2003; 125:4716. [PubMed: 12696883] b) Zhao GL, Shi M. Org Biomol Chem. 2005; 3:3686. [PubMed: 16211103] c) Baskar B, Dakas PY, Kumar K. Org Lett. 2011; 13:1988. [PubMed: 21417315]
8. Tran YS, Kwon O. J Am Chem Soc. 2007; 129:12632. [PubMed: 17914823]
9. a) Wurz RP, Fu GC. J Am Chem Soc. 2005; 127:12234. [PubMed: 16131196] b) Tran YS, Kwon O. Org Lett. 2005; 7:4289. [PubMed: 16146409]
10. a) Castellano S, Fiji HDG, Kinderman SS, Watanabe M, De Leon P, Tamanoi F, Kwon O. J Am Chem Soc. 2007; 129:5843. [PubMed: 17439124] b) Watanabe M, Fiji MHDF, Guo L, Chan L, Kinderman SS, Slamon DJ, Kwon O, Tamanoi F. J Biol Chem. 2008; 283:9571. [PubMed: 18351062]

- 18230616] c) Lu J, Chan L, Fiji HDF, Dahl R, Kwon O, Tamanai F. Mol Cancer Ther. 2009; 8:1218. [PubMed: 19417142] d) Wang Z, Castellano S, Kinder-man SS, Argueta CE, Beshir AB, Fenteany G, Kwon O. Chem Eur J. 2011; 17:649. [PubMed: 21207585] e) Cruz D, Wang Z, Kibbie J, Modlin R, Kwon O. Proc Natl Acad Sci U S A. 2011; 108:6769. [PubMed: 21383121] f) Choi J, Mouillesseaux K, Wang Z, Fiji HDG, Kinderman SS, Otto GW, Geisler R, Kwon O, Chen JN. Development. 2011; 138:1173. [PubMed: 21307094]
11. a) Ofial, A.; Mayr, H. [accessed February 2011] Reactivity Scales. <http://www.cup.uni-muenchen.de/oc/mayr/CDpublika.html> b) Brotzel F, Kempf B, Singer T, Zipse H, Mayr H. Chem Eur J. 2007; 13:336. [PubMed: 17009366] c) see Ref. [51].
12. Reichardt C. Angew Chem. 1979; 91:119. Angew Chem Int Ed Engl. 1979; 18:98.
13. It is possible that the initial addition of the intermediate **C** to the olefin **2** is the stumbling block; it might also be aided by the use of more-electrophilic Michael acceptors, such as arylidene malononitriles and arylidene malonates.
14. With mesitylene as a solvent, only traces of cyclohexenes were formed, presumably because the high temperature of refluxing mesitylene led to decomposition of the allenolate **1a**.
15. Unlike the cases for the [3+2] annulations of allenolates with alkenes or imines (see Refs. [16] and [17]), addition of water (25, 50, 100, or 400 mol%) under the optimized conditions for the [4+2] annulation of ethyl 2-methyl allenolate (**1a**) and the alkene **2a** was detrimental to the reaction.
16. Fang YQ, Jacobsen EN. J Am Chem Soc. 2008; 130:5660. [PubMed: 18393504]
17. a) Xia Y, Liang Y, Chen Y, Wang M, Jiao L, Huang F, Liu S, Li Y, Yu ZX. J Am Chem Soc. 2007; 129:3470. [PubMed: 17319666] b) Mercier E, Fonovic B, Henry CE, Kwon O, Dudding T. Tetrahedron Lett. 2007; 48:3617.
18. This result is consistent with our previous findings for allenolate/arylimine [4+2] annulations. See Ref. [7].
19. Compound **7** was obtained as a single diastereoisomer. The relative stereochemistry of the two stereogenic centers in compound **7** was not established.
20. a) Fernandez LS, Buchanan MS, Carroll AR, Feng YJ, Quinn RJ, Avery VM. Org Lett. 2009; 11:329. [PubMed: 19090698] b) Baldé AM, Pieters LA, Gergely A, Wray V, Claeys M, Vlietinck AJ. Phytochemistry. 1991; 30:997. c) Tillequin F, Koch M, Rabaron A. J Nat Prod. 1985; 48:120. d) Tillequin F, Rousselet R, Koch M, Bert M, Sevenet T. Ann Pharm Fr. 1979; 37:543. [PubMed: 546292] e) Tillequin F, Koch M, Bert M, Sevenet T. J Nat Prod. 1979; 42:92. f) Tillequin F, Koch M, Pousset JL, Cave A. J Chem Soc Chem Commun. 1978:826.
21. Yadav JS, Subba Reddy BV, Basak AK, Visali B, Narsaiah AV, Nagaih K. Eur J Org Chem. 2004:546.

Scheme 1.

Ethyl 2-methylallenoate functioning as 1,4-dipoles.

Scheme 2.

Proposed Mechanism for phosphine-catalyzed cyclohexene formation.

Scheme 3. Mechanistic rationale for the formation of cyclohexenes **4**.

Scheme 4.

Hydrolysis of the nitrile **3o** to the carboxamide **6**. DMSO=dimethyl sulfoxide.

Scheme 5.

Pyrroloindolinone formation from the cyclohexene gem-diester **4m**. TFA=trifluoroacetic acid.

Table 1Survey of phosphine catalysts for [4+2] annulation of the allenolate **1a** and the alkene **2a**.^[a]

Entry	Phosphine	Reaction Time [h]	Yield [%] ^[b]	3a:4a ^[c]
1	PPh ₃	96	32	>95:5
2	(4-Me ₂ NC ₆ H ₄) ₃ P	24	NR ^[d]	NA ^[e]
3	(4-MeOC ₆ H ₄) ₃ P	24	NR ^[d]	NA ^[e]
4	(4-FC ₆ H ₄) ₃ P	24	NR ^[d]	NA ^[e]
5	PBu ₃	48	NR ^[d]	NA ^[e]
6	PEt ₂ Ph	48	NR ^[d]	NA ^[e]
7	PEtPh ₂	48	NR ^[d]	NA ^[e]
8	(EtO) ₃ P	48	NR ^[d]	NA ^[e]
9	(EtO) ₂ PhP	48	NR ^[d]	NA ^[e]
10	(MeO) ₂ (Et ₂ N)P	48	NR ^[d]	NA ^[e]
11	(MeO)(Et ₂ N) ₂ P	48	2	ND ^[f]
12	(Et ₂ N) ₂ PhP	48	10	22:78
13	HMPT	24	63	28:72

^[a] Reaction conditions: **1a** (1.4–2.0 mmol), **2a** (1 mmol), and the phosphine (20 mol%) were heated under reflux in benzene (10 mL).^[b] Yield of isolated product.^[c] Regioisomeric ratio **3a:4a** was determined through analysis of the ¹H NMR spectrum of the crude reaction mixture.^[d] No reaction.^[e] Not applicable.^[f] Not determined.

Table 2

Survey of solvents for [4+2] annulation of the allenolate **1a** and the alkene **2a**.^[a]

Entry	Solvent	Reaction Temp. [°C]	Reaction Time [h]	Yield [%] ^[b]	3a:4a ^[c]
1	benzene	80	24	63	28:72
2	toluene	111	24	65	33:67
3	xylene	140	48	72	23:77
4	mesitylene	165	48	trace	ND ^[d]
5	hexanes	69	24	48	28:72
6	pentane	36	14	72	>5:95
7	CH ₂ Cl ₂ /pentane (1:1)	36	24	trace	55:45
8	CH ₂ Cl ₂	39	48	NR ^[e]	NA ^[f]
9	Et ₂ O	33	24	15	20:80
10	THF	66	24	64	55:45
11	dioxane	101	48	12	53:47
12	MeCN	81	15	22	60:40
13	benzene	80	24	89 ^[g]	28:72
14	benzene	80	24	90 ^[h]	28:72

^[a] Reaction conditions: **1a** (1.4–2.0 mmol), **2a** (1 mmol), and the phosphine (20 mol%) were heated under reflux in the designated solvent (10 mL).

^[b] Yield of isolated product.

^[c] Regioisomeric ratio **3a:4a** was determined through analysis of the ¹H NMR spectrum of the crude reaction mixture.

^[d] Not determined.

^[e] No reaction.

^[f] Not applicable.

^[g] All starting reagents were freshly distilled.

[*h*] 1 or 2 equivalents of HMPT was used.

NIH-PA Author Manuscript

NIH-PA Author Manuscript

NIH-PA Author Manuscript

Table 3Survey of arylidenemalonates for [4+2] annulations with the allenolate **1a**.^[a]

Entry	Ar	R	Yield [%] ^[b]	3a/4a ^[c]
1	Ph (2a)	Et	89	28:72
2	4-O ₂ NC ₆ H ₄ (2b)	Me	69	23:77
3	4-NCC ₆ H ₄ (2c)	Me	75	24:76
4	4-BrC ₆ H ₄ (2d)	Me	84	21:79
5	4-ClC ₆ H ₄ (2e)	Me	86	30:70
6	4-FC ₆ H ₄ (2f)	Me	82	26:74
7	4-MeC ₆ H ₄ (2g)	Me	23	22:78
8	4-MeC ₆ H ₄ (2g)	Me	84 ^[d]	22:78
9	3-MeC ₆ H ₄ (2h)	Me	81	50:50
10	3-BrC ₆ H ₄ (2i)	Me	83	39:61
11	3-MeOC ₆ H ₄ (2j)	Me	81	33:66
12	2-furyl (2k)	Me	72	16:84
13	2-thienyl (2l)	Me	71	29:71
14	<i>N</i> -Boc-2-indolyl (2m)	Me	87	20:80

^[a] Reaction conditions: A solution of the allenolate **1a** (1.4–3.0 mmol) in benzene (10 mL) was added to a solution of the malonate **2** (1 mmol) and HMPT (20 mol%) in benzene (5 mL) under reflux over 4 h and then the mixture was further heated under reflux in benzene until the disappearance (TLC) of the malonate.

^[b] Yield of isolated product.

^[c] Regioisomeric ratio **3a/4a** was determined through analysis of the ¹H NMR spectrum of the crude reaction mixture.

^[d] The allenolate **1a** was added over 12 h.

Table 4Survey of Arylidencyanoacetates for [4 + 2] Annulations with the Allenolate **1a**.^[a]

Entry	Ar	Product	Yield [%] ^[b]
1	Ph (2n)	3n	75
2	4-ClC ₆ H ₄ (2o)	3o	75
3	4-BrC ₆ H ₄ (2p)	3p	76
4	4-MeC ₆ H ₄ (2q)	3q	68
5	4-MeOC ₆ H ₄ (2r)	3r	65
6	3-MeOC ₆ H ₄ (2s)	3s	63
7	2-MeOC ₆ H ₄ (2t)	3t	30
8	2-ClC ₆ H ₄ (2u)	3u	60
9	3-BrC ₆ H ₄ (2v)	3v	63

^[a] Reaction conditions: A solution of the allenolate **1a** (1.4–3.0 mmol) in benzene (10 mL) was added to a solution of the cyanoacetate **2** (1 mmol) and HMPT (20 mol%) in benzene (5 mL) under reflux over 4 h and then the mixture was further heated under reflux.

^[b] Yield of isolated product.

Table 5

Survey of allenates **1** for [4+2] annulations with benzyldenecyanoacetate **2n**.^[a]

Entry	R	Product	Yield [%] ^[b]	cis/trans ^[c]
1	Ph (1b)	3w	81	88:12
2	4-ClC ₆ H ₄ (1c)	3x	74	89:11
3	CO ₂ Et (1d)	3y	78	77:23

^[a] Reaction conditions: A solution of the allenate **1** (1.4–3.0 mmol) in benzene (10 mL) was added to a solution of the olefin **2n** (1 mmol) and HMPT (20 mol%) in benzene (5 mL) under reflux over 4 h and then the mixture was further heated under reflux.

^[b] Yield of isolated product.

^[c] Determined through analysis of the ¹H NMR spectrum of the crude reaction mixture and by comparison with the spectrum of *cis*-ethyl 5,5-dicyano-4,6-diphenylcyclohex-1-enecarboxylate (for which a single-crystal X-ray structure was obtained). See Ref. [8].