Published in final edited form as:

Med Sci Sports Exerc. 2013 November ; 45(11): 2091–2097. doi:10.1249/MSS.0b013e318299afd4.

Physical Activity and Lower Limb Lymphedema among Uterine Cancer Survivors

Justin C. Brown¹, Gabriella M. John², Saya Segal³, Christina S. Chu⁴, and Kathryn H. Schmitz¹

¹Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania, PA

²Department of Epidemiology, Columbia University, New York, NY

³Division of Female Pelvic Medicine and Reconstructive Surgery; University of Medicine and Dentistry of New Jersey, New Brunswick, NJ

⁴Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, University of Pennsylvania, PA

Abstract

Purpose—Physical activity (PA) is known to provide physical and mental health benefits to uterine cancer survivors. However, it is unknown if PA associates with lower limb lymphedema (LLL), an accumulation of protein-rich fluid in the lower limbs. Therefore, we sought to examine the association between PA and LLL in uterine cancer survivors, with a focus on walking.

Methods—We conducted a cross-sectional study using mailed surveys among uterine cancer survivors who received care at a university-based cancer center. We asked about PA, walking, and LLL symptoms using validated self-report questionnaires. PA was calculated using metabolic equivalent hours per week (MET-hrs·wk⁻¹), and walking was calculated using blocks per day (blocks·d⁻¹).

Results—The response rate to our survey was 43%. Among the 213 uterine cancer survivors in our survey, 36% were classified as having LLL. Compared with participants who reported <3 MET-hrs·wk⁻¹ of PA, participants who reported ≥8.0 MET-hrs·wk⁻¹ of PA had an odds ratio of LLL of 0.32 (95% CI: 0.15–0.69; $P_{trend} = .003$). Stratified analyses suggested the association of PA and LLL existed only among women with a body mass index (BMI) <30 kg/m² ($P_{trend} = .007$), compared to women with a BMI ≥30 kg/m² ($P_{trend} = .47$). Compared with participants who reported <4.0 blocks·d⁻¹ of walking, participants who reported ≥12 blocks·d⁻¹ of walking had an odds ratio of LLL of 0.19 (95% CI: 0.09–0.43; $P_{trend} < .0001$). Stratified analyses suggested the association of walking and LLL was similar among women with a BMI <30 kg/m² ($P_{trend} = .007$) and women with a BMI ≥30 kg/m² ($P_{trend} = .03$).

Corresponding Author: Kathryn H. Schmitz, University of Pennsylvania School of Medicine, 423 Guardian Drive, 8th Floor, Blockley Hall, Philadelphia, PA 19104, Phone: 215-898-6604, Fax: 251-573-5311, schmitz@mail.med.upenn.edu.

Disclosure of funding received: None

Conflicts of Interest: All other authors declare there are no conflicts of interest.

Conclusion—Participation in higher levels of PA or walking is associated with reduced proportions of LLL in dose-response fashion. These findings should be interpreted as preliminary, and investigated in future studies.

Keywords

gynecologic cancer; exercise; mobility disability; edema; quality of life

INTRODUCTION

Lower limb lymphedema (LLL) affects 7–47% of the 40,000 women diagnosed with uterine cancer each year in the United States (15, 22). The prevalence of LLL varies depending on the modality and threshold used for diagnosis (8, 9, 41). LLL is characterized by swelling and accumulation of protein-rich fluid in the lower limbs (23, 27) which impairs physical function (16), and reduces quality of life (5, 14). LLL imparts long-term psychological burden due to limited efficacious therapies for symptom management (28, 43). Established risk factors for the development of LLL include treatment with adjuvant radiation therapy, removal of 31 lymph nodes, and dissection of the saphenous vein (9, 44, 46). The incidence of LLL will likely rise as the number of lymph nodes sampled among gynecologic cancer survivors has increased by as much as 350% in the past two decades (1). Therefore, efficacious interventions are needed to minimize the deleterious sequelae associated with LLL.

The American Cancer Society (35) suggests all uterine cancer survivors participate in physical activity (PA) following the guidelines set forth by the American College of Sports Medicine (ACSM) (38). However, LLL is associated with impaired leg blood flow (7), poor tissue oxygenation (42), and poor wound healing (13), such that the ACSM recommended uterine cancer survivors (or any gynecologic cancer survivor) with swelling or inflammation of the lower limbs be examined by a medical professional for clearance prior to engaging in PA (38). PA such as aerobic treadmill walking improves leg blood flow (19), tissue oxygenation (20, 30), wound healing (40) among people with peripheral artery disease (PAD) and venous leg ulcers. Further, PA has been shown to improve symptoms of the lower limbs known to impair quality of life (31). These symptoms are similar to those associated with LLL experienced by cancer survivors.

It is unknown if PA associates with LLL among uterine cancer survivors. Identifying an association between PA and LLL would provide a foundation for future research to examine the efficacy of PA to reduce LLL symptom frequency and severity among this group of cancer survivors. Alternatively, an association between PA and LLL may serve to identify cancer survivors whose symptoms preclude them from safely completing the standard prescription of PA for cancer survivors (38), and may require a tailored PA prescription to maximize the potential health-benefits gained through PA participation. Therefore, the primary goal of our study was to examine the association between PA and LLL.

METHODS

Participants and procedures

We conducted a cross-sectional survey of patients with uterine cancer who received care at the Abramson Cancer Center at the University of Pennsylvania (Philadelphia, PA). Potential participants included women ≥ 20 years old, with a history of uterine cancer. Potentially eligible participants were identified using fellow surgical case logs from 2008–2010, and ICD-9 diagnosis codes 179.0 and 182.0–182.8, from 2006–2010. ICD-9 codes 179.0 and 182.0–182.8 are the primary codes used to classify cancers of the uterus. ICD-9 codes are commonly used in medical settings to systematically classify diagnoses for billing purposes. Participants identified to meet inclusion criteria by study staff were sent a letter signed by their oncologist explaining the purpose of the study. Potentially eligible participants were provided with the option to decline participation within two-weeks of receiving the letter from their oncologist. Those who did not decline participation were sent a survey to complete. After two-weeks, a second survey was sent to those who did not reply to the first mailed survey (25). This protocol was approved by the University of Pennsylvania Institutional Review Board, and the University of Pennsylvania cancer center. We classified women who provided their informed consent as those who mailed back a completed survey.

Lower limb lymphedema assessment

The Gynecologic Cancer Lymphedema Questionnaire (GCLQ) was used to assess symptoms associated with LLL (10). The GCLQ is a validated self-report measure that assesses seven domains of symptoms in both lower extremities. The seven domains include heaviness, general swelling, limb-related swelling, infection, aching, numbness, and physical function, with one or more symptom questions per domain. Participants reporting ≥ 5 symptoms of the lower extremities within the seven above-listed domains were classified as having LLL. The GCLQ demonstrates good psychometric characteristics, with ≥ 5 symptoms maximizing the sensitivity, specificity, positive- and negative-predictive values compared to other cutoff scores (10). Our study group omitted one of the three questions in the swelling-general domain (question #20). In the validation study of the GCLQ there were no women that reported only having answered yes to the question we omitted (personal communication with J. Carter). We conducted sensitivity analyses assuming everyone responded “no” or “yes” to this question; our findings were consistent in all analyses with those reported herein.

Physical activity assessment

The Paffenbarger Physical Activity Questionnaire (PPAQ) was used to assess participation in PA (34). The PPAQ has been validated (3, 26) and used previously among cancer survivors (32, 33).

Participants were asked to “list any sports, leisure, or recreation activities you have participated in on a regular basis during the past year.” Participants were also asked to list the average number of sessions per week and the duration of each session for each PA listed. Trained research staff then coded each PA listed with a metabolic equivalent (MET) using the compendium of physical activities (2). For reference, one MET is the energy expended

when sitting quietly for one hour, and 3.5 METs is walking for pleasure. For each MET value, we calculated the weekly activity-specific MET-hrs-wk⁻¹, as the product of the MET value, the number of sessions per week, and the number of hours per session. For each participant, we summed the activity-specific MET-hrs-wk⁻¹ to generate an aggregate measure of MET-hrs-wk⁻¹. We created categories of MET-hrs-wk⁻¹, defined as <3.0, 3.0–8.9, 9.0–17.9, and ≥18.0 that correspond to <1.0, 1.0–2.9, 3.0–5.9, and ≥6.0 hours per week of moderate-intensity PA, consistent with prior analyses among cancer survivors (32, 33).

Participants were asked “how many city blocks or their equivalent have you walked on an average day during the past year.” It was noted on the questionnaire that “12 blocks equals one mile.” We then created categories of blocks per day (blocks·d⁻¹) of walking, defined as <4.0, 4.0–11.9, and ≥12 blocks·d⁻¹, which correspond to <¼, ¼ to <1, and ≥1 mile of walking per day, consistent with prior analyses among older adults (18).

Covariates

Information on covariates came from self-report or electronic medical records. Variables collected from self-report included age, marital status, race, education, employment, body mass index (BMI). We used the Charlson Comorbidity Index (11, 12) to identify comorbidities whose symptoms may be similar to LLL, such as congestive heart failure, PAD, and diabetes mellitus (45). We used the Gynecologic Oncology Group neurotoxicity sensory index to assess neuropathy from cancer treatment, as neuropathy may influence participation in PA (21). This self-administered questionnaire quantifies numbness and discomfort in the hands and feet, and ranges from 0 to 16 with higher scores representing more severe neuropathy. Variables collected from the electronic medical record included pathology type, stage of cancer, time since diagnosis, and cancer treatment history.

Statistical analysis

Response rates to our survey were calculated using the method described by the American Association for Public Opinion Research (25). We performed descriptive statistics and bivariate analyses on all study variables using the Wilcoxon Rank-Sum test for continuous variables and Fishers exact test for categorical variables. We used Logistic regression models to estimate the odds ratio (OR) of reporting LLL with 95% confidence intervals (95% CI). The *P* value for the linear trend test across categories (*P*_{trend}) was calculated using the median value for each category as a continuous variable in a logistic regression model. We examined unadjusted regression models, then adjusted for age and BMI, and subsequently built a multivariable regression model adjusting for demographic and clinical characteristics. Variables analyzed as potential covariates are shown in Tables 1 and 2. Statistical tests were two-sided and *P* < .05 was the threshold for statistical significance. All statistical analyses were conducted using Stata 12.0 (College Station, TX).

RESULTS

Mailed survey results

We identified 531 potentially eligible participants using the fellow surgical case logs and ICD-9 codes. Among the 531 mailed letters, we had a 43% response rate. Sixty-seven

potentially eligible participants were not interested in participating in our study, and 213 potentially eligible participants did not respond to either the letter or the mailed survey. There were 19 letters returned by the post office, labeled as undeliverable, and an additional seven people had died. A total of 225 participants returned surveys, and 12 were subsequently identified as not meeting inclusion criteria (i.e., 10 diagnosed with cancer before 2006, and two misclassified (diagnosed with other gynecologic cancers)). The remaining 213 eligible participants replied to our survey and contributed data to the study analyses.

Participant Characteristics

Demographic characteristics of the study participants are depicted in Table 1. The age of the 213 participants ranged from 29–94 years. The majority of participants reported being white, married, high school graduates, and retired or working full time. Clinical characteristics of the study participants are depicted in Table 2. Participants were commonly diagnosed with stage I endometroid adenocarcinoma, and treated with surgery. The BMI of study participants ranged from 14–67 kg/m². The neuropathy impairment index indicated women had minimal symptoms in their hands and feet. A small proportion of study participants were previously diagnosed with conditions that may clinically resemble LLL symptoms including congestive heart failure (4%), PAD (2%), and diabetes mellitus (14%). These prior comorbidities were not associated with LLL symptom classification (*P* values greater than .05).

Characteristics between participants with versus without LLL

Among the 213 participants, 77 (36%) reported 5 symptoms on the GCLQ and were classified as having LLL. There existed no significant difference in demographic and clinical characteristics between women classified as having LLL versus not having LLL.

LLL symptoms by level of physical activity and walking distance

Among the 213 study participants, 40%, 13%, 13%, and 35% reported participating in <3.0, 3.0–8.9, 9.0–17.9, and 18.0 MET-hrs·wk⁻¹ of PA, respectively (Table 3). The odds of LLL decreased as MET-hrs·wk⁻¹ of PA increased (*P*_{trend} = .003). Compared with participants who reported <3 MET-hrs·wk⁻¹ of PA, participants who reported 18.0 MET-hrs·wk⁻¹ of PA had an OR of 0.32 (95% CI: 0.15–0.69). The most common PA reported was walking (42%), aerobic gym-based activities including the recumbent bicycle and elliptical machine (11%), and swimming (8%).

Among the 213 study participants, 36%, 26%, and 38% reported walking <4.0, 4.0–11.9, and 12 blocks·d⁻¹, respectively (Table 3). The odds of LLL decreased as the blocks·d⁻¹ of walking increased (*P*_{trend} < .0001). Compared with participants who reported <4.0 blocks·d⁻¹ of walking, participants who reported 12 blocks·d⁻¹ of walking had a multivariable adjusted OR of 0.19 (95% CI: 0.09–0.43).

Among the 213 study participants, 26%, 22%, and 52% reported a BMI of <25.0, 25.0–29.9, and 30.0 kg/m², respectively (Table 3). The odds of LLL did not change as BMI increased (*P*_{trend} = .85).

We also assessed the joint effects of BMI with PA, and BMI with walking (Table 4). Although the interaction for BMI and PA was not statistically significant ($P_{interaction} = .27$), stratified analyses suggested the association of PA and LLL existed only among women with a BMI $<30 \text{ kg/m}^2$ ($P_{trend} = .007$), compared to women with a BMI $\geq 30 \text{ kg/m}^2$ ($P_{trend} = .47$). The interaction for BMI and walking was not statistically significant ($P_{interaction} = .56$), and stratified analyses suggested the association of walking and LLL was similar among women with a BMI $<30 \text{ kg/m}^2$ ($P_{trend} = .007$), compared to women with a BMI $\geq 30 \text{ kg/m}^2$ ($P_{trend} = .03$).

DISCUSSION

The major findings of this study are 36% of uterine cancer survivors in our sample have LLL based on symptom self-report, and uterine cancer survivors who participate in more PA and walking are less likely to report symptoms sufficient for a diagnosis of LLL. These findings, particularly PA participation, were more pronounced among women with a BMI $<30 \text{ kg/m}^2$. BMI was not independently associated with LLL. Our data now provide evidence linking PA, walking, and LLL. Among uterine cancer survivors who engage in the highest level of PA or walking, there were 68% and 81% reduced odds of reporting LLL, compared to uterine cancer survivors who engage in the lowest levels of PA or walking, respectively.

While these findings are promising, they should be interpreted as preliminary. The major limitation of this study is the cross-sectional design which it is impossible to determine the direction of any causal association. It is plausible that uterine cancer survivors who engage in more PA or walking subsequently experience fewer LLL-related symptoms. Conversely, it is plausible that uterine cancer survivors with more severe LLL symptoms may be physically or psychologically unable to engage in PA or walking. BMI in our study was self-reported, which may be subject to bias. However, BMI is highly correlated with objective measures of height and weight, and is appropriate for epidemiologic studies (29). PA in our study was self-reported. While self-reported PA is valid and correlated with objective measures of PA (47), 47% of participants in our study reported meeting physical activity guidelines (i.e., $9.0 \text{ MET-hrs-wk}^{-1}$), whereas only approximately 10% of US adults meet such guidelines (48). Therefore, it is plausible participants in our study may have over reported their PA, consistent with the hypothesis that uterine cancer survivors are less physically active and have a higher BMI. Another limitation to our study was the omission of question #20 in the swelling-general domain of the GCLQ survey. The current gold-standard method to diagnosis LLL is the circumferential measures of the lower limbs and a clinical examination. However, this method has not been adopted for use in routine clinical care (8). Our method to assess LLL relied on symptoms using a self-report questionnaire that was validated against circumferential measures of the lower limbs, and had excellent psychometric characteristics (10).

The optimal exercise modality to reduce LLL symptom burden is unknown. This is a result of a limited understanding of the physiologic complexities of LLL. People with PAD use treadmill walking as a rehabilitative modality to improve leg claudication symptoms (17). Resistance training has also been recommended as a useful adjunct to treadmill walking for

people with PAD (31). Among 156 people with PAD who were randomized to 24-weeks of aerobic treadmill walking, lower extremity resistance training, or control, people in the aerobic treadmill walking group had significantly improved 6-minute walking distance times, and improved time to onset of leg symptoms, compared to those in the control group (31). The weight-lifting group improved quality of life, and stair climbing ability compared to the control group (31). The three-arm design such as that described above may be the most efficient approach to test the safety and efficacy of various exercise modalities among uterine cancer survivors with LLL.

Recent evidence has emerged that weight-lifting exercise is a safe and efficacious modality for upper-extremity lymphedema among breast cancer survivors (36, 39). Among breast cancer survivors with upper-extremity lymphedema, weight-lifting resulted in a 50% risk reduction of lymphedema exacerbations requiring medical treatment, and a significant reduction in the severity of self-reported lymphedema symptoms over one-year (36). Despite physiologic differences between upper-extremity lymphedema and LLL, a hypothesis would be to examine the use of weight-lifting among cancer survivors with LLL. A small pilot study of weight-lifting has been conducted among 10 cancer survivors with LLL (24). After 5-months of weight-lifting, participants had improvements in muscular strength and physical functioning ($P < .05$) compared to baseline values. There were no significant changes in objective LLL leg volume when measured using perometry. Two of the 10 participants developed cellulitis, a bacterial infection requiring broad-spectrum antibiotics during the study. Therefore, the safety and efficacy of weight-lifting among cancer survivors with LLL remains to be elucidated (24).

Daily voluntary walking has been demonstrated to correlate with the distance women can walk without difficulty (6). Further, a program that increases voluntary walking may have implications for disability, mortality, and health care costs (18). Compared to those who could walk ¼ mile without difficulty, those who had difficulty or were unable to walk ¼ mile were at 1.57- or 2.73-fold higher risk for mortality among 5,895 community-dwelling older adults, respectively. A dose-response relationship between ability to walk ¼ mile and medical care costs and hospitalizations has been observed: as ability to walk ¼ mile decreases, medical care costs and hospitalizations may increase proportionally (18). Therefore, future intervention studies designed to improve functional mobility among uterine cancer survivors with LLL may provide a multitude of benefits including reduced disability, mortality, and health care costs. To this end, randomized controlled trials designed to reduce disability among uterine cancer survivors with LLL should consider these outcomes as being important on patient and policy levels (4).

As noted in the introduction, if LLL precludes women participating in the recommended levels of PA (i.e., if reverse-causality exists), then our findings have clinical implications. For example, it is important for clinicians to know that the ACSM guidelines for exercise among cancer survivors (38) may need to be tailored to allow women with LLL to engage in PA that is safe, feasible, and efficacious to promote their physical and mental health. Future cohort studies that measure PA after diagnosis and follow participants longitudinally until incident LLL diagnosis would help to delineate the temporal relationship of PA with LLL. In the absence of reverse-causality, PA may be an intervention to prevent LLL among those

at risk for developing LLL, and to reduce symptom burden among those with LLL. The potential utility of PA across the LLL spectrum—from prevention to treatment—is consistent with other forms of lymphedema such as upper-extremity lymphedema among breast cancer survivors. For example, weight-lifting reduces the risk of incident upper-extremity lymphedema by 36% among those at-risk for developing upper-extremity lymphedema (37), and reduces the need for complete decongestive treatment by 53% among those with upper-extremity lymphedema (36). Our hypothesis is that PA would have similar effects on LLL like that of upper-extremity lymphedema. The remaining questions to be answered include: 1) the temporal sequence of PA and incident LLL; 2) the efficacy and safety of PA to prevent and/or treat LLL; and 3) the type of PA modality (i.e., aerobic, resistance, neuromuscular) that should be prescribed to maximize LLL outcomes and promote the health and longevity of uterine cancer survivors.

Conclusion

Among uterine cancer survivors, approximately 36% have LLL using a validated symptom self-report (10). Participation in higher levels of PA or walking was associated with reduced proportions of LLL in dose-response fashion, such that the group who engaged in the highest levels of PA or walking reported the smallest proportion of LLL cases. These findings should be interpreted as preliminary and be subject to investigation in future studies.

Acknowledgments

The senior author, KH Schmitz, has received funding from the National Institutes of Health unrelated to this work. The results of this study do not constitute endorsement by ACSM. This work was completed without funding.

REFERENCES

1. Abu-Rustum NR, Alektiar K, Iasonos A, Lev G, Sonoda Y, Aghajanian C, Chi DS, Barakat RR. The incidence of symptomatic lower-extremity lymphedema following treatment of uterine corpus malignancies: a 12-year experience at Memorial Sloan-Kettering Cancer Center. *Gynecol Oncol*. 2006; 103(2):714–718. [PubMed: 16740298]
2. Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, O'Brien WL, Bassett DR Jr, Schmitz KH, Emplainscourt PO, Jacobs DR Jr, Leon AS. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc*. 2000; 32(9 Suppl):S498–S504. [PubMed: 10993420]
3. Ainsworth BE, Leon AS, Richardson MT, Jacobs DR, Paffenbarger RS Jr. Accuracy of the College Alumni Physical Activity Questionnaire. *J Clin Epidemiol*. 1993; 46(12):1403–1411. [PubMed: 8263567]
4. Cancer Program Standards 2012: Ensuring Patient-Centered Care [Internet]. Available from: www.facs.org/cancer/clp/cpstandards.pdf.
5. Bergmark K, Avall-Lundqvist E, Dickman PW, Henningsohn L, Steineck G. Lymphedema and bladder-emptying difficulties after radical hysterectomy for early cervical cancer and among population controls. *Int J Gynecol Cancer*. 2006; 16(3):1130–1139. [PubMed: 16803496]
6. Brach JS, Simonsick EM, Kritchevsky S, Yaffe K, Newman AB. Health, Aging and Body Composition Study Research Group. The association between physical function and lifestyle activity and exercise in the health, aging and body composition study. *J Am Geriatr Soc*. 2004; 52(4):502–509. [PubMed: 15066063]
7. Brorson H, Svensson H. Skin blood flow of the lymphedematous arm before and after liposuction. *Lymphology*. 1997; 30(4):165–172. [PubMed: 9476248]

8. Brown JC, Chu CS, Cheville AL, Schmitz KH. The prevalence of lymphedema symptoms among survivors of long-term cancer with or at risk for lower limb lymphedema. *Am J Phys Med Rehabil.* 2013; 92(3):223–231. [PubMed: 23069748]
9. Brown, JC.; Winters-Stone, K.; Lee, A.; Schmitz, KH. *Comprehensive Physiology.* John Wiley & Sons, Inc; 2012. Cancer, Physical Activity, and Exercise.
10. Carter J, Raviv L, Appollo K, Baser RE, Iasonos A, Barakat RR. A pilot study using the Gynecologic Cancer Lymphedema Questionnaire (GCLQ) as a clinical care tool to identify lower extremity lymphedema in gynecologic cancer survivors. *Gynecol Oncol.* 2010; 117(2):317–323. [PubMed: 20163847]
11. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *J Clin Epidemiol.* 1994; 47(11):1245–1251. [PubMed: 7722560]
12. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis.* 1987; 40(5):373–383. [PubMed: 3558716]
13. Daroczy J. Pathology of lymphedema. *Clin Dermatol.* 1995; 13(5):433–444. [PubMed: 8665454]
14. Ferrandina G, Mantegna G, Petrillo M, Fuoco G, Venditti L, Terzano S, Moruzzi C, Lorusso D, Marcellusi A, Scambia G. Quality of life and emotional distress in early stage and locally advanced cervical cancer patients: a prospective, longitudinal study. *Gynecol Oncol.* 2012; 124(3):389–394. [PubMed: 22035809]
15. Finnane A, Hayes SC, Obermair A, Janda M. Quality of life of women with lower-limb lymphedema following gynecological cancer. *Expert Rev Pharmacoecon Outcomes Res.* 2011; 11(3):287–297. [PubMed: 21671698]
16. Franks PJ, Moffatt CJ, Doherty DC, Williams AF, Jeffs E, Mortimer PS. Assessment of health-related quality of life in patients with lymphedema of the lower limb. *Wound Repair Regen.* 2006; 14(2):110–118. [PubMed: 16630098]
17. Gardner AW, Poehlman ET. Exercise rehabilitation programs for the treatment of claudication pain. A meta-analysis. *JAMA.* 1995; 274(12):975–980. [PubMed: 7674529]
18. Hardy SE, Kang Y, Studenski SA, Degenholtz HB. Ability to walk 1/4 mile predicts subsequent disability, mortality, and health care costs. *J Gen Intern Med.* 2011; 26(2):130–135. [PubMed: 20972641]
19. Hiatt WR, Regensteiner JG, Hargarten ME, Wolfel EE, Brass EP. Benefit of exercise conditioning for patients with peripheral arterial disease. *Circulation.* 1990; 81(2):602–609. [PubMed: 2404633]
20. Hiatt WR, Wolfel EE, Meier RH, Regensteiner JG. Superiority of treadmill walking exercise versus strength training for patients with peripheral arterial disease. Implications for the mechanism of the training response. *Circulation.* 1994; 90(4):1866–1874. [PubMed: 7923674]
21. Huang HQ, Brady MF, Cella D, Fleming G. Validation and reduction of FACT/GOG-Ntx subscale for platinum/paclitaxel-induced neurologic symptoms: a gynecologic oncology group study. *Int J Gynecol Cancer.* 2007; 17(2):387–393. [PubMed: 17362317]
22. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin.* 2011; 61(2):69–90. [PubMed: 21296855]
23. Jensen MR, Simonsen L, Karlsmark T, Bulow J. Lymphoedema of the lower extremities--background, pathophysiology and diagnostic considerations. *Clin Physiol Funct Imaging.* 2010; 30(6):389–398. [PubMed: 20718809]
24. Katz E, Dugan NL, Cohn JC, Chu C, Smith RG, Schmitz KH. Weight lifting in patients with lower-extremity lymphedema secondary to cancer: a pilot and feasibility study. *Arch Phys Med Rehabil.* 2010; 91(7):1070–1076. [PubMed: 20599045]
25. Kelly BJ, Frazee TK, Hornik RC. Response rates to a mailed survey of a representative sample of cancer patients randomly drawn from the Pennsylvania Cancer Registry: a randomized trial of incentive and length effects. *BMC Med Res Methodol.* 2010; 10:65. [PubMed: 20630101]
26. Kohl HW, Blair SN, Paffenbarger RS Jr, Macera CA, Kronenfeld JJ. A mail survey of physical activity habits as related to measured physical fitness. *Am J Epidemiol.* 1988; 127(6):1228–1239. [PubMed: 3369421]

27. Langbecker D, Hayes SC, Newman B, Janda M. Treatment for upper-limb and lower-limb lymphedema by professionals specializing in lymphedema care. *Eur J Cancer Care (Engl)*. 2008; 17(6):557–564. [PubMed: 18771539]
28. Lockwood-Rayermann S. Lymphedema in gynecologic cancer survivors: an area for exploration? *Cancer Nurs*. 2007; 30(4):E11–E18. [PubMed: 17666969]
29. McAdams MA, Van Dam RM, Hu FB. Comparison of self-reported and measured BMI as correlates of disease markers in US adults. *Obesity (Silver Spring)*. 2007; 15(1):188–196. [PubMed: 17228047]
30. McCully KK, Halber C, Posner JD. Exercise-induced changes in oxygen saturation in the calf muscles of elderly subjects with peripheral vascular disease. *J Gerontol*. 1994; 49(3):B128–B134. [PubMed: 8169330]
31. McDermott MM, Ades P, Guralnik JM, Dyer A, Ferrucci L, Liu K, Nelson M, Lloyd-Jones D, Van Horn L, Garside D, Kibbe M, Domanchuk K, Stein JH, Liao Y, Tao H, Green D, Pearce WH, Schneider JR, McPherson D, Laing ST, McCarthy WJ, Shroff A, Criqui MH. Treadmill exercise and resistance training in patients with peripheral arterial disease with and without intermittent claudication: a randomized controlled trial. *JAMA*. 2009; 301(2):165–174. [PubMed: 19141764]
32. Meyerhardt JA, Giovannucci EL, Holmes MD, Chan AT, Chan JA, Colditz GA, Fuchs CS. Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol*. 2006; 24(22):3527–3534. [PubMed: 16822844]
33. Meyerhardt JA, Heseltine D, Niedzwiecki D, Hollis D, Saltz LB, Mayer RJ, Thomas J, Nelson H, Whittom R, Hantel A, Schilsky RL, Fuchs CS. Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol*. 2006; 24(22):3535–3541. [PubMed: 16822843]
34. Paffenbarger RS Jr, Hyde RT, Wing AL. Physical activity and incidence of cancer in diverse populations: a preliminary report. *Am J Clin Nutr*. 1987; 45(1 Suppl):312–317. [PubMed: 3799521]
35. Rock CL, Doyle C, Demark-Wahnefried W, Meyerhardt J, Courneya KS, Schwartz AL, Bandera EV, Hamilton KK, Grant B, McCullough M, Byers T, Gansler T. Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin*. 2012
36. Schmitz KH, Ahmed RL, Troxel A, Cheville A, Smith R, Lewis-Grant L, Bryan CJ, Williams-Smith CT, Greene QP. Weight lifting in women with breast-cancer-related lymphedema. *N Engl J Med*. 2009; 361(7):664–673. [PubMed: 19675330]
37. Schmitz KH, Ahmed RL, Troxel AB, Cheville A, Lewis-Grant L, Smith R, Bryan CJ, Williams-Smith CT, Chittams J. Weight lifting for women at risk for breast cancer-related lymphedema: a randomized trial. *JAMA*. 2010; 304(24):2699–2705. [PubMed: 21148134]
38. Schmitz KH, Courneya KS, Matthews C, Demark-Wahnefried W, Galvao DA, Pinto BM, Irwin ML, Wolin KY, Segal RJ, Lucia A, Schneider CM, VON Gruenigen VE, Schwartz AL. American college of sports medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc*. 2010; 42(7):1409–1426. [PubMed: 20559064]
39. Schmitz KH, Troxel AB, Cheville A, Grant LL, Bryan CJ, Gross CR, Lytle LA, Ahmed RL. Physical Activity and Lymphedema (the PAL trial): assessing the safety of progressive strength training in breast cancer survivors. *Contemp Clin Trials*. 2009; 30(3):233–245. [PubMed: 19171204]
40. Simon DA, Dix FP, McCollum CN. Management of venous leg ulcers. *BMJ*. 2004; 328(7452):1358–1362. [PubMed: 15178615]
41. Spillane AJ, Saw RP, Tucker M, Byth K, Thompson JF. Defining lower limb lymphedema after inguinal or ilio-inguinal dissection in patients with melanoma using classification and regression tree analysis. *Ann Surg*. 2008; 248(2):286–293. [PubMed: 18650640]
42. Stanton AW, Levick JR, Mortimer PS. Cutaneous vascular control in the arms of women with postmastectomy oedema. *Exp Physiol*. 1996; 81(3):447–464. [PubMed: 8737078]
43. Symvoulakis EK, Anyfantakis DI, Lionis C. Primary lower limb lymphedema: a focus on its functional, social and emotional impact. *Int J Med Sci*. 2010; 7(6):353–357. [PubMed: 20975845]

44. Tada H, Teramukai S, Fukushima M, Sasaki H. Risk factors for lower limb lymphedema after lymph node dissection in patients with ovarian and uterine carcinoma. *BMC Cancer*. 2009; 9:47. [PubMed: 19193243]
45. Tiwari A, Cheng KS, Button M, Myint F, Hamilton G. Differential diagnosis, investigation, and current treatment of lower limb lymphedema. *Arch Surg*. 2003; 138(2):152–161. [PubMed: 12578410]
46. Todo Y, Yamamoto R, Minobe S, Suzuki Y, Takeshi U, Nakatani M, Aoyagi Y, Ohba Y, Okamoto K, Kato H. Risk factors for postoperative lower-extremity lymphedema in endometrial cancer survivors who had treatment including lymphadenectomy. *Gynecol Oncol*. 2010; 119(1):60–64. [PubMed: 20638109]
47. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008; 40(1):181–188. [PubMed: 18091006]
48. Tucker JM, Welk GJ, Beyler NK. Physical activity in U.S.: adults compliance with the Physical Activity Guidelines for Americans. *Am J Prev Med*. 2011; 40(4):454–461. [PubMed: 21406280]

Table 1

Demographic characteristics stratified by lower limb lymphedema (LLL) status

Variable	Total Sample (n=213)	LLL (n=77)	No LLL (n=136)	<i>P</i> ^a
Age — yr	63.6±10.6	62.7±10.2	64.0±10.9	.24
Marital status — no. (%)				.54
Never married	20 (9%)	8 (10%)	12 (9%)	
Married	128 (60%)	47 (61%)	81 (60%)	
Divorced or separated	31 (15%)	8 (10%)	23 (17%)	
Widowed	33 (16%)	14 (18%)	19 (14%)	
Self-reported race — no. (%)				.66
White	177 (84%)	66 (86%)	111 (82%)	
Black	28 (13%)	8 (10%)	20 (15%)	
Other	7 (3%)	3 (4%)	4 (3%)	
Education — no. (%)				.15
High school or less	46 (22%)	16 (21%)	30 (22%)	
Some college	51 (24%)	13 (17%)	38 (28%)	
College degree or more	114 (54%)	47 (62%)	67 (50%)	
Employment — no. (%)				.61
Retired	94 (45%)	35 (46%)	59 (44%)	
Unemployed	7 (3%)	4 (5%)	3 (2%)	
Homemaker	16 (8%)	6 (8%)	10 (7%)	
Other	14 (7%)	3 (4%)	11 (8%)	
Full time	80 (38%)	28 (37%)	52 (38%)	

^aBy Wilcoxon rank sum, or Fishers Exact test. Values may not sum to 213 or 100% due to rounding error and item non-response.

Table 2

Clinical characteristics stratified by lower limb lymphedema (LLL) status

Variable	Total Sample (n=213)	LLL (n=77)	No LLL (n=136)	<i>P</i> ^a
Pathology type — no. (%)				.73
Endometroid Adenocarcinoma	158 (75%)	57 (74%)	101 (75%)	
Papillary serous or Clear Cell	35 (17%)	14 (18%)	21 (16%)	
Sarcoma	8 (4%)	2 (3%)	6 (4%)	
Carcinosarcoma	8 (4%)	2 (3%)	6 (4%)	
Other (Undifferentiated)	3 (1%)	2 (3%)	1 (1%)	
Stage — no. (%)				.76
1	157 (74%)	57 (74%)	100 (74%)	
2	13 (6%)	5 (6%)	8 (6%)	
3	26 (12%)	9 (12%)	17 (13%)	
4	5 (2%)	3 (4%)	2 (1%)	
Unknown	12 (6%)	3 (4%)	9 (7%)	
Treatment Modalities — no. (%)				.51
Surgery	100 (47%)	40 (52%)	60 (44%)	
Surgery, Chemotherapy	37 (17%)	12 (16%)	25 (18%)	
Surgery, Radiation	47 (22%)	13 (17%)	34 (25%)	
Surgery, Chemotherapy, Radiation	22 (10%)	10 (13%)	12 (9%)	
None or Unknown	7 (3%)	2 (3%)	5 (4%)	
No. of nodes removed	8.9±10.2	7.6±9.8	9.7±10.4	.10
Neuropathy Impairment Index ^b	3.3±3.6	2.8±3.4	3.6±3.7	.17
Time since diagnosis — no. (%)				.74
0–2 yrs	69 (32%)	23 (30%)	46 (34%)	
3–4 yrs	94 (44%)	34 (44%)	60 (44%)	
5–6 yrs	50 (23%)	20 (26%)	30 (22%)	
BMI — kg/m ²	31.1±8.9	30.8±9.3	31.2±8.7	.66
Prior comorbidities ^c — no. (%)				
Congestive heart failure	8 (4%)	2 (3%)	6 (5%)	.50
Peripheral artery disease	5 (2%)	3 (4%)	2 (2%)	.27
Diabetes mellitus	29 (14%)	11 (15%)	18 (14%)	.82

^aBy Wilcoxon rank sum, or Fishers Exact test. Values may not sum to 213 or 100% due to rounding error and item non-response.^bThe neuropathy impairment index is scored from 0 to 16 with higher scores representing more severe neuropathy in the hands and feet.^cComorbidities associated with symptoms in the lower limbs that may resemble lower limb lymphedema.

Table 3

Cases of lower limb lymphedema (LLL) by level of physical activity, walking distance, and BMI

Physical Activity (MET-hrs-wk ⁻¹)	Total in Category	Cases of LLL	Model 1 ^a OR (95% CI)	Model 2 ^b OR (95% CI)	Model 3 ^c OR (95% CI)
<3.0	85	39 (46%)	1 — Referent	1 — Referent	1 — Referent
3.0–8.9	27	12 (44%)	0.94 (0.40–2.25)	0.92 (0.37–2.28)	0.75 (0.28–1.99)
9.0–17.9	27	9 (33%)	0.59 (0.24–1.46)	0.31 (0.24–1.52)	0.50 (0.17–1.45)
18.0	74	17 (23%)	0.35 (0.18–0.70)	0.36 (0.18–0.73)	0.32 (0.15–0.69)
<i>P</i> _{trend}	—	—	.002	.004	.003
Walking (blocks·d ⁻¹)	Total in Category	Cases of LLL	Model 1 ^a OR (95% CI)	Model 2 ^b OR (95% CI)	Model 3 ^c OR (95% CI)
<4.0	75	41 (55%)	1 — Referent	1 — Referent	1 — Referent
4.0–11.9	53	15 (28%)	0.33 (0.15–0.69)	0.31 (0.14–0.68)	0.24 (0.10–0.57)
12.0	78	18 (23%)	0.25 (0.12–0.50)	0.23 (0.11–0.48)	0.19 (0.09–0.43)
<i>P</i> _{trend}	—	—	<.0001	<.0001	<.0001
BMI (kg/m ²)	Total in Category	Cases of LLL	Model 1 ^a OR (95% CI)	Model 2 ^{b,d} OR (95% CI)	Model 3 ^{c,d} OR (95% CI)
<25.0	56	20 (36%)	1 — Referent	1 — Referent	1 — Referent
25.0–29.9	47	17 (36%)	1.02 (0.45–2.29)	1.01 (0.45–2.27)	1.38 (0.55–3.44)
30.0	110	40 (36%)	1.03 (0.53–2.01)	1.01 (0.52–1.98)	1.12 (0.52–2.43)
<i>P</i> _{trend}	—	—	.94	.97	.85

^aModel 1 is the crude (unadjusted) odds ratio and 95% confidence interval.

^bModel 2 is the age and BMI adjusted odds ratio and 95% confidence interval.

^cModel 3 is the fully adjusted (multivariable) odds ratio and 95% confidence interval, controlling for age, BMI, race, cancer stage, type of treatment received, years since diagnosis, and diabetes.

^dBMI excluded as a covariate in these analyses.

Table 4

Multivariable-adjusted cases of lower limb lymphedema (LLL) by level of physical activity and walking distance, stratified by BMI^a

Physical Activity (MET-hrs·wk ⁻¹)	BMI <30 kg/m ² (n = 103)	BMI ≥ 30 kg/m ² (n=110)	<i>P</i> _{interaction}
<3.0	1 — Referent	1 — Referent	.27
3.0–8.9	1.01 (0.20–5.10)	0.65 (0.16–2.70)	
9.0–17.9	0.24 (0.03–1.66)	0.84 (0.19–3.64)	
18.0	0.21 (0.06–0.70)	0.63 (0.21–1.92)	
<i>P</i> _{trend}	.007	.47	
Walking (blocks·d ⁻¹)	BMI <30 kg/m ² (n = 103)	BMI ≥ 30 kg/m ² (n=110)	<i>P</i> _{interaction}
<4.0	1 — Referent	1 — Referent	.56
4.0–11.9	0.20 (0.05–0.87)	0.26 (0.08–0.90)	
12.0	0.15 (0.04–0.61)	0.27 (0.09–0.83)	
<i>P</i> _{trend}	.007	.03	

^a Fully adjusted (multivariable) odds ratio and 95% confidence interval controlling for age, race, cancer stage, type of treatment received, years since diagnosis, and diabetes.