Published in final edited form as:

Gastroenterology. 2014 August ; 147(2): 473–484.e2. doi:10.1053/j.gastro.2014.04.042.

Interleukin 17-Producing $\gamma\delta$ T Cells Promote Hepatic Regeneration in Mice

Raghavendra Rao^{1,*}, Christopher S. Graffeo^{2,*}, Rishabh Gulati^{1,*}, Mohsin Jamal¹, Suchithra Narayan², Constantinos Zambirinis¹, Rocky Barilla², Michael Deutsch¹, Stephanie Greco¹, Atsuo Ochi¹, Lena Tomkötter¹, Reuven Blobstein¹, Antonina Avanzi¹, Daniel M. Tippens¹, Yisroel Gelbstein¹, Eliza Van Heerden¹, and George Miller^{1,2}

¹S. Arthur Localio Laboratory, Department of Surgery, New York University School of Medicine, 550 First Avenue, New York, NY 10016

²S. Arthur Localio Laboratory, Department of Cell Biology, New York University School of Medicine, 550 First Avenue, New York, NY 10016

Abstract

Background & Aims—Subsets of leukocytes synergize with regenerative growth factors to promote hepatic regeneration. $\gamma\delta$ T cells are early responders to inflammation-induced injury in a number of contexts. We investigated the role of $\gamma\delta$ T cells in hepatic regeneration using mice with disruptions in *Tcrd* (encodes the T cell receptor δ chain) and *Clec7a* (encodes C-type lectin domain family 7 member a, also known as *DECTIN1*).

Methods—We performed partial hepatectomies on wild-type C57BL/6, CD45.1, *Tcrd*^{−/−}, or *Clec7a*^{−/−} mice. Cells were isolated from livers of patients and mice via mechanical and enzymatic digestion. $\gamma\delta$ T cells were purified by fluorescence-activated cell sorting.

Results—In mice, partial hepatectomy upregulated expression of CCL20 and ligands of Dectin-1, associated with recruitment and activation of $\gamma\delta$ T cells and their increased production of interleukin (IL)17 family cytokines. Recruited $\gamma\delta$ T cells induced production of IL6 by antigen-presenting cells and suppressed expression of interferon γ by natural killer T cells, promoting hepatocyte proliferation. Absence of IL17-producing $\gamma\delta$ T cells or deletion of Dectin-1 prevented development of regenerative phenotypes in subsets of innate immune cells. This slowed liver

© 2014 The American Gastroenterological Association. Published by Elsevier Inc. All rights reserved.

Address correspondence to: George Miller, MD, Departments of Surgery and Cell Biology, New York University School of Medicine, 450 East 29th Street, East River Science Park, Room 843F, New York, NY 10016, Tel: (646) 501-2208, george.miller@nyumc.org.

*RR, CSG, and RG contributed equally to this work

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Author Contributions: Rao (acquisition of data; analysis and interpretation), Graffeo (analysis and interpretation; manuscript preparation; statistical analysis), Gulati (acquisition of data; analysis and interpretation), Narayan (acquisition of data), Zambirinis (analysis and interpretation; manuscript preparation), Barilla (acquisition of data; manuscript preparation), Deutsch (acquisition of data), Greco (acquisition of data), Ochi (acquisition of data; critical revision), Jamal (acquisition of data), Tomkötter (acquisition of data), Blobstein (acquisition of data), Avanzi (acquisition of data), Tippens (acquisition of data), Gelbstein (statistical analysis), Van Heerden (in vivo experiments; animal breeding), Miller (study concept and design; study supervision; critical revision).

Conflicts: The authors have no conflicts of interest to declare.

regeneration and was associated with reduced expression of regenerative growth factors and cell cycle regulators. Conversely, exogenous administration of IL17 family cytokines or Dectin-1 ligands promoted regeneration. More broadly, we found that $\gamma\delta$ T cells are required for inflammatory responses mediated by IL17 and Dectin-1.

Conclusions— $\gamma\delta$ T cells regulate hepatic regeneration by producing IL22 and IL17, which have direct mitogenic effects on hepatocytes and promote a regenerative phenotype in hepatic leukocytes, respectively. Dectin-1 ligation is required for $\gamma\delta$ T cells to promote hepatic regeneration.

Keywords

Inflammation; NKT cells; Kupffer cells; IL-22

Introduction

Liver regeneration has attracted intense scientific attention since the early 20th century owing to its importance in maintaining liver function after hepatic resection as well as in rescuing hepatic function in acute and chronic liver disease. Previous work has shown the importance of soluble mediators including TNF- α , IL-6, and IL-22 in driving hepatic regeneration¹. In particular, both IL-6 and IL-22 induce phosphorylation of STAT3 in hepatocytes, which is directly mitogenic and pro-proliferative. Conversely, IFN γ has inhibitory effects on hepatocyte proliferation. Macrophage secreted IL-6 is an early event in promoting liver regeneration¹. Dendritic cells regulate hepatic IFN γ levels and thereby also positively influence the rate of hepatic regeneration². Although NKT cells have been shown to promote liver regeneration at baseline^{3, 4}, the presence of activated NKT cells suppresses liver regeneration via their secretion of IFN γ ⁵. Nevertheless, a comprehensive understanding of the coordinated innate immune regulation of liver regeneration remains elusive.

$\gamma\delta$ T cells were first described nearly thirty years ago, yet much about their function, including their interactions with neighboring immune-competent and parenchymal cells, remains a mystery. $\gamma\delta$ T cells are distinct from $\alpha\beta$ T cells in that they are activated in a non-antigen-restricted manner via direct cellular or environmental stimuli. Hence, $\gamma\delta$ T cells are more closely aligned to innate rather than adaptive immunity. As such, they are one of the first cellular populations to be activated during an inflammatory response⁶.

We found that $\gamma\delta$ T cells are important orchestrators of hepatic regeneration via production of IL-22 and IL-17, which collectively have direct mitogenic effects on hepatocytes, promote a pro-regenerative phenotype in neutrophils, dendritic cells, and Kupffer cells, and suppress IFN γ production by NKT cells. We found that intra-hepatic Dectin-1 ligands are upregulated after partial hepatectomy, and Dectin-1 ligation is central to $\gamma\delta$ T cell activation and their induction of a pro-regenerative innate immune-phenotype. More generally, we discovered a novel role for Dectin-1 in non-pathogenic sterile inflammation, and revealed the centrality of the Dectin-1– $\gamma\delta$ T cell–IL-17 axis in modulating regenerative inflammation.

Materials & Methods

Animals and model of partial hepatectomy

C57BL/6, TCR $\delta^{-/-}$, and C57BL/6-Trdc^{tm1Mal} mice were purchased from Jackson (Bar Harbor, ME). For selected experiments, CD45.1 mice and C57BL/6 controls were purchased from Taconic (Germantown, NY). Dectin-1^{-/-} mice were a gift from Gordon Brown (University of Aberdeen). Bone marrow chimeric animals were created by irradiating mice (9 Gy) followed by i.v. bone marrow transfer (10^7 cells) as described by us⁷. The 70% partial hepatectomy procedure entailed ligation and removal of the left and median lobes of the liver in 8–12 week-old mice⁸. The sham operation consisted of a midline laparotomy alone. All animal procedures were approved by NYU School of Medicine IACUC.

Human and murine liver cell isolation

Murine hepatic non-parenchymal cells (NPC) were collected as previously described⁹. Briefly, the portal vein was cannulated and infused with 1% Collagenase IV (Sigma, Saint Louis, MO). The liver was then removed, minced, and filtered to obtain single cell suspensions. Hepatocytes were excluded with serial low speed (300 RPM) centrifugation followed by high-speed (1500 RPM) centrifugation to isolate the NPC, which were then further enriched over a 40% Optiprep (Sigma) gradient⁹. Human liver NPCs were isolated using a similar protocol as we have described¹⁰. All studies on human tissue were carried out under an IRB-approved protocol.

Statistics

Data is presented as mean $\pm$ standard error. Comparisons were made using student's t test for paired or unpaired samples. $p < 0.05$ was considered significant.

See supplemental information for additional methods

Results

Liver $\gamma\delta$ T cells express IL-17, IL-22, and Dectin-1 in both mice and humans

Before investigating the phenotypic and functional alterations in liver $\gamma\delta$ T cells during hepatic regeneration, we established their baseline characteristics. Compared with $\gamma\delta$ T cells in the spleen, liver $\gamma\delta$ T cells (Figure S1a) were more activated, expressing elevated NK1.1, CD44, CD54, FasL, CCR6 and lower CD62L (Figure S1b). Conversely, the CD27⁺ $\gamma\delta$ T cell population was more prominent in the spleen (Figure S1b). Liver $\gamma\delta$ T cells also expressed higher IL-17 and IL-22 than their splenic counterparts (Figure S1c) and higher Dectin-1 (Figure S1d). Further, the hepatic $\gamma\delta$ T cell population contained a higher fraction of V γ 4⁺ cells, which are associated with enhanced functional activation¹¹, whereas V γ 1.1⁺ $\gamma\delta$ T cells predominated in the spleen (Figure S1e).

We observed parallel findings in the human liver, where hepatic $\gamma\delta$ T cells were more mature than their counterparts in the peripheral blood, expressing higher CD122, Dectin-1, and CCR6, as well as down-regulating CD62L and CD27 (Figure S2a, b). Further, similar to their murine counterparts, human liver $\gamma\delta$ T cells produced markedly high levels of IL-17 and

IL-22 relative to $\gamma\delta$ T cells in PBMC and compared to other hepatic parenchymal and inflammatory cells (Figure S2c–e).

$\gamma\delta$ T cells expand after partial hepatectomy and are necessary for robust regeneration

To investigate a potential role for $\gamma\delta$ T cells in liver regeneration, we determined whether this population was altered after partial hepatectomy. The number of intra-hepatic $\gamma\delta$ T cells increased sharply within the first 3h after hepatectomy, but returned to baseline by 6h (Figure 1a). Correspondingly, there was a marked increase in hepatic expression of CCL20, a potent $\gamma\delta$ T cell chemokine¹², immediately following hepatectomy on analysis by PCR, immunofluorescence, and flow cytometry (Figure 1b, S3). CCL20 was expressed mostly from hepatic parenchymal cells (Figure 1b, S3). In addition, liver regeneration was associated with further $\gamma\delta$ T cell activation, as their population expressed higher CD44, CD54, and lower V γ 1.1 as compared to hepatic $\gamma\delta$ T cells from control liver (Figure 1c).

Since $\gamma\delta$ T cell populations expand and mature after partial hepatectomy, we postulated that they promote regeneration. To test this, we investigated the rate of liver regeneration in TCR $\delta^{-/-}$ mice. Hepatocyte proliferation was markedly lower in TCR $\delta^{-/-}$ animals compared to WT mice following partial hepatectomy, as measured by multiple proliferative indices including PCNA, Ki67, and BrdU incorporation (Figure 1d). These data suggest that $\gamma\delta$ T cells are necessary for vigorous hepatic regeneration. Accordingly, expression of immediate early genes including c-fos, c-jun, and c-myc, was markedly higher in WT liver compared with TCR $\delta^{-/-}$ liver within the first hour after hepatectomy (Figure 1e). Similarly, Cyclin D1 was expressed at substantially higher levels in WT liver compared to TCR $\delta^{-/-}$ liver at multiple post-hepatectomy time points (Figure 1e, f). Elevations in hepatic expression of hepatocyte growth factor (HGF) were also depressed in absence of $\gamma\delta$ T cells (Figure 1f). Consistent with a depressed rate of regeneration, weight of liver remnant and plasma levels of prothrombin were lower in TCR $\delta^{-/-}$ animals compared with WT at early time points after hepatectomy (Figure S4a, b). WT mice also exhibited lower elevations serum transaminases but elevated serum glucose (Figure S4c–e). WT mice made chimeric using bone marrow derived from TCR $\delta^{-/-}$ animals similarly exhibited a diminished rate of liver regeneration compared with WT mice transferred with WT bone marrow (Figure S5).

Regenerating TCR $\delta^{-/-}$ liver have a dampened inflammatory milieu

A dynamic interplay between inflammatory cytokines and pro-proliferative signaling pathways promote hepatic regeneration¹. To further investigate the mechanistic role of $\gamma\delta$ T cells in regeneration, we studied inflammatory signaling in regenerating TCR $\delta^{-/-}$ liver compared with WT. Liver from TCR $\delta^{-/-}$ mice expressed lower STAT3 and pSTAT3 and exhibited diminished phosphorylation of MAP kinase signaling intermediates within the first 3h after partial hepatectomy (Figure 2a). Similarly, there was absent elevation of Notch-1 expression in liver of regenerating TCR $\delta^{-/-}$ mice (Figure 2a). There was also dampened elevation in hepatic expression of the pro-regenerative cytokines IL-4, IL-6, IL-22, and IL-17 after partial hepatectomy in TCR $\delta^{-/-}$ mice (Figure 2b). Consistent with these findings, serum levels of IL-6 were markedly higher in hepatectomized WT mice compared with TCR $\delta^{-/-}$ (Figure 2c). Conversely, IFN γ , which is associated with inhibited hepatocyte proliferation¹³, was expressed at higher levels in TCR $\delta^{-/-}$ liver after hepatectomy (Figure

2d). Taken together, these data suggest that the absence of $\gamma\delta$ T cells after partial hepatectomy is associated with muted expression of soluble inflammatory mediators and signaling intermediates that promote hepatic regeneration, along with elevated expression of inhibitory cytokines.

$\gamma\delta$ T cells regulate intra-hepatic inflammatory cell recruitment during liver regeneration

Based on our data illuminating the muted inflammatory milieu in regenerating $\text{TCR}\delta^{-/-}$ liver, we postulated that $\gamma\delta$ T cells may affect liver regeneration by influencing the recruitment and activation of neighboring immune subsets. The influx of bulk CD45^{+} hepatic leukocytes was markedly diminished in $\text{TCR}\delta^{-/-}$ mice after partial hepatectomy (Figure 2e). Similarly, the numbers of B220^{+} cells, MPO^{+} cells, and CD68^{+} cells - which have been reported to contribute to hepatic regeneration¹⁴ - were lower in the livers of regenerating $\text{TCR}\delta^{-/-}$ mice compared with WT (Figure 2e). Conversely, the fractions of bulk NKT cells and iNKT cells - whose activation has been associated with inhibiting hepatic regeneration when activated^{5, 13} - were higher in $\text{TCR}\delta^{-/-}$ liver (Figure 2f). Taken together, these data suggest that $\gamma\delta$ T cells are required for the generation of a pro-regenerative intra-hepatic leukocyte composition, whereas absence of $\gamma\delta$ T cells results in a diminished influx of neutrophils, B cells, and Kupffer cells, paired with an expansion of NKT cell populations.

$\gamma\delta$ T cells affect inflammatory cell activation

We postulated that $\gamma\delta$ T cells may affect liver regeneration by not only affecting leukocyte recruitment, but also influencing their activation. Using WT and $\text{TCR}\delta^{-/-}$ mice, we tested the *in vivo* phenotypic activation of intra-hepatic leukocyte subsets with established roles in modulating liver regeneration. We found that NK and NKT cells were more activated in regenerating $\text{TCR}\delta^{-/-}$ liver compared with WT, expressing elevated CD69 and producing higher IFN- γ (Figure 3a, b). This is consistent with reports showing that NKT and NK cells can inhibit liver regeneration via their production of IFN- γ ¹³. To investigate whether the presence of activated NK or NKT cells in $\text{TCR}\delta^{-/-}$ mice definitively contributed to their retarded liver regeneration, we selectively depleted these cellular subsets using a mAb directed against NK1.1¹⁴. Consistent with our hypothesis, depletion of NK and NKT cells partially reversed the depressed rate of liver regeneration in $\text{TCR}\delta^{-/-}$ mice (Figure 3c). Interestingly, regeneration was depressed in WT mice following depletion of non-activated NK and NKT cell populations (Figure 3c), a finding consistent with a recent report suggesting that NK and NKT cells can accelerate hepatic regeneration by upregulating IL-6 and HGF⁴. Further, Kupffer cells and Dendritic cells - which are proregenerative^{1,2} - expressed higher IL-6 in regenerating WT liver as compared to $\text{TCR}\delta^{-/-}$ liver (Figure 3d, e). Taken together, these data suggest that the presence of $\gamma\delta$ T cells affects the activation of varied inflammatory cell subsets with critical roles in modulating liver regeneration.

$\gamma\delta$ T cells influence the activation of hepatic leukocyte subsets via IL-17

To test whether hepatic $\gamma\delta$ T cells can directly induce a pro-regenerative phenotype in neighboring hepatic leukocytes, we performed *in vitro* co-culture experiments. Hepatic $\gamma\delta$ T cells were purified by FACS and co-cultured with equal numbers of NKT cells, Kupffer

cells, DC, or neutrophils. Consistent with our *in vivo* data, $\gamma\delta$ T cells induced diminished activation of NKT cells, modestly lowering their expression of CD44 and CD69 (Figure S6a). Conversely, hepatic $\gamma\delta$ T cells modestly up-regulated expression of MHCII and CD86 on Kupffer cells (Figure S6b), and induced their production of IL-6 (Figure S6c). Both V γ 1.1⁺ and V γ 4⁺ subsets were equally effective activators of Kupffer cells (Figure S6c). Similarly, $\gamma\delta$ T cells moderately activated the surface phenotype of DC (Figure S6d) and neutrophils (Figure S6e). Taken together, these data suggest that liver $\gamma\delta$ T cells can directly influence the generation of a pro-regenerative phenotype in neighboring hepatic innate inflammatory subsets.

We found that hepatic $\gamma\delta$ T cells express elevated IL-17 at baseline in mice (Figure S1d) and humans (Figure S2c–e) which further increased markedly within 3h following hepatectomy (Figure 4a). Moreover, compared with hepatic CD3⁺TCR $\delta^{-/-}$ T cell subsets, CD3⁺CD45⁺ cells, and CD45⁺ cells, a higher proportion of $\gamma\delta$ T cells were IL-17⁺ cells by flow cytometry in human liver (Figure S2c, d) and in the regenerating mouse liver (Figure 4b, c). Since emerging data suggest that IL-17 can modulate intra-hepatic sterile inflammation^{15, 16}, we postulated that $\gamma\delta$ T cells induce a pro-regenerative hepatic inflammatory milieu via IL-17. To test this, leukocytes from WT mice and TCR $\delta^{-/-}$ mice were stimulated with PMA and ionomycin, in the presence or absence of an IL-17 mAb. WT leukocytes expressed higher levels of IL-6 at baseline compared with those from TCR $\delta^{-/-}$ mice (Figure 4d), consistent with our previous observations that $\gamma\delta$ T cells promote the production of pro-regenerative cytokines. Moreover, WT leukocyte concentrates down-regulated IL-6 transcript in the context of IL-17 inhibition, whereas TCR $\delta^{-/-}$ leukocyte suspensions upregulated IL-6 production after IL-17 blockade (Figure 4d). Further, in concert with our previous experiments, WT leukocyte concentrates expressed lower IFN γ compared with inflammatory cell suspensions from TCR $\delta^{-/-}$ mice (Figure 4e). Conversely, IFN γ mRNA was upregulated after IL-17 blockade in WT leukocyte concentrates, but not in inflammatory cell concentrates from TCR $\delta^{-/-}$ mice (Figure 4e). Taken together, these data suggest that IL-17 promotes inflammatory cell expression of high IL-6 and low IFN γ in $\gamma\delta$ T cell-rich leukocyte concentrates, but has the opposite effects in the absence of $\gamma\delta$ T cells.

To specifically investigate whether $\gamma\delta$ T cell-derived IL-17 influences the generation of a pro-regenerative NKT cell phenotype, we examined NKT populations in inflammatory cell suspensions derived from WT and TCR $\delta^{-/-}$ mice. We found that NKT cells produced higher IFN γ in the TCR $\delta^{-/-}$ suspensions (Figure 4f), consistent with our *in vivo* observations of greater NKT cell activation in regenerating TCR $\delta^{-/-}$ liver (Figure 3b). Additionally, blockade of IL-17 increased NKT cell IFN γ expression in $\gamma\delta$ T cell-rich cultures, but not in suspensions deficient in $\gamma\delta$ T cells (Figure 4f). These data suggest that $\gamma\delta$ T cells can inhibit NKT cell activation via IL-17.

IL-17 producing $\gamma\delta$ T cells are necessary for robust liver regeneration

The CD45.1⁺ congenic C57BL/6 mouse sub-strain is genetically identical to the WT strain, except that it carries the differential B cell antigen Ly5.1. However, a recent report found that CD45.1⁺ mice exhibit selective deficiency in IL-17 producing $\gamma\delta$ T cells due to a mutation in Sox13 resulting in their defective development in the neonatal thymus¹⁷. We

confirmed markedly lower hepatic $\gamma\delta$ T cell expression of IL-17 in regenerating CD45.1 mice (Figure S7a). Further, we showed that - consistent with absence of IL-17 producing $\gamma\delta$ T cells - there was almost no IL-17 production in regenerating liver of CD45.1 mice compared with WT (Figure S7b). Therefore, we postulated that CD45.1 mice would have deficient hepatic regeneration. We found that CD45.1 mice exhibited a diminished rate of hepatocyte proliferation after 70% hepatectomy (Figure S7c), reduced elevation in Cyclin D1 expression (Figure S7d), and lower serum levels of IL-6 (Figure S7e). Similarly, administration of a neutralizing IL-17 mAb to WT mice lowered the rate of hepatic regeneration (Figure S7f).

$\gamma\delta$ T cells accelerate liver regeneration by IL-22 production via Dectin-1

IL-22 is an IL-17-family cytokine that directly stimulates hepatocyte proliferation by inducing MAP kinase signaling¹⁸. Innate inflammatory cells reportedly generate IL-22 in a Dectin-1 dependent manner¹⁹. Therefore, we postulated that hepatic $\gamma\delta$ T cells, which express markedly elevated Dectin-1 (Figures S1c, S2b), influence hepatic regeneration via Dectin-1-dependent IL-22 production. We found that IL-22 expression was upregulated in hepatic $\gamma\delta$ T cells in the regenerating liver (Figure 5a). Further, in parallel with IL-17, compared with hepatic CD3⁺TCR $\delta^{-/-}$ T cell subsets, CD3⁻CD45⁺ cells, and CD45⁻ cells, a higher proportion of $\gamma\delta$ T cells were IL-22⁺ cells by flow cytometry (Figure 5b). Dectin-1 expression was also upregulated on $\gamma\delta$ T cells in the regenerating liver compared with control liver (Figure 5c). Co-expression analysis by both flow cytometry and confocal microscopy revealed that IL-22 was specifically expressed on Dectin-1⁺ $\gamma\delta$ T cells (Figure 5d) whereas hepatic $\gamma\delta$ T cells from Dectin-1^{-/-} mice exhibited an approximate 50% reduction in IL-22 production (Figure 5e). Moreover, Dectin-1 ligation induced IL-22 and IL-17 production in hepatic $\gamma\delta$ T cells (Figure 5e). In addition, using a human IgG Fc-Dectin-1-conjugated fusion protein, we found that hepatocytes and inflammatory cells over-express Dectin-1 ligands after partial hepatectomy (Figure S8a, b). Further, the marked elevations in IL-22 expression associated with liver regeneration were diminished in TCR $\delta^{-/-}$ liver (Figure 5f). Similarly, SOCS3 expression was lower in hepatectomized TCR $\delta^{-/-}$ mice compared with WT, consistent with reduced IL-22 signaling in the absence of $\gamma\delta$ T cells (Figure 5f).

To test whether IL-22 production plays a primary role in the capacity of $\gamma\delta$ T cells to accelerate liver regeneration, we administered recombinant mIL-22 to TCR $\delta^{-/-}$ mice coincident with partial hepatectomy. As predicted, IL-22 administration rescued the depressed hepatic regeneration in TCR $\delta^{-/-}$ mice (Figure S8c). Further, in concert with the dependence of IL-22 production on Dectin-1, we found retarded rates of liver regeneration in Dectin-1^{-/-} mice following partial hepatectomy along with a concomitant diminished upregulation of Cyclin D1 (Figure S8d, e).

The Dectin-1 – $\gamma\delta$ T cell – IL-17 axis regulates liver regeneration

To further investigate the link between Dectin-1-mediated IL-17-family cytokine production and $\gamma\delta$ T cell induction of pro-regenerative leukocytes, we stimulated leukocyte suspensions from WT or TCR $\delta^{-/-}$ mice with Zymosan depleted (Zy) to selectively activate Dectin-1 signaling. WT leukocyte concentrates produced higher IL-6 in response to Zy stimulation, as compared to $\gamma\delta$ T cell-depleted leukocyte suspensions (Figure 6a). Moreover, IL-17 blockade

diminished the activation of Zy-stimulated leukocytes in the presence of $\gamma\delta$ T cells, but had the opposite effect in their absence, as measured by both mRNA (Figure 6a) and protein analyses (Figure 6b). To determine whether Dectin-1 activation of $\gamma\delta$ T cells specifically results in a pro-regenerative phenotype in macrophages and DC in an IL-17-dependent manner, we gated on these cellular subsets in Zy-stimulated $\gamma\delta$ T cell-rich and $\gamma\delta$ T cell-depleted leukocyte concentrates and measured their production of IL-6 by intracellular cytokine analysis. In $\gamma\delta$ T cell-rich suspensions, both macrophages and DC upregulated expression of IL-6 in response to Dectin-1 ligation, the effect of which was abrogated by an IL-17 mAb (Figure 6c). Conversely, macrophages and DC did not become activated by Zy in TCR $\delta^{-/-}$ leukocyte concentrates (Figure 6c). By gating on NKT cells and measuring IFN γ expression, we further found that NKT cells markedly diminished their production of IFN γ in response to Zy in WT cultures (Figure 6d). Conversely, in TCR $\delta^{-/-}$ cultures, NKT cells increased IFN γ production in response to Dectin-1 ligand (Figure 6e). Taken together, these data show that Dectin-1-mediated IL-17-dependent inflammatory responses, and specifically their induction of a pro-regenerative phenotype in neighboring leukocyte subsets, is contingent on the presence of $\gamma\delta$ T cells.

To directly test the central role of the Dectin-1 – $\gamma\delta$ T cell – IL-17 axis *in vivo* in the regenerating liver, we administered Zy to WT and CD45.1 mice coincident with partial hepatectomy. We found that Dectin-1 ligation markedly increased Cyclin D1 expression in the liver of regenerating WT mice, but lowered Cyclin D1 expression in the CD45.1 liver (Figure 6f). Similarly, hepatocyte proliferation was accelerated in WT mice but diminished in CD45.1 mice following Zy administration (Figure 6f), consistent with our observation of paradoxical anti-inflammatory and anti-regenerative effects of Dectin-1 ligation in the absence of IL-17-producing $\gamma\delta$ T cells.

Discussion

Our work reveals a novel role for IL-17/IL-22-producing $\gamma\delta$ T cells in governing the inflammatory orchestration of hepatic regeneration by regulating the phenotype and recruitment of diverse hepatic leukocytes (Figure 7). While our correlative data in human systems suggests relative activation of human liver $\gamma\delta$ T cells and production of IL-17 family cytokines, the nomenclature and complexity of $\gamma\delta$ T cell subsets vary from mice to humans as do the specific $\gamma\delta$ T cell receptors for activating ligands suggesting caution before translating our data directly to human hepatic regeneration. We also describe a novel role for Dectin-1 and its cognate ligands in hepatic regeneration, through their influence on the production of IL-17-family cytokines in hepatic $\gamma\delta$ T cells.

Dectin-1 is a member of the C-type lectin receptor family of pattern recognition receptors and is necessary for innate immune responses to fungal pathogens. We found that Dectin-1 ligands are highly expressed on both parenchymal and inflammatory cells within hours after partial hepatectomy, and exogenous Dectin-1 ligand administration accelerates hepatocyte priming in a $\gamma\delta$ T cell- and IL-17-contingent manner. Further experiments using advanced proteomic techniques are necessary for definitive characterization of the precise Dectin-1 ligands expressed during liver regeneration. However, a novel report this year showed for the first time that intermediate filaments such as vimentin can ligate and activate Dectin-1²⁰.

Further, a recent analysis by mass spectrometry of proteins over-expressed in the regenerating liver compared with control showed that filamentous cytoskeletal proteins are markedly over-expressed in the liver during regeneration²¹. Therefore, these families of proteins seem likely candidate endogenous ligands for Dectin-1 activation during liver regeneration. Regardless, our data suggests that Dectin-1 activation may be an attractive strategy to promote hepatic regeneration after surgical resection or for patients with acute or chronic liver disease.

Downstream of Dectin-1, we ascribe a central role for IL-17-family cytokines in liver regeneration. We found that CD45.1 mice, whose $\gamma\delta$ T cells have diminished ability to produce IL-17, exhibit depressed liver regeneration and response to Dectin-1 ligation. Whereas CD4⁺ T helper cell subsets have frequently been considered primary sources of IL-17-related cytokines in neoplastic and inflammatory settings, we found minimal IL-22 or IL-17 secretion from CD3⁺TCR $\delta^{-/-}$ T cells after hepatectomy, while $\gamma\delta$ T cells robustly produced both cytokines in the regenerating liver.

Our experimental results suggest that IL-17 produced by $\gamma\delta$ T cells is critical in inducing a pro-regenerative phenotype in hepatic inflammatory cells including Kupffer cells, DC, and neutrophils, while simultaneously inhibiting NKT cell expansion or activation. Indeed, NKT cells are critical cellular targets of the Dectin-1 – $\gamma\delta$ T cell – IL-17 axis within the regenerating liver. This is exemplified by our findings that depletion of NKT cells *in vivo* accelerates liver regeneration and hepatocyte expression of Cyclin D1 in TCR $\delta^{-/-}$ mice, yet has the inverse effect of slowing liver regeneration in WT mice. This paradoxical role for NKT cells in liver regeneration - contingent on both their activation and their interaction with $\gamma\delta$ T cells - is consistent with recent reports showing that activated NKT cells which express IFN γ inhibit liver regeneration^{5,13}. Conversely, non-activated NK or NKT cells promote liver regeneration in WT mice⁴. It is conceivable that, in the presence of $\gamma\delta$ T cells, the reduced IFN γ synthesis from NKT cells shifts their functional properties from anti-regenerative to pro-regenerative owing to NKT cells simultaneously being sources of IL-4, a pro-regenerative cytokine³.

We found that in addition to IL-17, $\gamma\delta$ T cells are a vital source of IL-22 in the regenerating liver. IL-22 can directly induce hepatocyte proliferation¹⁸. However, the cellular sources of IL-22 during hepatic regeneration had not been previously demonstrated. Since innate inflammatory cells generate IL-22 in a Dectin-1 dependent manner¹⁹, we suspected that hepatic $\gamma\delta$ T cells - which express markedly elevated Dectin-1 - can influence hepatic regeneration via Dectin-1-dependent IL-22 production. We showed that IL-22 expression from $\gamma\delta$ T cells was Dectin-1 dependent. Further, administration of recombinant IL-22 impressively restored the sluggish rate of liver regeneration in TCR $\delta^{-/-}$ mice. IL-22 signals via the JAK/STAT pathway to upregulate numerous pro-regenerative genes, including TNF α and IL-6¹⁸. Accordingly, we observed diminished elevations in SOCS3 and MAP kinase intermediates - which serve as surrogate markers of IL-22 signaling²² - in the regenerating liver of TCR $\delta^{-/-}$ mice. Our observations of Dectin-1-mediated IL-22 expression in liver regeneration parallels its effects in pathogenic contexts, where Dectin-1 ligation has been associated with anti-fungal immunity in the lung via IL-22 production¹⁹.

A novel observation inherent to the mechanism underlying our *in vivo* findings, and emphasized by our *in vitro* studies, is that whereas IL-17 cytokine blockade in the presence of $\gamma\delta$ T cells has the effect of inhibiting a pro-regenerative phenotype in neighboring leukocyte subsets, including down-regulating IL-6 and upregulating IFN γ , it has diametrically opposite effects in the absence of $\gamma\delta$ T cells. These patterns are amplified by Dectin-1 ligation in both *in vitro* co-culture experiments and in our *in vivo* model of hepatic regeneration, suggesting that the IL-17-dependent inflammatory effects of Dectin-1 signaling are contingent on $\gamma\delta$ T cells. As such, we ascribe a novel role for the Dectin-1 – $\gamma\delta$ T cell – IL-17 axis in promoting liver regeneration, characterized by the induction of a pro-regenerative inflammatory milieu following hepatectomy. Our work further suggests a role for Dectin-1 in sterile inflammation and may have broader implications to the importance of $\gamma\delta$ T cells in directing Dectin-1 mediated IL-17-dependent inflammation in both sterile and pathogenic contexts.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Grant Support: This work was supported in-part by National Institute of Health Awards DK085278 (GM), and DK098303 (GM).

References

1. Amemiya H, Kono H, Fujii H. Liver regeneration is impaired in macrophage colony stimulating factor deficient mice after partial hepatectomy: the role of M-CSF-induced macrophages. *J Surg Res.* 2011; 165:59–67. [PubMed: 20031174]
2. Castellana A, Di Leo A, Francavilla R, et al. Functional modification of CD11c+ liver dendritic cells during liver regeneration after partial hepatectomy in mice. *Hepatology.* 2006; 43:807–816. [PubMed: 16557552]
3. DeAngelis RA, Markiewski MM, Kourtzelis I, et al. A complement-IL-4 regulatory circuit controls liver regeneration. *J Immunol.* 2012; 188:641–648. [PubMed: 22184721]
4. Hosoya S, Ikejima K, Takeda K, et al. Innate immune responses involving natural killer and natural killer T cells promote liver regeneration after partial hepatectomy in mice. *Am J Physiol Gastrointest Liver Physiol.* 2013; 304:G293–G299. [PubMed: 23086918]
5. Dong Z, Zhang J, Sun R, et al. Impairment of liver regeneration correlates with activated hepatic NKT cells in HBV transgenic mice. *Hepatology.* 2007; 45:1400–1412. [PubMed: 17523147]
6. Vantourout P, Hayday A. Six-of-the-best: unique contributions of gammadelta T cells to immunology. *Nat Rev Immunol.* 2013; 13:88–100. [PubMed: 23348415]
7. Ochi A, Graffeo CS, Zambirinis CP, et al. Toll-like receptor 7 regulates pancreatic carcinogenesis in mice and humans. *The Journal of clinical investigation.* 2012; 122:4118–4129. [PubMed: 23023703]
8. Mitchell C, Willenbring H. A reproducible and well-tolerated method for 2/3 partial hepatectomy in mice. *Nat Protoc.* 2008; 3:1167–1170. [PubMed: 18600221]
9. Connolly MK, Bedrosian AS, Mallen-St Clair J, et al. In liver fibrosis, dendritic cells govern hepatic inflammation in mice via TNF- α . *J Clin Invest.* 2009; 119:3213–3225. [PubMed: 19855130]
10. Ibrahim J, Nguyen AH, Rehman A, et al. Dendritic cell populations with different concentrations of lipid regulate tolerance and immunity in mouse and human liver. *Gastroenterology.* 2012; 143:1061–1072. [PubMed: 22705178]

11. Kisielow J, Kopf M, Karjalainen K. SCART scavenger receptors identify a novel subset of adult gammadelta T cells. *J Immunol.* 2008; 181:1710–1716. [PubMed: 18641307]
12. Li Z, Burns AR, Miller SB, et al. CCL20, gammadelta T cells, and IL-22 in corneal epithelial healing. *FASEB J.* 2011; 25:2659–2668. [PubMed: 21518851]
13. Sun R, Gao B. Negative regulation of liver regeneration by innate immunity (natural killer cells/interferon-gamma). *Gastroenterology.* 2004; 127:1525–1539. [PubMed: 15521020]
14. Selzner N, Selzner M, Odermatt B, et al. ICAM-1 triggers liver regeneration through leukocyte recruitment and Kupffer cell-dependent release of TNF-alpha/IL-6 in mice. *Gastroenterology.* 2003; 124:692–700. [PubMed: 12612908]
15. Wang X, Sun R, Wei H, et al. High-mobility group box 1 (HMGB1)-Toll-like receptor (TLR)4-interleukin (IL)-23-IL-17A axis in drug-induced damage-associated lethal hepatitis: Interaction of gammadelta T cells with macrophages. *Hepatology.* 2013; 57:373–384. [PubMed: 22821628]
16. Zhao N, Hao J, Ni Y, et al. Vgamma4 gammadelta T cell-derived IL-17A negatively regulates NKT cell function in Con A-induced fulminant hepatitis. *J Immunol.* 2011; 187:5007–5014. [PubMed: 21987663]
17. Gray EE, Ramirez-Valle F, Xu Y, et al. Deficiency in IL-17-committed Vgamma4(+) gammadelta T cells in a spontaneous Sox13-mutant CD45.1(+) congenic mouse substrain provides protection from dermatitis. *Nature immunology.* 2013; 14:584–592. [PubMed: 23624556]
18. Ren X, Hu B, Colletti LM. IL-22 is involved in liver regeneration after hepatectomy. *Am J Physiol Gastrointest Liver Physiol.* 2010; 298:G74–G80. [PubMed: 19875704]
19. Gessner MA, Werner JL, Lilly LM, et al. Dectin-1-dependent interleukin-22 contributes to early innate lung defense against *Aspergillus fumigatus*. *Infect Immun.* 2012; 80:410–417. [PubMed: 22038916]
20. Thiagarajan PS, Yakubenko VP, Elson DH, et al. Vimentin is an endogenous ligand for the pattern recognition receptor Dectin-1. *Cardiovascular research.* 2013; 99:494–504. [PubMed: 23674515]
21. Cao H, Yu J, Xu W, et al. Proteomic analysis of regenerating mouse liver following 50% partial hepatectomy. *Proteome science.* 2009; 7:48. [PubMed: 20040084]
22. Lejeune D, Dumoutier L, Constantinescu S, et al. Interleukin-22 (IL-22) activates the JAK/STAT, ERK, JNK, and p38 MAP kinase pathways in a rat hepatoma cell line. Pathways that are shared with and distinct from IL-10. *The Journal of biological chemistry.* 2002; 277:33676–33682. [PubMed: 12087100]

Figure 1. $\gamma\delta$ T cells expand and activate after partial hepatectomy and are necessary for robust liver regeneration

(a) Changes in the number of $\gamma\delta$ T cells in the remnant liver after partial hepatectomy compared with sham laparotomy were tested at 3, 6, and 24 hours. (b) Expression of CCL20 was compared at baseline and at 3h after partial hepatectomy by multicolor immunofluorescence microscopy. (c) Hepatic CD3⁺TCR $\gamma\delta$ ⁺ cells from partial hepatectomy and sham-operated liver were gated and tested for expression of CD44, CD54, and V γ 1.1⁺ at 3h. A representative experiment and averages of 3 replicates are shown. (d) Livers from WT

and TCR $\delta^{-/-}$ mice were harvested after 70% hepatectomy and tested for nuclear expression of PCNA, Ki67, and BrdU. Representative stained sections from the 36h time point are shown (n=5 mice/time point). **(e)** The relative increase in expression of c-fos, c-jun, and cmc at 1h in hepatectomy-treated WT and TCR $\delta^{-/-}$ livers compared with sham laparotomy-treated controls was tested by PCR. In addition, Cyclin D1 mRNA expression was tested at various time points in partial hepatectomy and sham-treated WT and TCR $\delta^{-/-}$ livers. **(f)** Hepatic expression of Cyclin D1 and HGF were assayed at 3h by Western blotting. Each experiment was repeated at least 3 times (*p<0.05, **p<0.01, ***p<0.001).

Figure 2. $\gamma\delta$ T cell depletion diminishes expression of pro-regenerative inflammatory mediators and modulates leukocyte expansion in the regenerating liver

(a) Expression of p-STAT3, STAT3, p-p38, p-Erk1/2, Erk1/2, Notch-1, and β -actin were tested by Western blotting within 3h after surgery in partial hepatectomy and sham-laparotomy treated WT and TCR $\delta^{-/-}$ mice. (b) Hepatic expression of IL-4, IL-6, IL-22, and IL-17 were tested in hepatectomized WT and TCR $\delta^{-/-}$ mice at 1 and 3 hours and compared with sham-laparotomy controls by PCR. (c) Serum IL-6 was measured in WT and TCR $\delta^{-/-}$ mice at 48h after partial hepatectomy. (d) IFN γ expression was compared at 3h in

regenerating WT and TCR $\delta^{-/-}$ liver by PCR. **(e)** WT and TCR $\delta^{-/-}$ mice were sacrificed 36h after partial hepatectomy. Liver sections were stained using antibodies directed against CD45 (40 $\times$), B220 (40 $\times$), MPO (40 $\times$), and CD68 (20 $\times$). The fraction of positive cells per HPF for each respective marker is shown. **(f)** The fraction of NK1.1⁺CD3⁻ NK cells, NK1.1⁺CD3⁺ bulk NKT cells, and NK1.1⁺CD1d tetramer⁺ iNKT cells were determined using flow cytometry in WT and TCR $\delta^{-/-}$ mice at 36h after partial hepatectomy. Representative data and averages of 3 replicates are shown. Experiments were repeated 3 times using 3–5 mice per group (*p<0.05, ***p<0.001).

Figure 3. $\gamma\delta$ T cells influence the pro-regenerative phenotype in hepatic inflammatory cells
(a) NK cells and **(b)** NKT cells harvested from the regenerating liver of WT and TCR $\delta^{-/-}$ mice at 36h were gated and tested for expression of CD69 and IFN γ . **(c)** Hepatocyte proliferation was tested by expression of PCNA and Ki67 at 36h in WT and TCR $\delta^{-/-}$ mice depleted of NK1.1 $^{+}$ cells or mock depleted (n=3/group). **(d)** CD11c $^{-}$ F480 $^{+}$ Kupffer cells and **(e)** CD11c $^{+}$ MHCII $^{+}$ DC were gated from the regenerating liver of WT and TCR $\delta^{-/-}$ mice at 36h and tested for expression of IL-6. Representative data and averages of replicates are shown. Each experiment was repeated at least 3 times (*p<0.05, ***p<0.001).

Figure 4. $\gamma\delta$ T cells induce a pro-regenerative phenotype in hepatic inflammatory cells in an IL-17 dependent manner

(a) IL-17 expression was compared at 3h by intracellular cytokine analysis in $\gamma\delta$ T cells from hepatectomy versus sham-operation liver in WT mice. Averages of quadruplicates is shown.

(b) CD3⁺ T cells from regenerating WT liver were gated and tested for co-expression of TCRγδ and IL-17. Similarly, CD3⁻ whole liver cellular concentrates were analyzed using mAbs directed against CD45 and IL-17. Based on triplicate experiments, the fraction of $\gamma\delta$ T cells, $\alpha\beta$ T cells, hepatic innate inflammatory cells, and hepatic parenchymal expression of

IL-17 is shown. **(c)** Liver NPC were harvested at 3h from regenerating murine liver of C57BL/6-Trdc^{tm1Mal}/J mice in which the $\gamma\delta$ T cells express GFP and stained for IL-17 (red) and DAPI (blue). A representative overlay image is shown. **(d, e)** Leukocyte concentrates from WT and TCR $\delta^{-/-}$ mice were stimulated with PMA-Ionomycin and tested for (d) IL-6 and (e) IFN γ expression by PCR, both alone or in the context of IL-17 blockade (*p<0.05, **p<0.01). **(f)** Leukocyte concentrates from WT and TCR $\delta^{-/-}$ mice were stimulated with PMA-Ionomycin. NKT cells were gated and tested for IFN γ expression by intra-cellular cytokine staining in the context of IL-17 blockade or control mAb. Representative data and average of triplicate repeats are shown. All experiments were performed at least 3 times with similar results (*p<0.05, **p<0.01, ***p<0.001).

Figure 5. Dectin-1 dependent IL-22 expression in $\gamma\delta$ T cells is vital for robust liver regeneration
(a) IL-22 mRNA expression was tested at 3h in partial hepatectomy and sham-laparotomy treated liver by PCR. **(b)** CD3⁺ T cells from regenerating WT liver were gated and tested for co-expression of TCR $\gamma\delta$ and IL-22. Similarly, CD3⁻ cells were gated and whole liver cellular concentrates were analyzed using mAbs directed against CD45 and IL-22. Based on quadruplicate experiments, the fraction of $\gamma\delta$ T cells, $\alpha\beta$ T cells, hepatic innate inflammatory cells, and hepatic parenchymal cell expression of IL-22 was calculated. **(c)** Expression of Dectin-1 in liver $\gamma\delta$ T cells was compared in partial hepatectomy and sham-laparotomy

treated mice. Representative data and the change in Dectin-1 expression based on MFI in 4 replicate experiments is shown. **(d)** Hepatic $\gamma\delta$ T cells were tested for co-expression of IL-22 and Dectin-1 by flow cytometry and confocal microscopy. **(e)** IL-22 expression was tested in $\gamma\delta$ T cells from WT and Dectin-1^{-/-} mice at 3h after hepatectomy by intracellular cytokine analysis. Summary data based on 3 replicates is shown. In addition, hepatic $\gamma\delta$ T cells were stimulated with the selective Dectin-1 ligand Zymosan depleted or saline and tested for expression of IL-17 and IL-22. **(f)** IL-22 was measured in the serum of WT and TCR δ ^{-/-} mice at 48h post-hepatectomy and SOCS3 mRNA expression was tested in WT and TCR δ ^{-/-} mice at multiple early time points post-hepatectomy (*p<0.05, ***p<0.001).

Figure 6. Dectin-1 mediated IL-17 cytokine production is necessary for the development of a pro-regenerative phenotype

(a, b) Leukocyte concentrates from WT and TCR $\delta^{-/-}$ mice were stimulated *in vitro* with the selective Dectin-1 ligand Zymosan depleted and tested for expression of (a) IL-6 mRNA or (b) protein in the context of IL-17 blockade or control mAb. (c) DC or macrophages were gated by flow cytometry in WT and TCR $\delta^{-/-}$ leukocyte suspensions and tested for expression of IL-6 by intracellular cytokine staining. (d, e) Similarly, bulk NKT cells were gated in (d) WT and (e) TCR $\delta^{-/-}$ leukocyte suspensions and tested for expression of IFN γ in

the context of IL-17 blockade or control. Experiments were repeated 3 separate times with similar results. (f) Cohorts of hepatectomy-treated WT or CD45.1⁺ mice were additionally treated with Zymosan depleted and tested at 24h for Cyclin D1 expression by PCR and hepatocyte proliferation by PCNA staining (n=4 mice/group; *p<0.05, **p<0.01, ***p<0.001).

Figure 7. Central position of the Dectin-1 – $\gamma\delta$ T cell – IL-17/IL-22 axis in promoting hepatic regeneration