MEETING ABSTRACT

Open Access

Perspectives of epidemiological control of the MRSA subpopulations in hospital patients with sepsis via molecular detection, genetic analysis and following typing of methicilin resistance-carrying mobile genetic elements *SCCmec* in Ukraine

Olena V Moshynets^{1*}, Svitlana U Rymar¹, Olga B Liutko², Nadiia M Oserjanskaja², Gennadii B Kolov², Nadiia P Kisyl³, Rostyslav V Bubnov⁴, Maria Boretska⁴, Dmytro M Irodov¹

From EPMA-World Congress 2013
Brussels, Belgium. 20-21 September 2013

Background

Sepsis and its after-effects are currently significant problems in hospital medicine. Successful treatment depends on the early detection of the etiological agent of the sepsis and of any antibiotic resistances. Bacteria that belong to *Staphylococcus* spp. are the most important Gram-positive causative agents of sepsis and are responsible for ~ 50% of sepsis and pyoinflammatory diseases in Ukraine. Up to 70% of Ukrainian staphylococci are also methicillin resistant which is determined by the genetic cassette *SCCmec* encoding the *mecA* β -lactam resistance gene. Besides *mecA*, some types of *SCCmec*s can also carry different genes of resistances to antibiotics like vancomycin etc. Until now nine types of *SCCmec* have been discovered based on the genetic construction, described and epidemiologically traced worldwide. It is now clear that new *SCCmec*s with enhanced antibiotic resistance characteristics are naturally appearing due to anthropogenic activities. Ukraine being geographically on a crossroads between Asia and Europe, having a big borderline with four countries-member of the EU and three non-EU European countries and exploiting a medical system where any antibiotic is easy to get without a doctor prescription can have a really dramatic impact on the *SCCmec* evolution in Europe. For this reason, a rapid screening assay with the following analysis of the

*SCCmec*s is very much needed for the early identification of new *SCCmec*s in Ukraine and prediction of new epidemiological waves in Europe.

The aim

was to elaborate a PCR rapid diagnostic test to reveal *SCCmec* in Ukrainian staphylococci.

Materials and methods

Four PCR-based assays [1-4] for methicillin-resistance identification were tested using forty *Staphylococcus* spp. isolates obtained from patients of the Institute of Traumatology and Orthopaedics of the NAMS of Ukraine.

Results

A new "in house" multiplex PCR test based on [3] for the direct simultaneous identification of *SCCmec* was developed and implemented.

Conclusions

A rapid PCR-based test for β -lactam resistance identification is now available in Ukraine. A collection of Ukrainian MRSA from different regions with PCR-determined presence of *SCCmec* has been now created and constantly supplemented.

Outlook and expert recommendations

We recommend to organize project based on direct bacteria identification using *SCCmec* for implementation

* Correspondence: moshynets@gmail.com

¹Institute of Molecular Biology and Genetics of NAS of Ukraine, Kyiv, Ukraine
Full list of author information is available at the end of the article

rapid screening in Ukraine. The epidemiological monitoring of Ukrainian *SCCmec*s with the detailed analysis of new *SCCmec* variants may impact prediction of new epidemiological waves in Europe.

Authors' details

¹Institute of Molecular Biology and Genetics of NAS of Ukraine, Kyiv, Ukraine.

²Institute of Traumatology and Orthopaedics of the NAMS of Ukraine, Kyiv, Ukraine. ³National Children Specialized Hospital "OCHMATDYT", Kyiv, Ukraine.

⁴Zabolotny Institute of Microbiology and Virology of NAS of Ukraine, Kyiv, Ukraine.

Published: 11 February 2014

References

1. Martineau, *et al*: *J. Antimic. Chemotherapy* 2000, **46**:527-533.
2. Strommenger, *et al*: *J Clin. Microbiol* 2003, **41**(9):4089-4094.
3. Huletsky, *et al*: *J. Clin. Microbiol* 2004, **42**(5):1875-1884.
4. Cuny, Witte: *Clinical Microbiology and Infection* 2005, **11**(10):834-837.

doi:10.1186/1878-5085-5-S1-A132

Cite this article as: Moshynets *et al*: Perspectives of epidemiological control of the MRSA subpopulations in hospital patients with sepsis via molecular detection, genetic analysis and following typing of methicillin resistance-carrying mobile genetic elements *SCCmec* in Ukraine. *EPMA Journal* 2014 **5**(Suppl 1):A132.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit