Published in final edited form as:

Org Lett. 2014 May 16; 16(10): 2630–2633. doi:10.1021/ol500792x.

Shagenes A and B, New Tricyclic Sesquiterpenes Produced by an Undescribed Antarctic Octocoral

Jacqueline L. von Salm[†], Nerida G. Wilson[‡], Brian A. Vesely[§], Dennis E. Kyle[§], Jason Cuce[†], and Bill J. Baker^{*†}

[†]Department of Chemistry and Center for Drug Discovery and Innovation, University of South Florida, Tampa, Florida 33620, United States

[§]Department of Global Health, University of South Florida, Tampa, Florida 33620, United States

[‡]Western Australia Museum, Perth, Western Australia, Australia

Abstract

The isolation and characterization of two new tricyclic sesquiterpenoids, shagenes A (**1**) and B (**2**) are presented. These compounds were isolated from an undescribed soft coral collected from the Scotia Arc in the Southern Ocean. One- and two-dimensional NMR spectroscopy and mass spectrometry provided the data necessary to characterize the compounds and their relative stereochemical configurations. Exploration of the bioactivity of shagenes A and B found **1** active against the visceral leishmaniasis causing parasite, *Leishmania donovani*, with no cytotoxicity against the mammalian host.

Sessile marine organisms such as sponges and corals are well-known for their chemically diverse natural products. It is common for these organisms to lack physical or behavioral defenses, therefore creating a necessity for the production of chemical defenses. From 1985 to 2008, cnidarians were second only to porifera in new marine natural products reported each year.¹ The largest class of cnidarians, Anthozoa, are almost solely responsible for

© 2014 American Chemical Society

*Corresponding Author: bjbaker@usf.edu.

Notes

The authors declare no competing financial interest.

Supporting Information

¹H, ¹³C, COSY, HSQC, HMQC, HMBC, and ROESY NMR spectra for structure elucidation of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

producing the natural products isolated from this phylum with greater than 80% of the compounds identified coming from the subclass Octocorallia.² Many of these secondary metabolites have been investigated for their ecological roles in feeding deterrence, allelopathy, and antifouling, providing evidence of widespread bioactivity.³ This has led many researchers to study their potential in drug discovery; however, the majority of published octocoral chemistry originates in tropical shallow waters with only a few dozen exceptions for deeper cold-water corals (>100 m).^{4,5}

Important metabolites for drug discovery that have been identified from soft corals include the potent anticancer diterpene glycosides eleutherobin⁶ and sarcodictyin A⁷ and the anti-inflammatory pseudopterosins.⁸ Terpenoids such as these comprise approximately 92% of all reported octocoral secondary metabolites.² Because terpenes are common in plant and algal extracts, it was once thought that the origin of octocoral terpenes was symbiotic microalgae called zooxanthellae.⁹ However, more recent results favor soft corals producing their own terpenoid metabolites;^{10,11} consider, for example, octocorals from depths of more than 100 m, where light has limited penetration through the water column and photosynthetic microalgal symbionts are absent, which are rich sources of terpenoids.¹² This paper presents the new terpenoid skeleton of shagene A and B (**1** and **2**)¹³ from a previously undescribed genus of octocoral collected via trawling near the South Georgia Islands.¹⁴ Shagenes are reminiscent of tamariscene, from the liverwort *Frullania tamarisci*, which bears the same fused 3/6/5 tricyclic ring system but displays a different methylation pattern on that ring system.¹⁵

Other cold-water sesquiterpenoids isolated from corals include the paesslerins,¹⁶ ainigmaptilonones,¹⁷ and alcyopterosins,¹⁸ all of which have shown moderate cytotoxicity toward either human cancer cell lines or microbial pathogens. Shagene A (**1**) demonstrates 5 μ M half maximal inhibitory concentration (IC_{50}) of the infected macrophage stage of the visceral leishmaniasis causing parasite,¹⁹ *Leishmania donovani*, showing a 10-fold increase in activity over the axenic amastigote form (54 μ M) with no cytotoxicity up to 100 μ g/mL (345 μ M) on J774.A1 macrophages.²⁰ The selectivity index (SI) of shagene A (**1**) is therefore greater than 70 as compared to the lower SI of the control drug miltefosine. Miltefosine shows an IC_{50} of 3 μ M against the axenic amastigote form, 1.4 μ M for the infected macrophage stage, but greater than 50 μ M cytotoxicity against mammalian cells. Shagene B (**2**) showed no activity (>20 μ g/mL) alluding to the importance of having the methoxy substituent at C-8.

The bright yellow and highly branched octocoral was collected during austral spring 2011 at depths of approximately 180 m via trawling. The location sampled was the sesquiterpenes namesake, Shag Rocks, which are a series of small islands along the Scotia Arc (S53° 26' 16.10", W42° 02' 15.52"). Specimens were frozen immediately due to noticeable oxidation of the yellow pigment. Once freeze-dried, the coral was processed with a Soxhlet extractor with refluxing CH₂Cl₂. The crude extract was mounted on silica gel and subsequently fractionated by normal-phase MPLC. Further purification of the nonpolar portion eluting in EtOAc-*n*-hexane (1:1) was accomplished by HPLC equipped with UV and ELSD detection. The more lipophilic colorless oil shagene A (**1**) (27 mg, 0.009% dry wt) was collected as the major component of the mixture with shagene B (**2**) (2 mg colorless oil, 0.0007% dry wt) considerably less abundant.

The ¹H NMR spectra of shagenes A (**1**), [α]²⁶_D -0.9, and B (**2**), [α]²⁶_D +0.2, displayed characteristic peaks indicative of smaller cyclic terpenes with 16 proton signals largely composed of single proton integrations, multiple downfield olefinic protons, and methyl substituents. The ¹³C NMR spectrum of **1** showed 18 carbon signals including five downfield signals of four olefinic carbons and the carbonyl carbon of the acetyl group (δ_C 170.9). As represented in Table 1, the spectra display functionality indicative of one acetate methyl (δ_H 2.09), one methoxy group (δ_H 3.34), one olefinic methylene (δ_H 5.05, 5.20), one trisubstituted olefin (δ_H 5.60), three vinyl or aliphatic methyl groups, six sp³ methines, one sp³ methylene, and two olefinic quaternary carbons. HREIMS of **1** provided the molecular formula C₁₈H₂₆O₃ (m/z calcd 290.1882, found m/z 290.1878) demonstrating six degrees of unsaturation and providing validation for the hypothesized 15 carbon skeleton of a tricyclic sesquiterpene bearing a quaternary carbon at the junction of all three rings. Further verification of this scaffold and relative stereochemistry were accomplished via 2D NMR experiments (COSY, gHMQC, gHMBCAD, and ROESY).

One-bond correlations from the HMQC experiment for **1** confirmed the olefinic methylene protons at C-12 (δ_C 115.9) and the quaternary nature of carbons C-5 and C-11 (δ_C 41.9, 140.6). Shared HMBC correlations (Figure 1) of the two protons at C-12 and H₃-13 (δ_H 5.05, 5.20, and 1.77) with C-10 (δ_C 31.5) and C-11 provided proof of the isopropenyl attachment to the cyclopropane at C-10. Placement of C-5 as a bridgehead carbon for this cyclopropane, the cyclohexane and cyclopentene fused rings, is verified by HMBC correlations with H-4, H-7, H-8, H-9, H-10, and H₃-15. Chemical shifts for C-4 (δ_C 22.6), C-5, and C-10 of shagene A are similar to those reported for the cyclopropane ring of the tricyclic skeleton comprising tamariscene (δ_C 30.1, δ_C 34.2, δ_C 33.6)¹⁵ with the added functionality of shagene A accounting for shift variance. Connectivity of the cyclopentene moiety can be completed on the basis of COSY coupling of the olefinic proton H-7 (δ_H 5.60) with vinyl methyl protons H₃-15 (δ_H 1.49) and the oxygen-bearing methine H-8 (δ_H 4.09). HMBC coupling of the methoxy protons with the oxygen bearing methine at C-8 verifies their connectivity. Further analysis of the 2D COSY spectrum shows H-1 (δ_H 1.19) to be vicinal to H-9 (δ_H 1.48), the methyl protons at H₃-14 (δ_H 1.01), and the diastereotopic methylene protons at H-2 (δ_H 1.25, δ_H 1.65). The acetate methyl protons provided HMBC correlations to the carbonyl C-16 (δ_C 170.9) and C-3 (δ_C 72.2) of the cyclohexane ring. COSY coupling of H-3 (δ_H 5.34) with neighboring methylene protons H-2 and methine H-4

(δ_{H} 1.65) confirms a large portion of the cyclohexane ring. HMBC correlations of H-4 with C-2, C-5, C-6, C-10, and C-11 provide the evidence of a third bridgehead carbon, this time between the cyclohexane and cyclopropane rings to complete the connectivity of the overall scaffold.

Relative stereochemistry of shagene A (**1**) was deducible upon analysis of strong, unanticipated ROESY correlations (Figure 2) between the exocyclic methylene protons on C-12 with the cyclopentane and cyclohexane bridgehead H-9, a methylene proton at C-2, and the methyl protons of the acetate. This places the isopropenyl group in the periphery of the acetate and methyl substituents of the cyclohexane ring essentially blocking that face of the molecule. Spatial placement of the methyl substituent at C-1 is a result of correlations between H₃-14 with H-9, and ROESY correlations between H-1 and H-8 place the methoxy group on the same face as H-9. Additional support for the β -methoxy is provided by conformational analysis of the H-9 coupling constants. Energy-minimized conformation of the α -methoxy configuration requires a large $^3J_{\text{HH}}$ coupling between H-8 and H-9 due to a nearly eclipsed conformation. On the other hand, the β -methoxy displays a roughly 118° dihedral angle resulting in a small $J_{\text{H8,H9}}$. The observed $J_{\text{H8,H9}}$ of 1.5 Hz indicates a β -methoxy substituent at C-8. H₃-18 and H₃-13 show weak ROESY correlation.

The obvious difference in the ^1H NMR spectra between compounds **1** and **2** was the absence of the proton signals associated with the methoxy functionality at δ_{H} 3.34 ppm and its oxymethine at δ_{H} 4.09 ppm as well as the downfield shift of multiple proton signals surrounding C-8. A carbon shift in **2**, but absent in **1** was observed at δ_{C} 208.5, establishing the oxidized α , β -unsaturated ketone moiety of **2**. The presence of a carbonyl at C-8 was further verified by the absence of H-8 COSY correlations with H-9 and H-7. Evidence of a conjugated ketone was indicated by absorbance at a wavelength of 257 nm in the UV spectrum and the extra carbonyl stretch on IR (1696 cm^{-1}). HREIMS of **2** revealed an expected mass similar to **1** save one carbon and four protons (m/z calcd 274.1569, found 274.1569) providing the molecular formula $\text{C}_{17}\text{H}_{22}\text{O}_3$. Validation of **2** having the same connectivity and stereochemistry as **1** is shown in the 2D NMR experiments (COSY, HSQC, HMBC, and ROESY).

The proposed biosynthetic pathway to **1** and **2** is shown in Scheme 1. Structural similarities to biosynthetic intermediates displayed in the biosynthesis of valerenic acid makes these molecules likely to undergo cyclization through a caryophyllene intermediate (**3**) via copalyl diphosphate synthase (CPS).²¹ Alternative pathways have been suggested with a concerted three-ring cyclization step; however, this would produce an undesirable stereochemical outcome.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the crew and research scientists on board the Nathaniel B. Palmer during the above-mentioned field season that kept us afloat and were always willing to lend a helping hand. The project was supported by National

Science Foundation awards ANT-1043749 (N.G.W.) and ANT-0838776 (B.J.B.) from the Antarctic Organisms and Ecosystems program, and National Institutes of Health grant AI103673 (B.J.B. and D.E.K.) from the National Institute of Allergy and Infectious Diseases. Greg W. Rouse, Scripps Institution of Oceanography, was instrumental in the logistical and conceptual implementation of this research. We extend special thanks to family and friends of The Baker Lab for their support, hard work, and dedication.

References

1. Hu GP, Yuan J, Sun L, She ZG, Wu JH, Lan XJ, Zhu X, Lin YC, Chen SP. Mar Drugs. 2011; 9:514–525. [PubMed: 21731546]
2. Harper, MK.; Bugni, TS.; Copp, BR.; James, RD.; Lindsay, BS.; Richardson, AD.; Schnabel, PC.; Tasdemir, D.; VanWagoner, RM.; Verbitski, SM.; Ireland, CM. Introduction to the Chemical Ecology of Marine Natural Products. In: McClintock, JB.; Baker, BJ., editors. Marine Chemical Ecology. CRC Press; Boca Raton, FL: 2001. p. 3-70.
3. Coll JC. Chem Rev. 1992; 92:613–631.
4. Skropeta D. Nat Prod Rep. 2008; 25:1131–1166. [PubMed: 19030606]
5. Lebar MD, Heimbegner JL, Baker BJ. Nat Prod Rep. 2007; 24:774–797. [PubMed: 17653359]
6. Lindel T, Jensen PR, Fenical W, Long BH, Casazza AM, Carboni J, Fairchild CR. J Am Chem Soc. 1997; 119:8744–8745.
7. Dambrosio M, Guerriero A, Pietra F. Helv Chim Acta. 1987; 70:2019–2027.
8. Look SA, Fenical W, Jacobs RS, Clardy J. Proc Natl Acad Sci USA. 1986; 83:6238–6240. [PubMed: 2875457]
9. Coll JC, Bowden BF, Tapiolas DM, Willis RH, Djura P, Streamer M, Trott L. Tetrahedron. 1983; 41:1085–1092.
10. Bandurraga MM, McKittrick B, Fenical W, Arnold E, Clardy J. Tetrahedron. 1982; 38:305–310.
11. Michalek-Wagner K, Bourne DJ, Bowden BF. Mar Biol. 2001; 138:753–760.
12. Scheuer PJ. Science. 1990; 248:173–177. [PubMed: 2183350]
13. Shagene A (1): UV (CH₂Cl₂) $\lambda_{\max}$ (ϵ) 240 (5945) nm; IR (thin film) 2959, 3050, 2933, 2877, 1734, 1655, 1458, 1369, 1246, 1089, 1026 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; 70 eV HREIMS m/z 290.1878 ([M⁺]) (C₁₈H₂₆O₃ calcd, 290.1882). Shagene B (2): UV (CH₂Cl₂) $\lambda_{\max}$ (ϵ) 257 (5425), 240 (6357) nm; IR (thin film) 3080, 2960, 2929, 2858, 1734, 1696, 1607, 1462, 1380, 1242, 1186, 1030 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; 70 eV HREIMS m/z 274.1569 ([M⁺]) (C₁₇H₂₂O₃ calcd, 274.1569).
14. This coral was assumed to be *Paragorgia* sp. upon collection; however, the bright yellow coloration and DNA sequence data from the mismatch repair (mtMutS) gene suggests this species is not related to any known species, genera, or families of Octocorallia. Further genetic analysis of the specimen is necessary for further resolution. A sample voucher has been accessioned in the Benthic Invertebrate Collection, Scripps Institution of Oceanography (S3528).
15. Paul C, Konig WA, Muhle H. Phytochemistry. 2001; 57:307–313. [PubMed: 11382249]
16. Rodriguez Brasco MF, Seldes AM, Palermo JA. Org Lett. 2001; 3:1415–1417. [PubMed: 11388830]
17. Iken KB, Baker BJ. J Nat Prod. 2003; 66:888–890. [PubMed: 12828486]
18. Palermo JA, Rodriguez Brasco MF, Spagnuolo C, Seldes AM. J Org Chem. 2000; 65:4482–4486. [PubMed: 10959848]
19. Siqueira-Neto JL, Moon S, Jang J, Yang G, Lee C, Moon HK, Chatelain E, Genovesio A, Cechetto J, Freitas-Junior LH. PLoS Negl Trop Dis. 2012; 6:e1671. [PubMed: 22720099]
20. The axenic amastigote bioassay for leishmaniasis was an MTS [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium] 72 h assay using 66000 amastigotes per well. IC₅₀ values were calculated from sigmoidal inhibition curves using SigmaPlot 11.0 software. The procedure for the J447.A-1 cytotoxicity assay is the same as the axenic amastigote bioassay with one exception, 50000 amastigotes are used per well.
21. Yeo YS, Nybo SE, Chittiboyina AG, Weerasooriya AD, Wang YH, Gongora-Castillo E, Vaillancourt B, Buell CR, DellaPenna D, Celiz MD, Jones AD, Wurtele ES, Ransom N, Dudareva

N, Shaaban KA, Tibrewal N, Chandra S, Smillie T, Khan IA, Coates RM, Watt DS, Chappell J. J Biol Chem. 2013; 288:3163–3173. [PubMed: 23243312]

Figure 1.
Key COSY and HMBC correlations of **1**.

Figure 2.
Key ROESY correlations for shagene A (**1**).

Scheme 1.
Proposed Biosynthetic Pathway for Sesquiterpenes 1 and 2

Table 1

¹H^a and ¹³C^b NMR Data for Shagenes A (1) and B (2) in CDCl₃

position	shagene A (1)			shagene B (2)		
	δ _C	δ _H (m, J, Hz)	HMBC	δ _C	δ _H (m, J, Hz)	HMBC
1	33.2	1.19 (1H, m)	2, 8, 9, 14	32.2	1.42 (1H, m)	2, 8, 9, 14
2	35.7	1.25 (1H, q, 11.8)	1, 3, 4, 9, 14	36.4	1.35 (1H, q, 11.8)	1, 3, 4, 9, 14
		1.65 (1H, m)	1, 3, 4, 9, 14		1.68 (1H, m)	1, 3, 4, 9, 14
3	72.2	5.34 (1H, dt, 11.7, 6.4)	2, 4, 10, 16	71.4	5.33 (1H, dt, 11.7, 6.4)	2, 4, 10, 16
4	22.6	1.65 (1H, m)	2, 5, 6, 10, 11	25.5	2.03 (1H, d, 6.1)	2, 3, 5, 6, 9, 10, 11
5	41.9			40.2		
6	147.3			178.0		
7	123.2	5.60 (1H, m)	5, 6, 8, 9, 15	129.7	5.94 (1H, m)	5, 6, 8, 9, 15
8	88.6	4.09 (1H, d, 0.6)	1, 5, 6, 7, 18	208.5		
9	48.4	1.48 (1H, dd, 1.5, 9)	1, 2, 5, 8, 10, 14	49.0	1.89 (1H, d, 10.9)	1, 2, 5, 6, 8, 10, 14
10	31.5	1.78 (1H, br s)	3, 4, 5, 6, 9	33.1	2.06 (1H, br s)	4, 5, 6, 9, 11, 12
11	140.6			139.4		
12	115.9	5.05 (1H, q, 1.6)	10, 13	116.9	5.11 (1H, q, 1.6)	10, 13
		5.20 (1H, br s)	10, 11, 13		5.20 (1H, br s)	10, 11, 13
13	24.1	1.77 (3H, s)	10, 11, 12	24.1	1.76 (3H, s)	10, 11, 12
14	19.5	1.01 (3H, d, 6.4)	1, 2, 3, 9	19.3	1.23 (3H, d, 6.1)	1, 2, 3, 9
15	11.6	1.49 (3H, t, 1.2)	5, 6, 7	13.2	1.84 (3H, d, 1.2)	5, 6, 7
16	170.9			170.8		
17	21.6	2.09 (3H, s)	3, 16	21.5	2.11 (3H, s)	3, 16
18	55.6	3.34 (3H, s)	8			

^a¹H NMR experiments at 500 MHz, recorded in ppm (integration, multiplicity, J-coupling in Hz);

^b¹³C NMR experiments at 125 MHz, recorded in ppm.