Published in final edited form as:

Am J Med. 2014 August ; 127(8): 739–743. doi:10.1016/j.amjmed.2014.02.033.

A prospective study of caffeine intake and risk of incident tinnitus

Jordan T. Glicksman, MD, MPH¹, Sharon G. Curhan, MD, MSc², and Gary C. Curhan, MD, ScD^{2,3,4}

¹Department of Otolaryngology—Head and Neck Surgery, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada

²Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, Massachusetts

³Renal Division, Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts

⁴Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts

Abstract

Background—Caffeine is a commonly consumed substance that has long been thought to play a role in the development of tinnitus, but prospective data are lacking. We prospectively evaluated the association between caffeine intake and self-reported tinnitus in a female cohort.

Methods—Participants were 65,085 women in the Nurses Health Study II, aged 30–44 years and without tinnitus at baseline in 1991, who completed questionnaires about lifestyle and medical history every two years and food frequency questionnaires every four years. Information on self-reported tinnitus and date of onset was obtained from the 2009 questionnaire, with cases defined as those reporting experiencing symptoms “a few days/week” or “daily.” Multivariable adjusted hazard ratios (HRs) were calculated using Cox proportional hazards regression models.

Results—At baseline the mean age of the cohort was 36.3 years and the mean caffeine intake was 242.3 mg/day. After 18 years of follow-up, 5,289 incident cases of tinnitus were reported. There was a significant inverse association between caffeine intake and the incidence of tinnitus. Compared with women with caffeine intake less than 150 mg/day (150 mg corresponds to approximately one 8 ounce cup of coffee), the multivariable adjusted hazard ratios were 0.85

© 2014 Elsevier Inc. All rights reserved.

Corresponding Author: Dr. Jordan Glicksman, Department of Otolaryngology—Head and Neck Surgery, University of Western Ontario, London Health Sciences Centre, 800 Commissioners Road East, PO Box 5010, London, Ontario N6A 5W9, jglicksm@uwo.ca.

Conflicts of Interest: None

Contribution Statement: All authors have met the criteria for authorship, including 1) substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; 2) drafting the article or revising it critically for important intellectual content; and 3) final approval of the version to be published.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

(95% CI 0.76–0.95) for those who consumed 450–599 mg/day and 0.79 (0.68–0.91) for those who consumed 600mg/day or more.

Conclusion—In this prospective study higher caffeine intake was associated with a lower risk of incident tinnitus in women.

Keywords

Caffeine; Tinnitus; Epidemiology

Introduction

Tinnitus is a highly prevalent condition estimated to afflict 50 million Americans, severely disabling 3 million.^[1, 2] In the majority of cases, the precise pathophysiology underlying tinnitus remains unknown and, in the absence of a reversible underlying condition, treatment is generally not highly effective.^[3, 4] Preventing the development of the condition may prove to be the best way to reduce its burden on the individual and society. While some risk factors such as hearing loss and trauma are well established, evidence for other risk factors is scarce.

Caffeine is the most commonly used psychoactive substance, frequently consumed in coffee.^[5] Over 50% of Americans drink coffee, and the per capita intake is 2 cups per day.^[6] Although caffeine has long been implicated in the development of tinnitus,^[7, 8] evidence to support caffeine as a risk factor or exacerbating factor for tinnitus is lacking in the medical literature. However, caffeine cessation or reduction in intake to reduce tinnitus symptoms has been recommended by national primary care and specialty organizations.^[9, 10] Notably, a recent randomized controlled crossover trial failed to demonstrate an improvement in tinnitus symptoms with cessation of caffeine intake.^[11]

With the high prevalence of caffeine intake and tinnitus, prospective evaluation of this relation is important. Therefore, we prospectively examined the association between caffeine intake and the risk of incident tinnitus in 65, 085 women in the Nurses' Health Study II.

Methods

Study Population

The Nurses' Health Study II cohort began in 1989 with questionnaires completed by 116,430 female registered nurses aged 25 to 42 years. Questionnaires have been mailed on a biennial basis to update exposure information and the incidence of a wide variety of medical conditions. The follow-up rate for this cohort exceeds 90% of eligible person-time.

We excluded individuals if they reported onset of tinnitus prior to 1991 (the first year that caffeine intake was assessed) or if they had a history of cancer (excluding non-melanoma skin cancer). Participants were eligible to contribute person-time to the study as of their 30th birthday, which is the youngest age for which we were able to estimate the onset of tinnitus.

Assessment of caffeine intake

Caffeine intake was assessed using detailed, extensively validated semiquantitative food frequency questionnaires that inquired about the average intake of over 130 foods and beverages over the preceding year in 1991 and at four year intervals thereafter. [12–14]

The relevant beverages on the questionnaire included low-calorie cola (e.g. Diet Coke or Diet Pepsi with caffeine), regular cola (e.g. Coke, Pepsi or other cola beverages with sugar), tea with caffeine, tea without caffeine, coffee with caffeine and decaffeinated coffee. In 2003 and 2007, a question about intake of dairy coffee drinks—hot or cold (e.g. cappuccino) was also included. The relevant foods included candy bars (e.g., Snickers, Milky Way, Reeses), dark chocolate (e.g. Hershey's Dark or Dove Dark), milk chocolate—bar or pack (e.g., Hershey's, M&M's) and brownies. The response options for specified serving sizes were: never or less than once per month; 1–3 per month; 1 per week; 2 to 4 per week; 5 to 6 per week; 1 per day; 2 to 3 per day; 4 to 5 per day; and 6 or more per day.

Total caffeine intake was calculated using US Department of Agriculture food composition sources. For example, the caffeine content of the specific items was considered to be 137 mg per cup of coffee, 47 mg per cup of tea, 46 mg per can or bottle of cola beverage and 7 mg per serving of chocolate candy.^[15] This method of determining caffeine intake was shown to be valid in prior studies.^[12–14]

Ascertainment of tinnitus

Participants were asked on the 2009 questionnaire “In the past 12 months, have you had ringing, roaring, or buzzing in your ears?” The response categories were never, once per week or less, a few days per week, or daily. To determine the date of onset, participants were asked “At what age did this first begin?” (less than 30, 30 to 39, 40 to 49, 50 to 59 and 60 years of age or older). Onset of tinnitus was defined to be the midpoint of an age interval (e.g. 35 for the age category of onset of 30–39). We defined a case as a participant who reported tinnitus occurring a few days per week or daily.

Assessment of covariates

Data on BMI and intake of tobacco were collected on each questionnaire. On the 2009 questionnaire, each participant was asked if they had a hearing problem and age when they first noticed a change. Data on other health conditions were ascertained on each questionnaire. Conditions of interest included diabetes, hypertension, multiple sclerosis and depression as these have been associated with tinnitus.^[2, 7, 16, 17] Data regarding use of medications proposed to be associated with tinnitus were obtained from the biennial questionnaires (analgesics (acetaminophen, aspirin, ibuprofen, other NSAIDs, and cyclooxygenase-2 inhibitors), thiazide diuretics, furosemide, and selective serotonin reuptake inhibitors/other antidepressants).

The Crown Crisp Questionnaire, used to assess phobic anxiety, was administered in 1993 and 2005 and was used to adjust for phobic anxiety based on score categories.^[18, 19]

Statistical Methods

The time of observation was between 1991 and 2009. Participants who did not return a questionnaire for one time period and who remained eligible contributed person-time for later time intervals. Participants were censored after the onset of tinnitus or diagnosis of cancer.

Cox proportional hazards regression models were used to estimate multivariable adjusted hazard ratios and 95% confidence intervals based on categories of cumulative average caffeine intake (in 150mg/day increments, which is approximately one cup of coffee). We used the Anderson-Gill data structure and stratified by age and calendar year. Information on caffeine intake on each subsequent questionnaire was used to update the cumulative average caffeine intake up to that point. Covariates considered in multivariable adjusted models included age (continuous), hearing loss, smoking status (current, past or never), comorbid conditions (diabetes, hypertension, multiple sclerosis, and depression), medications used at least one day per week (analgesics, diuretics and antidepressants), BMI and smoking status. We used stratified models to examine whether the relation between caffeine intake and incident tinnitus varied by age (less than or greater than/equal to 50 years) and self-reported hearing loss. We also evaluated the association between decaffeinated coffee and incident tinnitus, after excluding those who consumed caffeinated coffee, to explore whether any observed relations were due to coffee (the major source of caffeine in this cohort) or caffeine.

Covariate information was updated at the beginning of each time period using the most recently available responses. Participants with missing covariate data were assigned to a missing category for that specific time period.

Level of significance (α) for two-tailed p-values was <0.05 . We used SAS version 9.3 for UNIX statistical software package (SAS Institute Inc, Cary, NC).

Results

There were 65,085 women included in the analysis. The baseline characteristics of the cohort according to category of caffeine intake are shown in Table 1. At baseline the mean age of the cohort was 36.3 years, mean BMI was 24.5 kg/m² and the mean caffeine intake was 242.3 mg/day. Comorbidities present at baseline included hearing problems (prevalence = 3.3%), history of depression (13.7%), diabetes (0.8%), hypertension (6.1%) and multiple sclerosis (0.3%). With increasing caffeine intake, there were higher percentages of participants who reported smoking, depression, aspirin use and ibuprofen use. The incidence rate of tinnitus increased substantially with age, ranging from 104 per 100,000 person-years below age 40 to 1273 per 100,000 person-years for those 50 and older (Table 2).

The age and multivariable adjusted hazard ratios (HRs) of the association between caffeine intake and incident tinnitus are shown in Table 3. After adjusting for age, there was a significant inverse association between caffeine intake and the incidence of tinnitus at higher levels of caffeine intake (HR 0.89, 95% CI 0.80–0.99 for 450–599 mg/day and HR 0.83, 0.73–0.95 for 600 or more mg/day, p for trend <0.001). After adjusting for other potential

confounders, a slightly greater inverse association was observed between caffeine intake and tinnitus at higher levels of caffeine intake (HR 0.85 (0.76–0.95) for 450–599 mg/day and HR 0.79 (0.68–0.91) for 600 or more mg/day, p for trend <0.001). The association between caffeine intake and tinnitus did not vary by age (p for interaction, 0.5) or hearing loss (p for interaction, 0.8).

To exclude further the possibility that tinnitus onset might have led to decreased caffeine consumption, we compared caffeine intake in participants before and after the time they indicated they developed tinnitus, but found no differences. For example, between 1991 and 1999 there was no significant change in caffeine consumption category amongst participants who indicated their onset of tinnitus was between 1995 and 1999 ($p=0.53$ using Bowker's test of symmetry). We also examined changes in mean caffeine intake over time among those who did and did not develop tinnitus, in order to account for potential temporal trends. There was a similar slight decrease in total caffeine consumption between 1991 and 1999 among those who did develop tinnitus (19.1 mg, 95% CI = 8.5–29.7 mg) and those who did not develop tinnitus during follow-up (14.3 mg, 95% CI = 12.7–15.9 mg). The decreased consumption over time was not significantly different between groups.

There was no association between decaffeinated coffee intake and incident tinnitus. Compared with those who drank decaffeinated coffee less than once per month, the multivariable adjusted HR for tinnitus was 1.00 (0.89–1.13) for those who drank 2–3 cups/day and 0.85 (0.62–1.16) for 4 or more cups/day (p for trend = 0.9).

Discussion

Main Findings

In this large, prospective study among women we observed an inverse association between caffeine intake and incident tinnitus. The majority (approximately 70%) of caffeine intake in these individuals was from coffee consumption. We did not find an association between decaffeinated coffee intake and incident tinnitus. This suggests it is caffeine rather than another component of coffee that was associated with a decreased incidence of tinnitus.

Comparison with other studies

A recent randomized controlled crossover trial failed to demonstrate an improvement in tinnitus symptoms with cessation of caffeine intake while demonstrating harm in the way of withdrawal symptoms among participants.^[11] The investigation included men and women with a mean age of 59.2 years. A mean difference in tinnitus severity score of 0.04 (95% CI 1.99, 1.93) was observed between patients during caffeine-consuming and caffeine-abstaining states. Our study examined the risk of incident tinnitus, whereas the aforementioned study evaluated the effect of caffeine on existing tinnitus. Based on the results of these studies, there is not persuasive evidence that reducing caffeine intake would be beneficial with respect to tinnitus. We are not aware of other prospective studies evaluating the association between caffeine and incident tinnitus.

The precise pathophysiology underlying subjective tinnitus remains unknown. Purported mechanisms include hyperactivity of the ascending auditory pathways or a reduced

suppressive influence of the central nervous system.^[20] Caffeine is a central nervous system stimulant, and this biologic action may have led to the idea that it may play a role in the development of tinnitus. Noting a paucity of evidence for such an association, Anderson et al suggested that if such an association exists, the stimulatory effect of caffeine may lead to increased detection of tinnitus through increased arousal or anxiety.^[21]

In addition to general stimulatory effects of caffeine on the central nervous system, caffeine has been shown to have a direct effect on the inner ear in both in vitro and animal studies. Caffeine is thought to result in shortening of outer hair cells, and this effect may be mediated through the release of calcium from ryanodine receptors.^[22] At this time it is unclear whether this pathway is involved in the development or protection from tinnitus and therefore this may be a topic for future investigation.

Limitations

Our study has limitations. The presence of tinnitus was established based on self-report; however, there is no diagnostic test for tinnitus so it is not possible to validate a self-report of tinnitus. Hearing loss was also self-reported so there was likely some misclassification of this covariate. There does remain the possibility of misclassification due to recall with respect to the date of onset of tinnitus. Furthermore, the questionnaire does not distinguish between subjective and objective tinnitus, but the vast majority of cases of tinnitus are subjective. Finally, our cohort was composed of women who were predominantly Caucasian. Since there are no previous studies that have examined the association between caffeine and incidence tinnitus in men, it is unknown if the findings apply to men or other racial groups.

Conclusion

In this prospective study, higher caffeine intake was associated with a reduced risk of tinnitus. At present the mechanism underlying this association remains unknown. Given that our investigation evaluated the incidence of tinnitus and not the effect of caffeine amongst participants with existing tinnitus, we are unable to make a recommendation as to whether the addition of caffeine would improve symptoms.

Acknowledgments

This work was supported by grants UM1 CA176726 and DK91417 from the National Institutes of Health. We would like to gratefully acknowledge Elaine Coughlan-Gifford for the programming assistance she provided our research team.

References

1. Adams PF, Hendershot GE, Marano MA. Centers for Disease Control and Prevention/National Center for Health Statistics. Current estimates from the National Health Interview Survey, 1996. *Vital Health Stat.* 1999 Oct; 10(200):1–203.
2. Shargorodsky J, Curhan GC, Farwell WR. Prevalence and characteristics of tinnitus among US adults. *Am J Med.* 2010 Aug; 123(8):711–8. [PubMed: 20670725]
3. Heller AJ. Classification and epidemiology of tinnitus. *Otolaryngol Clin North Am.* 2003 Apr; 36(2):239–48. Review. [PubMed: 12856294]

4. Hoare DJ, Kowalkowski VL, Kang S, Hall DA. Systematic review and meta-analyses of randomized controlled trials examining tinnitus management. *Laryngoscope*. 2011 Jul; 121(7): 1555–64. [PubMed: 21671234]
5. Kendler KS, Myers J, Gardner OC. Caffeine intake, toxicity and dependence and lifetime risk for psychiatric and substance use disorders: an epidemiologic and co-twin control analysis. *Psychol Med*. 2006 Dec; 36(12):1717–25. [PubMed: 16893482]
6. Lundsberg, L. *Caffeine consumption in Caffeine*. Spiller, G., editor. CRC Press; Boca Raton: 1998.
7. Schleuning, AJ.; Shi, YJ.; Martin, WH. Tinnitus in Head and Neck Surgery—Otolaryngology. Bailey, BJ.; Johnson, JT.; Newlands, SD., editors. Lippincott Williams & Wilkins; Philadelphia, Pennsylvania: 2006. p. 2241
8. Mayo Clinic Staff. Tinnitus. 2010. [cited 2012 July 28]; Available from: <http://www.mayoclinic.com/health/tinnitus/DS00365/DSECTION=causes>
9. Insight into Causes and Treatments for Tinnitus. Dec. 2010 [cited 2012 July 8]; Available from: <http://www.entnet.org/HealthInformation/tinnitus.cfm>
10. Crummer RW, Hassan GA. Diagnostic approach to tinnitus. *Am Fam Physician*. 2004 Jan 1; 69(1): 120–6. Review. [PubMed: 14727828]
11. Claire LS, Stothart G, McKenna L, Rogers PJ. Caffeine abstinence: an ineffective and potentially distressing tinnitus therapy. *Int J Audiol*. 2010 Jan; 49(1):24–9. [PubMed: 20053154]
12. Willett WC, Sampson L, Stampfer MJ, et al. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol*. 1985 Jul; 122(1):51–65. [PubMed: 4014201]
13. Salvini S, Hunter DJ, Sampson L, et al. Food-based validation of a dietary questionnaire: the effects of week-to-week variation in food consumption. *Int J Epidemiol*. 1989 Dec; 18(4):858–67. [PubMed: 2621022]
14. Grobbee DE, Rimm EB, Giovannucci E, et al. Coffee, caffeine, and cardiovascular disease in men. *N Engl J Med*. 1990 Oct 11; 323(15):1026–32. [PubMed: 2215561]
15. Winkelmayr WC, Stampfer MJ, Willett WC, Curhan GC. Habitual caffeine intake and the risk of hypertension in women. *JAMA*. 2005 Nov 9; 294(18):2330–5. [PubMed: 16278361]
16. Loprinzi PD, Maskalick S, Brown K, Gilham B. Association between depression and tinnitus in a nationally representative sample of US older adults. *Aging Ment Health*. 2013 Aug; 17(6):714–7. [PubMed: 23461284]
17. Daugherty WT, Lederman RJ, Nodar RH, Conomy JP. Hearing loss in multiple sclerosis. *Arch Neurol*. 1983 Jan; 40(1):33–5. [PubMed: 6848085]
18. Crown S, Crisp AH. A short clinical diagnostic self-rating scale for psychoneurotic patients. The Middlesex Hospital Questionnaire (M.H.Q.). *Br J Psychiatry*. 1966 Sep; 112(490):917–23. [PubMed: 5970912]
19. Burgess PM, Mazzocco L, Campbell IM. Discriminant validity of the Crown-Crisp Experiential Index. *Br J Med Psychol*. 1987 Mar; 60(Pt 1):61–9. [PubMed: 3567112]
20. Nehlig A, Daval JL, Debry G. Caffeine and the central nervous system: mechanisms of action, biochemical, metabolic and psychostimulant effects. *Brain Res Brain Res Rev*. 1992 May-Aug; 17(2):139–70. Review. [PubMed: 1356551]
21. Bagueley, D.; Andersson, G.; McFerran, D.; McKenna, L. Tinnitus: A Multidisciplinary Approach (Second Edition). Wiley-Blackwell; London: 2013. Consequences and moderating factors; p. 100
22. Trinidad A, Robinson T, Phillips JS. The role of caffeine in otorhinolaryngology: guilty as charged? *Eur Arch Otorhinolaryngol*. 2013 Aug 11. (Ahead of Print).

Clinical Significance Bullet Points

- In women, the incidence rate of tinnitus increases with age and reaches a value greater than 1 per 100 per year in the 6th decade of life
- Higher caffeine intake is associated with a lower risk of incident tinnitus in women

Table 1

Baseline characteristics of women in 1991 by category of caffeine intake

	Energy-adjusted caffeine intake, mg/day				
	0-149 (n=29,037)	150-299 (n=13,049)	300-449 (n=13,188)	450-599 (n=4,453)	600+ (n=5,358)
Age, years*	35.7(4.7)	36.2(4.7)	37.0(4.3)	37.1(4.4)	37.7(4.2)
Body mass index, kg/m ²	24.4(5.3)	24.7(5.4)	24.2(4.8)	24.6(5.1)	24.7(4.9)
Smoking Status					
Never, %	77.8	68.0	55.0	52.5	39.5
Past, %	16.9	22.3	30.8	27.6	27.4
Current, %	5.2	9.5	14.0	19.7	33.0
Hearing problem, %	3.3	3.1	3.5	3.6	3.4
History of hypertension, %	5.9	6.6	5.6	6.4	6.4
History of diabetes, %	0.7	0.9	0.6	0.8	1.1
History of depression, 2003, %	12.9	13.7	14.2	14.8	16.0
History of multiple sclerosis, %	0.3	0.4	0.3	0.4	0.4
Thiazide diuretic use, %	1.6	1.8	1.7	1.9	1.8
Furosemide use, 1989, %	0.4	0.4	0.3	0.5	0.7
Aspirin use, %	3.6	4.8	4.5	5.0	6.2
Ibuprofen use, %	7.3	10.4	10.1	10.6	12.2
Acetaminophen use, %	6.7	8.2	7.6	8.2	8.1
COX-2 inhibitors use, 2001, %	8.6	9.4	9.3	10.3	10.6
SSRI use, 1997, %	7.2	7.9	8.3	9.0	9.1
Other antidepressant use, 1993, %	9.0	10.0	9.5	10.9	11.6
Anxiety Crown-Crisp Score					
2 or less, %	58.1	56.1	56.2	55.7	53.2
3 or more, %	31.8	32.7	32.2	32.1	35.3

The values in the table reflect responses from 1991 unless otherwise indicated. Use of medication refers to use at least one day per week. Values are means (SD) or percentages and are standardized to the age distribution of the study population.

SSRI- Selective Serotonin Reuptake Inhibitor

* indicates the value is not age adjusted

Table 2

Age-specific tinnitus incidence rates

Age (years)	Person-years	Cases	Incidence Rate per 100,000 person-years
<40	348,676	361	104
40–49	545,612	2,286	419
50	207,464	2,642	1,273
Total	1,101,753 *	5,289	475

* Total differs slightly due to rounding

Table 3

Age and multivariable adjusted hazard ratios for the association between caffeine intake and incident tinnitus

Caffeine intake in mg/day	No. of cases	Person-years	Age-adjusted HR (95%CI)	Multivariate Model * HR (95% CI)
0–149	2,268	470,228	1.00 (referent)	1.00(referent)
150–299	1,394	274,909	0.96 (0.89–1.02)	0.94 (0.88–1.00)
300–449	1,029	214,588	0.93 (0.87–1.01)	0.91 (0.84–0.98)
450–599	364	80,433	0.89 (0.80–0.99)	0.85 (0.76–0.95)
600	234	61,595	0.83 (0.73–0.95)	0.79 (0.68–0.91)
			$p_{\text{trend}} < 0.001$	$p_{\text{trend}} < 0.001$

* Covariates in Model included caffeine, age, hearing loss, medications, medical conditions, phobic anxiety, BMI and smoking status. Medications included acetaminophen, aspirin, ibuprofen, other NSAIDs, cyclooxygenase-2 inhibitors, thiazide diuretics, furosemide, and selective serotonin reuptake inhibitors/other antidepressants. Medical conditions included diabetes mellitus, hypertension, multiple sclerosis and depression.