Published in final edited form as:

Psychopharmacology (Berl). 2014 September ; 231(17): 3257–3272. doi:10.1007/s00213-014-3564-8.

Divergent neuroactive steroid responses to stress and ethanol in rat and mouse strains: Relevance for human studies

Patrizia Porcu¹ and A. Leslie Morrow²

¹Neuroscience Institute, National Research Council of Italy (CNR), Cagliari, Italy

²Departments of Psychiatry and Pharmacology, Bowles Center for Alcohol Studies, University of North Carolina School of Medicine, Chapel Hill, NC, USA

Abstract

Rationale—Neuroactive steroids are endogenous or synthetic steroids that rapidly alter neuronal excitability via membrane receptors, primarily GABA_A receptors. Neuroactive steroids regulate many physiological processes including hypothalamic-pituitary-adrenal (HPA) axis function, ovarian cycle, pregnancy, aging, and reward. Moreover, alterations in neuroactive steroid synthesis are implicated in several neuropsychiatric disorders.

Objectives—This review will summarize the pharmacological properties and physiological regulation of neuroactive steroids, with a particular focus on divergent neuroactive steroid responses to stress and ethanol in rats, mice and humans.

Results—GABAergic neuroactive steroids exert a homeostatic regulation of the HPA axis in rats and humans, whereby the increase in neuroactive steroid levels following acute stress counteracts HPA axis hyperactivity and restores homeostasis. In contrast, in C57BL/6J mice, acute stress decreases neurosteroidogenesis and neuroactive steroids exert paradoxical excitatory effects upon the HPA axis. Rats, mice and humans also differ in the neuroactive steroid responses to ethanol. Genetic variation in neurosteroidogenesis may explain the different neuroactive steroid responses to stress or ethanol.

Conclusions—Rats and mouse strains show divergent effects of stress and ethanol on neuroactive steroids in both plasma and brain. The study of genetic variation in the various processes that determine neuroactive steroids levels as well as their effects on cell signaling may underlie these differences and may play a relevant role for the potential therapeutic benefits of neuroactive steroids.

Keywords

Neuroactive steroids; allopregnanolone; stress; HPA axis; ethanol; rat; C57BL/6J mice; DBA/2J mice

Introduction

Neuroactive steroids are endogenous or synthetic steroids that rapidly alter neuronal excitability via membrane receptors. These steroids are derived from cholesterol, which is converted to pregnenolone that is further metabolized into several steroid hormones that include glucocorticoids, estrogens, progesterone, dehydroepiandrosterone (DHEA) as well as their metabolites. The $3\alpha,5\alpha$ - and $3\alpha,5\beta$ -reduced metabolites of progesterone, deoxycorticosterone (DOC), DHEA and testosterone are positive modulators of γ -aminobutyric acid type A ($GABA_A$) receptors. Among these steroids, the progesterone metabolite ($3\alpha,5\alpha$)-3-hydroxypregnan-20-one ($3\alpha,5\alpha$ -THP or allopregnanolone) and the DOC metabolite ($3\alpha,5\alpha$)-3,21-dihydroxypregnan-20-one ($3\alpha,5\alpha$ -THDOC or allotetrahydrodeoxy-corticosterone) are the most potent $GABA_A$ receptor modulators and the most studied neuroactive steroids.

This review will summarize the current literature on GABAergic neuroactive steroid responses to stress and ethanol in rat and mouse strains as well as humans. We highlight recent work on genetic variation in neuroactive steroid responses in mouse strains that may be particularly relevant for translational studies, since this genetic variation models the genetic diversity of the human population.

Pharmacological properties of neuroactive steroids

Systemic administration of GABAergic neuroactive steroids exerts a variety of pharmacological responses including anxiolytic, antidepressant, anticonvulsant, sedative, anesthetic, and analgesic effects in animal models and human studies (Belelli et al. 1989; Bitran et al. 1991; Hogskilde et al. 1987; Kavaliers and Wiebe 1987; Khisti et al. 2000), that are consistent with their GABAergic actions. However, there are also reports of paradoxical excitatory actions of GABAergic neuroactive steroids that have received less attention in the literature. For example, pregnanolone ($3\alpha,5\beta$ -THP) and alphaxalone were both shown to produce dose dependent myoclonic seizures in outbred Bantin and Kingman male mice (File and Simmonds 1988), pregnanolone induced seizures in male C57BL/6J mice (Shannon et al. 2005a), $3\alpha,5\alpha$ -THP enhanced the frequency and amplitude of postural tremor in $GABA_A$ receptor $\alpha 1$ subunit knockout mice (Osterman et al. 2005), and $3\alpha,5\alpha$ -THP enhanced aggression in outbred Swiss mice in various experimental settings (Fish et al. 2001; Miczek et al. 2003). The molecular mechanisms of these excitatory behavioral actions are unclear, but may involve disinhibition of neuronal circuits via $GABA_A$ receptors, alterations in membrane chloride gradient that result in depolarization by endogenous GABA or novel actions of these steroids. $3\alpha,5\alpha$ -THP also activates feminine sexual behavior of rodents (Frye et al. 1998), which is thought to involve dopamine transmission as well as pregnane xenobiotic receptors for steroidogenesis (see Walf et al., this issue). Moreover, GABAergic neuroactive steroids, like various drugs of abuse, exhibit rewarding properties in DBA/2J mice and Long-Evans rats (Finn et al. 1997a; Sinnott et al. 2002) and enhance brain stimulation reward in C57BL/6J mice, much like cocaine (Fish et al., this issue). $3\alpha,5\alpha$ -THP and other GABAergic neuroactive steroids exhibit biphasic modulation of ethanol intake in various rodent models of ethanol self-administration (Besheer et al. 2010; Finn et al. 2008; Ford et al. 2007; Ford et al. 2005; Janak et al. 1998; Morrow et al. 2001), suggesting

complex actions on neuronal circuitry mediating these behaviors. The complexity of these behavioral responses in rodents points to the possibility that genetic variation may play an important role in neuroactive steroid actions across species.

Mechanisms and sites of action

The 3 α ,5 α - and 3 α ,5 β -reduced metabolites of progesterone, DOC, DHEA and testosterone exert inhibitory actions that are mediated by synaptic and extrasynaptic GABA_A receptors. These neuroactive steroids interact with synaptic GABA_A receptors to produce phasic inhibition via specific binding sites on the TM1 region of the α subunits that allosterically modulate binding to GABA and benzodiazepine recognition sites (Hosie et al. 2006). GABA_A receptor function is enhanced by potentiation of GABA-mediated Cl⁻ conductance as well as direct stimulation of Cl⁻ conductance (Harrison et al. 1987; Majewska et al. 1986; Morrow et al. 1987). GABA_A receptors appear to have multiple neurosteroid recognition sites that likely reflect distinct recognition sites on GABA_A receptor subtypes (Morrow et al. 1990). Neuroactive steroids modulate both synaptic and extrasynaptic GABA_A receptors with lower potency at synaptic receptors that contain γ 2 subunits and higher potency at extrasynaptic receptors that contain subunits and mediate tonic inhibition (Belelli and Lambert 2005; Carver and Reddy 2013). The excitatory neuroactive steroids include the sulfated derivatives of pregnenolone and DHEA as well as the 3 α ,5 α - and 3 α ,5 β -reduced metabolites of cortisol. The excitatory actions of sulfated steroids are partially mediated by direct interactions with N-methyl-D-aspartate (NMDA) receptors, where they act as weak agonists (micromolar potency) (Park-Chung et al. 1997; Wu et al. 1991), as well as by a blocking action on GABA_A receptors (Akk et al. 2001). Pregnenolone sulfate also potently increases glutamate release (Meyer et al. 2002) and inhibits GABA release (Mtchedlishvili and Kapur 2003), effects that are blocked by sigma-1 receptor antagonists. Finally, pregnenolone sulfate enhances the trafficking of NMDA receptors, resulting in greater excitatory drive (Kostakis et al. 2013). In addition, 3 α ,5 α - and 3 α ,5 β -reduced cortisol have antagonist properties at both GABA and neurosteroid recognition sites of GABA_A receptors and these compounds are the most abundant metabolites of cortisol in human urine (Penland and Morrow 2004).

It is not clear how endogenous neuroactive steroids influence neurotransmission in the brain. The available data suggests that elevations of 3 α ,5 α -THP within a cell reduce the excitability of that particular cell (Akk et al. 2007; Saalman et al. 2006; Tokuda et al. 2011; Tokuda et al. 2010). 3 α ,5 α -THP may access the neuroactive steroid transmembrane binding sites on GABA_A receptors via intracellular (i.e. presynaptic) lateral diffusion through the cell membrane (Akk et al. 2007) or by a paracrine or autocrine mechanism (Herd et al. 2007), as no active release mechanism has been identified. There is also evidence for extracellular pockets of neuroactive steroid storage that allow for rapid and repeated access to extracellular binding sites on neurons (Akk et al. 2007; Akk et al. 2005). Figure 1 summarizes these putative mechanisms and sites of action of GABAergic neuroactive steroids.

Physiological regulation of neuroactive steroid levels in plasma and brain

Neuroactive steroid concentrations vary across physiological conditions mostly associated with reproduction and development such as puberty, ovarian cycling, pregnancy or aging. Brain levels of 3 α ,5 α -THP are elevated in the embryonic Sprague-Dawley rat, decrease immediately before birth, remain low throughout early life (Grobin and Morrow 2001) and increase immediately before puberty in Sprague-Dawley rats (Grobin and Morrow 2001), C57BL/6J mice (Shen et al. 2007) and humans (Fadalti et al. 1999). Circulating 3 α ,5 α -THP levels decline during menopause in women (Bixo et al. 1997). A similar decline has been observed in several rat brain areas and it was hypothesized it might contribute to cognitive impairment in aged Sprague-Dawley rats (George et al. 2010).

Interestingly, alterations in neuroactive steroid levels that occur *in vivo* under some physiological conditions are associated with changes in GABA_A receptor function and expression. These data are essential to understand the behavioral sequelae of changes in levels of these steroids. This work is reviewed in several other papers in this special issue and we refer the reader to those contributions for a complete review of neuroactive steroid regulation of GABA_A receptor gene expression (see, MacKenzie and Maguire, this issue).

GABAergic neuroactive steroids concentrations vary throughout the ovarian cycle in both rodents and humans. 3 α ,5 α -THP and progesterone levels vary throughout the estrus cycle in brain and plasma of HsdOla:Tuck-Ordinary mice (Corpechot et al. 1997). In female C57BL/6J mice, the diestrus phase is accompanied by elevated levels of progesterone and 3 α ,5 α -THP, and a subsequent increase in tonic inhibition and decreased seizure susceptibility and anxiety (Maguire et al. 2005). Moreover, GABA_A receptor plasticity throughout the ovarian cycle is accompanied to changes in sensitivity to exogenous 3 α ,5 α -THP; administration of 3 α ,5 α -THP potentiates tonic inhibition and exerts a protective action against hippocampus kindling epileptogenesis during the diestrus phase in female C57BL/6-129SV hybrid mice (Wu et al. 2013). Increased circulating levels of 3 α ,5 α -THP have been reported during the luteal phase of the menstrual cycle in women (Wang et al. 1996), and fluctuations in neuroactive steroid concentrations across the menstrual cycle correlate with symptoms of premenstrual dysphoric disorder (Girdler et al. 2001; Wang et al. 1996). Interestingly, treatment with hormonal contraceptives decreases plasma neuroactive steroids and prevents the increase in 3 α ,5 α -THP during the luteal phase in women (Follesa et al. 2002; Rapkin et al. 2006). The same treatment also dramatically decreased brain 3 α ,5 α -THP and progesterone concentrations, altered GABA_A receptor subunit expression and induced anxiety-like behavior in female Sprague-Dawley rats (Follesa et al. 2002; Porcu et al. 2012).

Neuroactive steroid concentrations increase dramatically during pregnancy in both rats and women (Concas et al. 1998; Gilbert Evans et al. 2005). Levels of progesterone and 3 α ,5 α -THP decrease immediately before parturition and return to baseline levels two days after parturition in Sprague-Dawley rats (Concas et al. 1998). These abrupt changes in steroid concentrations may contribute to post-partum depressive symptoms.

GABAergic neuroactive steroids and stress/HPA axis regulation

The hypothalamic-pituitary-adrenal (HPA) axis is regulated by numerous neurotransmitter systems and by negative feedback of steroid hormones. Activation of the HPA axis in response to acute stress increases the release of corticotrophin releasing hormone (CRH) from the hypothalamus that stimulates the release of adrenocorticotrophic hormone (ACTH) from the pituitary, which, in turn, stimulates the adrenal cortex to release glucocorticoids (cortisol in humans and corticosterone in rodents) as well as the GABAergic neuroactive steroids. The ability of these steroids to modulate HPA axis activation may play an important role in stress response, homeostasis and allostasis. In contrast, chronic stress leads to dysregulation of the HPA axis, a feature observed in several psychiatric and neurologic disorders, which are also associated with alterations in neuroactive steroid levels in plasma, cerebrospinal fluid or brain (Girdler and Klatzkin 2007; Morrow et al. 2006; Uzunova et al. 1998). The following sections will describe the effects of neuroactive steroids on the stress/HPA axis response in rats, mice and human subjects (see also Table 1 for a summary).

Homeostatic regulation of the HPA axis in rats

Acute swim stress increases brain and plasma concentrations of neuroactive steroids in Sprague-Dawley rats (Purdy et al. 1991). Likewise, other paradigms of acute stress also increase brain and plasma levels of GABAergic neuroactive steroids and their precursors in male Sprague-Dawley rats (Barbaccia et al. 1994; Barbaccia et al. 1996; Barbaccia et al. 1997; Vallee et al. 2000). The increase in neuroactive steroids is mediated by HPA activation as adrenalectomized rats show no circulating $3\alpha,5\alpha$ -THP and $3\alpha,5\alpha$ -THDOC levels, although brain levels of $3\alpha,5\alpha$ -THP are still detectable (Purdy et al. 1991). Thus, activation of the HPA axis ultimately leads to a release of glucocorticoids and neuroactive steroids from the adrenals. Glucocorticoids provide negative feedback upon the hypothalamus and pituitary. Likewise, GABAergic neuroactive steroids inhibit CRH production, release, ACTH release and subsequent corticosterone levels following stress in rats. $3\alpha,5\alpha$ -THP exerts a regulatory effect on HPA axis activity by reducing the neuroendocrine response to stress. While $3\alpha,5\alpha$ -THP per se appears not to be involved in basal CRH release, it counteracts the anxiety behavior induced by CRH administration in Wistar rats (Patchev et al. 1994). Moreover, $3\alpha,5\alpha$ -THP or $3\alpha,5\alpha$ -THDOC administration before stress attenuates the stress-induced increase in ACTH and corticosterone in Wistar rats (Owens et al. 1992; Patchev et al. 1996). In agreement, intracerebroventricular administration of $3\alpha,5\alpha$ -THP antiserum to male and female Wistar rats potentiated the corticosterone response to stress in prepubertal and adult rats, without affecting basal plasma corticosterone levels (Guo et al. 1995).

Under normal basal conditions, systemic administration of $3\alpha,5\alpha$ -THP to adult male non-stressed Sprague-Dawley rats increases hypothalamic CRH content as well as serum ACTH and corticosterone (Naert et al. 2007), supporting a regulatory role for this neuroactive steroid in HPA function, whereby $3\alpha,5\alpha$ -THP increases hormone levels in basal conditions and decreases them in stress-induced perturbations in order to restore homeostasis. Systemic administration of pregnenolone, DHEA and their sulfate metabolites also increased hypothalamic CRH and serum ACTH and corticosterone (Naert et al. 2007). All these

effects were rapid and likely mediated by a direct action of neuroactive steroids on GABA and glutamate neurotransmission in the hypothalamus that regulate HPA axis activation. Indeed, the hypothalamus contains the enzymes to locally synthesize $3\alpha,5\alpha$ -THP and is an area particularly rich in $3\alpha,5\alpha$ -THP immunoreactivity (Saalman et al. 2007).

Acute stress rapidly decreases GABAergic transmission in male Sprague-Dawley rats (Biggio et al. 2007), which leads to HPA axis activation. The elevations in neuroactive steroids levels that occur approximately 30 minutes after acute stress may represent a homeostatic mechanism to restore GABAergic inhibition upon the paraventricular nucleus of the hypothalamus, thus acting as a brake upon HPA axis activity.

GABAergic neuroactive steroid regulation of HPA axis function may differ across lifespan. In early development, administration of $3\alpha,5\alpha$ -THDOC to Wistar rat pups, exposed to repeated maternal separation, attenuated vocalizations and abolished the anxiety-like behavior observed in these rats during adulthood (Patchev et al. 1997). Moreover, pretreatment of Wistar rat pups with $3\alpha,5\alpha$ -THP or $3\alpha,5\alpha$ -THDOC exerts long-term protection against HPA responses to stress (Mitev et al. 2003; Patchev et al. 1997), suggesting that neuroactive steroids may protect the developing brain against adverse challenges.

A heightened HPA axis response to stress with prolonged glucocorticoid release has been reported in adolescent Sprague-Dawley rats (Lupien et al. 2009). However it is not clear whether alterations in neuroactive steroids may contribute to these changes in HPA responsivity since data is not available in adolescent animals and the effects of neuroactive steroids on the HPA axis appear to differ with genetic background.

Intracerebroventricular administration of $3\alpha,5\alpha$ -THP antiserum potentiated the stress response in prepubertal and adult Wistar rats, but not aged rats suggesting that the modulatory effects of $3\alpha,5\alpha$ -THP on the HPA response to stress are lost during aging (Guo et al. 1995). However, another study reported that aged male Sprague-Dawley rats show increased sensitivity to stress with greater elevations in cerebral cortical $3\alpha,5\alpha$ -THP and $3\alpha,5\alpha$ -THDOC levels following acute stress exposure, compared to adult rats (Barbaccia et al. 1998). These results may differ due to different stress paradigms or to the different rat strains.

Chronic stress exposure induces adaptations in HPA axis function that lead to hyper-responsiveness of the axis to stressful stimuli and to altered neuroactive steroid responses. Thus, adult male Sprague-Dawley rats subjected to social isolation, an animal model of anxiety disorders that generates chronic stress, show decreased basal levels of $3\alpha,5\alpha$ -THP and $3\alpha,5\alpha$ -THDOC in brain and plasma, but a greater neuroactive steroid response to a stress challenge, compared to group-housed rats (Serra et al. 2000). Alterations in HPA axis responsiveness and neurosteroidogenesis have also been found in adult male offspring of socially isolated Sprague-Dawley rats (Pisu et al. 2013) and in juvenile offspring of Long-Evans rats subjected to chronic gestational restraint stress (Paris and Frye 2011).

Orthodromic regulation of the HPA axis in C57BL/6J mice

In contrast to the findings in rats, acute swim stress decreases plasma concentrations of 3α , 5α -THP, measured by gas chromatography/mass spectrometry, 3α -hydroxysteroid dehydrogenase (3α -HSD) mRNA levels in the adrenal glands and 3α , 5α -THP immunoreactivity in various brain regions of male C57BL/6J mice (Maldonado-Devicci et al., this issue). Furthermore, recent studies have highlighted a novel action of GABAergic neuroactive steroids in HPA axis regulation. In C57BL/6J male mice, these steroids promote the physiological response to stress. Under basal conditions, local administration of 3α , 5α -THDOC into the paraventricular nucleus decreased circulating levels of corticosterone, consistent with the homeostatic regulation of the HPA axis by GABAergic neuroactive steroids. However, under stress conditions, administration of 3α , 5α -THDOC, 30 minutes before restraint stress, increased corticosterone levels, an effect abolished by finasteride pretreatment, suggesting that neurosteroidogenesis plays a critical role in HPA axis response to stress. Moreover, finasteride treatment before restraint stress decreases anxiety in the elevated plus maze test, suggesting that neurosteroidogenesis is required to mediate the anxiety-like behavior induced by stress in these mice (Sarkar et al. 2011). Under basal conditions, 3α , 5α -THDOC decreases the spontaneous firing of CRH neurons; in contrast, following restraint stress, 3α , 5α -THDOC increases the firing rate of CRH neurons. These excitatory actions of neuroactive steroids are mediated by GABA_A subunit-containing receptors, since that they are absent in subunit knock-out mice. Moreover, following stress, GABAergic transmission on CRH neurons switches from inhibitory to excitatory, due to a dephosphorylation and downregulation of the K⁺/Cl⁻ co-transporter KCC2, which may represent a novel target for stress and neuroactive steroid responses (Sarkar et al. 2011).

Paradoxical excitatory effects of neuroactive steroids have also been reported in mice during puberty. Thus, administration of 3α , 5β -THP to adolescent C57BL/6J female mice increased anxiety following restraint stress; this effect appears to be mediated by inhibition of the tonic GABAergic currents through $\alpha 4\beta 2\delta$ GABA_A receptors in the hippocampus (Shen et al. 2007). It is unknown if these excitatory actions of neuroactive steroids extend to other mouse strains, rats or humans.

The effects of HPA axis adaptations to chronic stress on neuroactive steroid actions are largely unexplored in mice. Similar to the above mentioned studies in rats, social isolation decreases 3α , 5α -THP content and 5α -reductase expression in frontal cortex, hippocampus, amygdala and olfactory bulb of adult male Swiss-Webster mice (Dong et al. 2001; Pibiri et al. 2008), suggesting a potential dysregulation at the HPA level. Moreover, it has recently been shown that poor maternal care, a model of early life adversity, blunts the 3α , 5α -THP-induced suppression of neuronal firing in CRH neurons in young pre-pubertal C57BL/6J-129SvSvJ mice (Gunn et al. 2013). Whether these changes are present in other mouse strains or persist in adult mice, remains to be determined.

Neuroactive steroid regulation of the HPA axis in mice may be more complex and not yet totally unraveled. As already mentioned, GABAergic neuroactive steroids have anticonvulsant properties in both rats (Reddy and Rogawski 2002) and mice [Swiss-Webster (Belelli et al. 1989; Gasior et al. 1997), NIH Swiss (Kokate et al. 1999; Kokate et al. 1994; Kokate et al. 1998; Reddy and Rogawski 2002) and C3HJ strains (Pericic et al. 2007)].

Interestingly, acute swim stress also exerts anticonvulsant actions. In male Sprague-Dawley rats, swim stress increases the seizure threshold along with circulating $3\alpha,5\alpha$ -THDOC levels; administration of the 5α -reductase inhibitor finasteride prevents the stress-induced increase in $3\alpha,5\alpha$ -THDOC levels and actually decreases the seizure threshold below that in non stressed rats (Reddy and Rogawski 2002). Acute swim stress also has anticonvulsant effects in CBA and C3HJ mice (Pericic et al. 2007; Pericic et al. 2000). Levels of GABAergic neuroactive steroids have not been measured in these mice, but they may likely play a role. Indeed, prior administration of finasteride does not alter seizure threshold in CBA mice (Pericic et al. 2000) and administration of $3\alpha,5\alpha$ -THP prior to swim stress potentiates its anticonvulsant effect (Pericic et al. 2007). Thus, acute swim stress might influence seizure susceptibility in both rats and mice through interactions with the GABAergic neuroactive steroids.

GABAergic neuroactive steroids and HPA axis in humans

HPA axis function can be tested in humans by measuring the hormonal response to pharmacological challenges with CRH, ACTH or dexamethasone. Thus, administration of CRH and ACTH to humans (mixed population of males and females) increased serum $3\alpha,5\alpha$ -THP, progesterone and DHEA concentrations, while administration of dexamethasone decreased serum levels of these steroids (Genazzani et al. 1998), suggesting a homeostatic regulation of the HPA axis that contributes to neuroactive steroid synthesis. In agreement with these results, we found that concentrations of DOC, the progesterone metabolite and precursor of the GABAergic neuroactive steroid $3\alpha,5\alpha$ -THDOC, were regulated by the HPA axis at hypothalamic, pituitary and adrenal levels in healthy men. Naloxone, CRH and ACTH challenges increased serum DOC levels, while dexamethasone challenge suppressed them. In contrast, concentrations of the excitatory neuroactive steroid pregnenolone sulfate were regulated by the adrenals but not the hypothalamus or pituitary; in fact, naloxone and CRH challenges failed to affect pregnenolone sulfate levels in healthy men, while ACTH and dexamethasone challenges increased and decreased such levels, respectively (Porcu et al. 2008).

Acute stress activates the HPA axis and increases serum $3\alpha,5\alpha$ -THP concentrations in human subjects (Droogleever Fortuyn et al. 2004; Girdler et al. 2001), similar to what has been shown in rats. However, others have also reported no change in serum $3\alpha,5\alpha$ -THP levels after acute stress, albeit different stress paradigms and time points were examined (Altemus et al. 2001; Childs and de Wit 2009; Childs et al. 2010; Girdler et al. 2006). Most of the studies in humans have focused on dysregulation of the HPA axis associated to several psychiatric and neurologic conditions. For instance, a blunted HPA axis function and altered neuroactive steroid levels are reported in premenstrual dysphoric disorder (PMDD), although the literature is controversial, with reports of increases, decreases, and no change in $3\alpha,5\alpha$ -THP plasma levels (Girdler and Klatzkin 2007). Following mental stress, $3\alpha,5\alpha$ -THP levels increased in control subjects, as expected, but decreased in PMDD women and this lack of $3\alpha,5\alpha$ -THP responsiveness to stress was related to the greater baseline $3\alpha,5\alpha$ -THP concentrations found in these subjects (Girdler et al. 2001). Moreover, women with a history of depression, regardless of PMDD symptoms, had reduced basal neuroactive steroid levels

and a blunted 3 α ,5 α -THP response to stress (Klatzkin et al. 2006). This work is reviewed in detail in another contribution to this Special Issue (Crowley and Girdler, this issue).

Decreased 3 α ,5 α -THP levels have been found in cerebrospinal fluid of women with posttraumatic stress disorder (Rasmusson et al. 2006) and chronic stress exposure may also affect neuroactive steroid sensitivity. A recent study has shown that women with work related psychosocial stress have similar baseline concentrations of 3 α ,5 α -THP compared to controls, but greater sensitivity to exogenous 3 α ,5 α -THP. In fact, 3 α ,5 α -THP administration decreased the saccadic eye velocity to a greater extent in patients compared to controls (Backstrom et al. 2013).

Alterations in HPA axis responsiveness are also found in alcoholism during drinking and abstinence (Morrow and Porcu 2009). Basal levels of 3 α ,5 α -THP and 3 α ,5 α -THDOC are reduced during ethanol withdrawal in humans (Romeo et al. 1996), when circulating cortisol levels are elevated. Their responses to stress have not been studied under these conditions, but are likely to be impacted since both human and animal studies show that neuroactive steroid responses to stress or HPA axis activation are blunted when ACTH or glucocorticoid responses are blunted (Morrow and Porcu 2009). Furthermore, abstinent alcoholics show a blunted pregnenolone sulfate response to adrenal stimulation and a delayed DOC response to CRH challenge, supporting the idea that blunting of the HPA axis also impacts the neuroactive steroid responses to stress in alcoholics (Porcu et al. 2008).

All this experimental evidence emphasizes the important link between HPA axis function and neuroactive steroid levels in the maintenance of homeostasis and healthy brain function.

GABAergic neuroactive steroids and ethanol interactions

The effects of ethanol involve various GABAergic mechanisms that contribute to many of its behavioral effects, including anxiolytic, anticonvulsant, sedative-hypnotic, cognitive-impairing and motor incoordinating actions (see Kumar et al. 2009, for review). These mechanisms are thought to include direct and indirect effects of ethanol on GABA_A receptors as well as effects on GABA release and the synthesis and availability of endogenous neuroactive steroids. This work was primarily based on studies in rat models, but certain effects of ethanol differ across various rat and mouse strains. Here, we focus on the ethanol regulation of GABAergic neuroactive steroid levels in rats as well as across mouse strains and demonstrate how genetic variation in neuroactive steroid responses can be linked to variation in the behavioral actions of ethanol.

Ethanol increases GABAergic neuroactive steroids in rats

Like stress, systemic administration of moderate ethanol doses (1-2.5 g/kg) increases plasma, cerebrocortical and hippocampal levels of 3 α ,5 α -THP and 3 α ,5 α -THDOC in Sprague-Dawley rats (Morrow et al. 1999; Morrow et al. 1998; Porcu et al. 2010; VanDoren et al. 2000) and Sardinian alcohol-preferring rats (Barbaccia et al. 1999). Recent studies using immunohistochemistry to determine ethanol-induced changes in cellular 3 α ,5 α -THP across brain show both regional and cell population specificity in this response in Wistar rats (Cook et al. 2014a). Ethanol increased 3 α ,5 α -THP immunoreactivity in the medial

prefrontal cortex, the hippocampal CA1 pyramidal cell layer, the polymorph cell layer of the dentate gyrus, the bed nucleus of the stria terminalis, and the paraventricular nucleus of the hypothalamus. In contrast, ethanol administration significantly reduced 3 α ,5 α -THP immunoreactivity in the nucleus accumbens and the central nucleus of the amygdala. No changes were observed in the ventral tegmental area, dorsomedial striatum, granule cell layer of the dentate gyrus, or the lateral or basolateral amygdala (Cook et al. 2014a). Therefore, ethanol produces divergent brain region and cell-type specific changes in 3 α ,5 α -THP concentrations.

The ethanol-induced increase in plasma neuroactive steroids is mediated by the HPA axis, since it is no longer observed in hypophysectomized and adrenalectomized Sprague-Dawley (Boyd et al. 2010; Khisti et al. 2003; Porcu et al. 2004) or Wistar (O'Dell et al. 2004) rats. However, studies of ethanol effects using immunohistochemistry reveal that ethanol-induced elevations of 3 α ,5 α -THP are independent of adrenal activation in the CA1 pyramidal cell layer, dentate gyrus polymorphic layer, bed nucleus of the stria terminalis, and paraventricular nucleus of the hypothalamus of Wistar rats. Furthermore, ethanol produced decreases in 3 α ,5 α -THP labeling in the nucleus accumbens shell and central nucleus of the amygdala that also occurred independent of adrenal activation. However, in the medial prefrontal cortex ethanol increased 3 α ,5 α -THP immunoreactivity after sham surgery, but there was no change of 3 α ,5 α -THP after adrenalectomy. These data indicate ethanol dynamically regulates local 3 α ,5 α -THP levels in several subcortical regions, but the adrenal glands contribute to 3 α ,5 α -THP elevations in the medial prefrontal cortex (Cook et al. 2014b).

These data in the hippocampal formation agree with the previous *in vitro* studies showing that ethanol induced local brain synthesis of 3 α ,5 α -THP. For example, it was first shown that incubation with ethanol (50 or 100 mM) can increase 3 α ,5 α -THP levels (measured by radioimmunoassay) and GABAergic transmission in hippocampal minces from intact Sprague-Dawley rats (Sanna et al. 2004) and those that had undergone adrenalectomy/gonadectomy (Follesa et al. 2006). More recently, it was shown that ethanol increases cellular 3 α ,5 α -THP in CA1 pyramidal cells in the slice preparation from juvenile Sprague-Dawley rats (Tokuda et al. 2011). Clearly, ethanol's ability to stimulate brain synthesis of 3 α ,5 α -THP in the hippocampal formation is isolated to specific cellular populations, since we did not previously observe ethanol-induced changes of 3 α ,5 α -THP in the granule cell layer of the dentate gyrus (Cook et al. 2014a). The CA1 pyramidal cells, and the polymorphic and granule cell layers of the dentate gyrus all exhibit dense 3 α ,5 α -THP staining. Therefore, the presence of this very specific effect of ethanol on cellular 3 α ,5 α -THP in the hippocampus is intriguing and may underlie neuron specific responses to ethanol in the hippocampal formation.

Ethanol-induced elevations in GABAergic steroids reach physiologically relevant concentrations that are capable of enhancing GABAergic transmission (Barbaccia et al. 1999; Morrow et al. 1999; Morrow et al. 1998; VanDoren et al. 2000). A large body of evidence from multiple laboratories suggests that ethanol-induced elevations of GABAergic neuroactive steroids contribute to many behavioral effects of ethanol in rodents. These steroids have been shown to modulate ethanol's anticonvulsant effects (VanDoren et al.

2000), sedation (Khisti et al. 2003), impairment of spatial memory (Matthews et al. 2002; Morrow et al. 2001), anxiolytic-like (Hirani et al. 2005) and antidepressant-like (Hirani et al. 2002) actions. Each of these behavioral responses is prevented by pretreatment with the neuroactive steroid biosynthesis inhibitor finasteride and/or by prior adrenalectomy. The hypnotic effect of ethanol is partially blocked by adrenalectomy. Importantly, administration of 5 α -dihydroprogesterone, the immediate precursor of 3 α ,5 α -THP, to adrenalectomized rats restores effects of ethanol, showing that brain synthesis of neuroactive steroids modulates effects of ethanol (Khisti et al. 2003). However, neuroactive steroids do not appear to influence the motor incoordinating effects of ethanol, since neither finasteride administration nor adrenalectomy diminish these actions (Khisti et al. 2004). Taken together, these studies suggest that elevations in neuroactive steroids influence many of the GABAergic effects of ethanol *in vivo* and contribute to sensitivity to behavioral effects of ethanol. We have suggested that ethanol-induced elevations of GABAergic neuroactive steroids protect against the risk for ethanol dependence (Morrow 2007; Morrow et al. 2006). Diminished elevations of GABAergic neuroactive steroids following ethanol exposure would result in reduced sensitivity to the anxiolytic, sedative, anticonvulsant, cognitive-impairing and discriminative stimulus properties of ethanol. Reduced sensitivity to ethanol is associated with greater risk for the development of alcoholism in individuals with genetic vulnerability to alcoholism (Schuckit 1994; 2009). Further studies are needed to test this hypothesis.

Strain dependent effects of ethanol on GABAergic neuroactive steroids in mice

Studies regarding the effects of ethanol on neuroactive steroids in mice have yielded different results depending on which strains have been used. Among the most studied mouse strains are C57BL/6J and DBA/2J. We recently reported that C57BL/6J mice have higher basal plasma neuroactive steroid levels compared to DBA/2J mice and that plasma 3 α ,5 α -THP levels are decreased in C57BL/6J mice and not altered in DBA/2J mice following injection of 2 g/kg ethanol (Porcu et al. 2010). Moreover, acute ethanol administration failed to alter cerebrocortical and hippocampal levels of 3 α ,5 α -THP and progesterone in naïve male C57BL/6J and DBA/2J mice (Porcu et al. 2014). Other studies have reported that injection of 2 g/kg ethanol increased whole brain 3 α ,5 α -THP levels in male DBA/2J (Gabriel et al. 2004) but not C57BL/6J mice (Finn et al. 2004b), while orally consumed ethanol increases whole brain 3 α ,5 α -THP levels in male C57BL/6J mice (Finn et al. 2004b).

These two strains differ in the behavioral response to neuroactive steroids: C57BL/6J mice are more sensitive to the anxiolytic, locomotor stimulant and anticonvulsant effects of 3 α ,5 α -THP compared to DBA/2J mice (Finn et al. 1997b). However, similar discriminative stimulus effects of its 5 isomer pregnanolone were reported for both strains (Shannon et al. 2005a; Shannon et al. 2005b). C57BL/6J and DBA/2J mice are also two of the most widely used inbred strains in addiction research because they show different sensitivity to several drugs of abuse, including ethanol (Belknap et al. 1993a; Belknap et al. 1993b). For instance, DBA/2J mice have more severe handling-induced convulsions after withdrawal from acute (Roberts et al. 1992) and chronic (Crabbe et al. 1983) ethanol, have greater acute functional tolerance (Gallaher et al. 1996), greater locomotor activity (Crabbe et al. 1983), and greater hypnosis (Linsenbardt et al. 2009) compared to C57BL/6J mice. These strains also differ in

ethanol preference and consumption. In the two-bottle choice paradigm, C57BL/6J mice drink alcohol, while DBA/2J mice avoid it (Belknap et al. 1977; Phillips et al. 1994). However, when allowed to self-administer ethanol intravenously, these two strains do not differ in the amount of ethanol self-administered, suggesting that ethanol's reinforcing properties are similar in both strains (Grahame and Cunningham 1997). It has been suggested that endogenous neuroactive steroids may play a role in ethanol sensitivity in these strains. In fact, 3 α ,5 α -THP modulates ethanol intake (Ford et al. 2007; Ford et al. 2005), reinstates ethanol seeking behavior in C57BL/6J mice (Finn et al. 2008) and affects ethanol withdrawal severity (Finn et al. 2004a; Gililand and Finn 2007).

Effects of ethanol on GABAergic neuroactive steroids in humans

The role of neuroactive steroids in alcohol action in humans is relatively unexplored and inconsistent. Studies in humans have shown that male and female adolescents seen in the emergency room for alcohol intoxication had elevated plasma levels of the neuroactive steroid 3 α ,5 α -THP (Torres and Ortega 2003; 2004). Although the peak ethanol concentrations were not determined in these studies, it is likely that high ethanol concentrations were achieved since the subjects were taken to the emergency room for treatment of alcohol intoxication. In contrast, laboratory administration of low or moderate doses of ethanol does not alter plasma levels of 3 α ,5 α -THP and other GABAergic neuroactive steroids (Holdstock et al. 2006; Porcu et al. 2010) or may decrease 3 α ,5 α -THP levels (Nyberg et al. 2005; Pierucci-Lagha et al. 2006). The basis of these conflicting results are unknown, but may involve pharmacologically different ethanol doses, different analytic methods to measure neuroactive steroids or environmental factors that influence neuroactive steroid synthesis in humans. Indeed, the same dose of ethanol consumed in the human laboratory studies above (~80 mg/dl) produced no effect in rats when administered systemically (Porcu et al. 2010), suggesting that dose might be a key factor in the difference between rat and human studies.

Despite these results, other studies suggest the importance of GABAergic neuroactive steroids in ethanol's actions in humans. In fact, various subjective effects of ethanol, measured during the rising phase of the blood alcohol curve, are diminished by prior administration of the neuroactive steroid biosynthesis inhibitor finasteride (Pierucci-Lagha et al. 2005). Finasteride prevents the formation of the 3 α ,5 α -reduced metabolites of progesterone, DOC, DHEA and testosterone, thus blocking 3 α ,5 α -THP synthesis (Azzolina et al. 1997). Indeed, finasteride pretreatment blocked subjective effects of alcohol using three different scales to measure the activating, sedating, anesthetic and peripheral dynamic aspects of alcohol actions. The ability of finasteride to reduce the subjective effects of alcohol was not observed in individuals carrying the GABA_A 2 subunit polymorphism associated with alcoholism, and individuals carrying this polymorphism had reduced sensitivity to alcohol as well as finasteride (Pierucci-Lagha et al. 2005). Covault and collaborators (this issue) further show that inhibition of 5 α -reductase with dutasteride also reduced the sedative and anesthetic effects of alcohol in men classified as moderate-heavy drinkers. Interestingly, this study, which was conducted in a larger population, did not find any association of dutasteride effects with the GABA_A α 2 subunit polymorphism. Indeed all these observations are consistent with the idea that GABAergic neuroactive steroid

responses contribute to ethanol sensitivity in humans and may be a risk factor for alcoholism.

Covault and collaborators (this issue) also report that dutasteride administration reduced subsequent alcohol consumption in those subjects classified as heavy drinkers. In agreement, it has recently been shown that male subjects, who took finasteride for treatment of male pattern hair loss, reported a decrease in alcohol consumption, which was greater in those subjects who consumed the most alcohol. These subjects also reported increased anxiety, tiredness, dizziness, and in few cases even intoxication, after alcohol exposure, all factors that contributed to their alcohol dislike (Irwig 2013). Taken together, these results further support the hypothesis that neuroactive steroids mediate subjective effects of alcohol in humans.

Studies on GABAergic neuroactive steroids in alcohol dependence have shown that 3 α ,5 α -THP and 3 α ,5 α -THDOC levels are decreased in serum of alcoholics during the peak of alcohol withdrawal and return to normal levels upon recovery (Romeo et al. 1996). Likewise, abstinent alcoholics exhibit diminished progesterone levels as well as a lowered ratio of progesterone to pregnenolone (Hill et al. 2005).

Some recent studies have investigated the role of genetic polymorphisms in alcohol dependence with a focus on neuroactive steroids. In alcohol dependent subjects, naltrexone administration, compared to placebo, increased 3 α ,5 α -THP serum levels among individuals with the Asp40 allele of the OPRM1 gene (AG/GG) but not among those carrying the Asn40 allele (Ray et al. 2010). In this population, no correlation was found between basal 3 α ,5 α -THP levels and average alcohol use frequency, quantity or "Alcohol Use Disorders Identification Test (AUDIT)" scores. Moreover, basal 3 α ,5 α -THP levels were not influenced by smoking status, sex, ethnicity, family history of alcoholism or OPRM1 status. These results highlight a potential contribution of neuroactive steroids to the pharmacological effects of naltrexone in alcohol dependent subjects that lack the OPRM1 polymorphism associated with therapeutic efficacy to naltrexone (Ray et al. 2010). Future studies will be required to further examine these relationships.

Risk of alcohol dependence was found to be associated with polymorphic variation in the enzymes 5 α -reductase and 3 α -HSD implicated in the conversion of progesterone and DOC to their neuroactive metabolites 3 α ,5 α -THP and 3 α ,5 α -THDOC. The minor C-allele for SRD5A1 exon 1 SNP rs248793, the gene encoding for 5 α -reductase, and the minor G-allele for AKR1C3*2 exon 1 SNP rs12529, the gene encoding for 3 α -HSD, were found to be more frequent in control subjects compared to alcohol dependent subjects. These results suggest that the minor allele for 5 α -reductase type I and 3 α -HSD type 2 genes may have a protective effect against alcohol dependence and further provide indirect evidence that neuroactive steroids may play a role in alcohol effects in humans (Milivojevic et al. 2011).

Genetic variation in neuroactive steroid levels across BXD recombinant inbred strains

Individual differences in vulnerability to alcoholism have a genetic component (Schuckit 2009). Studies in rodents indicate a shared genetic sensitivity to ethanol, anxiety and stress/HPA axis response (Boehm et al. 2002; Crabbe et al. 1999). We have identified the genetic regulation of basal levels of the neuroactive steroid DOC across the C57BL/6J (B6) × DBA/2J (D2) (BXD) recombinant inbred mouse strains, a reference population to study networks of phenotypes and their modulation by gene variants (Gora-Maslak et al. 1991; Williams et al. 2001). DOC is an endogenous neuroactive steroid, synthesized from progesterone mainly in the adrenal glands but also in the brain; it is the precursor of the GABAergic neuroactive steroid 3 α ,5 α -THDOC and of the glucocorticoid corticosterone. DOC levels are elevated in rat brain and mouse plasma following acute ethanol administration (Khisti et al. 2005; Porcu et al. 2010), are regulated by hypothalamic and pituitary activation of the HPA axis in both cynomolgus monkeys and humans and this regulation is altered following ethanol dependence (Porcu et al. 2006; Porcu et al. 2008). Across the BXD mouse population we found significant genetic variation in basal DOC levels. Cerebral cortical DOC levels ranged between 1.4 and 12.2 ng/g, resulting in a 8.7-fold genetic variation and heritability of 0.37. Basal plasma DOC levels ranged between 2.8 and 12.1 ng/ml, resulting in a 4.3-fold genetic variation and heritability of 0.32. We mapped variation in basal DOC levels using genetic and bioinformatics tools in GeneNetwork (www.genenetwork.org), a public repository of genetic and phenotypic data as well as a resource for multivariate genetic analysis of complex traits in genetic reference populations (Chesler et al. 2005; Wang et al. 2003). We identified quantitative trait loci (QTLs) on chromosomes 4 and 14, for basal DOC levels in cerebral cortex and plasma, respectively (Porcu et al. 2011). Moreover, variation in basal DOC levels in both plasma and cerebral cortex was linked to several ethanol and anxiety phenotypes previously characterized across the BXD strains by several groups and whose data is available in GeneNetwork. Thus, basal DOC levels are positively correlated with increased ethanol-induced sedation, ethanol-induced ataxia, and ethanol-induced corticosterone levels. These correlations are of particular importance because studies in rats and humans have suggested that neuroactive steroids may play a role in alcohol sensitivity (Morrow et al. 2006). The finding that mouse strains with higher basal DOC levels show a greater ethanol sensitivity is consistent with the hypothesis that elevated GABAergic neuroactive steroids may protect against the risk for alcohol dependence (Morrow and Porcu 2009; Morrow et al. 2006). Diminished elevations of neuroactive steroids following ethanol exposure would result in reduced sensitivity to the anxiolytic, sedative, anticonvulsant, cognitive-impairing, and discriminative stimulus properties of ethanol (Morrow et al. 2006). Reduced sensitivity to ethanol is associated with greater risk for the development of alcoholism in individuals with genetic vulnerability to alcoholism (Schuckit 1994; Wilhelmsen et al. 2003). Variation in basal DOC levels was also correlated to some anxiety phenotypes and seizure susceptibility, in agreement with the fact that neuroactive steroids exert anxiolytic and anticonvulsant properties (Belelli et al. 1989; Bitran et al. 1991; Reddy and Rogawski 2002) and that their levels are altered in several psychiatric diseases involving stress and anxiety (Eser et al. 2006; Girdler et al. 2001).

Given that many steroids, including DOC, function as gene regulatory molecules by acting on nuclear receptors (McEwen 1991), we used bioinformatics analysis to examine correlations between cerebral cortical DOC levels and whole brain gene expression profiles across the BXD strains. The genetic network analysis revealed the involvement of multiple genes regulating inflammation, vesicle trafficking and nuclear receptor signaling, supporting previous evidence for a role of neuroactive steroids in inflammation, neurotransmitter release and synaptic function (Belelli and Lambert 2005; Brinton 2013; Carver and Reddy 2013; Meyer et al. 2002; Mtchedlishvili and Kapur 2003; Schumacher et al. 2007).

We hypothesized that basal and ethanol-induced $3\alpha,5\alpha$ -THP levels may differ across BXD strains and may contribute to a network of alcohol-related phenotypes. Thus, based on the genetic variation in basal DOC levels (Porcu et al. 2011), we collected preliminary data on selected BXD strains that were chosen to capture the range of variation in neuroactive steroid content. We measured $3\alpha,5\alpha$ -THP levels in the cerebral cortex of these mice, 1 hour after acute ethanol administration (2 g/kg, i.p.) or saline (basal levels). We found that basal cerebral cortical $3\alpha,5\alpha$ -THP levels across strains ranged between 1.81 and 3.72 ng/g, equivalent to a 2.0-fold genetic variation [$F(9,79) = 6.27$, $p < 0.0001$] and heritability of 0.40 (Figure 2a). The ethanol-induced changes in cerebral cortical $3\alpha,5\alpha$ -THP levels ranged between +4% and +63% (Figure 2b).

We further examined genetic correlations with behavioral phenotypes previously determined in the BXD strains and available in GeneNetwork. Both basal and ethanol-induced cerebral cortical $3\alpha,5\alpha$ -THP levels were correlated with some ethanol phenotypes. Basal $3\alpha,5\alpha$ -THP levels were negatively correlated with consumption of 10% ethanol at 2 hours [Spearman $r = -0.82$, $p = 0.02$, $n = 7$, GeneNetwork ID 12733, (Cook et al., unpublished), Figure 3a], that is those strains with low basal $3\alpha,5\alpha$ -THP levels consumed more alcohol. Interestingly, the ethanol-induced changes in $3\alpha,5\alpha$ -THP levels were negatively correlated with ethanol consumption both of 3% [Spearman $r = -0.82$, $p = 0.02$, $n = 7$, GeneNetwork ID 10474, (Phillips et al. 1994), Figure 3b] and 10% [Spearman $r = -0.82$, $p = 0.02$, $n = 7$, GeneNetwork ID 10582, (Rodriguez et al. 1994), Figure 3c]. Those strains with increased $3\alpha,5\alpha$ -THP levels in response to acute ethanol consumed less alcohol. This finding, albeit observed in a small number of strains, supports the hypothesis that neuroactive steroid responses to ethanol may be associated with excessive alcohol consumption. Future studies are required to expand and validate these preliminary results.

Genetic variability in the neuroactive steroid response to ethanol has also been shown for corticosterone. DBA/2J mice display greater plasma ACTH and corticosterone responses to acute ethanol administration compared to C57BL/6J mice, suggesting that the HPA axis in DBA/2J mice is more responsive to ethanol (Roberts et al. 1992). In agreement, the increase in cerebral cortical corticosterone levels, induced by acute ethanol, was greater in DBA/2J mice compared to C57BL/6J mice (Porcu et al. 2014). The effects of acute ethanol administration on plasma corticosterone levels have been examined across the BXD strains: an effect of strain on plasma corticosterone was found 1 hour following ethanol administration, but not following saline injection; the genetic analysis suggested that a single gene may account for most of the genetic variability in the corticosterone response to ethanol at this time point (Roberts et al. 1995). Moreover, the corticosterone response to

ethanol was negatively correlated with consumption of 3% ethanol (Roberts et al. 1995), suggesting that those strains with an increased stress response to acute ethanol consumed less alcohol, similar to our preliminary observation on the ethanol-induced 3 α ,5 α -THP responses.

Thus, genetic variation may contribute to the differences in the neuroactive steroid responses to stress or ethanol challenges. These studies have mainly been conducted in mouse strains, but it is likely that genetic variation could also explain the different neuroactive steroid sensitivity observed in rats and humans, compared to C57BL/6J mice.

Summary and Conclusions

Rats and mouse strains clearly show divergent effects of stress and ethanol on GABAergic neuroactive steroids in both plasma and brain. The molecular mechanisms that underlie these differences are not known and have received very little attention by the field. Endocrine differences between rodents and humans, including the ovarian cycle or the predominant type of neuroactive steroid must be taken into account. Moreover, the amount and chronicity of alcohol exposure that underlies addiction to alcohol also differs between rodent and human studies. How then do we choose an appropriate animal model to better understand the potential role of these neuroactive steroids in various physiological processes in humans? At first glance, it may appear that GABAergic neuroactive steroid responses to stress in humans are more similar to rats than C57BL/6J mice. This may be true with respect to the populations of humans that have been included in laboratory studies. However, it is important to recognize that many individuals are excluded from human studies due to various neuropsychiatric diagnoses or even age restrictions. C57BL/6J mice are widely used in biomedical research, particularly due to their selection for transgenic and knock-out strategies. However, these mice exhibit phenotypes such as excessive drinking, aggression and other traits that would likely result in exclusion from laboratory studies in human populations. Perhaps C57BL/6J mice model some human populations while common laboratory rats better model other human populations. One approach to address these issues lies in the study of genetic variation in the various processes that determine neuroactive steroids levels as well as their effects on cell signaling. However, translation from mouse to human applications of GABAergic neuroactive steroids clearly presents major hurdles that will require further research.

Neuroactive steroids may have great potential for therapeutic development. It will be important however to capitalize on human genetic diversity with respect to neuroactive steroid levels and responses to stress in humans, moving closer to a personalized medicine approach in the field. We must consider human genetic diversity in this field, in order to be successful in our attempts to utilize neuroactive steroids for therapeutic benefits.

Acknowledgments

This work was supported by UO1-AA016672 to ALM, UO1-AA013641 to PP, and by the Bowles Center for Alcohol Studies. We acknowledge the myriad contributions of Robert H. Purdy to the field of neuroactive steroids in whose memory we dedicate this work.

References

- Akk G, Bracamontes J, Steinbach JH. Pregnenolone sulfate block of GABA_A receptors: mechanism and involvement of a residue in the M2 region of the alpha subunit. *J Physiol.* 2001; 532:673–684. [PubMed: 11313438]
- Akk G, Covey DF, Evers AS, Steinbach JH, Zorumski CF, Mennerick S. Mechanisms of neurosteroid interactions with GABA_A receptors. *Pharmacol Ther.* 2007; 116:35–57. [PubMed: 17524487]
- Akk G, Shu HJ, Wang C, Steinbach JH, Zorumski CF, Covey DF, Mennerick S. Neurosteroid access to the GABA_A receptor. *J Neurosci.* 2005; 25:11605–11613. [PubMed: 16354918]
- Altemus M, Redwine LS, Leong YM, Frye CA, Porges SW, Carter CS. Responses to laboratory psychosocial stress in postpartum women. *Psychosom Med.* 2001; 63:814–821. [PubMed: 11573030]
- Azzolina B, Ellsworth K, Andersson S, Geissler W, Bull HG, Harris GS. Inhibition of rat alphareductases by finasteride: evidence for isozyme differences in the mechanism of inhibition. *J Steroid Biochem Mol Biol.* 1997; 61:55–64. [PubMed: 9328210]
- Backstrom T, Bixo M, Nyberg S, Savic I. Increased neurosteroid sensitivity--an explanation to symptoms associated with chronic work related stress in women? *Psychoneuroendocrinology.* 2013; 38:1078–1089. [PubMed: 23177572]
- Barbaccia ML, Affricano D, Trabucchi M, Purdy RH, Colombo G, Agabio R, Gessa GL. Ethanol markedly increases "GABAergic" neurosteroids in alcohol-preferring rats. *Eur J Pharmacol.* 1999; 384:R1–R2. [PubMed: 10611449]
- Barbaccia ML, Concas A, Serra M, Biggio G. Stress and neurosteroids in adult and aged rats. *Exp Gerontol.* 1998; 33:697–712. [PubMed: 9951617]
- Barbaccia ML, Roscetti G, Trabucchi M, Cuccheddu T, Concas A, Biggio G. Neurosteroids in the brain of handling-habituated and naive rats: effect of CO₂ inhalation. *Eur J Pharmacol.* 1994; 261:317–320. [PubMed: 7813554]
- Barbaccia ML, Roscetti G, Trabucchi M, Mostallino MC, Concas A, Purdy RH, Biggio G. Time-dependent changes in rat brain neuroactive steroid concentrations and GABA_A receptor function after acute stress. *Neuroendocrinology.* 1996; 63:166–172. [PubMed: 9053781]
- Barbaccia ML, Roscetti G, Trabucchi M, Purdy RH, Mostallino MC, Concas A, Biggio G. The effects of inhibitors of GABAergic transmission and stress on brain and plasma allopregnanolone concentrations. *Br J Pharmacol.* 1997; 120:1582–1588. [PubMed: 9113382]
- Belelli D, Bolger MB, Gee KW. Anticonvulsant profile of the progesterone metabolite 5 α -pregnan-3 α -ol-20-one. *Eur J Pharmacol.* 1989; 166:325–329. [PubMed: 2792198]
- Belelli D, Lambert JJ. Neurosteroids: endogenous regulators of the GABA_A receptor. *Nat Rev Neurosci.* 2005; 6:565–575. [PubMed: 15959466]
- Belknap JK, Belknap ND, Berg JH, Coleman R. Preabsorptive vs. postabsorptive control of ethanol intake in C57BL/6J and DBA/2J mice. *Behav Genet.* 1977; 7:413–425. [PubMed: 603475]
- Belknap JK, Crabbe JC, Riggan J, O'Toole LA. Voluntary consumption of morphine in 15 inbred mouse strains. *Psychopharmacology (Berl).* 1993a; 112:352–358. [PubMed: 7871041]
- Belknap JK, Crabbe JC, Young ER. Voluntary consumption of ethanol in 15 inbred mouse strains. *Psychopharmacology (Berl).* 1993b; 112:503–510. [PubMed: 7871064]
- Besheer J, Lindsay TG, O'Buckley TK, Hodge CW, Morrow AL. Pregnenolone and ganaxolone reduce operant ethanol self-administration in alcohol-preferring p rats. *Alcohol Clin Exp Res.* 2010; 34:2044–2052. [PubMed: 20946297]
- Biggio G, Concas A, Follsea P, Sanna E, Serra M. Stress, ethanol, and neuroactive steroids. *Pharmacol Ther.* 2007; 116:140–171. [PubMed: 17555824]
- Bitran D, Hilvers RJ, Kellogg CK. Anxiolytic effects of 3 α -hydroxy-5 α [β]-pregnan-20-one: endogenous metabolites of progesterone that are active at the GABA_A receptor. *Brain Res.* 1991; 561:157–161. [PubMed: 1686744]
- Bixo M, Andersson A, Winblad B, Purdy RH, Backstrom T. Progesterone: 5 α -pregnane-3,20-dione and 3 α -hydroxy-5 α -pregnane-20-one in specific regions of the human female brain in different endocrine states. *Brain Res.* 1997; 764:173–178. [PubMed: 9295207]

- Boehm SL 2nd, Reed CL, McKinnon CS, Phillips TJ. Shared genes influence sensitivity to the effects of ethanol on locomotor and anxiety-like behaviors, and the stress axis. *Psychopharmacology (Berl)*. 2002; 161:54–63. [PubMed: 11967631]
- Boyd KN, Kumar S, O'Buckley TK, Porcu P, Morrow AL. Ethanol induction of steroidogenesis in rat adrenal and brain is dependent upon pituitary ACTH release and de novo adrenal StAR synthesis. *J Neurochem*. 2010; 112:784–796. [PubMed: 20021565]
- Brinton RD. Neurosteroids as regenerative agents in the brain: therapeutic implications. *Nat Rev Endocrinol*. 2013; 9:241–250. [PubMed: 23438839]
- Carver CM, Reddy DS. Neurosteroid interactions with synaptic and extrasynaptic GABA_A receptors: regulation of subunit plasticity, phasic and tonic inhibition, and neuronal network excitability. *Psychopharmacology (Berl)*. 2013; 230:151–188. [PubMed: 24071826]
- Chesler EJ, Lu L, Shou S, Qu Y, Gu J, Wang J, Hsu HC, Mountz JD, Baldwin NE, Langston MA, Threadgill DW, Manly KF, Williams RW. Complex trait analysis of gene expression uncovers polygenic and pleiotropic networks that modulate nervous system function. *Nat Genet*. 2005; 37:233–242. [PubMed: 15711545]
- Childs E, de Wit H. Hormonal, cardiovascular, and subjective responses to acute stress in smokers. *Psychopharmacology (Berl)*. 2009; 203:1–12. [PubMed: 18936915]
- Childs E, Dlugos A, De Wit H. Cardiovascular, hormonal, and emotional responses to the TSST in relation to sex and menstrual cycle phase. *Psychophysiology*. 2010; 47:550–559. [PubMed: 20070572]
- Concas A, Mostallino MC, Porcu P, Follesa P, Barbaccia ML, Trabucchi M, Purdy RH, Grisenti P, Biggio G. Role of brain allopregnanolone in the plasticity of γ -aminobutyric acid type A receptor in rat brain during pregnancy and after delivery. *Proc Natl Acad Sci USA*. 1998; 95:13284–13289. [PubMed: 9789080]
- Cook JB, Dumitru AM, O'Buckley TK, Morrow AL. Ethanol administration produces divergent changes in GABAergic neuroactive steroid immunohistochemistry in the rat brain. *Alcohol Clin Exp Res*. 2014a; 38:90–99. [PubMed: 23906006]
- Cook JB, Nelli SM, Neighbors MR, Morrow DH, O'Buckley TK, Maldonado-Devincci AM, Morrow AL. Ethanol alters local cellular levels of (3 α , 5 α)-3-hydroxypregnan-20-one (3 α , 5 α -THP) independent of the adrenals in subcortical brain regions. *Neuropsychopharmacology*. 2014b Epub Feb 25;
- Corpechot C, Collins BE, Carey MP, Tsouros A, Robel P, Fry JP. Brain neurosteroids during the mouse oestrous cycle. *Brain Res*. 1997; 766:276–280. [PubMed: 9359616]
- Crabbe JC, Kosobud A, Young ER. Genetic selection for ethanol withdrawal severity: differences in replicate mouse lines. *Life Sci*. 1983; 33:955–962. [PubMed: 6684200]
- Crabbe JC, Phillips TJ, Buck KJ, Cunningham CL, Belknap JK. Identifying genes for alcohol and drug sensitivity: recent progress and future directions. *Trends Neurosci*. 1999; 22:173–179. [PubMed: 10203855]
- Dong E, Matsumoto K, Uzunova V, Sugaya I, Takahata H, Nomura H, Watanabe H, Costa E, Guidotti A. Brain 5 α -dihydroprogesterone and allopregnanolone synthesis in a mouse model of protracted social isolation. *Proc Natl Acad Sci USA*. 2001; 98:2849–2854. [PubMed: 11226329]
- Droogleever Fortuyn HA, van Broekhoven F, Span PN, Backstrom T, Zitman FG, Verkes RJ. Effects of PhD examination stress on allopregnanolone and cortisol plasma levels and peripheral benzodiazepine receptor density. *Psychoneuroendocrinology*. 2004; 29:1341–1344. [PubMed: 15288714]
- Eser D, Schule C, Romeo E, Baghai TC, di Michele F, Pasini A, Zwanzger P, Padberg F, Rupprecht R. Neuropsychopharmacological properties of neuroactive steroids in depression and anxiety disorders. *Psychopharmacology (Berl)*. 2006; 186:373–387. [PubMed: 16247651]
- Fadalti M, Petraglia F, Luisi S, Bernardi F, Casarosa E, Ferrari E, Luisi M, Saggese G, Genazzani AR, Bernasconi S. Changes of serum allopregnanolone levels in the first 2 years of life and during pubertal development. *Pediatr Res*. 1999; 46:323–327. [PubMed: 10473049]
- File SE, Simmonds MA. Myoclonic seizures in the mouse induced by alphaxalone and related steroid anaesthetics. *J Pharm Pharmacol*. 1988; 40:57–59. [PubMed: 2896777]

- Finn DA, Ford MM, Wiren KM, Roselli CE, Crabbe JC. The role of pregnane neurosteroids in ethanol withdrawal: behavioral genetic approaches. *Pharmacol Ther.* 2004a; 101:91–112. [PubMed: 14761701]
- Finn DA, Mark GP, Fretwell AM, Gililand-Kaufman KR, Strong MN, Ford MM. Reinstatement of ethanol and sucrose seeking by the neurosteroid allopregnanolone in C57BL/6 mice. *Psychopharmacology (Berl).* 2008; 201:423–433. [PubMed: 18758755]
- Finn DA, Phillips TJ, Okorn DM, Chester JA, Cunningham CL. Rewarding effect of the neuroactive steroid 3 α -hydroxy-5 α -pregnan-20-one in mice. *Pharmacol Biochem Behav.* 1997a; 56:261–264. [PubMed: 9050083]
- Finn DA, Roberts AJ, Lotrich F, Gallaher EJ. Genetic differences in behavioral sensitivity to a neuroactive steroid. *J Pharmacol Exp Ther.* 1997b; 280:820–828. [PubMed: 9023296]
- Finn DA, Sinnott RS, Ford MM, Long SL, Tanchuck MA, Phillips TJ. Sex differences in the effect of ethanol injection and consumption on brain allopregnanolone levels in C57BL/6 mice. *Neuroscience.* 2004b; 123:813–819. [PubMed: 14751275]
- Fish EW, Faccidomo S, DeBold JF, Miczek KA. Alcohol, allopregnanolone and aggression in mice. *Psychopharmacology (Berl).* 2001; 153:473–483. [PubMed: 11243495]
- Follesa P, Biggio F, Talani G, Murru L, Serra M, Sanna E, Biggio G. Neurosteroids, GABA_A receptors, and ethanol dependence. *Psychopharmacology (Berl).* 2006; 186:267–280. [PubMed: 16163526]
- Follesa P, Porcu P, Sogliano C, Cinus M, Biggio F, Mancuso L, Mostallino MC, Paoletti AM, Purdy RH, Biggio G, Concas A. Changes in GABA_A receptor 2 subunit gene expression induced by long-term administration of oral contraceptives in rats. *Neuropharmacology.* 2002; 42:325–336. [PubMed: 11897111]
- Ford MM, Mark GP, Nickel JD, Phillips TJ, Finn DA. Allopregnanolone influences the consummatory processes that govern ethanol drinking in C57BL/6J mice. *Behav Brain Res.* 2007; 179:265–272. [PubMed: 17376546]
- Ford MM, Nickel JD, Phillips TJ, Finn DA. Neurosteroid modulators of GABA_A receptors differentially modulate ethanol intake patterns in male C57BL/6J mice. *Alcohol Clin Exp Res.* 2005; 29:1630–1640. [PubMed: 16205363]
- Frye CA, Bayon LE, Pursnani NK, Purdy RH. The neurosteroids, progesterone and 3 α ,5 α -THP, enhance sexual motivation, receptivity, and proceptivity in female rats. *Brain Res.* 1998; 808:72–83. [PubMed: 9795145]
- Gabriel KI, Cunningham CL, Finn DA. Allopregnanolone does not influence ethanol-induced conditioned place preference in DBA/2J mice. *Psychopharmacology (Berl).* 2004; 176:50–56. [PubMed: 15083256]
- Gallaher EJ, Jones GE, Belknap JK, Crabbe JC. Identification of genetic markers for initial sensitivity and rapid tolerance to ethanol-induced ataxia using quantitative trait locus analysis in BXD recombinant inbred mice. *J Pharmacol Exp Ther.* 1996; 277:604–612. [PubMed: 8627537]
- Gasior M, Carter RB, Goldberg SR, Witkin JM. Anticonvulsant and behavioral effects of neuroactive steroids alone and in conjunction with diazepam. *J Pharmacol Exp Ther.* 1997; 282:543–553. [PubMed: 9262314]
- Genazzani AR, Petraglia F, Bernardi F, Casarosa E, Salvestroni C, Tonetti A, Nappi RE, Luisi S, Palumbo M, Purdy RH, Luisi M. Circulating levels of allopregnanolone in humans: gender, age, and endocrine influences. *J Clin Endocrinol Metab.* 1998; 83:2099–2103. [PubMed: 9626145]
- George O, Vallee M, Vitiello S, Le Moal M, Piazza PV, Mayo W. Low brain allopregnanolone levels mediate flattened circadian activity associated with memory impairments in aged rats. *Biol Psychiatry.* 2010; 68:956–963. [PubMed: 20471631]
- Gilbert Evans SE, Ross LE, Sellers EM, Purdy RH, Romach MK. 3 α -reduced neuroactive steroids and their precursors during pregnancy and the postpartum period. *Gynecol Endocrinol.* 2005; 21:268–279. [PubMed: 16373246]
- Gililand KR, Finn DA. The impact of gonadectomy and adrenalectomy on acute withdrawal severity in male and female C57BL/6J and DBA/2J mice following a single high dose of ethanol. *Alcohol Clin Exp Res.* 2007; 31:1846–1857. [PubMed: 17850218]

- Girdler SS, Beth Mechlin M, Light KC, Morrow AL. Ethnic differences in allopregnanolone concentrations in women during rest and following mental stress. *Psychophysiology*. 2006; 43:331–336. [PubMed: 16916428]
- Girdler SS, Klatzkin R. Neurosteroids in the context of stress: implications for depressive disorders. *Pharmacol Ther*. 2007; 116:125–139. [PubMed: 17597217]
- Girdler SS, Straneva PA, Light KC, Pedersen CA, Morrow AL. Allopregnanolone levels and reactivity to mental stress in premenstrual dysphoric disorder. *Biol Psychiatry*. 2001; 49:788–797. [PubMed: 11331087]
- Gora-Maslak G, McClearn GE, Crabbe JC, Phillips TJ, Belknap JK, Plomin R. Use of recombinant inbred strains to identify quantitative trait loci in psychopharmacology. *Psychopharmacology (Berl)*. 1991; 104:413–424. [PubMed: 1780413]
- Grahame NJ, Cunningham CL. Intravenous ethanol self-administration in C57BL/6J and DBA/2J mice. *Alcohol Clin Exp Res*. 1997; 21:56–62. [PubMed: 9046373]
- Grobin AC, Morrow AL. 3 α -hydroxy-5 α -pregnan-20-one levels and GABA_A receptor-mediated ³⁶Cl⁻ flux across development in rat cerebral cortex. *Brain Res Dev Brain Res*. 2001; 131:31–39.
- Gunn BG, Cunningham L, Cooper MA, Corteen NL, Seifi M, Swinny JD, Lambert JJ, Belelli D. Dysfunctional astrocytic and synaptic regulation of hypothalamic glutamatergic transmission in a mouse model of early-life adversity: relevance to neurosteroids and programming of the stress response. *J Neurosci*. 2013; 33:19534–19554. [PubMed: 24336719]
- Guo AL, Petraglia F, Criscuolo M, Ficarra G, Nappi RE, Palumbo MA, Trentini GP, Purdy RH, Genazzani AR. Evidence for a role of neurosteroids in modulation of diurnal changes and acute stress-induced corticosterone secretion in rats. *Gynecol Endocrinol*. 1995; 9:1–7. [PubMed: 7793294]
- Harrison NL, Majewska MD, Harrington JW, Barker JL. Structure-activity relationships for steroid interaction with the gamma-aminobutyric acidA receptor complex. *J Pharmacol Exp Ther*. 1987; 241:346–353. [PubMed: 3033209]
- Herd MB, Belelli D, Lambert JJ. Neurosteroid modulation of synaptic and extrasynaptic GABA_A receptors. *Pharmacol Ther*. 2007; 116:20–34. [PubMed: 17531325]
- Hill M, Popov P, Havlikova H, Kancheva L, Vrbikova J, Kancheva R, Pouzar V, Cerny I, Starka L. Altered profiles of serum neuroactive steroids in premenopausal women treated for alcohol addiction. *Steroids*. 2005; 70:515–524. [PubMed: 15894035]
- Hirani K, Khisti RT, Chopde CT. Behavioral action of ethanol in Porsolt's forced swim test: modulation by 3 α -hydroxy-5 α -pregnan-20-one. *Neuropharmacology*. 2002; 43:1339–1350. [PubMed: 12527484]
- Hirani K, Sharma AN, Jain NS, Ugale RR, Chopde CT. Evaluation of GABAergic neuroactive steroid 3 α -hydroxy-5 α -pregnane-20-one as a neurobiological substrate for the anti-anxiety effect of ethanol in rats. *Psychopharmacology (Berl)*. 2005; 180:267–278. [PubMed: 15719223]
- Hogskilde S, Wagner J, Carl P, Sorensen MB. Anaesthetic properties of pregnanolone emulsion. A comparison with alphaxolone/alphadolone, propofol, thiopentone and midazolam in a rat model. *Anaesthesia*. 1987; 42:1045–1050. [PubMed: 3500653]
- Holdstock L, Penland SN, Morrow AL, de Wit H. Moderate doses of ethanol fail to increase plasma levels of neurosteroid 3 α -hydroxy-5 α -pregnan-20-one-like immunoreactivity in healthy men and women. *Psychopharmacology (Berl)*. 2006; 186:442–450. [PubMed: 16240164]
- Hosie AM, Wilkins ME, da Silva HM, Smart TG. Endogenous neurosteroids regulate GABA_A receptors through two discrete transmembrane sites. *Nature*. 2006; 444:486–489. [PubMed: 17108970]
- Irwig MS. Decreased alcohol consumption among former male users of finasteride with persistent sexual side effects: a preliminary report. *Alcohol Clin Exp Res*. 2013; 37:1823–1826. [PubMed: 23763349]
- Janak PH, Redfern JE, Samson HH. The reinforcing effects of ethanol are altered by the endogenous neurosteroid, allopregnanolone. *Alcohol Clin Exp Res*. 1998; 22:1106–1112. [PubMed: 9726282]
- Kavaliers M, Wiebe JP. Analgesic effects of the progesterone metabolite: 3 α -hydroxy-5 α -pregnan-20-one, and possible modes of action in mice. *Brain Res*. 1987; 415:393–398. [PubMed: 3111642]

- Khisti RT, Boyd KN, Kumar S, Morrow AL. Systemic ethanol administration elevates deoxycorticosterone levels and chronic ethanol exposure attenuates this response. *Brain Res.* 2005; 1049:104–111. [PubMed: 15936001]
- Khisti RT, Chopde CT, Jain SP. Antidepressant-like effect of the neurosteroid 3 α -hydroxy-5 α -pregnan-20-one in mice forced swim test. *Pharmacol Biochem Behav.* 2000; 67:137–143. [PubMed: 11113493]
- Khisti RT, VanDoren MJ, Matthews DB, Morrow AL. Ethanol-induced elevation of 3 α -hydroxy-5 α -pregnan-20-one does not modulate motor incoordination in rats. *Alcohol Clin Exp Res.* 2004; 28:1249–1256. [PubMed: 15318125]
- Khisti RT, VanDoren MJ, O'Buckley T, Morrow AL. Neuroactive steroid 3 α -hydroxy-5 α -pregnan-20-one modulates ethanol-induced loss of righting reflex in rats. *Brain Res.* 2003; 980:255–265. [PubMed: 12867266]
- Klatzkin RR, Morrow AL, Light KC, Pedersen CA, Girdler SS. Histories of depression, allopregnanolone responses to stress, and premenstrual symptoms in women. *Biol Psychol.* 2006; 71:2–11. [PubMed: 15951099]
- Kokate TG, Banks MK, Magee T, Yamaguchi S, Rogawski MA. Finasteride, a 5 α -reductase inhibitor, blocks the anticonvulsant activity of progesterone in mice. *J Pharmacol Exp Ther.* 1999; 288:679–684. [PubMed: 9918575]
- Kokate TG, Svensson BE, Rogawski MA. Anticonvulsant activity of neurosteroids: correlation with -aminobutyric acid-evoked chloride current potentiation. *J Pharmacol Exp Ther.* 1994; 270:1223–1229. [PubMed: 7932175]
- Kokate TG, Yamaguchi S, Pannell LK, Rajamani U, Carroll DM, Grossman AB, Rogawski MA. Lack of anticonvulsant tolerance to the neuroactive steroid pregnanolone in mice. *J Pharmacol Exp Ther.* 1998; 287:553–558. [PubMed: 9808680]
- Kostakis E, Smith C, Jang MK, Martin SC, Richards KG, Russek SJ, Gibbs TT, Farb DH. The neuroactive steroid pregnenolone sulfate stimulates trafficking of functional N-methyl D-aspartate receptors to the cell surface via a noncanonical, G protein, and Ca²⁺-dependent mechanism. *Mol Pharmacol.* 2013; 84:261–274. [PubMed: 23716622]
- Kumar S, Porcu P, Werner DF, Matthews DB, Diaz-Granados JL, Helfand RS, Morrow AL. The role of GABA_A receptors in the acute and chronic effects of ethanol: a decade of progress. *Psychopharmacology (Berl).* 2009; 205:529–564. [PubMed: 19455309]
- Linsenbardt DN, Moore EM, Gross CD, Goldfarb KJ, Blackman LC, Boehm SL 2nd. Sensitivity and tolerance to the hypnotic and ataxic effects of ethanol in adolescent and adult C57BL/6J and DBA/2J mice. *Alcohol Clin Exp Res.* 2009; 33:464–476. [PubMed: 19120054]
- Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci.* 2009; 10:434–445. [PubMed: 19401723]
- Maguire JL, Stell BM, Rafizadeh M, Mody I. Ovarian cycle-linked changes in GABA_A receptors mediating tonic inhibition alter seizure susceptibility and anxiety. *Nat Neurosci.* 2005; 8:797–804. [PubMed: 15895085]
- Majewska MD, Harrison NL, Schwartz RD, Barker JL, Paul SM. Steroid hormone metabolites are barbiturate-like modulators of the GABA receptor. *Science.* 1986; 232:1004–1007. [PubMed: 2422758]
- Matthews DB, Morrow AL, Tokunaga S, McDaniel JR. Acute ethanol administration and acute allopregnanolone administration impair spatial memory in the Morris water task. *Alcohol Clin Exp Res.* 2002; 26:1747–1751. [PubMed: 12436065]
- McEwen BS. Non-genomic and genomic effects of steroids on neural activity. *Trends Pharmacol Sci.* 1991; 12:141–147. [PubMed: 2063480]
- Meyer DA, Carta M, Partridge LD, Covey DF, Valenzuela CF. Neurosteroids enhance spontaneous glutamate release in hippocampal neurons. Possible role of metabotropic sigma 1-like receptors. *J Biol Chem.* 2002; 277:28725–28732. [PubMed: 12042305]
- Miczek KA, Fish EW, De Bold JF. Neurosteroids, GABA_A receptors, and escalated aggressive behavior. *Horm Behav.* 2003; 44:242–257. [PubMed: 14609546]

- Milivojevic V, Kranzler HR, Gelernter J, Burian L, Covault J. Variation in genes encoding the neuroactive steroid synthetic enzymes 5 α -reductase type 1 and 3 α -reductase type 2 is associated with alcohol dependence. *Alcohol Clin Exp Res*. 2011; 35:946–952. [PubMed: 21323680]
- Mitev YA, Darwish M, Wolf SS, Holsboer F, Almeida OF, Patchev VK. Gender differences in the regulation of 3 α -hydroxysteroid dehydrogenase in rat brain and sensitivity to neurosteroid-mediated stress protection. *Neuroscience*. 2003; 120:541–549. [PubMed: 12890523]
- Monteleone P, Luisi S, Tonetti A, Bernardi F, Genazzani AD, Luisi M, Petraglia F, Genazzani AR. Allopregnanolone concentrations and premenstrual syndrome. *Eur J Endocrinol*. 2000; 142:269–273. [PubMed: 10700721]
- Morrow AL. Recent developments in the significance and therapeutic relevance of neuroactive steroids--Introduction to the special issue. *Pharmacol Ther*. 2007; 116:1–6. [PubMed: 17531324]
- Morrow AL, Janis GC, VanDoren MJ, Matthews DB, Samson HH, Janak PH, Grant KA. Neurosteroids mediate pharmacological effects of ethanol: a new mechanism of ethanol action? *Alcohol Clin Exp Res*. 1999; 23:1933–1940. [PubMed: 10630613]
- Morrow AL, Pace JR, Purdy RH, Paul SM. Characterization of steroid interactions with γ -aminobutyric acid receptor-gated chloride ion channels: evidence for multiple steroid recognition sites. *Mol Pharmacol*. 1990; 37:263–270. [PubMed: 1689453]
- Morrow, AL.; Porcu, P. Neuroactive steroid biomarkers of alcohol sensitivity and alcoholism risk. In: Ritsner, MS., editor. *The Handbook of Neuropsychiatric Biomarkers, Endophenotypes, and Genes*. Dordrecht: Springer Science + Business Media B.V.; 2009. p. 47-57.
- Morrow AL, Porcu P, Boyd KN, Grant KA. Hypothalamic-pituitary-adrenal axis modulation of GABAergic neuroactive steroids influences ethanol sensitivity and drinking behavior. *Dialogues Clin Neurosci*. 2006; 8:463–477. [PubMed: 17290803]
- Morrow AL, Suzdak PD, Paul SM. Steroid hormone metabolites potentiate GABA receptor-mediated chloride ion flux with nanomolar potency. *Eur J Pharmacol*. 1987; 142:483–485. [PubMed: 2828079]
- Morrow AL, VanDoren MJ, Devaud LL. Effects of progesterone or neuroactive steroid? *Nature*. 1998; 395:652–653. [PubMed: 9790185]
- Morrow AL, VanDoren MJ, Penland SN, Matthews DB. The role of GABAergic neuroactive steroids in ethanol action, tolerance and dependence. *Brain Res Rev*. 2001; 37:98–109. [PubMed: 11744078]
- Mitchellshvili Z, Kapur J. A presynaptic action of the neurosteroid pregnenolone sulfate on GABAergic synaptic transmission. *Mol Pharmacol*. 2003; 64:857–864. [PubMed: 14500742]
- Naert G, Maurice T, Tapia-Arancibia L, Givalois L. Neuroactive steroids modulate HPA axis activity and cerebral brain-derived neurotrophic factor (BDNF) protein levels in adult male rats. *Psychoneuroendocrinology*. 2007; 32:1062–1078. [PubMed: 17928160]
- Nyberg S, Andersson A, Zingmark E, Wahlstrom G, Backstrom T, Sundstrom-Poromaa I. The effect of a low dose of alcohol on allopregnanolone serum concentrations across the menstrual cycle in women with severe premenstrual syndrome and controls. *Psychoneuroendocrinology*. 2005; 30:892–901. [PubMed: 15979810]
- O'Dell LE, Alomary AA, Vallee M, Koob GF, Fitzgerald RL, Purdy RH. Ethanol-induced increases in neuroactive steroids in the rat brain and plasma are absent in adrenalectomized and gonadectomized rats. *Eur J Pharmacol*. 2004; 484:241–247. [PubMed: 14744609]
- Osterman, JL.; Kralic, JE.; O'Buckley, TK.; Homanics, GE.; Morrow, AL. GABA_A receptor 1 subunit knockout mice: A novel model of essential tremor. In: LeDoux, M., editor. *Animal Models of Movement Disorders*. Oxford: Elsevier Academic Press; 2005. p. 369-375.
- Owens MJ, Ritchie JC, Nemeroff CB. 5 α -pregnane-3 α , 21-diol-20-one (THDOC) attenuates mild stress-induced increases in plasma corticosterone via a non-glucocorticoid mechanism: comparison with alprazolam. *Brain Res*. 1992; 573:353–355. [PubMed: 1504771]
- Paris JJ, Frye CA. Juvenile offspring of rats exposed to restraint stress in late gestation have impaired cognitive performance and dysregulated progesterone formation. *Stress*. 2011; 14:23–32. [PubMed: 21034292]

- Park-Chung M, Wu FS, Purdy RH, Malayev AA, Gibbs TT, Farb DH. Distinct sites for inverse modulation of N-methyl-D-aspartate receptors by sulfated steroids. *Mol Pharmacol*. 1997; 52:1113–1123. [PubMed: 9396781]
- Patchev VK, Hassan AH, Holsboer DF, Almeida OF. The neurosteroid tetrahydroprogesterone attenuates the endocrine response to stress and exerts glucocorticoid-like effects on vasopressin gene transcription in the rat hypothalamus. *Neuropsychopharmacology*. 1996; 15:533–540. [PubMed: 8946427]
- Patchev VK, Montkowski A, Rouskova D, Koranyi L, Holsboer F, Almeida OF. Neonatal treatment of rats with the neuroactive steroid tetrahydrodeoxycorticosterone (THDOC) abolishes the behavioral and neuroendocrine consequences of adverse early life events. *J Clin Invest*. 1997; 99:962–966. [PubMed: 9062354]
- Patchev VK, Shoaib M, Holsboer F, Almeida OF. The neurosteroid tetrahydroprogesterone counteracts corticotropin-releasing hormone-induced anxiety and alters the release and gene expression of corticotropin-releasing hormone in the rat hypothalamus. *Neuroscience*. 1994; 62:265–271. [PubMed: 7816204]
- Penland SN, Morrow AL. 3 α ,5 β -Reduced cortisol exhibits antagonist properties on cerebral cortical GABA_A receptors. *Eur J Pharmacol*. 2004; 506:129–132. [PubMed: 15588732]
- Pericic D, Strac DS, Vlainic J. Interaction of diazepam and swim stress. *Brain Res*. 2007; 1184:81–87. [PubMed: 17945202]
- Pericic D, Svob D, Jazvinscak M, Mirkovic K. Anticonvulsive effect of swim stress in mice. *Pharmacol Biochem Behav*. 2000; 66:879–886. [PubMed: 10973529]
- Phillips TJ, Crabbe JC, Metten P, Belknap JK. Localization of genes affecting alcohol drinking in mice. *Alcohol Clin Exp Res*. 1994; 18:931–941. [PubMed: 7978106]
- Pibiri F, Nelson M, Guidotti A, Costa E, Pinna G. Decreased corticolimbic allopregnanolone expression during social isolation enhances contextual fear: A model relevant for posttraumatic stress disorder. *Proc Natl Acad Sci USA*. 2008; 105:5567–5572. [PubMed: 18391192]
- Pierucci-Lagha A, Covault J, Feinn R, Khisti RT, Morrow AL, Marx CE, Shampine LJ, Kranzler HR. Subjective effects and changes in steroid hormone concentrations in humans following acute consumption of alcohol. *Psychopharmacology (Berl)*. 2006; 186:451–461. [PubMed: 16341848]
- Pierucci-Lagha A, Covault J, Feinn R, Nellissery M, Hernandez-Avila C, Oncken C, Morrow AL, Kranzler HR. GABRA2 alleles moderate the subjective effects of alcohol, which are attenuated by finasteride. *Neuropsychopharmacology*. 2005; 30:1193–1203. [PubMed: 15702134]
- Pisu MG, Garau A, Olla P, Biggio F, Utzeri C, Dore R, Serra M. Altered stress responsiveness and hypothalamic-pituitary-adrenal axis function in male rat offspring of socially isolated parents. *J Neurochem*. 2013; 126:493–502. [PubMed: 23600845]
- Porcu P, Grant KA, Green HL, Rogers LS, Morrow AL. Hypothalamic-pituitary-adrenal axis and ethanol modulation of deoxycorticosterone levels in cynomolgus monkeys. *Psychopharmacology (Berl)*. 2006; 186:293–301. [PubMed: 16133132]
- Porcu P, Mostallino MC, Sogliano C, Santoru F, Berretti R, Concas A. Long-term administration with levonorgestrel decreases allopregnanolone levels and alters GABA_A receptor subunit expression and anxiety-like behavior. *Pharmacol Biochem Behav*. 2012; 102:366–372. [PubMed: 22634062]
- Porcu P, O'Buckley TK, Alward SE, Song SC, Grant KA, de Wit H, Morrow AL. Differential effects of ethanol on serum GABAergic 3 α ,5 α /3 α ,5 β neuroactive steroids in mice, rats, cynomolgus monkeys, and humans. *Alcohol Clin Exp Res*. 2010; 34:432–442. [PubMed: 20028362]
- Porcu P, O'Buckley TK, Morrow AL, Adinoff B. Differential hypothalamic-pituitary-adrenal activation of the neuroactive steroids pregnenolone sulfate and deoxycorticosterone in healthy controls and alcohol-dependent subjects. *Psychoneuroendocrinology*. 2008; 33:214–226. [PubMed: 18096321]
- Porcu P, O'Buckley TK, Song SC, Harenza JL, Lu L, Wang X, Williams RW, Miles MF, Morrow AL. Genetic analysis of the neurosteroid deoxycorticosterone and its relation to alcohol phenotypes: identification of QTLs and downstream gene regulation. *PLoS One*. 2011; 6:e18405. [PubMed: 21494628]
- Porcu P, Sogliano C, Ibba C, Piredda M, Tocco S, Marra C, Purdy RH, Biggio G, Concas A. Failure of γ -hydroxybutyric acid both to increase neuroactive steroid concentrations in adrenalectomized-

- orchiectomized rats and to induce tolerance to its steroidogenic effect in intact animals. *Brain Res.* 2004; 1012:160–168. [PubMed: 15158173]
- Purdy RH, Morrow AL, Moore PH Jr, Paul SM. Stress-induced elevations of γ -aminobutyric acid type A receptor-active steroids in the rat brain. *Proc Natl Acad Sci USA.* 1991; 88:4553–4557. [PubMed: 1852011]
- Rapkin AJ, Morgan M, Goldman L, Brann DW, Simone D, Mahesh VB. Progesterone metabolite allopregnanolone in women with premenstrual syndrome. *Obstet Gynecol.* 1997; 90:709–714. [PubMed: 9351749]
- Rapkin AJ, Morgan M, Sogliano C, Biggio G, Concas A. Decreased neuroactive steroids induced by combined oral contraceptive pills are not associated with mood changes. *Fertil Steril.* 2006; 85:1371–1378. [PubMed: 16580672]
- Rasmusson AM, Pinna G, Paliwal P, Weisman D, Gottschalk C, Charney D, Krystal J, Guidotti A. Decreased cerebrospinal fluid allopregnanolone levels in women with posttraumatic stress disorder. *Biol Psychiatry.* 2006; 60:704–713. [PubMed: 16934764]
- Ray LA, Hutchison KE, Ashenhurst JR, Morrow AL. Naltrexone selectively elevates GABAergic neuroactive steroid levels in heavy drinkers with the Asp40 allele of the OPRM1 gene: a pilot investigation. *Alcohol Clin Exp Res.* 2010; 34:1479–1487. [PubMed: 20528823]
- Reddy DS, Rogawski MA. Stress-induced deoxycorticosterone-derived neurosteroids modulate GABA_A receptor function and seizure susceptibility. *J Neurosci.* 2002; 22:3795–3805. [PubMed: 11978855]
- Roberts AJ, Crabbe JC, Keith LD. Genetic differences in hypothalamic-pituitary-adrenal axis responsiveness to acute ethanol and acute ethanol withdrawal. *Brain Res.* 1992; 579:296–302. [PubMed: 1320985]
- Roberts AJ, Phillips TJ, Belknap JK, Finn DA, Keith LD. Genetic analysis of the corticosterone response to ethanol in BXD recombinant inbred mice. *Behav Neurosci.* 1995; 109:1199–1208. [PubMed: 8748968]
- Rodriguez LA, Plomin R, Blizard DA, Jones BC, McClearn GE. Alcohol acceptance, preference, and sensitivity in mice. I. Quantitative genetic analysis using BXD recombinant inbred strains. *Alcohol Clin Exp Res.* 1994; 18:1416–1422. [PubMed: 7695038]
- Romeo E, Brancati A, De Lorenzo A, Fucci P, Furnari C, Pompili E, Sasso GF, Spalletta G, Troisi A, Pasini A. Marked decrease of plasma neuroactive steroids during alcohol withdrawal. *Clin Neuropharmacol.* 1996; 19:366–369. [PubMed: 8829001]
- Saalmann YB, Kirkcaldie MT, Waldron S, Calford MB. Cellular distribution of the GABA_A receptor-modulating 3 α -hydroxy, 5 α -reduced pregnane steroids in the adult rat brain. *J Neuroendocrinol.* 2007; 19:272–284. [PubMed: 17355317]
- Saalmann YB, Morgan IG, Calford MB. Neurosteroids involved in regulating inhibition in the inferior colliculus. *J Neurophysiol.* 2006; 96:3064–3073. [PubMed: 16971675]
- Sanna E, Talani G, Busonero F, Pisu MG, Purdy RH, Serra M, Biggio G. Brain steroidogenesis mediates ethanol modulation of GABA_A receptor activity in rat hippocampus. *J Neurosci.* 2004; 24:6521–6530. [PubMed: 15269263]
- Sarkar J, Wakefield S, MacKenzie G, Moss SJ, Maguire J. Neurosteroidogenesis is required for the physiological response to stress: role of neurosteroid-sensitive GABA_A receptors. *J Neurosci.* 2011; 31:18198–18210. [PubMed: 22171026]
- Schmidt PJ, Purdy RH, Moore PH Jr, Paul SM, Rubinow DR. Circulating levels of anxiolytic steroids in the luteal phase in women with premenstrual syndrome and in control subjects. *J Clin Endocrinol Metab.* 1994; 79:1256–1260. [PubMed: 7962316]
- Schuckit MA. Low level of response to alcohol as a predictor of future alcoholism. *Am J Psychiatry.* 1994; 151:184–189. [PubMed: 8296886]
- Schuckit MA. An overview of genetic influences in alcoholism. *J Subst Abuse Treat.* 2009; 36:S5–S14. [PubMed: 19062348]
- Schumacher M, Guennoun R, Stein DG, De Nicola AF. Progesterone: therapeutic opportunities for neuroprotection and myelin repair. *Pharmacol Ther.* 2007; 116:77–106. [PubMed: 17659348]

- Serra M, Pisu MG, Littera M, Papi G, Sanna E, Tuveri F, Usala L, Purdy RH, Biggio G. Social isolation-induced decreases in both the abundance of neuroactive steroids and GABA_A receptor function in rat brain. *J Neurochem*. 2000; 75:732–740. [PubMed: 10899949]
- Shannon EE, Porcu P, Purdy RH, Grant KA. Characterization of the discriminative stimulus effects of the neuroactive steroid pregnanolone in DBA/2J and C57BL/6J inbred mice. *J Pharmacol Exp Ther*. 2005a; 314:675–685. [PubMed: 15857945]
- Shannon EE, Purdy RH, Grant KA. Discriminative stimulus effects of 5.6 mg/kg pregnanolone in DBA/2J and C57BL/6J inbred mice. *Alcohol*. 2005b; 37:35–45. [PubMed: 16472717]
- Shen H, Gong QH, Aoki C, Yuan M, Ruderman Y, Dattilo M, Williams K, Smith SS. Reversal of neurosteroid effects at 2 GABA_A receptors triggers anxiety at puberty. *Nat Neurosci*. 2007; 10:469–477. [PubMed: 17351635]
- Sinnott RS, Mark GP, Finn DA. Reinforcing effects of the neurosteroid allopregnanolone in rats. *Pharmacol Biochem Behav*. 2002; 72:923–929. [PubMed: 12062582]
- Tokuda K, Izumi Y, Zorumski CF. Ethanol enhances neurosteroidogenesis in hippocampal pyramidal neurons by paradoxical NMDA receptor activation. *J Neurosci*. 2011; 31:9905–9909. [PubMed: 21734282]
- Tokuda K, O'Dell KA, Izumi Y, Zorumski CF. Midazolam inhibits hippocampal long-term potentiation and learning through dual central and peripheral benzodiazepine receptor activation and neurosteroidogenesis. *J Neurosci*. 2010; 30:16788–16795. [PubMed: 21159950]
- Torres JM, Ortega E. Alcohol intoxication increases allopregnanolone levels in female adolescent humans. *Neuropsychopharmacology*. 2003; 28:1207–1209. [PubMed: 12700685]
- Torres JM, Ortega E. Alcohol intoxication increases allopregnanolone levels in male adolescent humans. *Psychopharmacology (Berl)*. 2004; 172:352–355. [PubMed: 14647956]
- Uzunova V, Sheline Y, Davis JM, Rasmusson A, Uzunov DP, Costa E, Guidotti A. Increase in the cerebrospinal fluid content of neurosteroids in patients with unipolar major depression who are receiving fluoxetine or fluvoxamine. *Proc Natl Acad Sci USA*. 1998; 95:3239–3244. [PubMed: 9501247]
- Vallee M, Rivera JD, Koob GF, Purdy RH, Fitzgerald RL. Quantification of neurosteroids in rat plasma and brain following swim stress and allopregnanolone administration using negative chemical ionization gas chromatography/mass spectrometry. *Anal Biochem*. 2000; 287:153–166. [PubMed: 11078595]
- VanDoren MJ, Matthews DB, Janis GC, Grobin AC, Devaud LL, Morrow AL. Neuroactive steroid 3 α -hydroxy-5 α -pregnan-20-one modulates electrophysiological and behavioral actions of ethanol. *J Neurosci*. 2000; 20:1982–1989. [PubMed: 10684899]
- Wang J, Williams RW, Manly KF. WebQTL: web-based complex trait analysis. *Neuroinformatics*. 2003; 1:299–308. [PubMed: 15043217]
- Wang M, Seippel L, Purdy RH, Backstrom T. Relationship between symptom severity and steroid variation in women with premenstrual syndrome: study on serum pregnenolone, pregnenolone sulfate: 5 α -pregnane-3,20-dione and 3 α -hydroxy-5 α -pregnan-20-one. *J Clin Endocrinol Metab*. 1996; 81:1076–1082. [PubMed: 8772579]
- Wilhelmsen KC, Schuckit M, Smith TL, Lee JV, Segall SK, Feiler HS, Kalmijn J. The search for genes related to a low-level response to alcohol determined by alcohol challenges. *Alcohol Clin Exp Res*. 2003; 27:1041–1047. [PubMed: 12878909]
- Williams RW, Gu J, Qi S, Lu L. The genetic structure of recombinant inbred mice: high-resolution consensus maps for complex trait analysis. *Genome Biol*. 2001; 2 RESEARCH0046.
- Wu FS, Gibbs TT, Farb DH. Pregnenolone sulfate: a positive allosteric modulator at the Nmethyl-D-aspartate receptor. *Mol Pharmacol*. 1991; 40:333–336. [PubMed: 1654510]
- Wu X, Gangisetty O, Carver CM, Reddy DS. Estrous cycle regulation of extrasynaptic - containing GABA_A receptor-mediated tonic inhibition and limbic epileptogenesis. *J Pharmacol Exp Ther*. 2013; 346:146–160. [PubMed: 23667248]

Figure 1.

Schematic representation of the putative mechanisms and sites of action of GABAergic neuroactive steroids. Neuroactive steroids, such as 3 α ,5 α -THP and 3 α ,5 α -THDOC, exert inhibitory actions mediated by synaptic and extrasynaptic GABA_A receptors. No active release mechanism has been identified for endogenous neuroactive steroids. They may access the neuroactive steroid transmembrane binding sites via paracrine (a) or autocrine (b) mechanisms or by intracellular lateral diffusion through the cell membrane (c). There is also evidence for extracellular pockets of neuroactive steroid storage that allow for rapid and

repeated access to extracellular binding sites on neurons. These effects are described and referenced in the text.

Figure 2.

Variation in basal and ethanol-induced 3α,5α-THP across selected BXD strains. Group-housed male mice (8 to 12 weeks old) were injected with saline (basal levels) or ethanol (2 g/kg, i.p.) at 9:00 am and were euthanized 1 hour later. 3α,5α-THP levels were measured in the cerebral cortex by radioimmunoassay. (a) Data are expressed as ng/g of tissue and are means ± SEM of values from 3-16 mice per strain. (b) Data are expressed as percent change of the average 3α,5α-THP levels in the ethanol-treated group vs. the average 3α,5α-THP levels in the respective saline-treated group (n = 3-16/group/strain).

Figure 3.

Genetic correlations between basal or ethanol-induced 3α,5α-THP levels and behavioral phenotypes for ethanol consumption previously characterized in the BXD strains. Group-housed male mice (8 to 12 weeks old) were injected with saline (basal levels) or ethanol (2 g/kg, i.p.) at 9:00 am and were euthanized 1 hour later. 3α,5α-THP levels were measured in the cerebral cortex by radioimmunoassay. Data are expressed as ng/g of tissue (basal levels) or as percent change of the average 3α,5α-THP levels in the ethanol-treated group vs. the average 3α,5α-THP levels in the respective saline-treated group (n = 3-16/group/strain).

Behavioral data for ethanol consumption has been collected by independent labs and has been obtained from GeneNetwork (www.genenetwork.org). (a) Consumption of 10% ethanol at 2 hours, Spearman's $r=-0.82$, $p=0.02$, $n=7$, GeneNetwork ID 12733 (Cook et al., unpublished). (b) Consumption of 3% ethanol, Spearman's $r=-0.82$, $p=0.02$, $n=7$, GeneNetwork ID 10474 (Phillips et al. 1994). (c) Consumption of 10% ethanol, Spearman's $r=-0.82$, $p=0.02$, $n=7$, GeneNetwork ID 10582 (Rodriguez et al. 1994).

Table 1

Summary of the neuroactive steroids effects on the stress/HPA axis response in adult rats, mice and human subjects.

Species	Neuroactive steroid levels	Neuroactive steroid administration
Rat		
Acute stress	↑ basal levels (Barbaccia et al. 1994; Barbaccia et al. 1996; Barbaccia et al. 1997; Purdy et al. 1991; Vallee et al. 2000)	↓ levels of CRH, ACTH and corticosterone (Owens et al. 1992; Patchev et al. 1996; Patchev et al. 1994)
Chronic stress	↓ basal levels (Serra et al. 2000) ↑ response to a stress challenge (Serra et al. 2000)	N.A.
Mouse		
Acute stress	↓ basal levels (Maldonado-Devincini et al. this issue)	↑ levels of corticosterone (Sarkar et al. 2011)
Chronic stress	↓ basal levels (Dong et al. 2001; Pibiri et al. 2008)	N.A.
Human		
Acute stress/HPA challenges	↑ basal levels (Droogleever Fortuyn et al. 2004; Genazzani et al. 1998; Girdler et al. 2001; Porcu et al. 2008) -- basal levels (Altemus et al. 2001; Childs and de Wit 2009; Childs et al. 2010; Girdler et al. 2006)	N.A.
Chronic stress	↑ basal levels (Girdler et al. 2001) -- basal levels (Rapkin et al. 1997; Schmidt et al. 1994; Wang et al. 1996) ↓ basal levels (Klatzkin et al. 2006; Monteleone et al. 2000; Rasmusson et al. 2006) ↓ response to a stress challenge (Klatzkin et al. 2006)	↑ sensitivity to 3α, 5α-THP (Backstrom et al. 2013)

These effects are described and referenced in the text.

↑ = increase;

↓ = decrease;

-- = unchanged;

N.A. = not assayed.