Complete Sequencing of IncI1 Sequence Type 2 Plasmid pJIE512b Indicates Mobilization of *bla*_{CMY-2} from an IncA/C Plasmid

Kaitlin A. Tagg, Jonathan R. Iredell, Sally R. Partridge

Centre for Infectious Diseases and Microbiology, Westmead Millennium Institute, and The University of Sydney, Westmead, New South Wales, Australia

Sequencing of pJIE512b, a 92.3-kb IncI1 sequence type 2 (ST2) plasmid carrying *bla*_{CMY-2}, revealed a *bla*_{CMY-2} context that appeared to have been mobilized from an IncA/C plasmid by the insertion sequence IS1294. A comparison with published plasmids suggests that *bla*_{CMY-2} has been mobilized from IncA/C to IncI1 plasmids more than once by IS1294-like elements. Alignment of pJIE512b with the only other available IncI1 ST2 plasmid revealed differences across the backbones, indicating variability within this sequence type.

Plasmid-mediated AmpC β -lactamases, particularly in *Escherichia coli* and in *Salmonella* species, represent a significant public health concern, as they confer resistance to the globally important cephalosporin antibiotics and β -lactamase inhibitors (1). The most prevalent AmpC β -lactamase, CMY-2, encoded by *bla*_{CMY-2}, was first reported in 1990 (1, 2). *bla*_{CMY-2} and its minor variants are now globally disseminated among a number of species, most likely through a combination of clonal expansion of strains bearing *bla*_{CMY-2}-carrying plasmids and horizontal transfer of the plasmids themselves (3–6).

Sequence comparisons indicate that a region including *bla*_{CMY-2} was mobilized from the *Citrobacter freundii* chromosome onto a plasmid (7) by the insertion sequence ISEcp1 (Fig. 1a and b) (1, 8). Variations in the context of *bla*_{CMY-2}-like genes are due to differences in the size of the *C. freundii* region mobilized and/or the result of recombination (3, 8, 9).

Various plasmid types, including IncA/C, IncI1, IncI2, and IncFII, are associated with the carriage of *bla*_{CMY-2} (10–15). However, IncA/C and IncI1 plasmids are often reported as the most common carriers of *bla*_{CMY-2} (10, 13–15), particularly IncI1 sequence type 2 (ST2), ST12, and ST23 (10, 13, 16), as defined by plasmid multilocus sequence typing (pMLST) (17). Despite their importance, no ST2 or ST23 plasmid sequences have been published. Here, we completely sequenced pJIE512b, an IncI1 ST2 plasmid carrying *bla*_{CMY-2}.

Clinical *E. coli* isolate JIE512b was identified as carrying *bla*_{CMY-2} during routine PCR screening for plasmid-borne *ampC* genes (see Table S1 in the supplemental material) at the Centre for Infectious Diseases and Microbiology at Westmead Hospital, New South Wales, Australia. Conjugation of JIE512b with DH5 α Rf, with selection on 16 μ g/ml cefoxitin plus 80 μ g/ml rifampin, performed as previously described (18), gave transconjugants carrying *bla*_{CMY-2} and a single plasmid that was named pJIE512b. pMLST (17) indicated that pJIE512b is ST2. Plasmid DNA was purified using the HiSpeed plasmid midikit (Qiagen, Germany), and 1 ng was used for library preparation with the Nextera XT DNA sample preparation kit (Illumina, Inc., USA). Sequencing by Illumina MiSeq technology was performed at the Australian Genome Research Facility (Melbourne, Australia), and sequence reads were processed using FLASH (http://ccb.jhu.edu/software/FLASH/) and were assembled into contigs with Velvet (http://www.ebi.ac.uk/~zerbino/velvet/), SPAdes (http://bioinf.spbau.ru/spades), and CLC Genomics Workbench (http://www.clcbio.com/products/clc-genomics-workbench/). Seven contigs (~900-fold

average coverage) were assembled using SeqMan (DNASTAR, Madison, WI, USA) with PCR amplification (see Table S1 in the supplemental material) and Sanger sequencing to confirm contig boundaries. Analysis and annotation of the resulting sequence were performed using the RAST server (19), BLASTn searches (http://blast.ncbi.nlm.nih.gov/Blast.cgi), ISfinder (https://www-is.biotoul.fr/), and the Gene Construction Kit program (Textco BioSoftware, Inc., USA).

Four contigs corresponded to components of the IncI1 shufflon. This multiple inversion system is responsible for generating variation in the C terminus of the PilV tip adhesin of the thin pilus, leading to differences in recipient cell specificity in liquid mating (20). The shufflon contains up to four segments separated by seven inverted repeats, and a shufflon-specific recombinase (encoded by *rci*) catalyzes recombination between these repeats (20). Sequencing of the amplicons obtained with primers in the *rci* and *pilV* genes flanking the shufflon (see Table S1 in the supplemental material) revealed mixed bases in this region, suggesting a population of plasmids harboring different arrangements. We assembled the four shufflon contigs according to the arrangement that appeared to be dominant.

The resulting 92,339-bp plasmid was similar in organization to other IncI1 plasmids, with a complete conjugal transfer region, including the *traABCD* gene cluster, thin pilus formation region *pilI* through *pilV*, *traL* through *traY*, and *trb* transfer regions, and DNA-processing *nikAB* genes (21). Genes involved in the inhibition of the bacterial SOS response (*psiAB*) and in plasmid addiction (*pndCA*) were also identified (16, 21). pJIE512b harbored a 4,831-bp insert containing 161 bp of the right end of ISEcp1 and a 2,823-bp region of the *C. freundii* chromosome, including *bla*_{CMY-2} (Fig. 1c). ISEcp1 was truncated by IS1294, and the *C. freundii* region was followed by a 159-bp fragment of IncA/C backbone. IS1294 transposes by a

Received 12 March 2014 Returned for modification 20 April 2014

Accepted 27 May 2014

Published ahead of print 2 June 2014

Address correspondence to Sally R. Partridge, sally.partridge@health.nsw.gov.au.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.02773-14>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.02773-14

FIG 1 Different *bla*_{CMY-2} contexts. *ISecp1* is indicated by dark blue, inverted repeats are indicated by vertical black bars, and direct repeats are labeled and indicated by vertical gray bars. *IS1294* and *IS1294*-like elements are depicted with green and stippled green, respectively, target sites are labeled with black arrows, *terIS* sequences are depicted with green dots, and *terIS* look-alike sequences are depicted with black dots. Plasmid backbones are indicated by dashed red (IncA/C), cyan (IncI1), and pink (IncFII) lines. IncA/C backbone regions that have been mobilized are indicated by solid red lines. The lengths of the *Citrobacter freundii* part of the insertions are shown. (a) *C. freundii* chromosomal region, including *bla*_{CMY-2} (GenBank accession no. [U21727](#) and [AY125469](#)). (b) The 2,823-bp *bla*_{CMY-2} region in IncA/C plasmid pAR060302 (GenBank accession no. [FJ621588](#)) is associated with a complete *ISecp1*, and the whole insert is flanked by direct repeats. *IS1294* is present elsewhere in the backbone. (c) *bla*_{CMY-2} context in pJIE512b. The insert is flanked by the *IS1294* target site CTTG at the 5' end and bounded by the *IS1294* *terIS* look-alike sequence GTTC at the 3' end. (d) IncA/C plasmid pSH696_135 (GenBank accession no. [JN983048](#)) contains a complex *ISecp1*-*bla*_{CMY-2} arrangement, part of which is shown. An *IS1294b* element is present elsewhere in the plasmid. (e) The *IS1294b*-*ΔISecp1*-*bla*_{CMY-2} segment in pR7AC (GenBank accession no. [KF434766](#)) and pTN38148 ([FM246883](#)) is flanked at the 5' end by the imperfect target site GTCC and bounded at the 3' end by the *terIS* look-alike sequence GTTC. (f) The shorter *bla*_{CMY-2} insert in the IncFII plasmid pEQ011 (GenBank accession no. [KF582523](#)) is flanked by the *IS1294* target site GTTC and the imperfect *terIS* look-alike sequence CTTG.

rolling-circle mechanism and can capture adjacent regions by inefficient replication through its termination site (*terIS*) to an alternative *terIS* look-alike sequence (22). The IncA/C fragment in pJIE512b ends in GTTC, which matches the last 4 bp of *terIS*. This implies that *IS1294* inserted into *ISecp1* in an IncA/C plasmid and subsequently mobilized the adjacent region, including *bla*_{CMY-2} plus 159 bp of the IncA/C backbone. An IncA/C plasmid such as pAR060302 (GenBank accession no. [FJ621588](#)) (23), which carries an intact *ISecp1*, the 2,823-bp *bla*_{CMY-2} region, and *IS1294*, might be the source (Fig. 1b).

A similar *bla*_{CMY-2} context was identified in five partially sequenced IncI1 plasmids isolated from *E. coli* in France, including pTN38148 (GenBank accession no. [FM246883](#)) (9), and in a recently sequenced IncI1 ST2 plasmid from Denmark, pR7AC ([KF434766](#)). However, in these plasmids, an *IS1294*-like element, designated *IS1294b* here, truncates *ISecp1* at a different position, leaving a longer 372-bp fragment. The IncA/C fragment at the 3' end of the insert (1,395 bp) is also substantially longer than that in pJIE512b and again ends with the *IS1294* *terIS* look-alike sequence GTTC (Fig. 1e). *IS1294b* is ~95% identical to much of *IS1294*, but a 332-bp region that overlaps the start of the transposase gene is unrelated. *IS1294b* appears to be mobile, as it is found in at least eight different contexts in plasmid sequences in GenBank, including in IncA/C plasmids that also carry *bla*_{CMY-2}. One of these, e.g., pSH696_135 (GenBank accession no. [JN983048](#); Fig. 1d) (24), may be the source of the insert transferred to pR7AC. An *IS1294b*-*ΔISecp1*-*bla*_{CMY-2} insert with the same 372 bp of *ISecp1* but a shorter (1,553-bp) *C. freundii* region and with no IncA/C frag-

ment is present in the recently published IncFII plasmid pEQ011 (GenBank accession no. [KF582523](#)) (25) (Fig. 1f). This insert ends in an imperfect *terIS* look-alike sequence (CTTG) and thus appears to have been mobilized by *IS1294b* from a pR7AC-like IncI1 plasmid or directly from the IncA/C progenitor, both of which have longer *C. freundii* regions.

In pJIE512b, the *bla*_{CMY-2} insertion is upstream from *yafB* (unknown function), and an adjacent deletion removed part of the *cia* (colicin) gene (Fig. 2). In pR7AC, the insertion is downstream of *yafB*, a larger adjacent region is deleted, and *cia* is intact. Despite high homology across other regions of the backbones and identity at the pMLST target sites, a number of regions display <90% identity, including a short region encoding a single-stranded-DNA binding protein and several hypothetical proteins. An ~1.2-kb region of pJIE512b (at ~kb 10.5 in Fig. 2) (unknown function) shows no similarity to pR7AC but displays 100% identity to two IncI1 ST19 plasmids (16). These differences between the backbones and the different *bla*_{CMY-2} inserts reflect variability within pMLST ST2 and indicate, for at least this sequence type, that a more discriminatory plasmid typing method may be necessary for epidemiological purposes.

The novel *bla*_{CMY-2} context identified in this study, in addition to the contexts reported in the literature, suggests at least two separate mobilization events of *bla*_{CMY-2} from an IncA/C plasmid to an IncI1 plasmid mediated by an *IS1294*-like element. A recent paper also reported the *ISecp1*-mediated transfer of a *bla*_{CMY-2} region that includes 4,276 bp of backbone from an IncA/C plasmid to an IncX1 plasmid during conjugation experiments (26).

FIG 2 Comparison of the pJIE512b and pR7AC backbones, with *bla*_{CMY-2} inserts removed. Percent identity is displayed on the right of the diagram. The lengths and directions of selected genes are indicated by horizontal black arrows labeled with the gene name. pMLST target genes are shown in bold type, and pMLST amplicon positions are indicated by vertical blue arrows. The positions of the *bla*_{CMY-2} inserts in pJIE512b and pR7AC are indicated by red arrows. Important regions are shaded in green (*tra* region), purple (shufflon), or blue (thin pilus formation region). The location of the 2.3-kb deletion in pJIE512b is indicated by a vertical black arrow, and the position of the 3.3-kb deletion in pR7AC is labeled. The shufflon region is not included in the pR7AC GenBank entry. This diagram was drawn using mVISTA (<http://genome.lbl.gov/vista/index.shtml>) (28) using a calculation window of 100 bp.

These examples demonstrate movement of *bla*_{CMY-2} from a broad-host-range (IncA/C) plasmid to narrow-host-range (IncI1, IncF, IncX) plasmids that are more likely to be well adapted to their ecological niches (27), shedding light on the possible pathways by which *bla*_{CMY-2} (and potentially other resistance genes) have been mobilized into *E. coli*.

Nucleotide sequence accession number. The complete nucleotide sequence of pJIE512b has been submitted to the European Nucleotide Archive (ENA) under the accession number [HG970648](https://www.ebi.ac.uk/ena/entry/HG970648).

ACKNOWLEDGMENTS

Funding was provided by grants from the Australian Society for Antimicrobials and the Australian National Health and Medical Research Council (Centre for Research Excellence in Critical Infection G1001021). K.A.T. was supported by an Australian Postgraduate Award from the University of Sydney and a Westmead Millennium Institute Top-Up Scholarship.

We are grateful to Justin Ellem for the initial isolation of pJIE512b and pMLST, Carola Venturini for technical advice, Hana Dobiasova for designing shufflon primers, Joey Lai for library preparation, and Grant Hill-Cawthorne and Michael Roper for help with sequence assembly.

REFERENCES

- Jacoby GA. 2009. AmpC beta-lactamases. *Clin. Microbiol. Rev.* 22:161–182. <http://dx.doi.org/10.1128/CMR.00036-08>.
- Bauernfeind A, Stempling I, Jungwirth R, Giamarellou H. 1996. Characterization of the plasmidic beta-lactamase CMY-2, which is responsible for cephamycin resistance. *Antimicrob. Agents Chemother.* 40:221–224.
- Kang MS, Besser TE, Call DR. 2006. Variability in the region downstream of the *bla*_{CMY-2} beta-lactamase gene in *Escherichia coli* and *Salmonella enterica* plasmids. *Antimicrob. Agents Chemother.* 50:1590–1593. <http://dx.doi.org/10.1128/AAC.50.4.1590-1593.2006>.
- Su LH, Chen HL, Chia JH, Liu SY, Chu C, Wu TL, Chiu CH. 2006. Distribution of a transposon-like element carrying *bla*_{CMY-2} among *Salmonella* and other *Enterobacteriaceae*. *J. Antimicrob. Chemother.* 57:424–429. <http://dx.doi.org/10.1093/jac/dki478>.
- Allen KJ, Poppe C. 2002. Occurrence and characterization of resistance to extended-spectrum cephalosporins mediated by beta-lactamase CMY-2 in *Salmonella* isolated from food-producing animals in Canada. *Can. J. Vet. Res.* 66:137–144.
- Carattoli A, Tosini F, Giles WP, Rupp ME, Hinrichs SH, Angulo FJ, Barrett TJ, Fey PD. 2002. Characterization of plasmids carrying CMY-2 from expanded-spectrum cephalosporin-resistant *Salmonella* strains isolated in the United States between 1996 and 1998. *Antimicrob. Agents Chemother.* 46:1269–1272. <http://dx.doi.org/10.1128/AAC.46.5.1269-1272.2002>.
- Barlow M, Hall BG. 2002. Origin and evolution of the AmpC beta-lactamases of *Citrobacter freundii*. *Antimicrob. Agents Chemother.* 46:1190–1198. <http://dx.doi.org/10.1128/AAC.46.5.1190-1198.2002>.
- Partridge SR. 2011. Analysis of antibiotic resistance regions in Gram-negative bacteria. *FEMS Microbiol. Rev.* 35:820–855. <http://dx.doi.org/10.1111/j.1574-6976.2011.00277.x>.
- Verdet C, Gautier V, Chachaty E, Ronco E, Hidri N, Decré D, Arlet G. 2009. Genetic context of plasmid-carried *bla*_{CMY-2}-like genes in *Enterobacteriaceae*. *Antimicrob. Agents Chemother.* 53:4002–4006. <http://dx.doi.org/10.1128/AAC.00753-08>.
- Bortolaia V, Hansen KH, Nielsen CA, Fritsche TR, Guardabassi L. 4 February 2014. High diversity of plasmids harbouring *bla*_{CMY-2} among clinical *Escherichia coli* isolates from humans and companion animals in the upper midwestern U. S. A. *J. Antimicrob. Chemother.* <http://dx.doi.org/10.1093/jac/dku011>.
- Mataseje LF, Baudry PJ, Zhanel GG, Morck DW, Read RR, Louie M, Mulvey MR. 2010. Comparison of CMY-2 plasmids isolated from human, animal, and environmental *Escherichia coli* and *Salmonella* spp. from Canada. *Diagn. Microbiol. Infect. Dis.* 67:387–391. <http://dx.doi.org/10.1016/j.diagmicrobio.2010.02.027>.
- Hiki M, Usui M, Kojima A, Ozawa M, Ishii Y, Asai T. 2013. Diversity of plasmid replicons encoding the *bla*_{CMY-2} gene in broad-spectrum cephalosporin-resistant *Escherichia coli* from livestock animals in Japan. *Foodborne Pathog. Dis.* 10:243–249. <http://dx.doi.org/10.1089/fpd.2012.1306>.
- Weissman SJ, Adler A, Qin X, Zerr DM. 2013. Emergence of extended-spectrum beta-lactam resistance among *Escherichia coli* at a US academic children's hospital is clonal at the sequence type level for CTX-M-15, but not for CMY-2. *Int. J. Antimicrob. Agents* 41:414–420. <http://dx.doi.org/10.1016/j.ijantimicag.2013.01.006>.
- Folster JP, Tolar B, Pecic G, Sheehan D, Rickert R, Hise K, Zhao S, Fedorka-Cray PJ, McDermott P, Whichard JM. 2014. Characterization of *bla*_{CMY} plasmids and their possible role in source attribution of *Salmonella enterica* serotype Typhimurium infections. *Foodborne Pathog. Dis.* 11:301–306. <http://dx.doi.org/10.1089/fpd.2013.1670>.
- Mata C, Miró E, Alvarado A, Garcillán-Barcia MP, Toleman M, Walsh TR, de la Cruz F, Navarro F. 2012. Plasmid typing and genetic context of AmpC beta-lactamases in *Enterobacteriaceae* lacking inducible chromosomal *ampC* genes: findings from a Spanish hospital 1999–2007. *J. Antimicrob. Chemother.* 67:115–122. <http://dx.doi.org/10.1093/jac/dkr412>.
- Johnson TJ, Shepard SM, Rivet B, Danzeisen JL, Carattoli A. 2011. Comparative genomics and phylogeny of the IncI1 plasmids: a common plasmid type among porcine enterotoxigenic *Escherichia coli*. *Plasmid* 66:144–151. <http://dx.doi.org/10.1016/j.plasmid.2011.07.003>.
- García-Fernández A, Chiaretto G, Bertini A, Villa L, Fortini D, Ricci A, Carattoli A. 2008. Multilocus sequence typing of IncI1 plasmids carrying extended-spectrum beta-lactamases in *Escherichia coli* and *Salmonella* of human and animal origin. *J. Antimicrob. Chemother.* 61:1229–1233. <http://dx.doi.org/10.1093/jac/dkn131>.
- Valenzuela JK, Thomas L, Partridge SR, van der Reijden T, Dijkshoorn L, Iredell J. 2007. Horizontal gene transfer in a polyclonal outbreak of carbapenem-resistant *Acinetobacter baumannii*. *J. Clin. Microbiol.* 45:453–460. <http://dx.doi.org/10.1128/JCM.01971-06>.

19. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
20. Gyohda A, Furuya N, Ishiwa A, Zhu S, Komano T. 2004. Structure and function of the shufflon in plasmid R64. *Adv. Biophys.* 38:183–213.
21. Sampei G, Furuya N, Tachibana K, Saitou Y, Suzuki T, Mizobuchi K, Komano T. 2010. Complete genome sequence of the incompatibility group II plasmid R64. *Plasmid* 64:92–103. <http://dx.doi.org/10.1016/j.plasmid.2010.05.005>.
22. Tavakoli N, Comanducci A, Dodd HM, Lett Albiger MCB, Bennett P. 2000. IS1294, a DNA element that transposes by RC transposition. *Plasmid* 44:66–84. <http://dx.doi.org/10.1006/plas.1999.1460>.
23. Call DR, Singer RS, Meng D, Broschat SL, Orfe LH, Anderson JM, Herndon DR, Kappmeyer LS, Daniels JB, Besser TE. 2010. *bla*_{CMY-2}-positive IncA/C plasmids from *Escherichia coli* and *Salmonella enterica* are a distinct component of a larger lineage of plasmids. *Antimicrob. Agents Chemother.* 54:590–596. <http://dx.doi.org/10.1128/AAC.00055-09>.
24. Han J, Lynne AM, David DE, Tang H, Xu J, Nayak R, Kaldhøne P, Logue CM, Foley SL. 2012. DNA sequence analysis of plasmids from multidrug resistant *Salmonella enterica* serotype Heidelberg isolates. *PLoS One* 7:e51160. <http://dx.doi.org/10.1371/journal.pone.0051160>.
25. Karczmarczyk M, Wang J, Leonard N, Fanning S. 2014. Complete nucleotide sequence of a conjugative IncF plasmid from an *Escherichia coli* isolate of equine origin containing *bla*_{CMY-2} within a novel genetic context. *FEMS Microbiol. Lett.* 352:123–127. <http://dx.doi.org/10.1111/1574-6968.12364>.
26. Wiesner M, Fernández-Mora M, Cevallos MA, Zavala-Alvarado C, Zaidi MB, Calva E, Silva C. 2013. Conjugative transfer of an IncA/C plasmid-borne *bla*_{CMY-2} gene through genetic rearrangements with an IncX1 plasmid. *BMC Microbiol.* 13:264. <http://dx.doi.org/10.1186/1471-2180-13-264>.
27. De Gelder L, Williams JJ, Ponciano JM, Sota M, Top EM. 2008. Adaptive plasmid evolution results in host-range expansion of a broad-host-range plasmid. *Genetics* 178:2179–2190. <http://dx.doi.org/10.1534/genetics.107.084475>.
28. Frazer K, Pachter L, Poliakov A, Rubin EM, Dubchak I. 2004. VISTA: computational tools for comparative genomics. *Nucleic Acids Res.* 1:W273–W279. <http://dx.doi.org/10.1093/nar/gkh458>.