Published in final edited form as:

Mol Cell Neurosci. 2014 July ; 0: 163–175. doi:10.1016/j.mcn.2014.06.007.

Cortical parvalbumin GABAergic deficits with $\alpha 7$ nicotinic acetylcholine receptor deletion: Implications for schizophrenia

Hong Lin¹, Fu-Chun Hsu¹, Bailey H. Baumann¹, Douglas A. Coulter^{1,3}, Stewart A. Anderson^{2,3}, and David R. Lynch^{1,3}

¹Departments of Pediatrics and Neurology, The Children's Hospital of Philadelphia, Philadelphia PA, 19104

²Departments of Pediatrics and Child Psychiatry, The Children's Hospital of Philadelphia, Philadelphia PA, 19104

³Perelman School of Medicine, University of Pennsylvania, Philadelphia PA, 19104

Abstract

Dysfunction of cortical parvalbumin (PV)-containing GABAergic interneurons has been implicated in cognitive deficits of schizophrenia. In humans microdeletion of the *CHRNA7* ($\alpha 7$ nicotinic acetylcholine receptor, nAChR) gene is associated with cortical dysfunction in a broad spectrum of neurodevelopmental and neuropsychiatric disorders including schizophrenia while in mice similar deletion causes analogous abnormalities including impaired attention, working-memory and learning. However, the pathophysiological roles of $\alpha 7$ nAChRs in cortical PV GABAergic development remain largely uncharacterized. In both *in vivo* and *in vitro* models, we identify here that deletion of the $\alpha 7$ nAChR gene in mice impairs cortical PV GABAergic development and recapitulates many of the characteristic neurochemical deficits in PV-positive GABAergic interneurons found in schizophrenia. $\alpha 7$ nAChR *null* mice had decreased cortical levels of GABAergic markers including PV, Glutamic Acid Decarboxylase 65/67 (GAD65/67) and the $\alpha 1$ subunit of GABA_A receptors, particularly reductions of PV and GAD67 levels in cortical PV-positive interneurons during late postnatal life and adulthood. Cortical GABAergic synaptic deficits were identified in the prefrontal cortex of $\alpha 7$ nAChR *null* mice and $\alpha 7$ nAChR *null* cortical cultures. Similar disruptions in development of PV-positive GABAergic interneurons and perisomatic synapses were found in cortical cultures lacking $\alpha 7$ nAChRs. Moreover, NMDA receptor expression was reduced in GABAergic interneurons, implicating NMDA receptor hypofunction in GABAergic deficits in $\alpha 7$ nAChR *null* mice. Our findings thus demonstrate impaired cortical PV GABAergic development and multiple characteristic neurochemical deficits reminiscent of schizophrenia in cortical PV-positive interneurons in $\alpha 7$ nAChR gene deletion

© 2014 Elsevier Inc. All rights reserved.

CORRESPONDING AUTHOR: David R. Lynch, M.D., Ph.D., Division of Neurology, The Children's Hospital of Philadelphia, 502 Abramson Research Center, 3615 Civic Center Blvd, Philadelphia, Pennsylvania 19104-4318, Tel: 215-590-2242, Fax: 215-590-3779, lynchd@mail.med.upenn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CONFLICT OF INTEREST: No conflict of interest exists in this paper for all authors.

models. This implicates crucial roles of $\alpha 7$ nAChRs in cortical PV GABAergic development and dysfunction in schizophrenia and other neuropsychiatric disorders.

Keywords

cortex; $\alpha 7$ nicotinic ACh receptor; parvalbumin; GABAergic; synapse; NMDA receptor

Introduction

Nicotinic receptor mediated neurotransmission has been implicated in many CNS processes including neuronal plasticity, learning and memory as well as in neurological, neurodevelopmental and psychiatric disorders (Dani and Bertrand, 2007; Lin et al., 2010; Ross et al., 2010; Miwa et al., 2011; Griguoli and Cherubini, 2012; Wallace and Bertrand, 2013; Freedman, 2014; Lin et al., 2014). Some of the most powerful evidence comes from genetic studies in which a homozygous microdeletion of human15q13.3, including the $\alpha 7$ nAChR gene, is associated with cortical circuit disorders including schizophrenia, autism, mental retardation, epilepsy, and intellectual disability (Freedman et al., 1997; Sharp et al., 2008; Helbig et al., 2009; Miller et al., 2009; Shinawi et al., 2009; Endris et al., 2010; Lepichon et al., 2010; Ancin et al., 2011; Liao et al., 2011; Spielmann et al., 2011; Yasui et al., 2011; Hoppman-Chaney et al., 2012; Freedman, 2014). Based on the size and the overlap of deletions in such individuals, loss of the $\alpha 7$ nAChR gene alone appears sufficient to cause the majority of clinical features in such patients (Shinawi et al., 2009; Endris et al., 2010; Liao et al., 2011; Hoppman-Chaney et al., 2012). In humans reduced levels of $\alpha 7$ nAChR have been found in the prefrontal cortex and hippocampus of schizophrenia patients in postmortem studies (Freedman et al., 1995; Leonard et al., 1996; Freedman et al., 1997; Freedman et al., 2000; Freedman and Leonard, 2001; Freedman and Goldowitz, 2010; Ross et al., 2010). In animals loss of the $\alpha 7$ nAChR causes analogous abnormalities including impaired attention, working-memory and learning (Fernandes et al., 2006; Young et al., 2007; Levin et al., 2009; Brown et al., 2010; Young et al., 2011; Hellier et al., 2012). These findings suggest crucial roles of $\alpha 7$ nAChR in cortical circuit function and dysfunction.

Cortical parvalbumin (PV)-containing GABAergic interneurons have also been implicated in cortical circuit function and dysfunction in schizophrenia (Lewis et al., 2012; Nakazawa et al., 2012; Inan et al., 2013; Jiang et al., 2013; Le Magueresse and Monyer, 2013). A major role of GABA-mediated neurotransmission is to synchronize network oscillations essential for normal cognition (Gonzalez-Burgos et al., 2011), and impaired GABAergic neurotransmission in schizophrenia may underlie impairments in cortical gamma oscillations associated with cognitive dysfunction (McNally et al., 2013). In particular, the PV-positive subtype of GABAergic interneurons have uniquely fast membrane and synaptic properties that are crucial for cortical circuit functions such as gamma oscillations (Lewis et al., 2012). The cortical PV-positive interneurons consist of two main types: chandelier and basket cells. Cortical gamma oscillations require strong inhibitory inputs to pyramidal neurons from the PV-positive basket cells and are mediated through postsynaptic $\alpha 1$ subunit-containing GABA_A receptors in pyramidal neurons (Lewis et al., 2012; Inan et al., 2013). Dysfunction of cortical PV-positive basket cells has been implicated in aberrant gamma oscillations and

cognitive deficits in schizophrenia due to reduced levels of PV and the 67-kDa isoform of GABA synthesizing enzyme glutamic acid decarboxylase (GAD67) as well as lower GABA_A receptor $\alpha 1$ subunit levels in pyramidal neurons postsynaptic to basket cells (Curley and Lewis, 2012; Lewis et al., 2012). These observations implicate cortical PV-positive basket cells and their postsynaptic pyramidal cells in cortical circuit function and dysfunction in schizophrenia.

Thus, both $\alpha 7$ nAChR and cortical PV GABAergic abnormalities are associated with dysfunction in schizophrenia. Apart from decreased immunoreactivities of GAD65 and GABA_A receptor binding in the hippocampus of heterozygous $\alpha 7$ nAChR knockout mice, the pathophysiological processes linking nicotinic receptor and cortical PV GABAergic abnormalities remain largely uncharacterized (Adams et al., 2012). In the present studies in $\alpha 7$ nAChR gene deletion models, we demonstrate impaired cortical PV GABAergic development and multiple characteristic neurochemical deficits that are reminiscent of cortical PV GABAergic deficits in schizophrenia. These include reduced PV and GAD67 levels in PV-positive GABAergic interneurons, reduced GABA_A $\alpha 1$ levels in pyramidal neurons as well as perisomatic GABAergic synaptic deficits. Our findings thus implicate $\alpha 7$ nAChR in cortical PV GABAergic circuit development and dysfunction in schizophrenia and other neuropsychiatric disorders.

Materials and Methods

Materials

Timed-pregnant C57BL/6 mice were purchased from Charles River Laboratories and bred from $\alpha 7$ nAChR *knockout* mice (B6.129S7-Chrna7^{tm1Bay}/J, Jackson Laboratory). Biochemicals and antibodies included kynurenic acid (Sigma), α -GABA_A $\alpha 1$ (Millipore, rabbit polyclonal), α -somatostatin (Millipore, rabbit polyclonal), α -parvalbumin (Millipore, mouse monoclonal; Swant, rabbit polyclonal), α -GAD65/67 (Millipore, rabbit polyclonal), α -GAD65 (Developmental Studies Hybridoma Bank, mouse monoclonal), α -VGAT (Millipore, rabbit polyclonal), α -VGLUT1 (Synaptic Systems, rabbit polyclonal), α -GAPDH (Novus Biologicals, mouse monoclonal), α -actin (Sigma, rabbit polyclonal).

Neuronal cultures

Primary cortical cultures from E17-19 *wild-type* or $\alpha 7$ nAChR *knockout* mice were prepared as described (Dong et al., 2004). Briefly, the cortex was dissected, gently minced, trypsinized (0.027%, 37 °C; 7% CO₂ for 20 min), and then washed with 1× HBSS. Neurons were seeded to a density of 3×10^5 viable cells in 35-mm culture dish with five 12-mm glass coverslips or a density of 1.6×10^6 viable cells in 60-mm culture dishes. The culture dishes were coated with poly-D-Lysine (100 μ g/ml) prior to seeding neurons. Neurons were maintained at 37°C with 5% CO₂ in Neurobasal medium with B27 supplement. Neurobasal medium contains choline chloride, a selective agonist at $\alpha 7$ nAChR (Zhang and Warren, 2002). In addition, cholinergic neurons are present in cortical cultures as identified by ChAT immunostaining as described by Abcam in their manufacturer's instructions (Lin et al., 2010). At 21–28 *days in vitro* (DIV), cultures were subject to western blotting analysis, immunocytochemistry or patch clamp recording.

For cell lysate preparation, cultures were lysed in lysis buffer (150 mM NaCl, 1 mM EDTA, 100 mM Tris-HCl, 1% Triton X-100, 1% sodium deoxycholate and 1% SDS, pH 7.4, supplemented the day of use with 1:500 EDTA-free protease inhibitor cocktail III (Calbiochem) for 1 hr at 4°C and collected. Debris was cleared by centrifugation at 16,100 × g for 20 mins at 4°C. Supernatants were stored at –80°C until use.

Tissue preparation

For tissue homogenate preparation, the age-matched *wild-type* (WT) and $\alpha 7$ nAChR *null* ($\alpha 7$ -KO) mouse littermates at postnatal day (P1-P56) and 9 months older of either sex were anesthetized with isoflurane before decapitation in accordance with protocols approved by The Children's Hospital of Philadelphia Animal Care and Use Committee. The mouse brain was rapidly removed, and the cortex was dissected under Leica EZ4 stereomicroscope and immediately transferred to dry ice. The cortex was homogenized in 20ml lysis buffer per 1 g weight, and lysed for 1 hr at 4°C. Lysis buffer contained 150 mM NaCl, 1 mM EDTA, 100 mM Tris-HCl, 1% Triton X-100, and 1% sodium deoxycholate, pH 7.4, supplemented the day of use with 1:500 EDTA-free protease inhibitor cocktail III (Calbiochem). Debris was cleared by centrifugation at 39,000 × g for 1 hr at 4°C. Supernatants were stored at –80°C until use.

For immunohistochemical studies, the age-matched male WT and $\alpha 7$ -KO mice at P56-P90 were anesthetized with isoflurane and ketamine/xylazine mixture, and cardiac perfused with 10 ml of PBS, followed by 20 ml of PBS containing 4% paraformaldehyde in accordance with protocols approved by The Children's Hospital of Philadelphia Animal Care and Use Committee. Brains were excised and immersed overnight in 4% paraformaldehyde, washed in PBS, dehydrated and embedded in paraffin. A series of brain coronal sections (5 μ m thick) were obtained using microtome at The Children's Hospital of Philadelphia Pathology Core Facility.

Western blotting analysis

Western blotting was performed as described previously (Dong et al., 2004). Protein content was determined using BCA Protein Assay (Thermo Scientific). Equal amounts of total protein (5 μ g cell lysate or 30 μ g cortex homogenate per lane) was subjected to 4–12% NuPAGE Gel for electrophoresis and transferred to nitrocellulose membranes. Membranes were blocked with 3% nonfat milk and incubated with primary antibody overnight at 4 °C. Blots were then incubated with appropriate horseradish peroxidase, HRP-conjugated secondary antibodies (Cell Signaling) for 2 hr at room temperature and then washed; Reaction bands were visualized using a luminol-enhanced chemiluminescence (ECL) HRP substrate (Thermo Scientific). Each blot was then incubated with stripping buffer (2% SDS, 50 mM Tris, pH 6.8, and 100 mM β -mercaptoethanol) for 45 mins at room temperature to remove the signals and reprobed for other proteins including actin and GAPDH as internal controls. Reaction product levels were quantified by scanning densitometry and normalized by the levels of actin or GAPDH using NIH Image J software. Since the cortical actin and GAPDH levels alter during the time course of postnatal development, the protein levels were normalized by internal control at the same time-point between WT and KO mice.

Immunocytochemistry and immunohistochemistry

Primary cultured cortical neurons were fixed for 20 min at 4°C with 4% paraformaldehyde in phosphate-buffered saline (PBS) (pH 7.4), and then subject to immunostaining procedure. The paraffin-embedded brain sections were deparaffined, rehydrated, and antigen-retrieved in antigen unmasking solution (Vector Laboratories) and then subject to immunostaining procedure. For immunostaining procedure, after blocking with 5% normal goat serum and 1 % bovine serum albumin in combination with 0.3% (vol/vol) Triton X-100 in PBS at room temperature for 1 hr, the coverslips or slides were incubated with primary antibodies at 4°C overnight and then secondary antibodies conjugated to Alexa fluor 488 or 594 (Invitrogen) at room temperature for 60–90 mins. Following several washes with PBS, cells or slides were mounted with Vectashield with DAPI (Vector Laboratories).

Fluorescence imaging

Fluorescence images were obtained with Olympus FluoView laser scanning confocal microscope. For mouse cortex, the brain sections were stained for GABAergic markers (PV, SST, GAD65/67, GABA_Aα1). The infralimbic, prelimbic, or anterior cingulate regions of the mouse medial frontal cortex are collectively referred to as mouse prefrontal cortex (Rotaru et al., 2011). Confocal scans were performed in layer 2–6 of mouse prefrontal cortex, and the imaging parameters were identical for *wild-type* and α7 nAChR *knockout* mice. *Wild-type* controls were always included in all experiments to normalize for expected variations in antibody staining intensity performed on different days. The confocal images were acquired at the focal plane with maximal number of synapses from at least 3 sections per animal and from at least 3 animals per group. NIH Image J software was used to quantify VGAT- or GAD65-positive GABAergic synapses in the acquired confocal images. Thresholds were set at 3 SDs above the mean staining intensity of six nearby regions in the same visual field. Thresholded images present a fixed intensity for all pixels above threshold after having removed all of those below and VGAT- or GAD65-labeled puncta in the thresholded images were quantified (Lozada et al., 2012). For quantification of the immunofluorescence intensity of PV and SST in the soma of PV- or SST-positive interneurons, the acquired confocal images were quantified using NIH image J software in the thresholded images from 6–8 sections per animal and from at least 3 animals per group. For quantification of the number of the PV-positive GABAergic interneurons in mouse prefrontal cortex, a blind observer counted the PV-positive interneurons in mouse prefrontal cortex under the epifluorescence microscopy from 7–8 sections per animal and from at least 3 animals per group. For cortical cultures, neurons were stained for GABAergic markers (VGAT, GAD65, GABA_Aα1 and PV). For quantification analysis, the confocal images were acquired under 63× objectives from 4–6 samples and 4–5 different cultures for quantification of VGAT- or PV-labeled puncta in *wild-type* and α7 nAChR *knockout* cortical cultures. NIH Image J software and the thresholded images were used to quantify the number of GABAergic synapses in cortical cultures.

Whole-cell patch-clamp recording

Cortical pyramidal neurons were identified based on their morphological properties and firing pattern. Specifically, besides the typical morphology of cultured cortical pyramidal

neurons, pyramidal neurons were also confirmed by their action potential firing patterns in respond to intracellular injection of depolarized current using whole cell patch-clamping techniques (Rotaru et al., 2011). Since spontaneous IPSC recording was recorded using a CsCl-based intrapipette solution and resting membrane potential of recorded cells would be affected under current clamp mode of recording, an initial screening recording was conducted to reduce the data contamination by interneurons using potassium gluconate-based intrapipette solution. The recording electrodes were filled with potassium gluconate-based intrapipette solution (in mM: 145 K-gluconate, 2 MgCl₂, 2 ATP·Mg, 0.5 GTP·Tris, 0.1 BAPTA, 2.5 KCl, 2.5 NaCl, and 10 HEPES), and the recording chamber was perfused with artificial CSF (aCSF) with composition (in mM): 130 NaCl, 3 KCl, 2 CaCl₂, 1 MgCl₂, 1.25 NaH₂PO₄, 26 NaHCO₃ and 10 glucose. Cultured cortical neurons were recorded under current clamp mode and action potentials were induced by intracellular injection of depolarized current. Matched with the morphology, 100% of selected cells (n = 15) for an initial screening recording were cortical pyramidal neurons judged by their action potential firing patterns (Rotaru et al., 2011). Also, an example of cortical pyramidal neuron responding to intracellular injection (700 msec) of depolarized current is shown in Supplemental Figure 1.

Spontaneous inhibitory postsynaptic current (sIPSC) recording was performed using standard whole-cell voltage clamp techniques in cultured mouse cortical neurons grown on coverslips for 21–28 DIV. Cells were transferred to a recording chamber at room temperature (23 ± 2°C) containing an extracellular solution composed of (mM): 155 NaCl, 3 KCl, 3 CaCl₂, 1 MgCl₂, 10 HEPES, pH 7.4. Recording electrodes were pulled on a Flaming-Brown micropipette puller (Model P-97, Sutter Instrument Co.) and filled with a solution containing (mM): 130 CsCl, 2 MgCl₂, 10 HEPES, 0.8 CsOH, 2 Mg-ATP, pH 7.35, 280–300 mOsm. In order to isolate the inhibitory synaptic neuronal events, the glutamate receptor-mediated sEPSCs were blocked by bath application of glutamate receptor antagonist kynurenic acid (2 mM) in extracellular solution. The holding potential for sIPSC recording was –70 mV. Recorded signals were amplified using an Axopatch-1D amplifier, filtered at 5 kHz and then saved as data files using pCLAMP 9.01 software (Axon Instrument Inc., Hamden, CT) for off-line analysis. The synaptic events were analyzed using Mini-Analysis software to calculate the sIPSC frequency, amplitude as well as current decay time constant.

Statistical analysis

Data were shown as the mean ± S.E.M. Experiments were analyzed using Mann-Whitney Rank Sum test or Student's *t*-test to compare two conditions or ANOVA followed by planned comparisons of multiple conditions. Significance was set at *P* < 0.05.

Results

Cortical levels of GABAergic markers were decreased in $\alpha 7$ nAChR *null* mice during late postnatal development and adulthood

Our previous studies have demonstrated that $\alpha 7$ nAChR gene deletion leads to specific loss of synaptic NMDA receptors and their coagonist, D-serine, as well as glutamatergic synaptic deficits in mouse cortex (Lin et al., 2014). In the present study, we explored whether $\alpha 7$

nAChR gene deletion affects GABAergic neuronal development by examining the expression of parvalbumin (PV), the $\alpha 1$ subunit of GABA_A receptor (GABA_A $\alpha 1$), and two isoforms of the GABA synthesizing enzyme, glutamic acid decarboxylase 65/67 (GAD65/67) as markers of GABAergic interneurons in the cortex of $\alpha 7$ nAChR *null* ($\alpha 7$ -KO) mice and *wild-type* (WT) littermates at P1-P56 and P9M. In WT mice, expression of PV remained low until P21, peaked at P56, and remained high in adulthood (Fig. 1A). Levels of GAD65 remained low until P14, increased by P21 and reached its maximum at P56 (Fig. 1B), while the levels of GABA_A $\alpha 1$ remained low until P8, began to increase by P14, peaked at P56 and remained high in adulthood (Fig. 1C). The results suggest that the neurochemical maturation of mouse cortical GABAergic circuitry occurs by P56, shortly after sexual maturation, similar to development of GABAergic circuitry in monkeys (Hoftman and Lewis, 2011).

Remarkably, in the neocortex of $\alpha 7$ -KO mice expression levels of these markers of GABAergic maturation were significantly reduced. PV was modestly but significantly reduced at P21, an effect that persisted into adulthood (Fig. 1A). For both GAD65 and GABA_A $\alpha 1$, there was a trend for a decrease during the second postnatal week and in adults, with a statistically significant decrease present at P21 and P56 (Fig. 1B and 1C). Similar to GAD65, GAD67 expression levels were also decreased in the cortex of $\alpha 7$ -KO mice compared with WT littermates (Fig. 1B). These findings suggest that $\alpha 7$ nAChR deletion alters cortical GABA synthesis, PV regulation, and GABA_A $\alpha 1$ subunit-containing receptor presence during late postnatal development.

Neurochemical deficits of cortical PV-positive GABAergic interneurons in the prefrontal cortex of $\alpha 7$ nAChR *null* mice

We then examined the effects of $\alpha 7$ nAChR gene deletion on cortical PV-positive GABAergic interneurons in mouse prefrontal cortex by examining PV, GAD65/67 and GABA_A $\alpha 1$ immunoreactivities microscopically at P56-P90. In the prefrontal cortex of WT mice, the PV-positive GABAergic interneurons were scattered in layers 2–6 (Fig. 2A, 2G). GAD67 was localized in the soma of PV-positive GABAergic interneurons, and GAD65 in the GABAergic synapses. GAD65-positive GABAergic synapses were widely and extensively distributed in the neuropil surrounding the PV-positive GABAergic neurons and pyramidal neurons (Fig. 2B). The GABA_A $\alpha 1$ -containing receptors were found in both PV-positive GABAergic neurons and pyramidal neurons (Fig. 2H). In $\alpha 7$ -KO mice, the overall levels of PV, GAD65/67 and GABA_A $\alpha 1$ immunoreactivities (Fig. 2D–2F and 2J–2L) were decreased compared with WT littermates (Fig. 2A–2C and 2G–2I), demonstrating that deletion of the $\alpha 7$ nAChR gene reduced PV, GAD65/67 and GABA_A $\alpha 1$ expression during late postnatal development. We further examined the specificity of PV alteration in $\alpha 7$ -KO mice by examining somatostatin (SST) immunoreactivity, a marker of a different subclass of GABAergic interneurons, in the mouse cortex. In the prefrontal cortex of WT mice, the SST-positive GABAergic interneurons were scattered in all layers (Fig. 3B), while the PV-positive GABAergic interneurons were scattered in layers 2–6 (Fig. 3A). The overall level of somatostatin was not altered in the prefrontal cortex of $\alpha 7$ -KO mice (Fig. 3D–3F and 3J–3L), whereas that of PV was markedly reduced compared with WT littermates (Fig. 3A–3C and 3G–3I). Quantification analysis showed significant reduction of PV levels in PV-

positive GABAergic interneurons (Fig. 3M) and similar levels of SST in SST-positive GABAergic interneurons (Fig. 3N) in the prefrontal cortex of $\alpha 7$ -KO mice compared with WT mice. Although PV levels were decreased in the prefrontal cortex, the number of PV-positive GABAergic neurons in the prefrontal cortex of $\alpha 7$ -KO mice was not significantly decreased compared with that of WT littermates (Fig. 3O). Our findings suggest that deletion of the $\alpha 7$ nAChR gene reduces cortical PV levels but not PV interneuron number, reflecting a reduction of PV synthesis rather than a failure of PV-positive neurons to develop.

Moreover, PV and GAD67 levels were reduced in the soma of PV-positive GABAergic interneurons (Fig. 4D–4F, white arrows) as were the number of GAD65-positive GABAergic synapses surrounding the pyramidal neurons (Fig. 4D–4F grey arrows) in the prefrontal cortex of $\alpha 7$ -KO mice compared with WT littermates (Fig. 4A–4C). The GABA_A $\alpha 1$ -containing receptors were localized in both PV-positive GABAergic interneurons (Fig. 4G–4I, white arrows) and pyramidal neurons (Fig. 4G–4I, grey arrows), with higher levels in PV-positive GABAergic interneurons in the prefrontal cortex of WT mice. GABA_A $\alpha 1$ levels were reduced in both PV-positive GABAergic interneurons (Fig. 4J–4L, white arrows) and pyramidal neurons (Fig. 4J–4L, grey arrows) in the prefrontal cortex of $\alpha 7$ -KO mice compared with WT littermates (Fig. 4G–4I). Our findings thus demonstrated multiple characteristic neurochemical deficits reminiscent of schizophrenia in PV-positive GABAergic interneurons in mouse prefrontal cortex with $\alpha 7$ nAChR deletion.

Cortical GABAergic synapse formation was impaired in prefrontal cortex and in cortical cultures of $\alpha 7$ nAChR null mice

We then examined whether $\alpha 7$ nAChR gene deletion affects GABAergic synapse formation by examining the immunoreactivity of a GABAergic specific presynaptic marker, the vesicular GABA transporter (VGAT), in the prefrontal cortex. In the prefrontal cortex of WT mice, similar to GAD65-positive GABAergic presynaptic terminals, VGAT-positive GABAergic presynaptic terminals were extensively distributed in the neuropil surrounding the PV-positive GABAergic interneurons and pyramidal neurons (Fig. 5A–5C). In $\alpha 7$ -KO mice, the overall levels of VGAT immunoreactivity (Fig. 5D–5F) were reduced compared with WT littermates (Fig. 5A–5C), and the number of VGAT-positive GABAergic presynaptic terminals surrounding the pyramidal neurons (Fig. 5J–5L, grey arrows) were reduced in the prefrontal cortex of $\alpha 7$ -KO mice compared with WT littermates (Fig. 5G–5I, grey arrows). Quantification of VGAT-positive GABAergic presynaptic terminals identified a significant decrease in the prefrontal cortex of $\alpha 7$ -KO mice compared with WT littermates (Fig. 5M). Thus, deletion of the $\alpha 7$ nAChR decreased cortical GABAergic synapse formation in mouse prefrontal cortex.

We further examined the effects of $\alpha 7$ nAChR gene deletion on GABAergic synaptic development in cortical cultures. Double immunostaining with GABAergic presynaptic marker GAD65 and postsynaptic marker GABA_A $\alpha 1$ antibodies identified GAD65- and GABA_A $\alpha 1$ -positive GABAergic synapses and demonstrated a reduction in the number of GABAergic synapses and level of GABA_A $\alpha 1$ in $\alpha 7$ -KO cultures (Fig. 6D–6F) compared with WT cultures (Fig. 6A–6C). However, high magnification of confocal images showed

that 100% of GAD65-positive puncta were colocalized with or adjacent to one or more GABA α 1-positive puncta in the dendrites of cortical pyramidal neurons as shown in Fig. 6A–6F, the number of GABAergic specific presynaptic marker VGAT-positive puncta thus accurately reflects the number of GABAergic synapses on the dendrites of pyramidal neurons. Immunostaining with VGAT antibody identified specific presynaptic terminals of GABAergic synapses in WT (Fig. 6G) and α 7-KO (Fig. 6H) cultures and demonstrated a significant reduction in the number of GABAergic synapses on the dendrites of pyramidal neurons in α 7-KO cortical cultures compared with WT cultures based on quantification (Fig. 6I). Whole-cell voltage clamp recording showed significant reduction in sIPSC frequency and decay time (Fig. 6J–6L), but not amplitude (Fig. 6M), in cultured cortical pyramidal neurons, consistent with the reduction of GABAergic synapses in α 7-KO cortical cultures. The findings thus demonstrate that deletion of the α 7 nAChR gene in mice severely impairs cortical GABAergic synapse formation *in vivo* and *in vitro*.

The development and maturation of cortical PV-positive basket cells were impaired in α 7 nAChR null cortical cultures

We further explored whether deletion of the α 7 nAChR impaired the development of cortical PV-positive GABAergic interneurons by examining PV and GAD65 immunoreactivities in cortical cultures at 21–28 DIV. In WT cortical cultures, PV immunoreactivity was widely distributed in the soma, dendrites, axons and synapses of cortical PV-positive GABAergic interneurons (Fig. 7A–7C and 7G–7I). The axons of an individual PV-positive basket cells formed perisomatic GABAergic synapses surrounding the multiple pyramidal neurons. The PV-positive perisomatic GABAergic synapses were positive for GAD65 immunoreactivity (Fig. 7A–7C and 7G–7I, white arrows). In α 7-KO cultures, the development of PV-positive basket cells and the formation PV- and GAD65-positive perisomatic GABAergic synapses were impaired (Fig. 7D–7F and 7J–7L, white arrows) compared with WT cultures (Fig. 7A–7C and 7G–7I, white arrows). The overall PV immunoreactivity in the soma of PV-positive GABAergic interneurons was significantly decreased in α 7-KO cultures compared with WT cultures (Fig. 7M–7O), matching the PV reduction in prefrontal cortex of α 7-KO mice.

We further examined the maturation of PV-positive basket cells by examining the formation of PV-positive perisomatic GABAergic synapses surrounding the pyramidal neurons in cortical cultures. In WT cultures, the PV- and GAD65-positive perisomatic synapses formed well surrounding the soma and proximal dendrites of pyramidal neurons (Fig. 8A–8C and 8G–8I). In α 7-KO cultures, the formation of PV- and GAD65-positive perisomatic synapses was markedly impaired (Fig. 8D–8F and 8J–8L), and there was a significant reduction of the number of PV-positive perisomatic synapses in α 7-KO cultures compared with WT cultures (Fig. 8M). The findings thus suggest that deletion of the α 7 nAChR impaired the development and the perisomatic synapse formation of cortical PV-positive GABAergic interneurons in cortical cultures.

NMDA receptor expression was reduced in cortical GABAergic interneurons in $\alpha 7$ -KO cultures

As our previous findings demonstrated that NMDA receptor (NMDAR) expression and function was reduced in cortical $\alpha 7$ -KO pyramidal neurons (Lin et al., 2014), we further examined if NMDAR was reduced in cortical GABAergic interneurons by examining NR1 and GABAergic neuronal marker GABA immunoreactivities in cultures. Double immunostaining with α -GABA and α -NR1 antibodies identified GABA-positive interneurons and NR1-positive puncta in the soma and dendrites of cortical GABAergic interneurons in WT and $\alpha 7$ -KO cultures (Fig. 9A–9L, white arrows). The levels of NR1 immunoreactivity in GABAergic neurons were reduced in $\alpha 7$ -KO cultures (Fig. 9D–9F and 9J–9L) compared with WT cultures (Fig. 9A–9C and 9G–9I), suggesting a reduction and hypofunction of NMDAR in cortical GABAergic interneurons.

Discussion

The present study uses both *in vivo* and *in vitro* models to identify multiple deficits of cortical parvalbuminergic/GABAergic circuits from $\alpha 7$ nAChR deletion. The loss of $\alpha 7$ nAChR decreases cortical levels of PV and GABAergic markers, and alters the development of GABAergic synapses and circuits, particularly the development of PV-containing GABAergic interneurons and perisomatic synapses during late postnatal life and adulthood. Moreover, multiple neurochemical deficits in cortical PV-containing GABAergic interneurons with $\alpha 7$ nAChR deletion are reminiscent of characteristic features in schizophrenia. This implicates $\alpha 7$ nAChR-mediated control of cortical PV GABAergic development as crucial components in cortical circuit function and dysfunction in schizophrenia and other neuropsychiatric disorders. The present results thus provide a rationale for the association of $\alpha 7$ nAChR deletion in humans with a spectrum of neurodevelopmental and neuropsychiatric disorders. Relative loss of $\alpha 7$ receptors by genetic or epigenetic regulation could represent a fundamental pathogenic mechanism in such disorders.

In our studies, cortical GAD65/67, PV and GABA_A $\alpha 1$ levels were decreased in $\alpha 7$ -KO mice during postnatal development. GABA-mediated neurotransmission is crucial for shaping the functional maturation of cortical circuits (Le Magueresse and Monyer, 2013). GABA is synthesized by two isoforms of glutamic acid decarboxylase (GAD), GAD65 and GAD67. Both isoforms are present in most GABAergic interneurons, with GAD67 distributed throughout the soma and GAD65 in GABAergic synapses. The development of GAD65/67 expression is thus a reliable indicator of the maturation of cortical GABAergic interneurons' ability to produce GABA. Reduced cortical GAD65/67 levels and decreased spontaneous IPSC frequency suggest deficits in cortical GABA synthesis and maturation of GABAergic circuits with $\alpha 7$ nAChR deletion. As GAD67 is a key regulatory sensor of cortical circuit activity (McNally et al., 2013), reduced GAD67 levels in PV-containing interneurons thereby further imply cortical PV GABAergic dysfunction with $\alpha 7$ nAChR deletion. Moreover, since PV expression is essential for synchronizing GABA release and network gamma oscillatory activities (Volman et al., 2011; McNally et al., 2013), reduced cortical PV levels could thus contribute to abnormal synchronization and oscillations in the cortex

with $\alpha 7$ nAChR deletion. GABA_A $\alpha 1$ -containing receptors are crucial for mediating strong and fast inhibition of PV-positive basket cells onto pyramidal neurons and gamma oscillations (Hajos and Paulsen, 2009; Curley and Lewis, 2012; Lewis et al., 2012). Reduced GABA_A $\alpha 1$ levels in $\alpha 7$ -KO pyramidal neurons, along with reduced PV and GAD67 levels in PV-positive GABAergic interneurons may contribute to the compromised inhibitory control of network activity and could lead to abnormalities in gamma oscillations, working-memory, attention, sensory gating and learning.

Cortical PV levels in PV-positive GABAergic interneurons are specifically affected with $\alpha 7$ nAChR deletion, while the somatostatin immunoreactivity remains unaltered, suggesting that the development of neocortical PV-positive interneurons may be specifically regulated by $\alpha 7$ nAChR. In the developing brain, $\alpha 7$ nAChRs are expressed at high levels during the first postnatal week and reach peak levels at P7 in the prefrontal cortex and hippocampus, when GABAergic signaling changes from excitatory to inhibitory (Adams et al., 2002). $\alpha 7$ nAChR *null* mice indeed exhibited delays in the switch over from GABA-mediated excitation to inhibition during brain development (Liu et al., 2006). In addition, $\alpha 7$ nAChRs colocalize with GABA_A receptors in hippocampal interneurons and are critical in hippocampal gamma oscillations (Song et al., 2005; Le Magueresse et al., 2006; Liu et al., 2006; Bruel-Jungerman et al., 2011; Adams et al., 2012). The PV-positive, fast-spiking basket cells have uniquely fast membrane and synaptic properties that provide strong and fast inhibitory control of pyramidal neurons and gamma oscillations (Hajos and Paulsen, 2009; Curley and Lewis, 2012; Lewis et al., 2012). Furthermore, loss of the $\alpha 7$ nAChR impaired the development of PV-positive GABAergic interneurons and perisomatic synapses in cortical cultures. Indeed, $\alpha 7$ nAChRs have been found in mouse hippocampal interneurons and rat cortical PV GABAergic interneurons (Adams et al., 2001; Murakami et al., 2013). Our findings, along with other studies, thus suggest an intrinsic role of $\alpha 7$ nAChR in mediating cortical PV GABAergic neuronal and circuit development. Moreover, NMDAR expression in cortical GABAergic interneurons was reduced, suggesting NMDAR hypofunction in cortical GABAergic interneurons of $\alpha 7$ -KO mice. Ablation of NMDARs in parvalbumin-positive interneurons of the cerebral cortex results in diminished theta oscillations, impaired spatial representation, and defective memory (Korotkova et al., 2010; Carlen et al., 2012). Similarly, postnatal ablation of NR1 in cortical and hippocampal GABAergic neurons leads to schizophrenia-related symptoms after adolescence, whereas a conditional NR1 knockout in which NMDAR deletion occurring after adolescence does not result in such abnormalities (Belforte et al., 2010). Loss or reduction of $\alpha 7$ nAChRs may lead to such profound glutamatergic hypofunction in PV-positive interneurons through neuregulin 1, NRG1/ErbB4 pathways in schizophrenia, since both $\alpha 7$ nAChRs and NRG1 are schizophrenia risk genes that associate with NMDARs and excitatory synapse formation on PV-positive interneurons (Hahn et al., 2006; Hancock et al., 2008; Ting et al., 2011). Our findings thus implicate NMDAR hypofunction in GABAergic interneurons, which may contribute to cortical dysfunction leading to impaired attention working-memory and learning in $\alpha 7$ nAChR deletion models.

$\alpha 7$ nAChR *null* mice show many neurochemical findings that are analogous to features observed in schizophrenia. First, $\alpha 7$ nAChR levels are reduced in multiple brain regions

including the prefrontal cortex in schizophrenia patients (Freedman et al., 1995; Breese et al., 1997; Freedman et al., 2000; Mathew et al., 2007; Freedman and Goldowitz, 2010; Ross et al., 2010; Olincy and Freedman, 2012). . Though the transient upregulation of another nicotinic receptor subtype $\alpha 4\beta 2$ in $\alpha 7$ nAChR *null* mouse cortex and hippocampus during postnatal development may have some compensatory effects on brain development (Orr-Urtreger et al., 1997; Yu et al., 2007), the changes of $\alpha 4\beta 2$ nAChRs and their association with schizophrenia patients remain less clear and inconsistent in the literatures (Adams and Stevens, 2007). In addition, $\alpha 7$ nAChR selective agonists have demonstrated efficacy in clinical trials of schizophrenia patients and in animal models of positive and negative symptoms and cognitive deficits of schizophrenia (Hajos and Rogers, 2010; Tregellas et al., 2011; Wallace and Porter, 2011; Cannon et al., 2013; Freedman, 2014). Finally, cortical PV GABAergic circuitry is altered in schizophrenia patients in a manner similar to that observed in $\alpha 7$ -KO mice. Deficient cortical GABA synthesis is a conserved feature and loss of GAD67 is prominent in the PV-positive GABAergic interneurons in schizophrenia patients (Fung et al., 2010; Lewis et al., 2012). Postsynaptic GABA_A $\alpha 1$ expression in cortical layer 2/3 pyramidal neurons is reduced in schizophrenia patient dorsolateral prefrontal cortex (Lewis et al., 2012). Furthermore, the level of PV in PV-positive interneurons, but not the number of PV interneurons, is reduced in the prefrontal cortex of schizophrenia patients (Lewis et al., 2012). Thus the neurochemical abnormalities of the cortical PV GABAergic circuit in $\alpha 7$ -KO mice mirror the pathological alterations in schizophrenia patients. Our findings thus demonstrate impaired cortical PV GABAergic development and multiple neurochemical deficits in $\alpha 7$ nAChR gene deletion models that are reminiscent of PV GABAergic deficits in schizophrenia, thus implicating crucial roles of $\alpha 7$ nAChR in cortical PV GABAergic circuit development and dysfunction in schizophrenia and other neuropsychiatric disorders.

$\alpha 7$ nAChR heterozygous and *null* mutants in different mouse strains showed similar GABAergic deficits in the cortex and hippocampus (Adams et al., 2008; Adams et al., 2012). While our findings showed cortical PV GABAergic deficits in the prefrontal cortex of C57BL/6 strain of $\alpha 7$ nAChR *null* mutant mice, we also observed the reduction of PV and GABAergic markers in other cortical areas such as somatosensory cortex of $\alpha 7$ nAChR *null* mice (Supplemental Fig.2). Other lines of investigation have shown decreased levels of GAD65 and GABAA receptors in the hippocampus as well as abnormal hippocampal auditory gating in C3H $\alpha 7$ nAChR heterozygous mutant mice (Adams et al., 2008; Adams et al., 2012). Moreover, both the $\alpha 7$ nAChR null mutant and an inbred mouse strain that has lower expression of $\alpha 7$ nAChRs showed deficits in hippocampal (P20/N40) inhibitory sensory gating in DBA/2 mice (Stevens et al., 1998; Freedman, 2014). The P50 auditory sensory gating deficit has been considered a leading endophenotype in schizophrenia, and polymorphisms in the 5' promoter of *CHRNA7* and in a second gene *CHRFAM7A* with a partial duplication of *CHRNA7* are associated with schizophrenia and the P50 sensory gating deficit (Leonard et al., 2002; Sinkus et al., 2009; Ross et al., 2010; Freedman, 2014). Taken together, loss or reduction of $\alpha 7$ nAChRs may contribute to the totality of deficits expected in GABA-mediated inhibition both in the PV-positive interneurons on prefrontal cortex executive function deficits as well as hippocampal auditory gating deficits in schizophrenia patients.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by CHOP Foerderer Grant and NIH grants R21NS072842, R01 NS45986 and P30HD026979 to Drs. Hong Lin and David R. Lynch. We thank Ms. Margaret Maronski for preparation of cortical neuronal cultures, Dr. Hajime Takano for aid in confocal imaging, and Dr. Jessica Panzer for helpful discussion.

Abbreviations

nAChR	nicotinic acetylcholine receptor
PV	parvalbumin
SST	somatostatin
GAD65/67	Glutamic Acid Decarboxylase 65/67
GABA_Aα1	α1 subunit of GABA _A receptors
VGAT	vesicular GABA transporter
NMDAR	N-methyl-D-aspartate receptor
VGLUT1	vesicular glutamate transporter 1
WT	<i>wild-type</i>
α7-KO	α7 nAChR <i>null</i>

References

- Adams CE, Stevens KE. Evidence for a role of nicotinic acetylcholine receptors in schizophrenia. *Frontiers in bioscience : a journal and virtual library*. 2007; 12:4755–4772. [PubMed: 17485411]
- Adams CE, Stitzel JA, Collins AC, Freedman R. Alpha7-nicotinic receptor expression and the anatomical organization of hippocampal interneurons. *Brain Res*. 2001; 922:180–190. [PubMed: 11743948]
- Adams CE, Yonchek JC, Zheng L, Collins AC, Stevens KE. Altered hippocampal circuit function in C3H alpha7 null mutant heterozygous mice. *Brain Res*. 2008; 1194:138–145. [PubMed: 18199426]
- Adams CE, Broide RS, Chen Y, Winzer-Serhan UH, Henderson TA, Leslie FM, Freedman R. Development of the alpha7 nicotinic cholinergic receptor in rat hippocampal formation. *Brain research Developmental brain research*. 2002; 139:175–187. [PubMed: 12480132]
- Adams CE, Yonchek JC, Schulz KM, Graw SL, Stitzel J, Teschke PU, Stevens KE. Reduced ChRNA7 expression in mice is associated with decreases in hippocampal markers of inhibitory function: implications for neuropsychiatric diseases. *Neuroscience*. 2012; 207:274–282. [PubMed: 22314319]
- Ancin I, Cabranes JA, Santos JL, Sanchez-Morla E, Vazquez-Alvarez B, Rodriguez-Moya L, Pousada-Casal A, Fernandez C, Aparicio A, Barabash A. CHRNA7 haplotypes are associated with impaired attention in euthymic bipolar disorder. *J Affect Disord*. 2011; 133:340–345. [PubMed: 21550667]
- Belforte JE, Zsiros V, Sklar ER, Jiang Z, Yu G, Li Y, Quinlan EM, Nakazawa K. Postnatal NMDA receptor ablation in corticolimbic interneurons confers schizophrenia-like phenotypes. *Nat Neurosci*. 2010; 13:76–83. [PubMed: 19915563]
- Breese CR, Adams C, Logel J, Drebing C, Rollins Y, Barnhart M, Sullivan B, Demasters BK, Freedman R, Leonard S. Comparison of the regional expression of nicotinic acetylcholine receptor alpha7 mRNA and [125I]-alpha-bungarotoxin binding in human postmortem brain. *J Comp Neurol*. 1997; 387:385–398. [PubMed: 9335422]

- Brown KL, Comalli DM, De Biasi M, Woodruff-Pak DS. Trace eyeblink conditioning is impaired in alpha7 but not in beta2 nicotinic acetylcholine receptor knockout mice. *Front Behav Neurosci.* 2010; 4:166. [PubMed: 20976039]
- Bruel-Jungerman E, Lucassen PJ, Francis F. Cholinergic influences on cortical development and adult neurogenesis. *Behav Brain Res.* 2011; 221:379–388. [PubMed: 21272598]
- Cannon CE, Puri V, Vivian JA, Egbertson MS, Eddins D, Uslander JM. The nicotinic alpha7 receptor agonist GTS-21 improves cognitive performance in ketamine impaired rhesus monkeys. *Neuropharmacology.* 2013; 64:191–196. [PubMed: 22659472]
- Carlen M, Meletis K, Siegle JH, Cardin JA, Futai K, Vierling-Claassen D, Ruhlmann C, Jones SR, Deisseroth K, Sheng M, Moore CI, Tsai LH. A critical role for NMDA receptors in parvalbumin interneurons for gamma rhythm induction and behavior. *Mol Psychiatry.* 2012; 17:537–548. [PubMed: 21468034]
- Curley AA, Lewis DA. Cortical basket cell dysfunction in schizophrenia. *J Physiol.* 2012; 590:715–724. [PubMed: 22219337]
- Dani JA, Bertrand D. Nicotinic acetylcholine receptors and nicotinic cholinergic mechanisms of the central nervous system. *Annu Rev Pharmacol Toxicol.* 2007; 47:699–729. [PubMed: 17009926]
- Dong YN, Waxman EA, Lynch DR. Interactions of postsynaptic density-95 and the NMDA receptor 2 subunit control calpain-mediated cleavage of the NMDA receptor. *J Neurosci.* 2004; 24:11035–11045. [PubMed: 15590920]
- Endris V, Hackmann K, Neuhaus TM, Grasshoff U, Bonin M, Haug U, Hahn G, Schallner JC, Schrock E, Tinschert S, Rappold G, Moog U. Homozygous loss of CHRNA7 on chromosome 15q13.3 causes severe encephalopathy with seizures and hypotonia. *Am J Med Genet A.* 2010; 152A:2908–2911. [PubMed: 20979196]
- Fernandes C, Hoyle E, Dempster E, Schalkwyk LC, Collier DA. Performance deficit of alpha7 nicotinic receptor knockout mice in a delayed matching-to-place task suggests a mild impairment of working/episodic-like memory. *Genes Brain Behav.* 2006; 5:433–440. [PubMed: 16923147]
- Freedman R. alpha7-nicotinic acetylcholine receptor agonists for cognitive enhancement in schizophrenia. *Annual review of medicine.* 2014; 65:245–261.
- Freedman R, Leonard S. Genetic linkage to schizophrenia at chromosome 15q14. *American journal of medical genetics.* 2001; 105:655–657. [PubMed: 11803510]
- Freedman R, Goldowitz D. Studies on the hippocampal formation: From basic development to clinical applications: Studies on schizophrenia. *Prog Neurobiol.* 2010; 90:263–275. [PubMed: 19853005]
- Freedman R, Adams CE, Leonard S. The alpha7-nicotinic acetylcholine receptor and the pathology of hippocampal interneurons in schizophrenia. *Journal of chemical neuroanatomy.* 2000; 20:299–306. [PubMed: 11207427]
- Freedman R, Hall M, Adler LE, Leonard S. Evidence in postmortem brain tissue for decreased numbers of hippocampal nicotinic receptors in schizophrenia. *Biol Psychiatry.* 1995; 38:22–33. [PubMed: 7548469]
- Freedman R, et al. Linkage of a neurophysiological deficit in schizophrenia to a chromosome 15 locus. *Proceedings of the National Academy of Sciences of the United States of America.* 1997; 94:587–592. [PubMed: 9012828]
- Fung SJ, Webster MJ, Sivagnanasundaram S, Duncan C, Elashoff M, Weickert CS. Expression of interneuron markers in the dorsolateral prefrontal cortex of the developing human and in schizophrenia. *Am J Psychiatry.* 2010; 167:1479–1488. [PubMed: 21041246]
- Gonzalez-Burgos G, Fish KN, Lewis DA. GABA neuron alterations, cortical circuit dysfunction and cognitive deficits in schizophrenia. *Neural Plast.* 2011; 2011:723184. [PubMed: 21904685]
- Griguoli M, Cherubini E. Regulation of hippocampal inhibitory circuits by nicotinic acetylcholine receptors. *J Physiol.* 2012; 590:655–666. [PubMed: 22124144]
- Hahn CG, Wang HY, Cho DS, Talbot K, Gur RE, Berrettini WH, Bakshi K, Kamins J, Borgmann-Winter KE, Siegel SJ, Gallop RJ, Arnold SE. Altered neuregulin 1-erbB4 signaling contributes to NMDA receptor hypofunction in schizophrenia. *Nature medicine.* 2006; 12:824–828.
- Hajos M, Rogers BN. Targeting alpha7 nicotinic acetylcholine receptors in the treatment of schizophrenia. *Curr Pharm Des.* 2010; 16:538–554. [PubMed: 19909231]

- Hajos N, Paulsen O. Network mechanisms of gamma oscillations in the CA3 region of the hippocampus. *Neural networks : the official journal of the International Neural Network Society*. 2009; 22:1113–1119. [PubMed: 19683412]
- Hancock ML, Canetta SE, Role LW, Talmage DA. Presynaptic type III neuregulin1-ErbB signaling targets {alpha}7 nicotinic acetylcholine receptors to axons. *The Journal of cell biology*. 2008; 181:511–521. [PubMed: 18458158]
- Helbig I, et al. 15q13.3 microdeletions increase risk of idiopathic generalized epilepsy. *Nat Genet*. 2009; 41:160–162. [PubMed: 19136953]
- Hellier JL, Arevalo NL, Smith L, Xiong KN, Restrepo D. alpha7-Nicotinic acetylcholine receptor: role in early odor learning preference in mice. *PLoS One*. 2012; 7:e35251. [PubMed: 22514723]
- Hoftman GD, Lewis DA. Postnatal developmental trajectories of neural circuits in the primate prefrontal cortex: identifying sensitive periods for vulnerability to schizophrenia. *Schizophr Bull*. 2011; 37:493–503. [PubMed: 21505116]
- Hoppman-Chaney N, Wain K, Seger P, Superneau D, Hodge J. Identification of single gene deletions at 15q13.3: further evidence that CHRNA7 causes the 15q13.3 microdeletion syndrome phenotype. *Clin Genet*. 2012
- Inan M, Petros TJ, Anderson SA. Losing your inhibition: linking cortical GABAergic interneurons to schizophrenia. *Neurobiol Dis*. 2013; 53:36–48. [PubMed: 23201207]
- Jiang Z, Cowell RM, Nakazawa K. Convergence of genetic and environmental factors on parvalbumin-positive interneurons in schizophrenia. *Front Behav Neurosci*. 2013; 7:116. [PubMed: 24027504]
- Korotkova T, Fuchs EC, Ponomarenko A, von Engelhardt J, Monyer H. NMDA receptor ablation on parvalbumin-positive interneurons impairs hippocampal synchrony, spatial representations, and working memory. *Neuron*. 2010; 68:557–569. [PubMed: 21040854]
- Le Magueresse C, Monyer H. GABAergic interneurons shape the functional maturation of the cortex. *Neuron*. 2013; 77:388–405. [PubMed: 23395369]
- Le Magueresse C, Safiulina V, Changeux JP, Cherubini E. Nicotinic modulation of network and synaptic transmission in the immature hippocampus investigated with genetically modified mice. *J Physiol*. 2006; 576:533–546. [PubMed: 16901939]
- Leonard S, Adams C, Breese CR, Adler LE, Bickford P, Byerley W, Coon H, Griffith JM, Miller C, Myles-Worsley M, Nagamoto HT, Rollins Y, Stevens KE, Waldo M, Freedman R. Nicotinic receptor function in schizophrenia. *Schizophr Bull*. 1996; 22:431–445. [PubMed: 8873294]
- Leonard S, Gault J, Hopkins J, Logel J, Vianzon R, Short M, Drebing C, Berger R, Venn D, Sirota P, Zerbe G, Olincy A, Ross RG, Adler LE, Freedman R. Association of promoter variants in the alpha7 nicotinic acetylcholine receptor subunit gene with an inhibitory deficit found in schizophrenia. *Arch Gen Psychiatry*. 2002; 59:1085–1096. [PubMed: 12470124]
- Lepichon JB, Bittel DC, Graf WD, Yu S. A 15q13.3 homozygous microdeletion associated with a severe neurodevelopmental disorder suggests putative functions of the TRPM1, CHRNA7, and other homozygously deleted genes. *Am J Med Genet A*. 2010; 152A:1300–1304. [PubMed: 20425840]
- Levin ED, Petro A, Rezvani AH, Pollard N, Christopher NC, Strauss M, Avery J, Nicholson J, Rose JE. Nicotinic alpha7- or beta2-containing receptor knockout: effects on radial-arm maze learning and long-term nicotine consumption in mice. *Behav Brain Res*. 2009; 196:207–213. [PubMed: 18831991]
- Lewis DA, Curley AA, Glausier JR, Volk DW. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci*. 2012; 35:57–67. [PubMed: 22154068]
- Liao J, DeWard SJ, Madan-Khetarpal S, Surti U, Hu J. A small homozygous microdeletion of 15q13.3 including the CHRNA7 gene in a girl with a spectrum of severe neurodevelopmental features. *Am J Med Genet A*. 2011; 155A:2795–2800. [PubMed: 21990074]
- Lin H, Hsu FC, Baumann BH, Coulter DA, Lynch DR. Cortical synaptic NMDA receptor deficits in alpha7 nicotinic acetylcholine receptor gene deletion models: implications for neuropsychiatric diseases. *Neurobiol Dis*. 2014; 63:129–140. [PubMed: 24326163]
- Lin H, Vicini S, Hsu FC, Doshi S, Takano H, Coulter DA, Lynch DR. Axonal alpha7 nicotinic ACh receptors modulate presynaptic NMDA receptor expression and structural plasticity of

- glutamatergic presynaptic boutons. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107:16661–16666. [PubMed: 20817852]
- Liu Z, Neff RA, Berg DK. Sequential interplay of nicotinic and GABAergic signaling guides neuronal development. *Science*. 2006; 314:1610–1613. [PubMed: 17158331]
- Lozada AF, Wang X, Goukko NV, Massey KA, Duan J, Liu Z, Berg DK. Glutamatergic synapse formation is promoted by alpha7-containing nicotinic acetylcholine receptors. *J Neurosci*. 2012; 32:7651–7661. [PubMed: 22649244]
- Mathew SV, Law AJ, Lipska BK, Davila-Garcia MI, Zamora ED, Mitkus SN, Vakkalanka R, Straub RE, Weinberger DR, Kleinman JE, Hyde TM. Alpha7 nicotinic acetylcholine receptor mRNA expression and binding in postmortem human brain are associated with genetic variation in neuregulin 1. *Hum Mol Genet*. 2007; 16:2921–2932. [PubMed: 17884806]
- McNally JM, McCarley RW, Brown RE. Impaired GABAergic neurotransmission in schizophrenia underlies impairments in cortical gamma band oscillations. *Current psychiatry reports*. 2013; 15:346. [PubMed: 23400808]
- Miller DT, et al. Microdeletion/duplication at 15q13.2q13.3 among individuals with features of autism and other neuropsychiatric disorders. *J Med Genet*. 2009; 46:242–248. [PubMed: 18805830]
- Miwa JM, Freedman R, Lester HA. Neural systems governed by nicotinic acetylcholine receptors: emerging hypotheses. *Neuron*. 2011; 70:20–33. [PubMed: 21482353]
- Murakami K, Ishikawa Y, Sato F. Localization of alpha7 nicotinic acetylcholine receptor immunoreactivity on GABAergic interneurons in layers I–III of the rat retrosplenial granular cortex. *Neuroscience*. 2013; 252:443–459. [PubMed: 23985568]
- Nakazawa K, Zsiros V, Jiang Z, Nakao K, Kolata S, Zhang S, Belforte JE. GABAergic interneuron origin of schizophrenia pathophysiology. *Neuropharmacology*. 2012; 62:1574–1583. [PubMed: 21277876]
- Olinic A, Freedman R. Nicotinic mechanisms in the treatment of psychotic disorders: a focus on the alpha7 nicotinic receptor. *Handb Exp Pharmacol*. 2012:211–232. [PubMed: 23027417]
- Orr-Urtreger A, Goldner FM, Saeki M, Lorenzo I, Goldberg L, De Biasi M, Dani JA, Patrick JW, Beaudet AL. Mice deficient in the alpha7 neuronal nicotinic acetylcholine receptor lack alpha-bungarotoxin binding sites and hippocampal fast nicotinic currents. *J Neurosci*. 1997; 17:9165–9171. [PubMed: 9364063]
- Ross RG, Stevens KE, Proctor WR, Leonard S, Kisley MA, Hunter SK, Freedman R, Adams CE. Research review: Cholinergic mechanisms, early brain development, and risk for schizophrenia. *Journal of child psychology and psychiatry, and allied disciplines*. 2010; 51:535–549.
- Rotaru DC, Yoshino H, Lewis DA, Ermentrout GB, Gonzalez-Burgos G. Glutamate receptor subtypes mediating synaptic activation of prefrontal cortex neurons: relevance for schizophrenia. *J Neurosci*. 2011; 31:142–156. [PubMed: 21209199]
- Sharp AJ, et al. A recurrent 15q13.3 microdeletion syndrome associated with mental retardation and seizures. *Nat Genet*. 2008; 40:322–328. [PubMed: 18278044]
- Shinawi M, Schaaf CP, Bhatt SS, Xia Z, Patel A, Cheung SW, Lanpher B, Nagl S, Herding HS, Nevinsky-Stickel C, Immken LL, Patel GS, German JR, Beaudet AL, Stankiewicz P. A small recurrent deletion within 15q13.3 is associated with a range of neurodevelopmental phenotypes. *Nat Genet*. 2009; 41:1269–1271. [PubMed: 19898479]
- Sinkus ML, Lee MJ, Gault J, Logel J, Short M, Freedman R, Christian SL, Lyon J, Leonard S. A 2-base pair deletion polymorphism in the partial duplication of the alpha7 nicotinic acetylcholine gene (CHRFAM7A) on chromosome 15q14 is associated with schizophrenia. *Brain Res*. 2009; 1291:1–11. [PubMed: 19631623]
- Song C, Murray TA, Kimura R, Wakui M, Ellsworth K, Javedan SP, Marxer-Miller S, Lukas RJ, Wu J. Role of alpha7-nicotinic acetylcholine receptors in tetanic stimulation-induced gamma oscillations in rat hippocampal slices. *Neuropharmacology*. 2005; 48:869–880. [PubMed: 15829257]
- Spielmann M, Reichelt G, Hertzberg C, Trimborn M, Mundlos S, Horn D, Klopocki E. Homozygous deletion of chromosome 15q13.3 including CHRNA7 causes severe mental retardation, seizures, muscular hypotonia, and the loss of KLF13 and TRPM1 potentially cause macrocytosis and

- congenital retinal dysfunction in siblings. *Eur J Med Genet.* 2011; 54:e441–e445. [PubMed: 21596161]
- Stevens KE, Kem WR, Mahnir VM, Freedman R. Selective alpha7-nicotinic agonists normalize inhibition of auditory response in DBA mice. *Psychopharmacology (Berl).* 1998; 136:320–327. [PubMed: 9600576]
- Ting AK, Chen Y, Wen L, Yin DM, Shen C, Tao Y, Liu X, Xiong WC, Mei L. Neuregulin 1 promotes excitatory synapse development and function in GABAergic interneurons. *J Neurosci.* 2011; 31:15–25. [PubMed: 21209185]
- Tregellas JR, Tanabe J, Rojas DC, Shatti S, Olincy A, Johnson L, Martin LF, Soti F, Kem WR, Leonard S, Freedman R. Effects of an alpha 7-nicotinic agonist on default network activity in schizophrenia. *Biol Psychiatry.* 2011; 69:7–11. [PubMed: 20728875]
- Volman V, Behrens MM, Sejnowski TJ. Downregulation of parvalbumin at cortical GABA synapses reduces network gamma oscillatory activity. *J Neurosci.* 2011; 31:18137–18148. [PubMed: 22159125]
- Wallace TL, Porter RH. Targeting the nicotinic alpha7 acetylcholine receptor to enhance cognition in disease. *Biochem Pharmacol.* 2011; 82:891–903. [PubMed: 21741954]
- Wallace TL, Bertrand D. Importance of the nicotinic acetylcholine receptor system in the prefrontal cortex. *Biochem Pharmacol.* 2013; 85:1713–1720. [PubMed: 23628449]
- Yasui DH, Scoles HA, Horike S, Meguro-Horike M, Dunaway KW, Schroeder DI, Lasalle JM. 15q11.2-13.3 chromatin analysis reveals epigenetic regulation of CHRNA7 with deficiencies in Rett and autism brain. *Hum Mol Genet.* 2011; 20:4311–4323. [PubMed: 21840925]
- Young JW, Meves JM, Tarantino IS, Caldwell S, Geyer MA. Delayed procedural learning in alpha7-nicotinic acetylcholine receptor knockout mice. *Genes Brain Behav.* 2011; 10:720–733. [PubMed: 21679297]
- Young JW, Crawford N, Kelly JS, Kerr LE, Marston HM, Spratt C, Finlayson K, Sharkey J. Impaired attention is central to the cognitive deficits observed in alpha 7 deficient mice. *Eur Neuropsychopharmacol.* 2007; 17:145–155. [PubMed: 16650968]
- Yu WF, Guan ZZ, Nordberg A. Postnatal upregulation of alpha4 and alpha3 nicotinic receptor subunits in the brain of alpha7 nicotinic receptor-deficient mice. *Neuroscience.* 2007; 146:1618–1628. [PubMed: 17434683]
- Zhang L, Warren RA. Muscarinic and nicotinic presynaptic modulation of EPSCs in the nucleus accumbens during postnatal development. *J Neurophysiol.* 2002; 88:3315–3330. [PubMed: 12466449]

Figure 1. Cortical levels of GABAergic markers were decreased in $\alpha 7$ nAChR null mice during late postnatal development and adulthood

Western blotting analysis of whole cortex homogenates (30 μ g per lane) showing PV (A), GAD65/67 (B), GABA α 1 (C) expression in the cortex of *wild-type* (WT) and $\alpha 7$ nAChR *null* (KO) mice at postnatal days P1, P4, P8, P14, P21, P56 and P9M (>9 months old) (* p <0.05 ** p <0.01 KO vs. WT; n =6; t -test).

Figure 2. The levels of PV, GAD65/67 and GABA_Aα1 were decreased in the prefrontal cortex of α7 nAChR *null* mice during late postnatal development and adulthood

A–F. Confocal images of PV (green), GAD65/67 (red) immunofluorescence and merged images with DAPI-stained nucleus showing a reduction in the overall level of PV and GAD65/67 immunoreactivities in the prefrontal cortex of α7 nAChR *null* (α7-KO) mice (**D–F**) compared with WT littermates at P56-P90 (**A–C**). **G–L.** Confocal images of PV (green), GABA_Aα1 (red) immunofluorescence and merged images with DAPI-stained nucleus showing a reduction in the overall level of PV and GABA_Aα1 immunoreactivities in the prefrontal cortex of α7-KO mice (**J–L**) compared with WT littermates at P56-P90. Scale bars as indicated.

Figure 3. The levels of PV, but not somatostatin, immunoreactivities were decreased in the prefrontal cortex of $\alpha 7$ nAChR null mice during late postnatal development and adulthood
A–L. Confocal images of PV (green), somatostatin (red) immunofluorescence and merged images with DAPI-stained nucleus showing a reduction in the overall level of PV, but not somatostatin, immunoreactivities in the prefrontal cortex of $\alpha 7$ -KO mice (**D–F, J–L**) compared with WT littermates (**A–C, G–I**) at P56-P90. **M.** Quantification analysis showing reduced PV levels in the soma of PV-positive GABAergic interneurons of $\alpha 7$ -KO mouse prefrontal cortex compared with WT littermates. (** $p < 0.001$ KO vs. WT; 6–8 sections per animal, 3 animals each group; t -test). **N.** Quantification analysis showing similar SST levels in the soma of SST-positive GABAergic interneurons of $\alpha 7$ -KO mouse prefrontal cortex compared with WT littermates. ($p > 0.05$ KO vs. WT; 6–8 sections per animal, 3 animals each group; t -test). **O.** Quantification analysis showing that the number of PV-positive GABAergic interneurons in the prefrontal cortex was not significantly reduced in $\alpha 7$ -KO mice compared with WT littermates at P56-P90 ($p > 0.05$; 7–8 sections per animal, 3 animals per group; t -test). Scale bars as indicated.

Figure 4. PV and GAD67 levels in PV-positive GABAergic interneurons as well as GABA_Aα1 levels in pyramidal neurons were reduced in mouse prefrontal cortex with α7 nAChR deletion during late postnatal development and adulthood

A–F. Confocal images of PV (green), GAD65/67 (red) immunofluorescence and merged images with DAPI-stained nucleus showing reduced PV and GAD67 levels in PV-positive GABAergic neurons (white arrows) and reduced GAD65-positive GABAergic synapses surrounding pyramidal neurons (grey arrows) in the prefrontal cortex of α7-KO mice (**D–F**) compared with WT littermates at P56-P90 (**A–C**). **G–L.** Confocal images of PV (green), GABA_Aα1 (red) immunofluorescence and merged images with DAPI-stained nucleus showing reduced GABA_Aα1 levels in pyramidal neurons (**J–L**, grey arrows) and in PV-positive GABAergic interneurons (**J–L**, white arrows) in the prefrontal cortex of α7-KO mice compared with WT littermates at P56-P90 (**G–I**). Scale bars as indicated.

Figure 5. GABAergic synapse formation was impaired in the prefrontal cortex of $\alpha 7$ nAChR null mice during late postnatal development and adulthood

A–F. Confocal images of PV (green), VGAT (red) immunofluorescence and merged images with DAPI-stained nucleus showing a reduction in the overall levels of PV and VGAT in the prefrontal cortex of $\alpha 7$ -KO mice (**D–F**) compared with WT littermates at P56-P90 (**A–C**).

G–L. The PV levels in PV-positive GABAergic neurons (white arrows) and VGAT-positive GABAergic presynaptic terminals surrounding pyramidal neurons (grey arrows) were reduced in the prefrontal cortex of $\alpha 7$ -KO mice (**J–L**) compared with WT littermates at P56-P90 (**G–I**). **M.** Quantification of the number of VGAT-positive terminals per 1000 μm^2 showing a significant decrease in the number of GABAergic presynaptic terminals in the prefrontal cortex of $\alpha 7$ KO mice compared with WT littermates at P56-P90 (** $p < 0.001$ KO vs. WT; 4–5 sections per animal, 3 animals per group; t -test). Scale bars as indicated.

Figure 6. GABAergic synapse formation and neurotransmission were reduced in α7-KO cortical cultures

A–F. Confocal images of GAD65 (green), GABAAα1 (red) immunofluorescence and merged images showing reduced number of GABAergic synapses and levels of GABAAα1 in α7-KO cortical cultures (**D–F**) compared with WT cultures (**A–C**). High magnification of confocal images showing that 100% of GAD65-positive puncta (**A, D**) were colocalized with or adjacent to one or more GABAAα1 puncta (**B, E**) on the dendrites of cortical neurons in WT and α7-KO cultures (**C, F**). **G, H.** Confocal images of VGAT (red) immunofluorescence showing reduced GABAergic synapses in α7-KO cortical cultures (**H**) compared with WT cultures (**G**). **I.** Quantification of the number of VGAT-positive puncta per 10μm length dendrite of cortical neurons showing a significant decrease in the number of GABAergic synapses on the dendrites of pyramidal neurons in α7 KO cultures compared with WT cultures. (***) $p < 0.001$ KO vs. WT; 5–6 neurons per culture, 4 different cultures each group; t -test). **J–M.** Spontaneous IPSC (sIPSC) properties for WT and α7-KO cultured cortical pyramidal neurons (21–28 DIV). Cells were voltage-clamped at -70 mV and whole

cell recording of sIPSCs was performed. **J.** Representative traces for sIPSC for both WT and $\alpha 7$ -KO cultured cortical pyramidal neurons. sIPSC analysis demonstrated that the sIPSC frequency was decreased (**K**), current decay time constant was faster (**L**), but no change in amplitude (**M**) in $\alpha 7$ -KO cultures compared with those in WT cultures. Box-Whisker graphs were plotted due to the non-parametric data distribution. The data were from 4–5 independent experiments. (** $p < 0.01$ and *** $p < 0.001$ vs. WT; $n = 25$; Mann-Whitney Rank Sum test). Scale bars as indicated.

Figure 7. The development and maturation of PV-positive basket cells were impaired in $\alpha 7$ -KO cortical cultures

A–L. Confocal images of PV (red) and GAD65 (green) immunofluorescence and merged images showing PV immunoreactivities in the soma, dendrites, axons and synapses of PV-positive basket cells in WT and $\alpha 7$ -KO cultures (A, D, G and J). The axons of an individual PV-positive basket cell formed perisomatic GABAergic synapses surrounding the multiple pyramidal neurons (A–L, white arrows). The development of PV-positive basket cells and perisomatic GABAergic synapses was impaired in $\alpha 7$ -KO cortical cultures (D–F and J–L)

compared with WT cultures (**A–C** and **G–I**). **M, N**. Confocal images of PV (red) immunofluorescence showing PV immunoreactivity in PV-positive GABAergic interneurons in WT (**M**) and $\alpha 7$ -KO (**N**) cortical cultures. **O**. Quantification analysis showing reduced PV levels in the soma of PV-positive GABAergic interneurons in $\alpha 7$ -KO cultures compared with WT cultures. (** $p < 0.001$ KO *vs.* WT; 5–6 neurons per culture, 4 different cultures; *t*-test). Scale bars as indicated.

Figure 8. The PV-positive perisomatic GABAergic synapses were reduced in $\alpha 7$ -KO cortical cultures

A–L. Confocal images of PV (red) and GAD65 (green) immunofluorescence and merged images showing the reduced PV- and GAD65-positive perisomatic synapses of PV-positive basket cells surrounding the soma and proximal dendrites of pyramidal neurons in $\alpha 7$ -KO cultures (**D–F** and **J–L**) compared with WT cultures (**A–C** and **G–I**). **M.** Quantification of the number of PV-positive perisomatic GABAergic synapses surrounding the soma and proximal dendrites of pyramidal neurons showed significant reduction of PV-positive perisomatic synapses in $\alpha 7$ -KO cultures compared with WT cultures (*** $p < 0.001$ KO vs. WT; 5–6 neurons per culture, 4 different cultures; t -test). Scale bars as indicated.

Figure 9. Reduction of NMDAR in GABAergic interneurons in $\alpha 7$ -KO cortical cultures compared with WT cultures

A–F, Confocal imaging of NR1 (green) and GABA (red) immunofluorescence showing NR1 reduction in the soma and dendrites of GABA-positive interneurons (white arrows) in $\alpha 7$ -KO cortical cultures (**D–F**) compared with WT cultures (**A–C**). **G–L**, High magnification of confocal imaging of NR1 (green) and GABA (red) immunofluorescence showing reduced NR1 puncta in the soma and dendrites of GABAergic interneurons in $\alpha 7$ -KO cortical cultures (**J–L**) compared with WT cultures (**G–I**). Scale bar as indicated.