Published in final edited form as:

J Neurochem. 2014 July ; 130(2): 313–323. doi:10.1111/jnc.12720.

Evidence for an Angiotensin-(1–7) Neuropeptidase Expressed in the Brain Medulla and CSF of Sheep

Allyson C. Marshall¹, Nancy T. Pirro¹, James C. Rose², Debra I. Diz¹, and Mark C. Chappell¹

¹Hypertension and Vascular Research Center, Wake Forest School of Medicine, Winston Salem, NC

²Department of Obstetrics and Gynecology, Wake Forest School of Medicine, Winston Salem, NC

Abstract

Angiotensin-(1–7) [Ang-(1–7)] is an alternative product of the brain renin-angiotensin system (RAS) that exhibits central actions to lower blood pressure and improve baroreflex sensitivity. We previously identified a peptidase that metabolizes Ang-(1–7) to the inactive metabolite product Ang-(1–4) in CSF of adult sheep. The current study purified the peptidase 1445-fold from sheep brain medulla and characterized this activity. The peptidase was sensitive to the chelating agents o-phenanthroline and EDTA, as well as the mercury compound p-chloromercuribenzoic acid (PCMB). Selective inhibitors to angiotensin-converting enzyme, neprilysin, neurolysin, and thimet oligopeptidase did not attenuate activity; however, the metallopeptidase agent JMV-390 was a potent inhibitor of Ang-(1–7) hydrolysis ($K_i = 0.8$ nM). Kinetic studies using ¹²⁵I-labeled Ang-(1–7), Ang II, and Ang I revealed comparable apparent K_m values (2.6, 2.8 and 4.3 μ M, respectively), but a higher apparent V_{max} for Ang-(1–7) (72 vs. 30 and 6 nmol/min/mg, respectively; $P < 0.01$). HPLC analysis of the activity confirmed the processing of unlabeled Ang-(1–7) to Ang-(1–4) by the peptidase, but revealed $< 5\%$ hydrolysis of Ang II or Ang I, and no hydrolysis of neurotensin, bradykinin or apelin-13. The unique characteristics of the purified neuropeptidase may portend a novel pathway to influence actions of Ang-(1–7) within the brain.

Keywords

Renin-angiotensin system; Ang-(1–7); neuropeptidase; metabolism

INTRODUCTION

Angiotensin-(1–7) [Ang-(1–7)] is a bioactive hormone of the renin-angiotensin system (RAS) that we originally identified in the brain and circulation of the rat twenty-five years ago (Chappell *et al.* 1989). Since that time, accumulating scientific evidence suggests that Ang-(1–7) opposes the actions of Ang II and may constitute an endogenous buffering

Address for Correspondence: Mark C. Chappell, Professor of Surgical Sciences – Hypertension Center, Hanes Building 6th Floor, Medical Center Blvd, Winston Salem, NC, 27157, Telephone: 336-716-9236, Fax: 336-716-2456, mchappel@wakehealth.edu.

CONFLICT OF INTEREST DISCLOSURE

No conflicts of interest, financial or otherwise, are declared by the authors.

system to the ACE-Ang II-Ang II type 1 (AT₁) receptor axis in the regulation of autonomic function, blood pressure, and blood pressure-independent metabolic pathologies (Chappell 2007, Xu *et al.* 2011, Chappell *et al.* 2014). Indeed, administration of the Ang-(1–7) antagonist D-Ala⁷-Ang-(1–7) (A779) decreases baroreflex sensitivity (Kar *et al.* 2011, Arnold *et al.* 2011) and increases inflammation (Capettini *et al.* 2012), implying a significant degree of endogenous Ang-(1–7) tone in the brain. Moreover, the Ang-(1–7) receptor antagonist reduces the actions of either angiotensin-converting enzyme (ACE, EC 3.4.15.1) or AT₁ receptor inhibition, further suggesting a key role of the peptide in the therapeutic benefits of RAS blockade (Chappell 2007).

Ang II is primarily formed through the direct processing of Ang I by the dipeptidyl carboxypeptidase ACE, although other peptidases including chymase (EC 3.4.21.39) and cathepsin G (EC 3.4.21.20) may contribute under pathological conditions (Sadjadi *et al.* 2005, Rykl *et al.* 2006). In contrast, Ang-(1–7) can be processed directly from Ang I by several endopeptidase including neprilysin (EC 3.4.24.11), thimet oligopeptidase (TOP, EC 3.4.24.15), and prolyl endopeptidase (PE, EC 3.4.21.26) (Campbell *et al.* 1998, Allred *et al.* 2000, Alzayadneh & Chappell 2013, Welches *et al.* 1991). Additionally, Ang-(1–7) is efficiency generated directly from Ang II by the mono-carboxypeptidases ACE2 (EC 3.4.15.1) and prolyl carboxypeptidase (EC 3.4.16.2) (Pereira *et al.* 2013, Grobe *et al.* 2013). Although ACE constitutes the major Ang II-forming pathway, the peptidase also degrades a number of biological active peptides including Ang-(1–7) (Chappell *et al.* 1998). The impact of ACE inhibitors to increase circulating levels of Ang-(1–7) reflects the marked reduction in Ang-(1–7) metabolism to Ang-(1–5), as well as the alternative processing of Ang I by the metallopeptidase neprilysin (Chappell *et al.* 1998, Yamada *et al.* 1998, Campbell *et al.* 1998). We recently observed that ACE activity was higher in CSF from adult offspring of betamethasone-exposed pregnant ewes, a model of glucocorticoid-induced fetal programming characterized by higher blood pressure and reduced baroreflex function (Marshall *et al.* 2013b). Although the hydrolysis of Ang-(1–7) in sheep CSF *ex vivo* was attenuated by ACE inhibition, the majority of the peptide-degrading activity in the CSF was due to a thiol-sensitive endopeptidase that cleaved Ang-(1–7) to Ang-(1–4) (Marshall *et al.* 2013b). The Ang-(1–7) peptidase activity was significantly higher in the CSF of the betamethasone-exposed group and the CSF content of Ang-(1–7) was inversely correlated to peptidase activity (Marshall *et al.* 2013b). Selective inhibitors against the endopeptidases neprilysin, TOP, and neurolysin (EC 3.4.24.16) did not attenuate the hydrolysis of Ang-(1–7) to Ang-(1–4) in the CSF possibly suggesting a unique Ang-(1–7)-degrading activity in brain. To address this possibility, the present study sought to obtain a sufficient amount of the purified activity from brain medullary tissue to achieve a more complete characterization of the peptidase. A 1445-fold enrichment of the peptidase was achieved from the brain medulla of sheep and the purified activity evaluated against angiotensins and other neuropeptide substrates. We report that the medullary peptidase appears similar to the activity in the CSF to metabolize Ang-(1–7) to Ang-(1–4), exhibits marked sensitivity to mercury-based inhibitors, chelating agents, and metalloendopeptidase agent JMV-390, and hydrolyzes Ang-(1–7) to a greater extent than Ang II and Ang I, while other bioactive peptides including bradykinin, neurotensin and apelin-13 were not cleaved.

METHODS

Animals

Mixed breed sheep (obtained from a private local vendor) were delivered at term, farm raised, and weaned at 3-months of age. At 10–12-months of age, male offspring were brought to our Association for Assessment and Accreditation of Laboratory Animals Care (AAALAC) approved facility, where they were maintained on a normal diet with free access to tap water and a 12-hour light/dark cycle (lights on 7 AM to 7 PM). Sheep were anesthetized with ketamine and isoflurane and euthanized by exsanguination. Brain medullae were removed and immediately covered with optimal cryosection media and stored at -80°C . CSF (~3 ml per animal) was extracted, taking care to avoid contamination with blood, and tubes were stored at -80°C . All procedures were approved by the Wake Forest University School of Medicine ACUC for animal care.

Homogenization of Sheep Brain

Brain medullae were cut 4 mm rostral and 2 mm caudal to the obex and divided in half along the dorsoventral axis to isolate the dorsal medulla including the NTS. The dorsal medullae from two animals were pooled (2.5 g) for each purification and homogenized in HEPES buffer (25 mM Na^{+} free HEPES, 10 μM ZnCl_2 , 0.05% Triton, pH 7.0) using a Power Gen 1000 tissue grinder (Fisher Scientific) on setting 5 for 60 sec and centrifuged at $25,000\times g$ for 30 min at 4°C . Supernatant was retained for the subsequent purification steps.

Cerebrospinal Fluid Concentration

CSF (5 ml) was pooled and concentrated from five animals 1:5 using molecular weight filtration tubes to remove small proteins and endogenous angiotensin peptides (30 kDa, Millipore Bedford, MA). Concentrated CSF was resuspended in a final volume of 5 ml HEPES buffer (25 mM HEPES, 125 mM NaCl , 10 μM ZnCl_2 , pH = 7.4) and protein concentration was measured using a Bradford protein assay.

DEAE-Cellulose Chromatography

Diethylaminoethyl Sepharose (DEAE, Sigma-Aldrich, St Louis MO) was equilibrated with HEPES buffer (25 mM Na^{+} free HEPES, 10 μM ZnCl_2 , 0.05% Triton, pH 7.0) and incubated with the medullary supernatant for 60 min at room temperature. The DEAE gel was pelleted at $1,000\times g$ for 60 sec, and supernatant was removed. The gel was subsequently washed in HEPES buffer with an increasing step gradient of NaCl (100, 250, 500 mM, 1M for 30 mins) and the gel pelleted at $1,000\times g$. The eluted material from the gradient was assayed for protein content and enzyme activity.

Cibacron Blue Chromatography

Fractions containing the highest activity from the DEAE gel were combined (250 and 500 mM NaCl), concentrated on 50 kDa molecular weight concentrations tubes (Millipore Bedford, MA), washed with Na^{+} free HEPES buffer, and incubated with equilibrated Cibacron Blue Sepharose Fast Flow (GE Healthcare Life Sciences) for 60 min. The flow-

through fractions from the Cibacron gel contained the majority of activity and were applied directly to the Q-Sepharose column.

Q-Sepharose Chromatography

Q-Sepharose Fast Flow (Sigma-Aldrich, St Louis MO) was equilibrated with Na-free HEPES buffer in a 2.5×10 cm column. The unbound Cibacron fraction was applied directly to the Q-Sepharose column and washed with a step-gradient of increasing NaCl (0, 100, 200, 400, 600 mM and 1 M NaCl). Fractions were collected and assayed for protein content and enzyme activity.

Gel Electrophoresis

Purified protein eluted from the Q-Sepharose column was diluted 1:1 in Laemmli buffer containing β -mercaptoethanol and boiled for 5 mins to induce denaturing conditions. Proteins were separated on 10% Mini-PROTEAN TGX gels for 80 min at 120 V in Tris-glycine buffer. Gels were stained with ImperialTM Blue Stain (Thermo Fisher Scientific, Rockford, IL) or silver stain (PlusOneTM Silver Stain Kit, GE Healthcare, Sweden) according to manufacturer's instructions.

Inhibitor and pH Profile

Using aliquots of partially purified peptidase, various inhibitors were tested for their ability to inhibit ^{125}I -Ang-(1–7) metabolism including PCMB, E-64, EDTA, o-phenanthroline, SCH 39370 (SCH), 1-carboxy-3-phenyl-propyl (Ala-Ala-Phe-4-Abz-OH, CFP), Z-prolyl prolinal (ZPP), dithiothreitol (DTT), the dipeptide Pro-Ile, phenylmethylsulfonyl fluoride (PMSF), aprotinin, soybean trypsin inhibitor (SBTI), and the metallopeptidase inhibitor *N*-[3-[(hydroxyamino)carbonyl]-1-oxo-2(*R*)-benzylpropyl]-L-leucine (JMV-390). Each reaction was conducted in the presence of 0.5 nM ^{125}I -Ang-(1–7), 100 nM Ang-(1–7), 92 ng of the Q-Sepharose fraction or 25 μl of CSF, and a cocktail of inhibitors (metabolism cocktail) containing the aminopeptidase inhibitors amastatin (AM, 2 μM) and bestatin (BS, 10 μM), the chymase inhibitor chymostatin (CHYM, 10 μM), the carboxypeptidase A inhibitor benzy succinate (BSC 10 μM), and the ACE inhibitor lisinopril (LIS, 10 μM). All inhibitors were obtained from Sigma (St. Louis, MO) except SCH 39370 (gift from Schering Plow), CFP (Bachem, King of Prussia, PA), and JMV-390 (Tocris bioscience, UK). A pH profile was conducted as previously described (Marshall et al. 2013c). Purified peptidase (10 μl) was added to buffers ranging from pH 3.5 to 5.5 using 25 mM MES, 125 mM NaCl, and pH 7.5–9.5 using 25 mM HEPES, 125 mM NaCl. Iodinated products generated by the incubation of peptidase and ^{125}I -Ang-(1–7) were separated using HPLC.

Peptidase kinetics

Kinetic or competition assays were performed with ^{125}I -Ang-(1–7), -Ang II or -Ang I as substrates, and increasing concentrations of corresponding unlabeled peptides as previously described (Shaltout *et al.* 2009, Marshall et al. 2013b). Reaction velocities for generation of ^{125}I -Ang-(1–4) from the radiolabeled peptides were expressed as pmol/min/mg protein. The kinetic assays were performed with the metabolism cocktail to prevent the contribution of other peptidases and preserve the Ang-(1–4) product. Apparent kinetic constants (K_m),

V_{max}' were determined using Michaelis-Menten kinetics in the Prism 5 statistical program. Peptides were obtained from Bachem (Torrance CA) or custom synthesized (Genscript, Piscataway, NJ).

HPLC Separation

Metabolism reactions were conducted at 37°C in reaction buffer using 1445-fold purified peptidase (92 ng in a final volume of 250 µl) or 25 µl CSF. Each reaction included 0.5 nM ¹²⁵I-Ang-(1–7) and 0.1 µM non-iodinated Ang-(1–7) (Shaltout et al. 2009). The reactions were stopped after 60 min by addition of ice-cold 1.0% phosphoric acid and centrifuged at 16,000×g. The supernatant was immediately filtered for separation by reverse-phase HPLC on a Shimadzu equipped with an Aeris Peptide XB-C18 3.6 µm (2.1×100 mm, Phenomenex, Torrance CA). The ¹²⁵I-products were monitored by a Bioscan flow-through γ detector as described (Marshall et al. 2013b). Products were identified by comparison of their retention times to ¹²⁵I-standard peptides and sensitivity to peptidase inhibitors. Peptides were iodinated by the chloramine T method and purified by HPLC to a specific activity >2,000 Ci/mmol (Chappell et al. 1989).

UV HPLC Separation

Metabolism reactions were conducted at 37°C in reaction buffer using 1445-fold enriched peptidase (185 ng) and 50 nmol of unlabeled Ang I, Ang II, Ang-(1–7), Ang-(1–6) or Ang-(1–5), neurotensin, bradykinin, bradykinin-(1–7), bradykinin-(1–5), or apelin-13 in a final volume of 125 µl (final concentration of 0.4 mM). The reactions were stopped after 24 hours by addition of 1.0% phosphoric acid and separated by HPLC using a 0.1% phosphoric acid solvent system performed under a gradient of 0–15–25% phase B (80% acetonitrile) at a flow rate of 0.35 ml/min at ambient temperature. Peptides were monitored at 220 nm and the products identified by the retention time of standard peptides.

Statistics

Data are expressed as mean ± SEM. Unpaired t tests and one-way repeated measures ANOVA with Bonferroni posttests were used for the statistical analysis of the data with GraphPad Prism 5.01 (GraphPad Software, San Diego, CA). The criterion for statistical significance was set at **P* < 0.05.

RESULTS

Comparison of CSF and medullary metabolism of Ang-(1–7)

We initially compared the metabolism of ¹²⁵I-Ang-(1–7) in the cytosolic fraction of the brain medulla to metabolism in the CSF. As shown in Figure 1, ¹²⁵I-Ang-(1–7) was metabolized primarily to ¹²⁵I-Ang-(1–4) with a minor peak corresponding to ¹²⁵I-Ang-(3–4) in the medullary supernatant fraction; the ¹²⁵I-Ang-(1–4) peak was abolished by addition of the thiol inhibitor PCMB (10 µM). Comparison of the Ang-(1–7) saturation curves revealed similar apparent K_m (K_m') values for the CSF and brain medulla activities, but a 6-fold higher apparent V_{max} (V_{max}') for the medullary peptidase (Figure 1, right panel).

Purification of Ang-(1–7) peptidase activity from sheep medulla

In lieu of the higher tissue activity, the soluble fraction of the brain medulla was used as the initial source for the purification of peptidase activity. Soluble activity from 2.5 g of dorsal brain medulla was subjected to consecutive steps on DEAE, Cibacron Blue, and Q-Sepharose chromatography (Table 1). Peptidase activity was monitored by the conversion of ^{125}I -Ang-(1–7) to ^{125}I -Ang-(1–4) and sensitivity to the metalloendopeptidase inhibitor JMV-390 (see inhibitor sensitivity, below). The peptidase activity eluted in 250 mM NaCl on DEAE, was not retained on the Cibacron Blue gel, and eluted at 200 mM NaCl on Q-Sepharose following extensive washing with the 100 mM NaCl buffer. Fractions were collected in 15 ml increments from the 200 mM NaCl elution on the Q-Sepharose column; fraction 3 had the highest specific activity and was used to calculate purity and yield. We attained a 1445-fold purification with a 25% yield of the active peptidase using this protocol (Table 1). Peptidase activity apparently increased following the Cibacron Blue column (89 to 128%) which may reflect the separation of inhibitory substances that compete for the metabolism of Ang-(1–7). A second purification of the medullary soluble fraction obtained from the medullary tissue of different animals resulted in a similar degree of enrichment (1970-fold) and enzyme yield (31%) following the Q-Sepharose fractionation (preparation 2). SDS gels of the Q-Sepharose purified fraction stained with ImperialTM Protein Stain or the silver stain revealed approximately five protein bands in the 50 to 80 kDa range indicating that complete purification of the Ang-(1–7) peptidase was not achieved. Since the denatured gels abolished activity, it remains unclear what band corresponds to the peptidase.

Inhibitor Sensitivity

The sensitivity of the enriched peptidase from the Q-Sepharose column or concentrated CSF to various inhibitors was assessed by the conversion of ^{125}I -Ang-(1–7) to ^{125}I -Ang-(1–4). Similar to the CSF activity, the medullary peptidase was insensitive to specific inhibitors against neprilysin (10 μM , SCH 39370), TOP (10 μM , CFP), neurolysin (1 mM, Pro-Ile), the general cysteine peptidase inhibitor E-64 (10 μM), and the serine protease inhibitors PMSF (100 μM), aprotinin (80 μM), and SBTI (100 μM) (Figure 2A, Table 2). PCMB (10 μM) and o-phenanthroline (1 mM) abolished peptidase activity while DTT (5 mM) and EDTA (5 mM) exhibited partial inhibition (Figure 2B). The optimal pH for the peptidase purified from medullary tissue was pH 7.5 (Figure 2C). As compared to control conditions (Figure 3A), 1 nM JMV-390 reduced the Ang-(1–4) peak and 100 nM JMV-390 essentially abolished activity (Figure 3B and 3C, respectively). A dose-response curve for JMV-390 inhibition revealed an IC_{50} of 0.8 ± 0.2 nM with a R value of 0.996 for a one-site competition curve (Figure 3D). This was not significantly different than the IC_{50} for JMV-390 of the CSF activity (1.1 ± 0.4 nM) (Table 2).

Kinetic Properties

Saturation studies on the purified peptidase activity were performed using ^{125}I -labelled Ang-(1–7), Ang II and Ang I to compare the apparent kinetic constants among the three peptides. The kinetic analysis revealed apparent K_m (K_m') and V_{max} (V_{max}') values of 2.6 ± 0.3 μM and 72 ± 2.2 nmol/min/mg for Ang-(1–7), 2.8 ± 0.7 μM and 30 ± 3 nmol/min/mg for Ang II, and 4.3 ± 0.7 μM and 6 ± 0.3 nmol/min/mg for Ang I (Figure 4A–C, respectively). Although the

K_m' values were similar for all 3 peptides, Ang-(1–7) exhibited the highest V_{max}' as compared to Ang II or Ang I (p<0.01 vs. Ang I or Ang II, N=3). Apparent kinetic values for ¹²⁵I-labelled Ang-(1–7), Ang II and Ang I were also derived from the second batch of purification. While the K_m' values for Ang-(1–7) (2.0 μM), Ang II (7.4 μM) and Ang I (1.7 μM) were similar to the first purification, the V_{max}' values were >2-fold higher (174, 69, and 20 nmol/min/mg protein for Ang-(1–7), Ang II, and Ang I, respectively) and likely reflect the greater degree of purity in preparation 2 that yields a higher V_{max}'.

Substrate Specificity

We compared the metabolism of ¹²⁵I-Ang-(1–7) in the purified medullary fraction to the initial tissue supernatant in the absence of any peptidase or protease inhibitors. Essentially complete metabolism of ¹²⁵I-Ang-(1–7) to ¹²⁵I-Ang-(3–4) was evident in the medullary supernatant fraction that likely reflects dipeptidyl aminopeptidase activity (Figure 5A). In contrast, ¹²⁵I-Ang-(1–4) was the primary product following a 60 or 120 minute incubation with the purified peptidase (Figure 5B and C respectively). Addition of the inhibitor JMV-390 essentially abolished the ¹²⁵I-Ang-(1–4) peak, and did not reveal additional metabolites suggesting the absence of other activities that metabolize Ang-(1–7) (Figure 5D). Given the apparent lack of other peptidase activities in the purified preparation, we determined its specificity against various unlabeled angiotensins, as well as the bioactive peptides neurotensin, bradykinin, and apelin-13 (Figures 6 and 7). Ang-(1–7) was exclusively hydrolyzed to Ang-(1–4) and addition of JMV-390 abolished the peak of Ang-(1–4) (Figure 6A and 6B, respectively). Ang II and Ang I were metabolized to Ang-(1–4); however, the extent of metabolism was markedly less than that of Ang-(1–7) (Figure 6C and 6D). Ang-(1–6) and Ang-(1–5) were also metabolized to Ang-(1–4), but at lower rates than Ang-(1–7) (Table 3). Neurotensin, bradykinin, and the bradykinin metabolites bradykinin-(1–7) and bradykinin-(1–5) were not hydrolyzed by the peptidase as reflected by the absence of intermediate peaks distinct from the parent peptides (Figure 7 A–D). Based on the extent of hydrolysis of the unlabeled peptides, we calculated the rate of metabolism for each peptide substrate (Table 3). Ang-(1–7) was hydrolyzed to a greater extent than Ang II (11-fold) or Ang I (27-fold). The rate of Ang-(1–7) metabolism was 2-fold higher than Ang-(1–6) and 26-fold greater than the pentapeptide Ang-(1–5). There was no apparent metabolism of neurotensin, bradykinin or bradykinin metabolites, or apelin-13 (chromatograph not shown) following the 24 hour incubation conditions.

DISCUSSION

In the present study, we isolated and characterized an Ang-(1–7) metabolizing activity in the brain medulla of sheep that extends the original identification of the peptidase activity in the CSF (Marshall et al. 2013c). With respect to the similar kinetic parameters, substrate specificity, inhibitor sensitivity, and optimal pH, it is likely that the medullary peptidase and the CSF peptidase activities are identical (Table 2). Using a combination of DEAE Sepharose, Cibacron Blue, and Q-Sepharose chromatography, the peptidase was purified 1445- and 1950-fold from supernatant fraction of the brain medulla in two preparations using tissues pooled from multiple animals. The isolated activity is optimally active at pH 7.5 and sensitive to the mercury compounds PCMB and aminophenylmercuric acetate

(APMA), and chelating agents o-phenanthroline, EDTA, and DTT; however peptidase activity is not reduced by selective inhibitors against the metalloendopeptidases neprilysin, neurolysin, and TOP nor the serine protease inhibitors PMSF, aprotinin, or SBTI. The peptidase appears to preferentially cleave the Tyr⁴-Ile⁵ bond of Ang-(1–7) in comparison to other angiotensin peptides or the bioactive peptides neurotensin, bradykinin, and apelin-13. Kinetic analyses using [¹²⁵I]-labeled angiotensins revealed similar Km' values (2.6 ± 0.3 , 2.8 ± 0.7 , and 4.3 ± 0.7 μ M, for Ang-(1–7), Ang II, and Ang I, respectively), but a significantly higher Vmax' for Ang-(1–7) (72 ± 2.2 nmol/min/mg) versus Ang II (30 ± 3 nmol/min/mg) or Ang I (6 ± 0.3 nmol/min/mg). These data likely represent a novel pathway for specific regulation of central Ang-(1–7) levels.

Metallopeptidases are the major class of enzymes involved in extracellular peptide metabolism (Shrimpton *et al.* 2002). ACE, ACE2, neprilysin, TOP, and neurolysin belong to the same class of thermolysin-like zinc-dependent peptidases that hydrolyze peptide substrates (<40 amino acids) and are maximally active at a neutral pH (Rawlings & Barrett 1993, Shrimpton *et al.* 2002). The characterization of a soluble metalloendopeptidase in the CSF and brain medulla is consistent with this class of peptidases, yet appears to be more restricted regarding the substrate length and the site of hydrolysis. For example, angiotensin peptides larger than Ang-(1–7) such as Ang II and Ang I exhibited an 11- and 27-fold lower rate of hydrolysis to generate Ang-(1–4). Interestingly, neurotensin contains a Tyr-Ile site at positions 12 and 13 of the peptide but was not cleaved by the peptidase; this may reveal a more constrained effect of peptides greater than 10 residues in length. In this regard, both neurolysin and TOP exhibit restriction on the length of their peptide substrates that may reflect the limited access to the active sites of these peptidases (Shrimpton & Smith 2000). Peptides that lack the Tyr-Ile site such as bradykinin and apelin-13 were not cleaved by this peptidase. Moreover, the seven and five amino acid metabolites of bradykinin were not hydrolyzed by the medullary peptidase. The shorter fragments of Ang-(1–7) including Ang-(1–6) and Ang-(1–5) that are generally considered biologically inactive were hydrolyzed at a lower rate (2- and 25-fold, respectively) than Ang-(1–7), again suggesting an optimal length for recognition by the peptidase. Finally, we note that while the velocity rates for the hydrolysis of ¹²⁵I-Ang-(1–7) and Ang-(1–7) were similar (72 versus 54 nmol/min/mg), the velocity rate for ¹²⁵I-Ang II was 6-fold higher than that of Ang II (30 versus 5 nmol/min/mg) and the rate for ¹²⁵I-Ang I was 3 fold higher than Ang I (6 versus 2 nmol/min/mg). These data suggest an apparent influence of the addition of the ¹²⁵I-group on the Tyr⁴ residue, particularly for Ang II that may enhance the rate of hydrolysis of the peptide. Although additional kinetic studies are required to elucidate the mechanism for this effect on the Tyr residue, the use of ¹²⁵I-labeled peptides other than Ang-(1–7) may not accurately reflect the substrate characteristics for the peptidase.

The Ang-(1–7) peptidase was isolated from the supernatant fraction of the medullary tissue, consistent with the soluble activity detected in the CSF (Marshall *et al.* 2013c). Metallopeptidases such as ACE, ACE2, neprilysin and neurolysin are predominantly membrane-associated, thus they are situated to hydrolyze bioactive peptides in the extracellular space. In contrast, TOP is a soluble enzyme that is primarily intracellular with no hydrophobic membrane-spanning domain (Shrimpton *et al.* 2003, Acker *et al.* 1987,

Woulfe *et al.* 1992, Jeske *et al.* 2004). Several studies suggest that TOP localizes to the nucleus, consistent with the identification of a nuclear localization sequence for the peptidase (Healy & Orłowski 1992, Alzayadneh & Chappell 2013, Massarelli *et al.* 1999, Thompson *et al.* 1995). TOP reportedly undergoes both stimulated secretion and constitutive release from neuronal cells and is known to associate with lipid rafts in the plasma membrane (Jeske *et al.* 2004, Ferro *et al.* 1999, Jeske *et al.* 2003). Once in the extracellular space, TOP participates in the extracellular metabolism of neuropeptides such as gonadotropin-releasing hormone, bradykinin, and neurotensin (Ferro *et al.* 1999). We have not established the intracellular distribution of the Ang-(1–7) peptidase nor the specific cell type (ie. neuronal vs. glia) that express the peptidase within the medulla. Increasing evidence supports an intracellular RAS including the expression of intracellular Ang II and Ang-(1–7), as well as their respective receptors (Gwathmey *et al.* 2012, Chappell *et al.* 2014, Ellis *et al.* 2012, Singh *et al.* 2007). Indeed, we quantified the peptide content of Ang-(1–7), Ang II and Ang I in the brain medulla of sheep (Marshall *et al.* 2013a). Krob *et al.* demonstrated intense Ang-(1–7) immunostaining in hypothalamic neurons of the mRen2(27) transgenic rats suggesting the intracellular localization of the peptide (Krob *et al.* 1998). Furthermore, Gironacci and colleagues reported intracellular Ang-(1–7) expression in primary neuronal cells cultured from the hypothalamic-brainstem areas (Verrilli *et al.* 2009). In this regard, the peptidase may potentially influence the local processing of Ang-(1–7) within the cells. Alternatively, the peptidase may be secreted or released from medullary tissue to degrade extracellular Ang-(1–7) or perhaps other substrates that are unidentified at this time. Indeed, the presence of the soluble peptidase in the CSF may reflect the secretion of the enzyme into this compartment. Additional studies are necessary to identify the cell types that contain the peptidase and address whether the enzyme activity is actively secreted from medullary tissues.

The current chromatographic approach was not intended to achieve a preparation of pure enzyme, but to obtain sufficient activity to characterize the peptidase; thus, we cannot establish the identity of the protein with this preparation at the present time. Nevertheless, the studies of the enriched enzyme suggest a metalloendopeptidase-like activity to convert Ang-(1–7) to Ang-(1–4). Although the peptidase activity was not blocked by selective inhibitors to neprilysin, ACE, neurolysin or TOP, the enzyme exhibits marked sensitivity to the inhibitor JMV-390 ($IC_{50} = 0.80$ nM). In contrast, the reported IC_{50} values for JMV-390 against neprilysin, neurolysin, TOP, and leucine aminopeptidase (EC 3.4.11.1) are 30 to 60 nM, and a far higher value of 70 μ M against ACE (Doulut *et al.* 1993). JMV-390 was originally synthesized to inhibit metalloendopeptidase activity thereby extending the functional benefits of neurotensin and neuromedin in brain (Doulut *et al.* 1993). The molecular design of JMV-390 was based on the recognition of the leucine-leucine sequence that corresponds to neurotensin, and the amino-terminal hydroxy amino group for an interaction with the zinc cofactor in the active site (Doulut *et al.* 1993).

ACE constitutes a prominent pathway for the metabolism of Ang-(1–7) in the circulation (Chappell 2007). ACE inhibition significantly increases the half-life of Ang-(1–7) in the circulation by blocking the hydrolysis of Ang-(1–7) to Ang-(1–5) (Yamada *et al.* 1998, Chappell *et al.* 1998). The marked increase in circulating Ang-(1–7) following

administration of ACE inhibitors also reflects an increase in neprilysin-dependent processing of Ang I to Ang-(1–7) (Chappell et al. 1998, Campbell et al. 1998). We reported that soluble ACE contributes to Ang-(1–7) metabolism in sheep CSF *ex vivo*; however, the majority of the Ang-(1–7)-degrading activity was ACE independent and this activity inversely correlated to Ang-(1–7) content in the CSF (Marshall et al. 2013c). The presence of an Ang-(1–7) peptidase in the CSF and brain medulla may represent an alternative pathway to metabolize the peptide, particularly if the local expression of ACE is low or there is restricted access to ACE. Additional studies are necessary to compare the relative expression of ACE and the Ang-(1–7) peptidase in other areas of the brain, as well as the presence of the metallopeptidase in the circulation and peripheral tissues such as the heart and kidney that express Ang-(1–7).

Perspective and Significance

Central Ang-(1–7) is critically involved in blood pressure control and autonomic regulation. Low levels of Ang-(1–7) are associated with hypertension, reduced baroreflex sensitivity for control of heart rate, inflammation, cell proliferation, and oxidative stress (Averill & Diz 2000, Diz *et al.* 2008, Diz *et al.* 2011). The peptidase characterized in this study may potentially represent a novel Ang-(1–7) metabolizing pathway involved in the pathological reduction of central Ang-(1–7) levels (Marshall et al. 2013c). Moreover, we identified JMV-390 as a potent inhibitor of the Ang-(1–7) peptidase that exhibits an IC₅₀ value far below that reported for neprilysin, neurolysin, TOP, leucine aminopeptidase, and ACE. JMV-390, or more selective analogs, may constitute a novel selective inhibitor against the Ang-(1–7) peptidase to prevent metabolism of Ang-(1–7) in brain. Enhanced peptidase activity in CSF of animals exposed *in utero* to glucocorticoids, where Ang-(1–7) levels were significantly lower than control animals, further suggests that the peptidase participates in physiological regulation of the brain RAS.

Acknowledgments

Support for this study was provided by National Institutes of Health Grants HD-047584, HD-017644, and HL-51952; the Groskert Heart Fund and the Wake Forest Venture Fund. The authors gratefully acknowledge Ellen Tommasi and Eric LeSaine for their technical and surgical support.

Abbreviations

ang	Angiotensin
ACE	angiotensin converting enzyme
DEAE	diethylaminoethyl
DTT	dithiothreitol
JMV-390	<i>N</i> -[3-[(hydroxyamino) carbonyl]-1-oxo-2(<i>R</i>)-benzylpropyl]-L-leucine
PCMB	p-chloromercuribenzoic acetate
TOP	thimet oligopeptidase

REFERENCES

- Acker GR, Molineaux C, Orlowski M. Synaptosomal membrane-bound form of endopeptidase-24.15 generates Leu-enkephalin from dynorphin1–8, alpha- and beta-neoendorphin, and Met-enkephalin from Met-enkephalin-Arg6-Gly7-Leu8. *J Neurochem.* 1987; 48:284–292. [PubMed: 2878974]
- Allred AJ, Diz DI, Ferrario CM, Chappell MC. Pathways for angiotensin-(1–7) metabolism in pulmonary and renal tissues. *Am J Physiol Renal Physiol.* 2000; 279:F841–F850. [PubMed: 11053044]
- Alzayadneh EM, Chappell MC. Nuclear expression of components of renin-angiotensin system in the NRK52E cells. *Journal of the Renin-Angiotensin-Aldosterone System.* 2013 In press.
- Arnold AC, Shaltout HA, Gilliam-Davis S, Kock ND, Diz DI. Autonomic control of the heart is altered in Sprague-Dawley rats with spontaneous hydronephrosis. *Am J Physiol Heart Circ Physiol.* 2011; 300:H2206–H2213. [PubMed: 21460193]
- Averill DB, Diz DI. Angiotensin peptides and baroreflex control of sympathetic outflow: pathways and mechanisms of the medulla oblongata. *Brain Res Bull.* 2000; 51:119–128. [PubMed: 10709957]
- Campbell DJ, Anastasopoulos F, Duncan AM, James GM, Kladis A, Briscoe TA. Effects of neutral endopeptidase inhibition and combined angiotensin converting enzyme and neutral endopeptidase inhibition on angiotensin and bradykinin peptides in rats. *J Pharmacol Exp Ther.* 1998; 287:567–577. [PubMed: 9808682]
- Capettini LS, Montecucco F, Mach F, Stergiopoulos N, Santos RA, da Silva RF. Role of renin-angiotensin system in inflammation, immunity and aging. *Curr Pharm Des.* 2012; 18:963–970. [PubMed: 22283774]
- Chappell MC. Emerging evidence for a functional angiotensin-converting enzyme 2-angiotensin-(1–7)-Mas receptor axis: more than regulation of blood pressure? *Hypertension.* 2007; 50:596–599. [PubMed: 17785634]
- Chappell MC, Brosnihan KB, Diz DI, Ferrario CM. Identification of angiotensin-(1–7) in rat brain. Evidence for differential processing of angiotensin peptides. *J Biol Chem.* 1989; 264:16518–16523. [PubMed: 2777795]
- Chappell MC, Marshall AC, Alzayadneh EM, Shaltout HA, Diz DI. Update on the angiotensin converting enzyme 2-angiotensin (1–7)-Mas receptor axis: fetal programming, sex differences, and intracellular pathways. *Front Endocrinol (Lausanne).* 2014; 4:1–13.
- Chappell MC, Pirro NT, Sykes A, Ferrario CM. Metabolism of angiotensin-(1–7) by angiotensin-converting enzyme. *Hypertension.* 1998; 31:362–367. [PubMed: 9453329]
- Diz DI, Arnold AC, Nautiyal M, Isa K, Shaltout HA, Tallant EA. Angiotensin peptides and central autonomic regulation. *Curr Opin Pharmacol.* 2011; 11:131–137. [PubMed: 21367658]
- Diz DI, Garcia-Espinosa MA, Gallagher PE, Ganten D, Ferrario CM, Averill DB. Angiotensin-(1–7) and baroreflex function in nucleus tractus solitarii of (mRen2)27 transgenic rats. *J Cardiovasc Pharmacol.* 2008; 51:542–548. [PubMed: 18475201]
- Doulut S, Dubuc I, Rodriguez M, et al. Synthesis and analgesic effects of N-[3-[(hydroxyamino) carbonyl]-1-oxo-2(R)-benzylpropyl]-L-isoleucyl-L-leucine, a new potent inhibitor of multiple neurotensin/neuromedin N degrading enzymes. *J Med Chem.* 1993; 36:1369–1379. [PubMed: 8496905]
- Ellis B, Li XC, Miguel-Qin E, Gu V, Zhuo JL. Evidence for a functional intracellular angiotensin system in the proximal tubule of the kidney. *Am J Physiol Regul Integr Comp Physiol.* 2012; 302:R494–R509. [PubMed: 22170616]
- Ferro ES, Tullai JW, Glucksman MJ, Roberts JL. Secretion of metalloendopeptidase 24.15 (EC 3.4.24.15). *DNA Cell Biol.* 1999; 18:781–789. [PubMed: 10541437]
- Grobe N, Weir NM, Leiva O, Ong FS, Bernstein KE, Schmaier AH, Morris M, Elased KM. Identification of prolyl carboxypeptidase as an alternative enzyme for processing of renal angiotensin II using mass spectrometry. *Am J Physiol-Cell Ph.* 2013; 304:C945–C953.
- Gwathmey TM, Alzayadneh EM, Pendergrass KD, Chappell MC. Novel roles of nuclear angiotensin receptors and signaling mechanisms. *Am J Physiol Regul Integr Comp Physiol.* 2012; 302:R518–R530. [PubMed: 22170620]

- Healy DP, Orlowski M. Immunocytochemical localization of endopeptidase 24.15 in rat brain. *Brain Res.* 1992; 571:121–128. [PubMed: 1377082]
- Jeske NA, Glucksman MJ, Roberts JL. EP24.15 is associated with lipid rafts. *Journal of Neuroscience Research.* 2003; 74:468–473. [PubMed: 14598323]
- Jeske NA, Glucksman MJ, Roberts JL. Metalloendopeptidase EC 3.4.24.15 is constitutively released from the exofacial leaflet of lipid rafts in GT1–7 cells. *J Neurochem.* 2004; 90:819–828. [PubMed: 15287887]
- Kar S, Gao L, Belatti DA, Curry PL, Zucker IH. Central angiotensin (1–7) enhances baroreflex gain in conscious rabbits with heart failure. *Hypertension.* 2011; 58:627–634. [PubMed: 21844487]
- Krob HA, Vinsant SL, Ferrario CM, Friedman DP. Angiotensin-(1–7) immunoreactivity in the hypothalamus of the (mRen-2d)27 transgenic rat. *Brain Res.* 1998; 798:36–45. [PubMed: 966069]
- Marshall AC, Shaltout HA, Nautiyal M, Rose JC, Chappell MC, Diz DI. Fetal betamethasone exposure attenuates angiotensin-(1–7)-Mas receptor expression in the dorsal medulla of adult sheep. *Peptides.* 2013a; 44:25–31. [PubMed: 23538211]
- Marshall AC, Shaltout HA, Pirro NT, Rose JC, Diz DI, Chappell MC. Antenatal betamethasone exposure is associated with lower ANG-(1–7) and increased ACE in the CSF of adult sheep. *Am J Physiol Regul Integr Comp Physiol.* 2013b; 305:R679–R688. [PubMed: 23948771]
- Marshall AC, Shaltout HA, Pirro NT, Rose JC, Diz DI, Chappell MC. Enhanced activity of an angiotensin-(1–7) neuropeptidase in glucocorticoid-induced fetal programming. *Peptides.* 2013c; 52C:74–81. [PubMed: 24355101]
- Massarelli EE, Casatti CA, Kato A, Camargo AC, Bauer JA, Glucksman MJ, Roberts JL, Hirose S, Ferro ES. Differential subcellular distribution of neurolysin (EC 3.4.24.16) and thimet oligopeptidase (EC 3.4.24.15) in the rat brain. *Brain Res.* 1999; 851:261–265. [PubMed: 10642854]
- Pereira MG, Souza LL, Becari C, et al. Angiotensin II-independent angiotensin-(1–7) formation in rat hippocampus: involvement of thimet oligopeptidase. *Hypertension.* 2013; 62:879–885. [PubMed: 24041943]
- Rawlings ND, Barrett AJ. Evolutionary families of peptidases. *Biochem J.* 1993; 290(Pt 1):205–218. [PubMed: 8439290]
- Rykl J, Thiemann J, Kurzawski S, Pohl T, Gobom J, Zidek W, Schluter H. Renal cathepsin G and angiotensin II generation. *J Hypertens.* 2006; 24:1797–1807. [PubMed: 16915029]
- Sadjadi J, Kramer GL, Yu CH, Burrell Welborn M, Chappell MC 3rd, Gregory Modrall J. Angiotensin converting enzyme-independent angiotensin ii production by chymase is up-regulated in the ischemic kidney in renovascular hypertension. *J Surg Res.* 2005; 127:65–69. [PubMed: 15869764]
- Shaltout HA, Figueroa JP, Rose JC, Diz DI, Chappell MC. Alterations in circulatory and renal angiotensin-converting enzyme and angiotensin-converting enzyme 2 in fetal programmed hypertension. *Hypertension.* 2009; 53:404–408. [PubMed: 19047579]
- Shrimpton CN, Smith AI. Soluble neutral metallopeptidases: physiological regulators of peptide action. *J Pept Sci.* 2000; 6:251–263. [PubMed: 10912905]
- Shrimpton CN, Smith AI, Lew RA. Soluble metalloendopeptidases and neuroendocrine signaling. *Endocr Rev.* 2002; 23:647–664. [PubMed: 12372844]
- Shrimpton CN, Wolfson AJ, Smith AI, Lew RA. Regulators of the neuropeptide-degrading enzyme, EC 3.4.24.15 (thimet oligopeptidase), in cerebrospinal fluid. *J Neurosci Res.* 2003; 74:474–478. [PubMed: 14598324]
- Singh VP, Le B, Bhat VB, Baker KM, Kumar R. High-glucose-induced regulation of intracellular ANG II synthesis and nuclear redistribution in cardiac myocytes. *Am J Physiol-Heart C.* 2007; 293:H939–H948.
- Thompson A, Huber G, Malherbe P. Cloning and functional expression of a metalloendopeptidase from human brain with the ability to cleave a beta-APP substrate peptide. *Biochem Biophys Res Commun.* 1995; 213:66–73. [PubMed: 7639763]

- Verrilli MAL, Pirola CJ, Pascual MM, Dominici FP, Turyn D, Gironacci MM. Angiotensin-(1–7) through AT(2) receptors mediates tyrosine hydroxylase degradation via the ubiquitin-proteasome pathway. *Journal of Neurochemistry*. 2009; 109:326–335. [PubMed: 19183250]
- Welches WR, Santos RAS, Chappell MC, Brosnihan KB, Greene LJ, Ferrario CM. Evidence that prolyl endopeptidase participates in the processing of brain angiotensin. *Journal of Hypertension*. 1991; 9:631–638. [PubMed: 1653799]
- Woulfe J, Checler F, Beaudet A. Light and electron microscopic localization of the neutral metalloendopeptidase EC 3.4.24.16 in the mesencephalon of the rat. *Eur J Neurosci*. 1992; 4:1309–1319. [PubMed: 12106394]
- Xu P, Sriramula S, Lazartigues E. ACE2/ANG-(1–7)/Mas pathway in the brain: the axis of good. *Am J Physiol Regul Integr Comp Physiol*. 2011; 300:R804–R817. [PubMed: 21178125]
- Yamada K, Iyer SN, Chappell MC, Ganten D, Ferrario CM. Converting enzyme determines plasma clearance of angiotensin-(1–7). *Hypertension*. 1998; 32:496–502. [PubMed: 9740616]

Figure 1.

Ang-(1-7) metabolizing peptidase activity in the brain medulla and CSF. **A:** ^{125}I -Ang-(1-7) [A7] is metabolized to ^{125}I -Ang-(1-4) [A4] and ^{125}I -Ang-(3-4) [A3-4]. **B:** A7 metabolism is abolished with the addition of PCMB (10 μM). **C:** Comparison of apparent kinetics of ^{125}I -Ang-(1-7) metabolizing peptidase in brain medulla and CSF. Reactions were run in the presence of an inhibitor cocktail (AM, BS, CHYM, BSC, LIS) for 60 minutes at 37°C.

Figure 2.

Inhibitor and pH profiles of the enriched Ang-(1-7) brain medullary peptidase. **A:** Specific inhibitors SCH 39370 (10 μ M, SCH), Pro-Ile (1 mM), CFP (10 μ M), E-64 (10 μ M), or PMSF (100 μ M) did not inhibit activity. **B:** Peptidase was sensitive to the chelating agents PCMB (10 μ M), o-phenanthroline (1 mM, PHEN), dithiothreitol (5 mM, DTT), and EDTA (5 mM). **C:** pH profile reveals an optimal pH of 7.5 for peptidase activity. Reactions were carried out in the presence of an inhibitor cocktail (AM, BS, CHYM, BSC, LIS) for 60

minutes at 37°C. N=3 separate determinations from the Q-Sepharose purified fraction of preparation 1, ***P<0.0001 vs. control.

Figure 3. JMV-390 inhibits the Ang-(1-7) peptidase activity. **A–C:** ^{125}I -Ang-(1-7) was incubated with purified enzyme in the presence of 0, 1 nM, or 100 nM JMV-390. **D:** A dose-response curve was conducted to determine the IC_{50} for JMV-390. Reactions were carried out in the presence of an inhibitor cocktail (AM, BS, CHYM, BSC, LIS). N=3 separate determinations from the Q-Sepharose purified fraction of preparation 1, *** $P < 0.0001$ vs. control.

Figure 4.

^{125}I -Ang-(1-7) exhibits the highest apparent V_{max} for the brain medullary peptidase. **A–C:** Apparent kinetic values were determined for ^{125}I -Ang-(1-7), ^{125}I -Ang II, and ^{125}I -Ang I. Kinetic values were determined based on the conversion of radiolabeled peptide to ^{125}I -Ang-(1-4) in the presence of an inhibitor cocktail (AM, BS, CHYM, BSC, LIS). N=3 separate determinations from the Q-Sepharose purified fraction of preparation 1.

Figure 5.

Purified peptidase lacks other Ang-(1-7) metabolizing enzyme activity. **A:** Medullary supernatant incubated with ^{125}I -Ang-(1-7) and no inhibitors for 60 min. **B:** Q-Sepharose purified fraction incubated with ^{125}I -Ang-(1-7) and no inhibitors for 60 min. **C:** Q-Sepharose purified fraction incubated with ^{125}I -Ang-(1-7) and no inhibitors for 120 min. **D:** Q-Sepharose purified fraction incubated with ^{125}I -Ang-(1-7) and 1 nM JMV for 120 min.

Figure 6. Purified medullary peptidase preferentially hydrolyzes Ang-(1-7) to Ang-(1-4). Purified peptidase was incubated with 50 nmol (final concentration of 400 μ M) of **A:** Ang-(1-7), **B:** Ang-(1-7) + 10 μ M JMV-390, **C:** Ang II, **D:** Ang I for 24 hours at 37°C in the presence of no inhibitors.

Figure 7.

Purified medullary peptidase does not hydrolyze other selected neuropeptides. Purified peptidase was incubated with 50 nmol (final concentration of 400 μ M) of **A:** Neurotensin (NT), **B:** Bradykinin (BK)-(1-9), **C:** BK-(1-7), or **D:** BK-(1-5) for 24 hours at 37°C in the presence of no inhibitors.

Table 1

Four-step purification of brain peptidase. A combination of centrifugation, DEAE, Cibacron Blue, and Q-Sepharose chromatography were used to achieve 1445-fold purification over starting material.

Stage	Activity (pmol/ml)	Protein (mg/ml)	Specific Activity (pmol/mg)	Yield (%)	Purification Factor (fold)
Supernatant (25,000×g)	142	244	0.58	100%	1
DEAE	127	17	7.5	89%	13
Cibacron Blue	183	4	44.5	128%	77
Q-Sepharose	40	.047	831.9	25%	1445

Table 2

Comparison of CSF and the purified medullary peptidase activities for hydrolysis of ^{125}I -Ang-(1–7). Values for each inhibitor represent percent of control. N=3 separate determinations from the Q-Sepharose purified fraction.

	CSF	Medullary Tissue
Ang-(1–7) Affinity		
K _m ' (μM)	8.5 ± 0.5	3.0 ± 0.6
V _{max} ' (pmol/min/mg)	84 ± 1.5	495 ± 29
Inhibitor Sensitivity (% of control)		
JMV-390	0 (IC ₅₀ = 1.1 ± 0.3 nM)	0 (IC ₅₀ = 0.8 ± 0.2 nM)
PCMB (10 μM)	0	0
APMA (10 μM)	0	0
EDTA (5 mM)	61 ± 2	79 ± 5
DTT (5 mM)	50 ± 4	73 ± 3
o-phenanthroline (1 mM)	1 ± 2	4 ± 0.3
E-64 (10 μM)	100 ± 1	100
SCH (10 μM)	100	99 ± 0.6
CFP (10 μM)	102 ± 3	98 ± 3
Pro-Ile (1 mM)	101 ± 2	101 ± 0.5
Aprotinin (80 μM)	100	103 ± 0.9
SBTI (100 μM)	101 ± 2	101 ± 0.3
Lisinopril (10 μM)	100	100
Optimal pH	7.5	7.5

Table 3

Comparison of the metabolism velocities for angiotensins and other peptides. Velocity is an approximation of V_{max} , as reactions are carried out in the presence of excess peptide substrate. NC = not cleaved within 24 hours of digestion.

Peptide	Amino Acid Sequence	Velocity (nmol/min/mg)
Ang I	DRVYIHPFHL	2.0
Ang II	DRVYIHPF	4.9
Ang-(1-7)	DRVYIHP	54
Ang-(1-6)	DRVYIH	27
Ang-(1-5)	DRVYI	2.1
Neurotensin	ELYENLPRRPYIL	NC
Bradykinin-(1-9)	RPPRFSPFR	NC
Bradykinin-(1-7)	RPPRFSP	NC
Bradykinin-(1-5)	RPPRF	NC
Apelin 13	KFRRQRPRLSHKG	NC