Published in final edited form as:

Curr Opin HIV AIDS. 2014 July ; 9(4): 302–308. doi:10.1097/COH.0000000000000072.

HIV and Ageing: Emerging Research Issues

Peter W. Hunt, MD

SFGH HIV/AIDS Division, University of California, San Francisco

Abstract

Purpose of the review—It is now widely accepted that HIV-infected individuals remain at higher risk for mortality and age-related morbidities than the general population, but several unresolved issues need to be addressed by the research community in the coming years to further improve the health of HIV-infected individuals in the modern treatment era.

Recent findings—While recent studies have helped better define the contribution of HIV to life expectancy and morbidity in the modern ART era, questions remain about the generalizability of these findings to a future HIV-infected population that is expected to be much older. Furthermore, while a consensus has emerged that the persistent inflammatory state contributes to morbidity and mortality in this setting, the relative contributions of this process, health-related behaviors, co-morbidities, and medication toxicities remain incompletely understood. Lastly, significant uncertainty remains over the root causes of the persistent inflammatory state, the specific immunologic pathways to target with interventions, and the most appropriate biomarkers to use for surrogate outcomes in pilot trials of immune-based interventions.

Summary—Each of these issues will be addressed in this review, highlighting recently published and presented studies that inform the discussion, and recommendations will be made for prioritizing the future research agenda.

Keywords

HIV-1; inflammation; aging; multi-morbidity; mortality; antiretroviral therapy

Introduction

HIV-infected individuals have experienced a dramatic improvement in life expectancy in the modern antiretroviral therapy (ART) era, particularly those who initiated therapy at early disease stages [1, 2]. Nevertheless, millions of HIV-infected individuals around the world – and millions more to come – will have initiated ART at advanced disease stages and appear to be at substantially higher risk of morbidity and mortality than the general population, even when sustained viral suppression is achieved [3]. There is now an emerging consensus that the persistent inflammatory state contributes to this increased risk [4]. This inflammation-associated risk of morbidity and mortality is also superimposed on an aging HIV epidemic in both resource-rich and resource-limited settings as individuals live longer

on ART, potentially amplifying the risk of age-related morbidity. Much progress has been made in understanding the probable causes of excess morbidity and mortality in treated HIV infection, but several unresolved issues remain to be addressed to further improve the health of HIV-infected individuals receiving ART.

The scope of the problem

A broad consensus now exists that HIV-infected individuals in the modern treatment era have a higher risk of mortality and age-associated multi-morbidity than the general population [5–9]. This excess HIV-associated risk of morbidity and mortality will almost certainly become a greater public health issue as more HIV-infected individuals enter their sixth and seventh decades of life. Indeed, half of all HIV-infected individuals in the United States are expected to be over the age of 50 within the next few years [10], as will many more millions of HIV-infected individuals in resource-limited settings around the world [11, 12]. Nevertheless, significant uncertainty remains over whether inferences from recent cohort studies will accurately reflect a future HIV-infected population that is expected to be much older.

For example, several recent cohort studies have suggested that the life expectancy of HIV-infected individuals may approach that of the general population, particularly among those who initiated ART at earlier disease stages [1, 13–15]. These projections assume that the risk of mortality in the relatively small proportion of older individuals in contemporary cohort studies will reflect the risk actually observed in the future population. This may not be a safe assumption. As antiretroviral treatment regimens have become more effective and less toxic, and more patients initiate ART at earlier disease stages, it is possible that the risk of morbidity and mortality might be lower for older HIV-infected individuals in the future. On the other hand, most of the current older population of HIV-infected individuals survived the pre-ART and early ART eras and may well be enriched for favorable host genetics and healthier lifestyles than the general population. Such a survivorship bias might result in a significant underestimation of the influence of HIV on morbidity and mortality in older populations. Indeed, given the approximately 50% increased risk of coronary artery disease in the HIV-infected population after adjustment for traditional cardiovascular risk factors [16, 17], as well as a potentially even greater relative risk of sudden cardiac death [18], it seems likely that recent life expectancy projections for the HIV-infected population may well be overly optimistic. This may be particularly true if the impact of HIV on multi-morbidity and life expectancy increases with age, though most clinical cohorts currently have too few HIV-infected individuals in their sixth and seventh decades to definitively assess whether this might be the case. It will also take time before the leading edge of older HIV-infected patients in most cohorts is no longer strongly influenced by survivorship bias issues.

Does HIV Cause Accelerated Aging or Increased Co-Morbidity?

While it is well established that HIV-infected individuals have a higher risk of several age-related co-morbidities than the general population, there has been considerable debate as to whether HIV causes accelerated aging [7]. This is in part because “aging” itself is difficult

to define, but also because the framing of the debate depends on one's perspective. For example, if one wants to improve the overall health and quality of life of a frail HIV-infected individual with multiple co-morbidities, then the recognition that this clinical phenotype is similar to that observed in much older geriatric populations is helpful. Minimizing pill burden, overlapping toxicities, and drug interactions; avoiding destructive health-related behaviors; and focusing on interventions like moderate exercise that holistically address overall function rather than specific disease states are all valuable lessons from the geriatrics literature that are likely to be helpful for frail HIV-infected individuals. On the other hand, if the goal is to identify targets for interventions to prevent multi-morbidity and frailty in HIV infection, then understanding the specific diseases that are increased as well as the mechanisms that drive them is critical. Here, specificity is important. For example, while many non-AIDS cancers are in fact increased in HIV-infected individuals with a history of AIDS, not all age-related cancers are increased, and some cancers (i.e., breast, prostate, and colon) may even be decreased [19]. While surveillance bias may have led to under-reporting of some cancers in patients with AIDS, it is notable that a similar pattern of cancer incidence – both those that are increased and decreased – is observed in other inflammatory diseases like rheumatoid arthritis [20]. Here, recognizing the differences between the effects of HIV and aging on cancer incidence may actually provide clues as to the pathophysiology and highlight targets for interventions. Similarly, our group recently demonstrated that the phenotypic CD8+ T cell defects observed in treated HIV infection are actually quite distinct from those observed in aging and have qualitatively different effects on mortality risk [21, 22]. Thus, while the clinical features of multi-morbidity in HIV and aging may be similar, their root causes may be different and understanding those differences may help identify more appropriate interventional targets to prevent multi-morbidity from developing in the first place. A better understanding of the shared and distinct determinants of multi-morbidity and frailty in HIV and aging is thus clearly needed to identify the most appropriate interventional targets.

Relative importance of inflammation as a target for interventions

While there is now a broad consensus that immune activation and inflammation persist in the majority of HIV-infected individuals maintaining long-term ART-mediated viral suppression – even in those that restore normal CD4+ T cell counts [23–26], and that immune activation and inflammatory markers strongly predict morbidity and mortality and even frailty in this setting [27–37], the degree to which inflammation is a direct cause of morbidity and mortality in this setting remains controversial. A prominent role of the inflammatory state in explaining the excess morbidity and mortality observed in treated HIV infection is supported by several observations. First, the relationship between inflammatory markers and mortality is much stronger in treated HIV infection than it is in the general population [32, 38], consistent with a more central role of the inflammatory state in driving mortality risk in this setting. Second, while confounding medication toxicities may play a role, it is notable that other inflammatory diseases like rheumatoid arthritis and psoriasis are associated with similar magnitude increases in cardiovascular disease, osteoporosis, and lymphoma [20, 39–43], consistent with a role of systemic inflammation in promoting chronic disease. Indeed, each of the surrogate markers of end-organ disease (i.e.,

hemoglobin, creatinine, and transaminase levels) that contribute to the prognostic utility of the Veterans Aging Cohort Study (VACS) index is strongly associated with markers of inflammation, monocyte activation, and coagulation [31], underscoring the link between the inflammatory state and dysfunction of multiple organ systems in HIV infection.

Nevertheless, it remains unclear how much of the increased morbidity and mortality in treated HIV infection is attributable to the inflammatory state as opposed to confounding lifestyle factors, co-morbidities, or antiretroviral medication toxicities. Often, studies comparing HIV-infected and –uninfected individuals lack sufficient granularity in confounding risk factor assessment and/or are too small to completely exclude confounding by these factors. That said, some potential confounding co-morbidities like diabetes and hypertension may also be in part a consequence of the HIV-induced inflammatory state [44, 45], so adjustment for these factors may underestimate the true effect of HIV-associated inflammation on clinical events [46]. Furthermore, the effects of lifestyle factors on clinical outcomes might even be modified by HIV infection. For example, one recent study suggested that the relationship between smoking and atherosclerosis is stronger in HIV-infected than HIV–uninfected individuals [47]. Smoking may also cause monocyte activation, inflammation, and hypercoagulability, so it is perhaps not surprising that HIV may amplify its effects on cardiovascular risk [48]. HIV also tended to increase the impact of hypertension on myocardial infarction risk in the VACS, though this interaction did not reach statistical significance [49]. Similarly, moderate exercise has long been known to decrease inflammation and improve both cognitive function and overall health in elderly populations, and has recently been shown to decrease inflammation in a pilot trial of much younger HIV-infected individuals [50]. Thus, lifestyle changes may have an even more important impact on health in HIV-infected individuals than in the general population, in part because they decrease the inflammatory state.

Definitive evidence that inflammation causes excess morbidity and mortality in treated HIV infection will only come from a large clinical trial of a specific immune-based intervention that reduces clinical events in this setting. Unfortunately, few candidate interventions to date are both scalable and have promising enough immunologic effects in pilot studies to warrant investing in expensive clinical endpoint trials. A clinical endpoint trial of statins is being considered in treated HIV infection based on observational studies suggesting benefit [51–53], and clinical trials demonstrating favorable immunologic effects in HIV infection [54, 55]. Nevertheless, even if statins are shown to decrease cardiovascular events in HIV-infected individuals who don't otherwise meet criteria for statin therapy, questions might remain about the extent to which reductions in clinical events are due to declines in inflammation or cholesterol, as they have following statin trials in the general population [56]. The inferences might be more compelling if non-cardiovascular endpoints are also decreased by statins (e.g., cancer, bone fractures, etc). Indeed, rosuvastatin improved bone mineral density in a recent clinical trial of ART-suppressed individuals, though it was not clear whether this was mediated by immunologic mechanisms [57]. Other commonly used medications with anti-inflammatory properties (e.g., aspirin, angiotensin converting enzyme inhibitors, etc) have also shown some promise in HIV-infected individuals in small pilot studies [58, 59], but given cost constraints and research capacity issues, the threshold for advancing these interventions to clinical endpoint trials is likely to be high.

Earlier initiation of ART in the course of HIV infection is also likely to reduce the inflammatory set-point during ART-mediated viral suppression [22, 60, 61], and a clinical trial of early vs. later ART initiation (the Strategic Timing of AntiRetroviral Treatment, START, NCT00867048) is underway. This trial might also help inform whether reducing inflammation (via earlier ART initiation) decreases the risk of non-AIDS events. Nevertheless, inflammation continues to predict morbidity and mortality for at least a decade during treated HIV infection [62], so it is possible that the effects of chronic inflammation on morbidity and mortality may not be evident until trial subjects reach older ages, years after the trial ends. Indeed, the rate of non-AIDS events in the enrolled population is apparently lower than originally anticipated when START was planned [63].

Identifying the most appropriate targets for interventions

While the above interventional strategies currently being pursued are promising, it is unlikely that any of them are likely to completely reverse the persistent inflammatory state of treated HIV infection. Thus, developing more effective interventional strategies is a major focus of the field. Identifying the most appropriate specific targets for interventions to pursue - even in small pilot trials - remains a challenge, however, particularly in an era of constrained resources. Studies linking biomarkers to clinical events have therefore taken on increasing importance to identify those pathways that are most likely to drive disease. Two recent studies determined that soluble markers of innate immune activation predicted non-AIDS events and mortality during ART-mediated viral suppression much more strongly than T cell activation and senescence [64, 65]. These studies suggest that innate immune activation may be a more relevant target for interventions than T cell activation or senescence, but there remains a myriad of potential innate immunologic pathways that could be targeted [27–37].

To better prioritize potential interventional targets, it may be helpful to consider not just the individual relationships between specific biomarkers and clinical outcomes, but also how they fit together in biologic systems in vivo. Just because a particular inflammatory marker is strongly associated with clinical outcomes, it may not necessarily be the most appropriate direct target for interventions. For example, the inflammation reflected by IL-6 levels and coagulation reflected by D-dimer levels may be along the final common pathway linking immune activation to disease, but specifically blocking IL-6 signaling or coagulation might not reverse morbidity and mortality if the underlying causes of the inflammatory state are not addressed as parallel inflammatory pathways may persist or even increase (the “whack-a-mole” theory). Mediation and pathway modeling approaches may well prove useful in drawing inferences from these studies about the most promising interventional targets. Indeed, targeting the root causes of the inflammatory state may prove to be a more sensible strategy, even if the biomarkers that reflect those root causes tend to be less strongly associated with clinical outcomes.

Several root causes of the persistent inflammatory state during suppressive ART have been proposed including HIV itself (either from residual low-level replication [66–69] or passive release from infected cells), co-infections like cytomegalovirus (CMV) or hepatitis C virus (HCV) [70, 71], and microbial translocation [72]. Unfortunately, except for anti-HCV

therapy, we currently lack safe and effective interventions to potentially block these root causes. While ART intensification may decrease the extent of low-level HIV replication in a subset of patients (particularly those for whom penetration of antiretroviral medications into lymphoid tissues may be incomplete [66–69]), its impact on systemic immune activation is modest at best, and no currently available medications block HIV release from infected cells. While drug development for CMV is quite active, we currently lack safe therapies that can be given to ART-suppressed individuals long-term. Furthermore, microbial translocation has proven to be difficult to reverse in HIV-infected individuals [73–76], perhaps since HIV-associated microbial translocation involves defects at multiple levels (e.g., gut epithelial barrier integrity and function, immune surveillance in lamina propria, lymphoid tissues, etc) [77]. To overcome this challenge, one might need to combine synergistic interventions that address each one of these defects simultaneously. Alternatively, simply treating individuals with anti-inflammatory interventions for much longer durations of time may be required to reverse microbial translocation. Indeed, chronic immune activation and inflammation may not only impair immune function, but they may also directly cause gut epithelial barrier defects [78–80]. Just as it appears to take at least six months for systemic microbial translocation to fully ramp up during primary HIV infection [81], it may take several months of reducing inflammation before gut barrier integrity is noticeably restored. Thus, further research is required to develop more effective interventions to reverse the underlying causes of the inflammatory state in treated HIV infection.

Choosing the most appropriate surrogate markers in clinical trials

Just as there remains uncertainty as to the most appropriate interventional targets, the most appropriate surrogate markers to use in pilot trials of immune-based interventions also remains uncertain. The lack of a single stable, reproducible, and reliable surrogate marker for the persistent inflammatory state in treated HIV infection also limits the possibility for novel drug development in this area. Many of the biomarkers that most strongly predict mortality in treated HIV infection (e.g., IL-6, D-dimer, etc.) fluctuate greatly within individuals and typically require excessively large sample sizes for adequate power to detect clinically relevant effects in pilot trials. Conversely, T cell activation, which is much more stable within individuals and has been effectively used as a surrogate outcome in many prior pilot trials, fails to reliably predict clinical outcomes in treated HIV infection [64, 65]. Considering biomarkers in the context of in vivo systems may also be helpful here as some soluble markers of monocyte activation (e.g., sCD14, sCD163) reflect a key source of inflammation and coagulation in HIV infection [82], and predict mortality [83], but are also relatively stable within individuals and can be measured reliably in small pilot trials [84]. Thus, these biomarkers will likely be used more commonly in future pilot trials of immune-based interventions. An alternative approach would be to create an index of several biomarkers predictive of mortality in this setting, as has recently been proposed by the INSIGHT network [62], though more information is needed on the within-subject variability over time for such indices for these approaches to be effectively compared.

Conclusion

As the HIV epidemic ages, the impact of HIV on multi-morbidity and premature mortality is likely to take on increasing public health importance. Understanding the similarities between the multi-morbidity associated with HIV and aging may help frame the clinical management of frail HIV-infected patients, while understanding the differences in the root causes of multi-morbidity in HIV and aging may help better focus our attention on targets for interventions to prevent future morbidity and mortality in HIV-infected individuals. The persistent inflammatory state of treated HIV infection is likely to be an important determinant of morbidity and mortality, and it may even interact with health related behaviors, such that smoking cessation, diet, and exercise may be even more important interventions for HIV-infected patients than the general population. Better understanding the root causes of the persistent inflammatory state in treated HIV infection and developing more effective interventions for these targets will continue to be an important research agenda for the next several years.

Acknowledgments

The author is a consultant for Merck, Bristol Myers Squibb, Gilead, and Tobira Pharmaceuticals, and has received honoraria from Gilead. Dr. Hunt currently receives the following relevant research funding from the NIH (R56 AI100765, R01 MH54907-1S1, BRS-ACURE-S-11-000035-001312, R24 AI067039).

References

- 1*. Samji H, Cescon A, Hogg RS, et al. Closing the Gap: Increases in Life Expectancy among Treated HIV-Positive Individuals in the United States and Canada. *PLoS ONE*. 2013; 8:e81355. This large study suggests that the life expectancy of HIV-infected individuals in the modern era may be approaching that of the general population, but survivorship bias may underestimate the contribution of HIV to mortality. [PubMed: 24367482]
2. van Sighem A, Gras L, Reiss P, Brinkman K, de Wolf F. Life expectancy of recently diagnosed asymptomatic HIV-infected patients approaches that of uninfected individuals. *Aids*. 2010; 24:1527. [PubMed: 20467289]
3. Life expectancy of individuals on combination antiretroviral therapy in high-income countries: a collaborative analysis of 14 cohort studies. *Lancet*. 2008; 372:293–9. [PubMed: 18657708]
4. Lederman MM, Funderburg NT, Sekaly RP, Klatt NR, Hunt PW. Residual Immune Dysregulation Syndrome in Treated HIV infection. *Adv Immunol*. 2013; 119:51–83. [PubMed: 23886064]
5. Schouten, J.; Wit, FW.; Stolte, IG., et al. Comorbidity and ageing in HIV-1 infection: the AGEHIV Cohort Study. In: in the Program and Abstracts from the XIX International AIDS Conference; Washington DC. Abstract #THAB0205
6. Kendall CE, Wong J, Taljaard M, et al. A cross-sectional, population-based study measuring comorbidity among people living with HIV in Ontario. *BMC Public Health*. 2014; 14:161. [PubMed: 24524286]
7. High KP, Brennan-Ing M, Clifford DB, et al. HIV and aging: state of knowledge and areas of critical need for research. A report to the NIH Office of AIDS Research by the HIV and Aging Working Group. *Journal of acquired immune deficiency syndromes*. 2012; 60 (Suppl 1):S1–18. [PubMed: 22688010]
8. Kim DJ, Westfall AO, Chamot E, et al. Multimorbidity patterns in HIV-infected patients: the role of obesity in chronic disease clustering. *Journal of acquired immune deficiency syndromes*. 2012; 61:600–5. [PubMed: 23023101]
9. Salter ML, Lau B, Go VF, Mehta SH, Kirk GD. HIV infection, immune suppression, and uncontrolled viremia are associated with increased multimorbidity among aging injection drug

users. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2011; 53:1256–64. [PubMed: 21976463]

10. Brooks JT, Buchacz K, Gebo KA, Mermin J. HIV infection and older Americans: the public health perspective. *American journal of public health*. 2012; 102:1516–26. [PubMed: 22698038]
11. Mills EJ, Barnighausen T, Negin J. HIV and aging--preparing for the challenges ahead. *The New England Journal of Medicine*. 2012; 366:1270–3. [PubMed: 22475591]
12. Wester CW, Koethe JR, Shepherd BE, et al. Non-AIDS-defining events among HIV-1-infected adults receiving combination antiretroviral therapy in resource-replete versus resource-limited urban setting. *Aids*. 2011; 25:1471–9. [PubMed: 21572309]
13. Lewden C, Bouteloup V, De Wit S, et al. All-cause mortality in treated HIV-infected adults with CD4 $\geq$ 500/mm³ compared with the general population: evidence from a large European observational cohort collaboration{dagger}. *Int J Epidemiol*. 2012; 41:433–445. [PubMed: 22493325]
14. van Sighem AI, Gras LA, Reiss P, Brinkman K, de Wolf F. Life expectancy of recently diagnosed asymptomatic HIV-infected patients approaches that of uninfected individuals. *Aids*. 2010; 24:1527–35. [PubMed: 20467289]
- 15*. Rodger AJ, Lodwick R, Schechter M, et al. Mortality in well controlled HIV in the continuous antiretroviral therapy arms of the SMART and ESPRIT trials compared with the general population. *Aids*. 2013; 27:973–979. This study suggests that mortality rates among ART-suppressed patients who initiated therapy at earlier disease stages may approach that of the general population, but as these are clinical trial subjects, may be selected for a healthier population. [PubMed: 23698063]
- 16**. Freiberg MS, Chang CC, Kuller LH, et al. HIV Infection and the Risk of Acute Myocardial Infarction. *JAMA Intern Med*. 2013;1–9. This is perhaps the strongest study to date to establish that HIV increases the risk of cardiovascular disease independent of traditional cardiovascular risk factors.
17. Triant VA, Lee H, Hadigan C, Grinspoon SK. Increased acute myocardial infarction rates and cardiovascular risk factors among patients with human immunodeficiency virus disease. *J Clin Endocrinol Metab*. 2007; 92:2506–12. [PubMed: 17456578]
18. Tseng ZH, Secemsky EA, Dowdy D, et al. Sudden cardiac death in patients with human immunodeficiency virus infection. *Journal of the American College of Cardiology*. 2012; 59:1891–6. [PubMed: 22595409]
19. Shiels MS, Pfeiffer RM, Engels EA. Age at cancer diagnosis among persons with AIDS in the United States. *Ann Intern Med*. 2010; 153:452–60. [PubMed: 20921544]
20. Smitten AL, Simon TA, Hochberg MC, Suissa S. A meta-analysis of the incidence of malignancy in adult patients with rheumatoid arthritis. *Arthritis Research and Therapy*. 2008; 10:R45. [PubMed: 18433475]
- 21*. Lee SA, Sinclair E, Hatano H, et al. Impact of HIV on CD8+ T Cell CD57 Expression Is Distinct from That of CMV and Aging. *PLoS ONE*. 2014; 9:e89444. This study demonstrated that the classic T cell senescence marker CD57, which is typically increased on effector CD8+ T cells in aging and CMV infection, is actually inappropriately low on effector CD8+ T cells in HIV infection, highlighting immunologic differences between HIV and aging. [PubMed: 24586783]
- 22**. Lee SA, Sinclair E, Jain V, et al. Low Proportions of CD28- CD8+ T cells Expressing CD57 Can Be Reversed by Early ART Initiation and Predict Mortality in Treated HIV Infection. *The Journal of Infectious Diseases*. 2014 This study demonstrated that low –not high - CD57 expression on effector CD8+ T cells predicts mortality in treated HIV infection, a qualitatively different association than observed in elderly HIV-negative populations, suggesting distinct immunologic pathways driving morbidity and mortality in these settings.
23. Lederman MM, Calabrese L, Funderburg NT, et al. Immunologic failure despite suppressive antiretroviral therapy is related to activation and turnover of memory CD4 cells. *J Infect Dis*. 2011; 204:1217–26. [PubMed: 21917895]
24. Hunt PW, Martin JN, Sinclair E, et al. T cell activation is associated with lower CD4+ T cell gains in human immunodeficiency virus-infected patients with sustained viral suppression during antiretroviral therapy. *J Infect Dis*. 2003; 187:1534–43. [PubMed: 12721933]

25. Neuhaus J, Jacobs DR Jr, Baker JV, et al. Markers of inflammation, coagulation, and renal function are elevated in adults with HIV infection. *J Infect Dis.* 2010; 201:1788–95. [PubMed: 20446848]
26. French MA, King MS, Tschampa JM, da Silva BA, Landay AL. Serum immune activation markers are persistently increased in patients with HIV infection after 6 years of antiretroviral therapy despite suppression of viral replication and reconstitution of CD4+ T cells. *J Infect Dis.* 2009; 200:1212–5. [PubMed: 19728788]
27. Boulware DR, Hullsiek KH, Puroon CE, et al. Higher levels of CRP, D-dimer, IL-6, and hyaluronic acid before initiation of antiretroviral therapy (ART) are associated with increased risk of AIDS or death. *J Infect Dis.* 2011; 203:1637–46. [PubMed: 21592994]
- 28*. Byakwaga H II, YB, Huang Y, et al. The Kynurenine Pathway of Tryptophan Catabolism, CD4+ T Cell Recovery and Mortality in HIV-infected Ugandans Initiating ART. *J Infect Dis.* 2014 in press. This study is the first in the world to link the kynurenine pathway of tryptophan catabolism to mortality in treated HIV infection and may highlight a novel interventional target.
29. Duprez DA, Neuhaus J, Kuller LH, et al. Inflammation, coagulation and cardiovascular disease in HIV-infected individuals. *PLoS ONE.* 2012; 7:e44454. [PubMed: 22970224]
30. Ford ES, Greenwald JH, Richerman AG, et al. Traditional risk factors and D-dimer predict incident cardiovascular disease events in chronic HIV infection. *AIDS.* 2010; 24:1509–17. [PubMed: 20505494]
31. Justice AC, Freiberg MS, Tracy R, et al. Does an Index Composed of Clinical Data Reflect Effects of Inflammation, Coagulation, and Monocyte Activation on Mortality Among Those Aging With HIV? *Clin Infect Dis.* 2012; 54:984–994. [PubMed: 22337823]
32. Kuller LH, Tracy R, Belloso W, et al. Inflammatory and coagulation biomarkers and mortality in patients with HIV infection. *PLoS Med.* 2008; 5:e203. [PubMed: 18942885]
33. Ledwaba L, Tavel JA, Khabo P, et al. Pre-ART levels of inflammation and coagulation markers are strong predictors of death in a South African cohort with advanced HIV disease. *PLoS ONE.* 2012; 7:e24243. [PubMed: 22448211]
34. Tien PC, Choi AI, Zolopa AR, et al. Inflammation and mortality in HIV-infected adults: analysis of the FRAM study cohort. *J Acquir Immune Defic Syndr.* 2010; 55:316–22. [PubMed: 20581689]
- 35**. Erlandson KM, Allshouse AA, Jankowski CM, et al. Association of Functional Impairment with Inflammation and Immune Activation in HIV-1-Infected Adults on Effective Antiretroviral Therapy. *The Journal of Infectious Diseases.* 2013 This study demonstrated that the frailty phenotype is strongly associated with immune activation and inflammation in treated HIV infection.
- 36**. Borges AH, Silverberg MJ, Wentworth D, et al. Predicting risk of cancer during HIV infection: the role of inflammatory and coagulation biomarkers. *Aids.* 2013; 27:1433–41. This study provides the strongest evidence to date linking inflammation to subsequent non-AIDS cancer risk in HIV infection. [PubMed: 23945504]
- 37*. Bjerk SM, Baker JV, Emery S, et al. Biomarkers and bacterial pneumonia risk in patients with treated HIV infection: a case-control study. *PLoS ONE.* 2013; 8:e56249. This is the first study to link inflammatory biomarkers to the risk of subsequent bacterial pneumonia in treated HIV infection, one of the more common morbidities affecting this population. [PubMed: 23457535]
38. Harris TB, Ferrucci L, Tracy RP, et al. Associations of elevated interleukin-6 and C-reactive protein levels with mortality in the elderly. *Am J Med.* 1999; 106:506–12. [PubMed: 10335721]
39. Solomon DH, Karlson EW, Rimm EB, et al. Cardiovascular morbidity and mortality in women diagnosed with rheumatoid arthritis. *Circulation.* 2003; 107:1303–1307. [PubMed: 12628952]
40. Han C, Robinson DW, Hackett MV, Paramore LC, Fraeman KH, Bala MV. Cardiovascular disease and risk factors in patients with rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis. *The Journal of rheumatology.* 2006; 33:2167–2172. [PubMed: 16981296]
41. Gelfand JM, Berlin J, Van Voorhees A, Margolis DJ. Lymphoma rates are low but increased in patients with psoriasis: results from a population-based cohort study in the United Kingdom. *Archives of dermatology.* 2003; 139:1425–1429. [PubMed: 14623702]
42. Deodhar A, Woolf A. Bone mass measurement and bone metabolism in rheumatoid arthritis: a review. *Rheumatology.* 1996; 35:309–322.

43. Dreiherr J, Weitzman D, Cohen AD. Psoriasis and Osteoporosis: A Sex-Specific Association? *Journal of Investigative Dermatology*. 2009; 129:1643–1649. [PubMed: 19158845]
44. Brown TT, Tassiopoulos K, Bosch RJ, Shikuma C, McComsey GA. Association between systemic inflammation and incident diabetes in HIV-infected patients after initiation of antiretroviral therapy. *Diabetes Care*. 2010; 33:2244–9. [PubMed: 20664016]
45. Savoia C, Schiffrin EL. Inflammation in hypertension. *Current opinion in nephrology and hypertension*. 2006; 15:152–158. [PubMed: 16481882]
46. Armah KA, McGinnis K, Baker J, et al. HIV status, burden of comorbid disease, and biomarkers of inflammation, altered coagulation, and monocyte activation. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2012; 55:126–36. [PubMed: 22534147]
- 47*. Fitch KV, Looby SE, Rope A, et al. Effects of aging and smoking on carotid intima-media thickness in HIV-infection. *Aids*. 2013; 27:49–57. This study suggests that the effect of smoking on atherosclerosis may be even greater in HIV infection than it is in the general population, highlighting the importance of smoking cessation for HIV-infected individuals. [PubMed: 22874518]
48. Yang S-R, Chida AS, Bouter MR, et al. Cigarette smoke induces proinflammatory cytokine release by activation of NF- κ B and posttranslational modifications of histone deacetylase in macrophages. *American Journal of Physiology-Lung Cellular and Molecular Physiology*. 2006; 291:L46–L57. [PubMed: 16473865]
- 49*. Armah KA, Chang CC, Baker JV, et al. Prehypertension, hypertension, and the risk of acute myocardial infarction in HIV-infected and -uninfected veterans. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2014; 58:121–9. While not formally statistically significant, this study appears to suggest that the influence of hypertension on morbidity and mortality may be even greater in HIV infection than it is in uninfected individuals. [PubMed: 24065316]
- 50*. Longo, V.; Bonato, M.; Bossolasco, S., et al. Brisk Walking Improves Inflammatory Markers in cART-Treated Patients. In: in the Program and Abstracts of the 2014 Conference on Retroviruses and Opportunistic Infections; Boston, MA. Abstract #763. While uncontrolled, this trial suggests that even moderate exercise may lead to significant declines in inflammation in treated HIV infection
- 51*. Overton ET, Kitch D, Benson CA, et al. Effect of Statin Therapy in Reducing the Risk of Serious Non-AIDS-Defining Events and Nonaccidental Death. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2013 This study suggested a potential positive impact of statins on morbidity and mortality in treated HIV infection.
52. Moore RD, Bartlett JG, Gallant JE. Association between use of HMG CoA reductase inhibitors and mortality in HIV-infected patients. *PLoS ONE*. 2011; 6:e21843. [PubMed: 21765919]
53. Chao C, Xu L, Abrams DI, et al. HMG-CoA reductase inhibitors (statins) use and risk of non-Hodgkin lymphoma in HIV-positive persons. *Aids*. 2011; 25:1771–7. [PubMed: 21681055]
- 54*. Funderburg NT, Jiang Y, Debanne SM, et al. Rosuvastatin treatment reduces markers of monocyte activation in HIV-infected subjects on antiretroviral therapy. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2014; 58:588–95. This study provides the strongest evidence to date that statins decrease monocyte activation in treated HIV infection. [PubMed: 24253250]
55. Ganesan A, Crum-Cianflone N, Higgins J, et al. High Dose Atorvastatin Decreases Cellular Markers of Immune Activation Without Affecting HIV-1 RNA Levels: Results of a Double-blind Randomized Placebo Controlled Clinical Trial. *J Infect Dis*. 2011; 203:756–64. [PubMed: 21325137]
56. Ridker PM. Testing the inflammatory hypothesis of atherothrombosis: scientific rationale for the cardiovascular inflammation reduction trial (CIRT). *J Thromb Haemost*. 2009; 7 (Suppl 1):332–9. [PubMed: 19630828]
- 57*. McComsey, GA.; Jiang, Y.; Erlandson, KM.; Debanne, SM. Rosuvastatin Improves Hip Bone Mineral Density But Worsens Insulin Resistance. In: in the Program and Abstracts of the 2014 Conference on Retroviruses and Opportunistic Infections; Boston, MA. Abstract #134. This study suggests that statins may improve bone mineral density in addition to reducing immune

activation and inflammation in treated HIV infection, but it remains unclear whether these effects are linked

- 58*. O'Brien M, Montenont E, Hu L, et al. Aspirin attenuates platelet activation and immune activation in HIV-infected subjects on antiretroviral therapy: A Pilot Study. *Journal of acquired immune deficiency syndromes*. 2013 While uncontrolled, this study suggests substantially beneficial effects of low-dose aspirin on immune activation in treated HIV infection and provides strong rationale for a randomized controlled trial in this setting.
59. Baker JV, Huppler Hullsiek K, Prosser R, et al. Angiotensin converting enzyme inhibitor and HMG-CoA reductase inhibitor as adjunct treatment for persons with HIV infection: a feasibility randomized trial. *PLoS ONE*. 2012; 7:e46894. [PubMed: 23082133]
60. Burdo TH, Lentz MR, Autissier P, et al. Soluble CD163 made by monocyte/macrophages is a novel marker of HIV activity in early and chronic infection prior to and after anti-retroviral therapy. *J Infect Dis*. 2011; 204:154–63. [PubMed: 21628670]
- 61*. Jain V, Hartogensis W, Bacchetti P, et al. Antiretroviral Therapy Initiated within Six Months of HIV Infection is Associated with Lower T-cell Activation and Smaller HIV Reservoir Size. *The Journal of Infectious Diseases*. 2013 This study suggests that earlier initiation of ART in the course of HIV infection may result in a lower immune activation setpoint once viral suppression is achieved.
- 62*. Grund, B.; Baker, J.; Deeks, S., et al. Combined Effect of Interleukin-6 and D-dimer on the Risk of Serious Non-AIDS Conditions: Data from 3 Prospective Cohorts. In: in the Program and Abstracts from the 20th Conference on Retroviruses and Opportunistic Infections; Atlanta, GA. Abstract #60. This study suggests that IL-6 and D-dimer levels continue to predict morbidity and mortality for at least 10 years of ART-mediated viral suppression, suggesting that the inflammatory setpoint during ART may have long-lasting effects, well into the future
63. Babiker AG, Emery S, Fätkenheuer G, et al. Considerations in the rationale, design and methods of the Strategic Timing of AntiRetroviral Treatment (START) study. *Clinical Trials*. 2013; 10:S5–S36. [PubMed: 22547421]
64. Hunt, P.; Rodriguez, B.; Shive, C., et al. Gut Epithelial Barrier Dysfunction, Inflammation, and Coagulation Predict Higher Mortality during Treated HIV/AIDS. In: in the Program and Abstracts from the 19th Conference on Retroviruses and Opportunistic Infections; Seattle, WA. Abstract #278
- 65*. Tenorio, A.; Zheng, E.; Bosch, R., et al. Soluble Markers of Inflammation & Coagulation, but not T-Cell Activation, Predict Non-AIDS-Defining Events During Suppressive Antiretroviral Therapy. In: in the Program and Abstracts of the 20th Conference on Retroviruses and Opportunistic Infections; 2013; Atlanta, GA. Abstract #790. This study suggested that innate immune activation markers are stronger predictors of mortality than T cell activation or senescence in treated HIV infection
66. Buzon MJ, Massanella M, Llibre JM, et al. HIV-1 replication and immune dynamics are affected by raltegravir intensification of HAART-suppressed subjects. *Nat Med*. 2010; 16:460–5. [PubMed: 20228817]
- 67*. Hatano H, Strain MC, Scherzer R, et al. Increase in 2-Long Terminal Repeat Circles and Decrease in D-dimer After Raltegravir Intensification in Patients With Treated HIV Infection: A Randomized, Placebo-Controlled Trial. *The Journal of Infectious Diseases*. 2013; 208:1436–1442. This study confirmed an earlier study by Buzon et al, demonstrating that low-level re-infection of CD4+ T cells is likely occurring in many HIV-infected patients despite apparently undetectable plasma HIV RNA levels on clinical assays, potentially re-seeding the viral reservoir and contributing to immune activation. [PubMed: 23975885]
68. Yukl SA, Shergill AK, McQuaid K, et al. Effect of raltegravir-containing intensification on HIV burden and T-cell activation in multiple gut sites of HIV-positive adults on suppressive antiretroviral therapy. *Aids*. 2010; 24:2451–60. [PubMed: 20827162]
- 69**. Fletcher, CV.; Staskus, K.; Wietgreffe, SW., et al. Persistent HIV-1 replication is associated with lower antiretroviral drug concentrations in lymphatic tissues. *Proceedings of the National Academy of Sciences of the United States of America*; 2014. This study suggests that many commonly used antiretroviral medications may not adequately penetrate lymphoid tissues, where

most HIV replication is thought to occur, providing a mechanism to explain persistent low-level re-infection or replication even when plasma HIV RNA levels are clinically undetectable

70. Hunt PW, Martin JN, Sinclair E, et al. Valganciclovir reduces T cell activation in HIV-infected individuals with incomplete CD4+ T cell recovery on antiretroviral therapy. *J Infect Dis.* 2011; 203:1474–83. [PubMed: 21502083]
71. Gonzalez VD, Falconer K, Blom KG, et al. High levels of chronic immune activation in the T-cell compartments of patients coinfectd with hepatitis C virus and human immunodeficiency virus type 1 and on highly active antiretroviral therapy are reverted by alpha interferon and ribavirin treatment. *J Virol.* 2009; 83:11407–11. [PubMed: 19710147]
72. Brenchley JM, Price DA, Schacker TW, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med.* 2006; 12:1365–1371. [PubMed: 17115046]
73. Somsouk, M.; Dunham, R.; Cohen, M., et al. Mesalamine to Reduce Immune Activation during HIV Infection: A Randomized Controlled Trial. In: in the Program and Abstracts of the 2014 Conference on Retroviruses and Opportunistic Infections; Boston, MA. March 3–6, 2014; Abstract #841
74. Sandler NG.; Zhang, X.; Bosch, RJ., et al. Sevelamer Does Not Decrease Plasma LPS or sCD14 But Does Decrease Soluble Tissue Factor and LDL. In: in the Program and Abstracts from the 2014 Conference on Retroviruses and Opportunistic Infections; Boston, MA. Abstract #337
75. Tenorio, AR.; Wilson, CC.; Chan, ES., et al. Rifaximin Has Marginal Impact On Immune Activation in Immune Non-Responders To ART - ACTG 5286. In: in the Program and Abstracts from the 2014 Conference on Retroviruses and Opportunistic Infections; Boston, MA. Abstract #339
76. Stiksrud, B.; Nowak, P.; Kvale, D., et al. Decreased Levels of D-Dimer After Probiotic Supplementation in Patients Receiving ART. In: in the Program and Abstracts from the 2014 Conference on Retroviruses and Opportunistic Infections; Boston, MA. Abstract #342
77. Sandler NG, Douek DC. Microbial translocation in HIV infection: causes, consequences and treatment opportunities. *Nature reviews Microbiology.* 2012; 10:655–66.
78. Favre D, Mold J, Hunt PW, et al. Tryptophan catabolism by indoleamine 2,3-dioxygenase 1 alters the balance of TH17 to regulatory T cells in HIV disease. *Science Translational Medicine.* 2010; 2:32ra36.
79. Feng Y, Teitelbaum DH. Tumour necrosis factor--induced loss of intestinal barrier function requires TNFR1 and TNFR2 signalling in a mouse model of total parenteral nutrition. *J Physiol.* 2013; 591:3709–23. [PubMed: 23753529]
80. Petecchia L, Sabatini F, Usai C, Caci E, Varesio L, Rossi GA. Cytokines induce tight junction disassembly in airway cells via an EGFR-dependent MAPK/ERK1/2-pathway. *Laboratory investigation; a journal of technical methods and pathology.* 2012; 92:1140–8.
81. Chevalier MF, Petitjean G, Dunyach-Remy C, et al. The Th17/Treg ratio, IL-1RA and sCD14 levels in primary HIV infection predict the T-cell activation set point in the absence of systemic microbial translocation. *PLoS pathogens.* 2013; 9:e1003453. [PubMed: 23818854]
82. Funderburg NT, Mayne E, Sieg SF, et al. Increased tissue factor expression on circulating monocytes in chronic HIV infection: relationship to in vivo coagulation and immune activation. *Blood.* 2010; 115:161–7. [PubMed: 19828697]
83. Sandler NG, Wand H, Roque A, et al. Plasma levels of soluble CD14 independently predict mortality in HIV infection. *The Journal of Infectious Diseases.* 2011; 203:780–90. [PubMed: 21252259]
84. Hunt PW, Shulman NS, Hayes TL, et al. The immunologic effects of maraviroc intensification in treated HIV-infected individuals with incomplete CD4+ T-cell recovery: a randomized trial. *Blood.* 2013; 121:4635–46. [PubMed: 23589670]

Key Points

- HIV is associated with an increased risk of multi-morbidity and mortality, which is likely to take on increasing public health importance as the HIV epidemic gets older.
- While the clinical phenotype of HIV-associated multi-morbidity shares many features with aging-related frailty, its root causes may in fact be distinct, requiring different interventions for prevention.
- Persistent inflammation is likely to be a major driver of morbidity and mortality in treated HIV infection, and a better understanding of its root causes may help prioritize targets for interventions.
- Recent studies have also helped shape our understanding of the most appropriate surrogate biomarkers to employ in pilot studies of immune-based interventions.