Published in final edited form as:

J Int Neuropsychol Soc. 2014 August ; 20(7): 756–763. doi:10.1017/S1355617714000423.

Relationship Between Depressive Symptoms and Cognition in Older, Non-demented African Americans

Jamie L. Hamilton^{1,2}, Adam M. Brickman^{1,3,4}, Rosalyn Lang⁵, Goldie S. Byrd⁶, Jonathan L. Haines⁷, Margaret A. Pericak-Vance⁸, and Jennifer J. Manly^{1,3,4}

¹Taub Institute for Research on Alzheimer's Disease and the Aging Brain, College of Physicians and Surgeons, Columbia University, New York, New York

²Department of Psychiatry, College of Physicians and Surgeons, Columbia University, New York, New York

³G.H. Sergievsky Center, College of Physicians and Surgeons, Columbia University, New York, New York

⁴Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, New York

⁵North Carolina Agricultural and Technical State University, Department of Biology, Greensboro, North Carolina

⁶North Carolina Agricultural and Technical State University, Dean, College of Arts and Sciences, Greensboro, North Carolina

⁷Center for Human Genetics Research, Vanderbilt University, Nashville, Tennessee

⁸John P. Hussman Institute for Human Genomics, Miller School of Medicine, University of Miami, Miami, Florida

Abstract

Knowledge of the relationship between depressive symptoms and cognition in older adults has primarily come from studies of clinically depressed, functionally impaired or cognitively impaired individuals, and in predominately White samples. Limited minority representation in depression research exposes the need to examine these associations in more ethnic/racially diverse populations. We sought to examine the relationship between depressive symptoms and cognition in a sample of non-demented older African Americans recruited from surrounding U.S. cities of New York, Greensboro, Miami, and Nashville ($N = 944$). Depressive symptoms were evaluated with the Geriatric Depression Scale (GDS). Cognition was evaluated with a comprehensive neuropsychological battery. Test scores were summarized into attention, executive function, memory, language, and processing speed composites. Controlling for age, education, reading level, and sex, African American older adults who endorsed more symptoms obtained significantly lower scores on measures of memory, language, processing speed, and executive

Copyright © INS. Published by Cambridge University Press, 2014.

Correspondence and reprint requests to: Jennifer Manly, Columbia University, Taub Institute for Research on Alzheimer's Disease, and the Aging Brain, College of Physicians and Surgeons, PH Building, Suite 19-321, 630 W 168th Street, New York, NY 10032. jjm71@columbia.edu.

functioning. Further investigation of the causal pathway underlying this association, as well as potential mediators of the relationship between depressive symptoms and cognitive test performance among older African Americans, such as cardiovascular and cerebrovascular disease, may offer potential avenues for intervention.

Keywords

Aging; Ethnic groups; Depression; Executive function; Memory; Language

INTRODUCTION

The relationship between depressive symptoms and cognition has received increased attention in recent cognitive aging literature. Depressive symptoms are less frequent in older adults compared with younger adults (Fiske, Wetherell, & Gatz, 2009; Hasin, Goodwin, Stinson, & Grant, 2005). However, at higher levels of endorsement, they are linked to cognitive decline, greater functional disability, and mortality (Blazer, 2003; Ganguli, Snitz, Vander Bilt, & Chang, 2009; Lockwood, Alexopoulos, & van Gorp, 2002; Wilson et al., 2002). It remains unclear whether the association of depressive symptoms and cognitive function is due to a common cause, such as underlying cerebrovascular pathology; whether depressive symptoms appear in a reaction to the onset of cognitive decline (Steffens et al., 2006), or whether depressive symptoms simply exacerbate cognitive impairment (Horwath, Johnson, Klerman, & Weissman, 1992; Jorm, 2001; Ownby, Crocco, Acevedo, John, & Loewenstein, 2006). Causal and temporal explanations of the relationship between depression and cognition have spurred interest in examining depressive symptoms at varying levels of severity, particularly among non-depressed people who endorse few or very mild depressive symptoms.

In depressed older adults, poorer cognition has been reported in the domains of executive functioning, verbal memory, learning, and processing speed in comparison to healthy controls (Austin, Mitchell, & Goodwin, 2001; Boone et al., 1994; Comijs, Jonker, Beekman, & Deeg, 2001; Nebes et al., 2000; Stoudemire, Hill, Morris, Martino-Saltzman, & Lewison, 1993). Depressive symptoms that fall below clinical diagnostic criteria are often described as subthreshold or subsyndromal (Cohen, Magai, Yaffee, & Walcott-Brown, 2005; Pincus, Davis, & McQueen, 1999) and are shown to predict cognition (Dotson, Resnick, & Zonderman, 2008; Elderkin-Thompson et al., 2003; Lichtenberg, Ross, Millis, & Manning, 1995; Ravdin, Mattis, & Lachs, 2004). Subsyndromal and subthreshold phenomenology studies with significant or primarily African American representation have provided some cross-sectional and longitudinal evidence of depressive symptoms and cognition associations yet vary in the extent of neuropsychological measures used (Adams & Moon, 2009; Burt, Zembar, & Niederehe, 1995; Cohen et al., 2005; Wilson, Mendes De Leon, Bennett, Bienias, & Evans, 2004).

Despite evidence of higher lifetime major depression disorder (MDD) prevalence rates among Whites in comparison to African Americans and Hispanics in epidemiological and community-based samples (Blazer, Landerman, Hays, Simonsick, & Saunders, 1998;

Kessler et al., 2003), African Americans tend to have greater chronicity of depressive symptoms (Williams et al., 2007). Greater chronicity of depressive symptoms coupled with higher prevalence of vascular disease among African Americans over 65 in comparison to other racial groups (Roger et al., 2012) could amplify susceptibility to cognitive decline, cognitive impairment, and dementia risk (Barnes, Alexopoulos, Lopez, Williamson, & Yaffe, 2006; Fillenbaum, Peterson, Welsh-Bohmer, Kukull, & Heyman, 1998; Ogunniyi et al., 2006; Tang et al., 2001; Unverzagt et al., 2001).

It is not well established whether minimal endorsement of depressive symptoms, routinely recognized as non-clinically relevant, is related to performance in specific cognitive domains in older African Americans. Because depressive symptoms often go undetected in African Americans (Gallo, Rabins, Lyketsos, Tien, & Anthony, 1997; Lebowitz et al., 1997; Miller et al., 2004), it is important to examine whether low-level endorsement of depressive symptoms is predictive of cognition among older African Americans. The aim of this study was to examine whether depressive symptoms are related to cognition in a large, cross-sectional cohort of non-demented older African Americans. Our study is unique among prior studies of African American older adults (Espiritu et al., 2001; Gitlin, Hauck, Dennis, & Schulz, 2007; Livney et al., 2011; Lopez et al., 2003; McDougall, Morgan, & Vaughan, 2012; Rovner, Casten, & Leiby, 2012) in that we examined the relationship between depressive symptoms and cognition in a large sample without significant cognitive or functional impairment and used a comprehensive neuropsychological assessment battery. We hypothesized that depressive symptoms would be independently associated with lower scores on measures of executive functioning, memory and attention.

METHODS

Individuals were recruited for a case-control study to identify genes associated with Alzheimer's disease (AD) among African Americans. This multi-site study included research teams from Columbia University, North Carolina A&T State University, Vanderbilt University, and the University of Miami. Participants were recruited from surrounding areas of New York City, Greensboro, Miami, and Nashville. Self-reported ethnicity and race were based on U.S. Census criteria (U.S. Census Bureau, 2001). All study-related procedures, including recruitment, informed consent, diagnostic, medical, neuropsychological, and neurological evaluation, were approved by the Institutional Review Boards of participating sites. Eligibility for the current study was restricted to those who were identified as non-demented and had available neuropsychological and self-reported depression data across all participating sites.

Subjects

Eligible participants were aged 60 and over, self-identified as non-Hispanic and Black or African American, and endorsed English as their primary language. Individuals with a past history of psychosis, epilepsy, electroconvulsive therapy, Parkinson's disease, or Huntington's disease were excluded. Older adults with a history of MDD, as determined from a medical and psychiatric history questionnaire of symptoms, clinical diagnosis, and treatment, were not excluded. Diagnoses of dementia (American Psychiatric Association,

1994), Alzheimer's Disease (McKhann et al., 2011), vascular dementia (Roman et al., 1993), mild cognitive impairment (Petersen et al., 1999), and cognitive impairment (not MCI), and healthy control status were made according to standard research criteria and determined by consensus case conference that considered the information gathered from neurological, neuropsychological, medical, and functional evaluations.

A total of 1468 individuals were reviewed at consensus conference. Of those, 300 were categorized as AD, 3 were demented with undetermined etiology, and 3 were diagnosed as probable vascular dementia. Of the 1162 non-demented cases, 217 were not administered the 15-item Geriatric Depression Scale (GDS) (Yesavage 1983) due to differences in procedures across performance sites. Nevertheless, there was no significant difference across those who were and were not administered the GDS in age ($M = 69.42$, $SD = 6.9$ with GDS vs. $M = 69.40$; $SD = 7.63$ without GDS; $t = .0398$; $p = .96$), or years of school ($M = 13.87$; $SD = 2.99$ with GDS vs. $M = 13.35$; $SD = 6.904$ without GDS; $t = 1.718$; $p = .09$). Of the remaining 945 participants, 1 person was excluded because they did not complete enough of the neuropsychological test battery to be able to compute at least one composite score.

Depressive symptoms were ascertained from the 15-item self-rated GDS. The yes/no response format was adapted from the original, 30-item scale and is a frequently used, reliable, and valid measure of depression symptoms in cognitively intact (Norris, Gallagher, Wilson, & Winograd, 1987), cognitively impaired (Gerety et al., 1994), and African American (Pedraza, Dotson, Willis, Graff-Radford, & Lucas, 2009) older adults.

Neuropsychological Evaluation

A comprehensive neuropsychological battery that evaluated memory, language, processing speed, executive functioning, and attention was administered to participants. The battery comprised the following tests: California Verbal Learning Test-2 (CVLT-II; Trials 1–5 free recall, short delayed free recall, long delayed free recall, recognition hits-false positives) (Delis, Kramer, Kaplan, & Ober, 2000); 30-item Boston Naming Test (Kaplan, Goodglass, & Weintraub, 1983; Weintraub et al., 2009); Digit Symbol subtest of the Wechsler Adult Intelligence Scale – Revised (Wechsler, 1997); Trail-Making Test Parts A and B (Reitan, 1978); Digit Span Forward and Backward (Wechsler, 1997); Category Fluency Test (animal and vegetable naming) (Goodglass & Kaplan, 1983; Morris et al., 1989).

Exploratory and factor analysis of the neuropsychological test battery revealed five factors, and confirmatory factor analysis revealed structural and metric invariance of these factors across age, education, and diagnostic groups (Miloyan et al., in preparation). To create composite scores for each cognitive domain, Z scores were calculated and averaged from the raw neuropsychological data for each test within each factor. Domain composite scores were grouped as follows: California Verbal Learning Test-2 (CVLT-II; Trials 1–5 free recall, short delayed free recall, long delayed free recall, recognition hits-false positives) comprised the memory composite; 30-item Boston Naming Test, Letter and Category Fluency Test (animal and vegetable naming) were included in the language composite; the processing speed composite included time to complete Trail-Making Test Parts A and Digit Symbol Forward; the attention composite included Digit Span Forward and Backward; and the

executive functioning score was the residual of time to complete Trails A regressed onto time to complete Trails B. For all composites, higher scores reflect better performance.

Statistical Analysis

All analyses were conducted using SPSS version 19 (IBM Corp., Amonk, NY). To compare demographics across diagnostic groups (normal control vs. cognitively impaired, not demented) analysis of variance was used for continuous variables and χ^2 analysis for proportional data. We used five separate linear regression analyses to evaluate whether depressive symptoms were related to cognitive performance with each of the cognitive domains. In the unadjusted models, GDS scores were entered as a continuous predictor of cognitive test performance. The adjusted regression models included age, sex, number of years of education, and reading level WRAT-3 (WRAT-3; Wilkinson, 1993), as covariates. A value of 0 was assigned to men and a value of 1 was assigned to women for the categorical variable of sex. A heterogeneity of slopes analysis was performed to determine if the relationships between depressive symptoms and cognitive test performance differed across those with and without cognitive impairment.

RESULTS

On the GDS, 415 (44%) endorsed no symptoms, 218 (23.1%) endorsed 1 symptom, 109 (11.5%) endorsed 2 symptoms, and 202 (21.4%) endorsed 3 or more symptoms. Descriptive statistics across diagnostic categories are reported in Table 1. There was no significant difference in the proportion of women across groups. Individuals with cognitive impairment were older and endorsed more depressive symptoms compared to normal controls. Normal controls had more years of formal education compared to individuals with cognitive impairment.

In the unadjusted models, participants with more depressive symptomatology obtained lower scores on measures of attention, language, memory, and processing speed. Because of the skewed distribution of GDS scores, we used Spearman's correlation to confirm these associations. Total GDS scores were negatively and significantly correlated with language ($\rho = -.139$; $p < .001$), memory ($\rho = -.089$; $p < .001$), and processing speed ($\rho = -.071$; $p < .05$) but not executive functioning ($\rho = -.040$; $p = .220$) or attention ($\rho = -.065$; $p = .052$).

In the adjusted models, higher GDS scores were associated with lower performance on measures of language, memory, and processing speed (Table 2). The association between GDS scores and executive function approached significance ($p = .05$) once the covariates were included in our model. In contrast, the relationship between depressive symptoms and attention was no longer significant after accounting for the covariates. Reading level was significantly related to all domains of cognitive performance. Women obtained higher scores on all the cognitive composites than men.

A heterogeneity of slopes analysis was performed to determine whether the relationship between depressive symptoms and cognition varied between diagnostic groups. The interaction between diagnostic group and GDS total score was not significant in each of the five analyses for the cognitive domains ($p > .05$ for all). We repeated the adjusted regression

models analyses in samples stratified by cognitive impairment and found that there was not a significant relationship between depressive symptoms and cognitive test performance in any of the stratified groups ($p > .05$ for all).

DISCUSSION

We found that, in a large sample of non-demented older African Americans, depressive symptoms were associated with lower scores on measures of memory, language, attention, and processing speed. Attention was no longer related to depressive symptoms after adjusting for age, sex, years of education, and reading level; however, depressive symptoms remained significantly related to lower scores on measures of memory, language, and processing speed. The relationship of depressive symptoms with a measure of executive function approached significance in the context of the covariates. When we examined these relationships within subgroups divided by presence of cognitive impairment, we found the relationship between depressive symptoms and cognition was no longer significant in any domain or subgroup. This suggests that the relationship between depressive symptoms and cognitive function cannot be detected when the sample does not include the full range of cognitive function within non-demented African Americans. Our study is unique because it is the first to examine depressive symptoms as a continuous predictor of multiple cognitive domains among older non-demented, community-dwelling African Americans. Prior studies have focused on internal consistency study of the GDS self-report measure in older African Americans (Pedraza, Dotson, Willis, Graff-Radford, & Lucas, 2009) and have included medical populations (Kane, Yochim, & Lichtenberg, 2010).

Consistent with prior studies of depressed and remitted depressed older adults (Bassuk, Berkman, & Wypij, 1998; Dal Forno et al., 2005; Dufouil, Fuhrer, Dartiques, & Alperovitch, 1996; Feng, Yap, & Ng, 2013; Godin et al., 2007), we found a significant association between depressive symptoms and language, memory, attention, executive function, and processing speed. One longitudinal study of cognitively intact older women showed that baseline depressive symptoms were associated with later, accelerated declines in global cognition and memory function (Goveas et al., 2014). Cross-sectional associations of depressive symptoms and faster rate of cognitive decline have also been reported in cognitively impaired and AD diagnosed individuals (Devanand et al., 1996; Geerlings et al., 2000; Yaffe, Blackwell, Gore, Sands, Reus, & Browner, 1999; Zahodne, Devanand, & Stern, 2013). Poorer memory, language, and executive function performance in the presence of depressive symptoms may be indicative of greater AD risk within this population.

Several limitations should be noted. First, our study was a study of non-demented African American volunteers for a case-control genetic study. We were unable to compare our results to other ethnic and racial groups, and our participants are not representative of all older African Americans. Second, although the GDS has been shown to be reliable among older African Americans (Pedraza et al., 2009), it is still a self-rated screening instrument that was not based on a structured clinical interview. Third, our findings may not generalize to broader samples with more significant depressive symptoms or more severe cognitive or functional impairment. The cross-sectional nature of our sample is a limitation in that the cause-effect associations between depressive symptoms and cognitive function could not be

determined. Furthermore, we did not examine whether vascular risk factors mediated the relationship between depressive symptoms and cognition. Cardiovascular risk factors such as hypertension, diabetes mellitus, obesity, and diet have been implicated in poorer cognition among African Americans (Antonogeorgos et al., 2012; Sachs-Ericsson et al., 2007; Sims, Madhere, Callender, & Campbell, 2008; Wessels et al., 2011) and may be a common cause of depressive symptoms and cognitive impairment among African Americans. Future work will examine the role of vascular disease within the relationship between depressive symptoms and cognition within this population.

This is the largest study of depressive symptoms and cognition using a comprehensive neuropsychological battery and a well-characterized sample of non-demented African American older adults. It is important to examine possible contributing factors that may alter the course of cognitive aging and may be potential avenues for targeted behavioral and cognition interventions for African Americans, a group that is at higher risk than non-Hispanic Whites for the development of dementia.

Acknowledgments

This research study was supported by National Institutes of Health, National Institute of Aging (grant R01 AG028786) and in part by Columbia University's CTSA grant No. UL1TR000040 from NCATS/NIH and T32 MH020004 training grant in Late Lifespan Neuropsychiatric Disorders Program. Aspects of the data were presented at the 2013 International Neuropsychological Society's meeting in Kona, Hawaii. The authors extend their gratitude to the research assistants and coordinators Raquel Cabo, Josina Hector, Karmen Louie, and Steven Philemond for data collection and processing assistance.

REFERENCES

- Adams KB, Moon H. Subthreshold depression: characteristics and risk factors among vulnerable elders. *Aging Ment Health*. 2009; 13(5):682–692. [PubMed: 19882406]
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th ed. Washington, DC: American Psychiatric Association; 1994.
- Antonogeorgos G, Panagiotakos DB, Pitsavos C, Papageorgiou C, Chrysoshoou C, Papadimitriou GN, Stefanadis C. Understanding the role of depression and anxiety on cardiovascular disease risk, using structural equation modeling; the mediating effect of the Mediterranean diet and physical activity: The ATTICA study. *Annals of Epidemiology*. 2012; 22(9):630–637. [PubMed: 22831995]
- Austin MP, Mitchell P, Goodwin GM. Cognitive deficits in Depression: Possible implications for functional neuropathology. *British Journal of Psychiatry*. 2001; 178:200–206. [PubMed: 11230029]
- Barnes DE, Alexopoulos GS, Lopez OL, Williamson JD, Yaffe K. Depressive symptoms, vascular disease, and mild cognitive impairment: Findings from the Cardiovascular Health Study. *Archives of General Psychiatry*. 2006; 63(3):273–279. [PubMed: 16520432]
- Bassuk SS, Berkman LF, Wypij D. Depressive symptomatology and incident cognitive decline in an elderly community sample. *Archives of General Psychiatry*. 1998; 55:1073–1081. [PubMed: 9862549]
- Blazer DG. Depression in late life: Review and commentary. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*. 2003; 58(3):249–265.
- Blazer DG, Landerman LR, Hays JC, Simonsick EM, Saunders WB. Symptoms of depression among community-dwelling elderly African-American and white older adults. *Psychological Medicine*. 1998; 28(6):1311–1320. [PubMed: 9854272]
- Boone KB, Lesser I, Miller B, Wohl M, Berman N, Lee A, Palmer B. Cognitive functioning in a mildly to moderately depressed geriatric sample: Relationship to chronological age. *The Journal of Neuropsychiatry and Clinical Neurosciences*. 1994; 6(3):267–272. [PubMed: 7950350]

- Burt DB, Zembar MJ, Niederehe G. Depression and memory impairment: A meta-analysis of the association, its pattern, and specificity. *Psychol Bull.* 1995; 117(2):285–305. [PubMed: 7724692]
- Cohen CI, Magai C, Yaffee R, Walcott-Brown L. Racial differences in syndromal and subsyndromal depression in an older urban population. *Psychiatric Services.* 2005; 56(12):1556–1563. [PubMed: 16339618]
- Comijs HC, Jonker C, Beekman AT, Deeg DJ. The association between depressive symptoms and cognitive decline in community-dwelling elderly persons. *International Journal of Geriatric Psychiatry.* 2001; 16(4):361–367. [PubMed: 11333422]
- Dal Forno G, Palermo MT, Donohue JE, Karagiozis H, Zonderman AB, Kawas CH. Depressive symptoms, sex, and risk for Alzheimer's disease. *Annals of Neurology.* 2005; 57:381–387. [PubMed: 15732103]
- Delis, DC.; Kramer, JH.; Kaplan, E.; Ober, BA. California Verbal Learning Test. (2nd ed.). San Antonio, TX: Psychological Corporation; 2000.
- Devanand DP, Sano M, Tang MX, Taylor S, Gurland BJ, Wilder D, Mayeux R. Depressed mood and incidence of Alzheimer's disease in the elderly living in the community. *Archives of General Psychiatry.* 1996; 53:175–182. [PubMed: 8629893]
- Dotson VM, Resnick SM, Zonderman AB. Differential association of concurrent, baseline, and average depressive symptoms with cognitive decline in older adults. *The American Journal of Geriatric Psychiatry.* 2008; 16(4):318–330. [PubMed: 18378557]
- Dufouil C, Fuhrer R, Dartigues JF, Alperovitch A. A Longitudinal analysis of the association between depressive symptomatology and cognitive deterioration. *Journal of Epidemiology.* 1996; 144:634–641.
- Elderkin-Thompson V, Kumar A, Bilker WB, Dunkin JJ, Mintz J, Moberg PJ, Gur RE. Neuropsychological deficits among patients with late-onset minor and major depression. *Archives of Clinical Neuropsychology.* 2003; 18:529–549.
- Espiritu DA, Rashid H, Mast BT, Fitzgerald J, Steinberg J, Lichtenberg PA. Depression, cognitive impairment and function in Alzheimer's disease. *International Journal of Geriatric Psychiatry.* 2001; 16(11):1098–1103. [PubMed: 11746657]
- Feng L, Yap KB, Ng TP. Depressive symptoms in older adults with chronic kidney-disease: Mortality, quality of life outcomes, and correlates. *American Journal of Geriatric Psychiatry.* 2013; 21:570–579. [PubMed: 23567405]
- Fillenbaum GG, Peterson B, Welsh-Bohmer KA, Kukull WA, Heyman A. Progression of Alzheimer's disease in black and white patients: The CERAD experience, part XVI. Consortium to Establish a Registry for Alzheimer's Disease. *Neurology.* 1998; 51:154–158. [PubMed: 9674795]
- Fiske A, Wetherell JL, Gatz M. Depression in older adults. *Annual Review of Clinical Psychology.* 2009; 5:363–389.
- Gallo JJ, Rabins PV, Lyketsos CG, Tien AY, Anthony JC. Depression without sadness: Functional outcomes of nondysphoric depression in later life. *Journal of the American Geriatrics Society.* 1997; 45(5):570–578. [PubMed: 9158577]
- Ganguli M, Snitz B, Vander Bilt J, Chang CC. How much do depressive symptoms affect cognition at the population level? The Monongahela-Youghiogheny Healthy Aging Team (MYHAT) study. *International Journal of Geriatric Psychiatry.* 2009; 24(11):1277–1284. [PubMed: 19340894]
- Geerlings MI, Shoeve RA, Beekman AT, Jonker C, Deeg DJ, Schmand B, Van Tilburg W. Depression and risk of cognitive decline and Alzheimer's disease. Results of two prospective community-based studies in the Netherlands. *The British Journal of Psychiatry.* 2000; 176:568–575. [PubMed: 10974964]
- Gerety MB, Williams JW Jr, Mulrow CD, Cornell JE, Kadri AA, Rosenberg J, Long M. Performance of case-finding tools for depression in the nursing home: Influence of clinical and functional characteristics and selection of optimal threshold scores. *Journal of the American Geriatric Society.* 1994; 42(10):1103–1109.
- Gitlin LN, Hauck WW, Dennis MP, Schulz R. Depressive symptoms in older African-American and white adults with functional difficulties: The role of control strategies. *Journal of the American Geriatric Society.* 2007; 55(7):1023–1030.

- Godin O, Dufouil C, Ritchie K, Dartigues JF, Tzourio C, Peres K, Alperovitch A. Depressive symptoms, major depressive episode and cognition in the elderly: The three-city study. *Neuroepidemiology*. 2007; 28:101–108. [PubMed: 17409771]
- Goodglass, H.; Kaplan, E. The assessment of aphasia and related disorders. Philadelphia: Lea & Febiger; 1983.
- Goveas JS, Espeland MA, Hogan PE, Tindle HA, Shih RA, Kotchen JM, Resnick SM. Depressive symptoms and longitudinal changes in cognition: Women's health initiative study of cognitive aging. *Journal of Geriatric Psychiatry and Neurology*. 2014
- Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: Results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Archives of General Psychiatry*. 2005; 62(10):1097–1106. [PubMed: 16203955]
- Horwath E, Johnson J, Klerman GL, Weissman MM. Depressive symptoms as relative and attributable risk factors for first-onset major depression. *Archives of General Psychiatry*. 1992; 49(10):817–823. [PubMed: 1417435]
- Jorm AF. History of depression as a risk factor for dementia: An updated review. *The Australian and New Zealand Journal of Psychiatry*. 2001; 35(6):776–781. [PubMed: 11990888]
- Kane KD, Yochim BP, Lichtenberg PA. Depressive symptoms and cognitive impairment predict all-cause mortality in long-term care residents. *Psychology and Aging*. 2010; 25(2):446–452. [PubMed: 20545428]
- Kaplan, E.; Goodglass, H.; Weintraub, S. Boston Naming Test. Philadelphia: Lea and Febiger; 1983.
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, National Comorbidity Survey R. The epidemiology of major depressive disorder: Results from the National Comorbidity Survey Replication (NCS-R). *Journal of the American Medical Association*. 2003; 289(23):3095–3105. [PubMed: 12813115]
- Lebowitz BD, Pearson JL, Schneider LS, Reynolds CF, Alexopoulos GS III, Bruce ML, Parmelee P. Diagnosis and treatment of depression in late life. Consensus statement update. *Journal of the American Medical Association*. 1997; 278(14):1186–1190. [PubMed: 9326481]
- Lichtenberg PA, Ross T, Millis SR, Manning CA. The relationship between depression and cognition in older adults: A cross-validation study. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*. 1995; 50(1):P25–P32.
- Livney MG, Clark CM, Karlawish JH, Cartmell S, Negron M, Nunez J, Arnold SE. Ethnoracial differences in the clinical characteristics of Alzheimer's disease at initial presentation at an urban Alzheimer's disease center. *American Journal of Geriatric Psychiatry*. 2011; 19(5):430–439. [PubMed: 21522051]
- Lockwood KA, Alexopoulos GS, van Gorp WG. Executive dysfunction in geriatric depression. *American Journal of Psychiatry*. 2002; 159:1119–1126. [PubMed: 12091189]
- Lopez OL, Jagust WJ, Dulberg C, Becker JT, DeKosky ST, Fitzpatrick A, Kuller LH. Risk factors for mild cognitive impairment in the Cardiovascular Health Study Cognition Study: Part 2. *Archives of Neurology*. 2003; 60(10):1394–1399. [PubMed: 14568809]
- McDougall GJ Jr, Morgan S, Vaughan PW. Sixteen-month evaluation of depressive symptomatology in older adults. *Archives of Psychiatric Nursing*. 2012; 26(2):e13–e21. [PubMed: 22449566]
- McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, Kawas CH, Phelps CH. The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers & Dementia*. 2011; 7(3):263–269.
- Miller DK, Malmstrom TK, Joshi S, Andresen EM, Morley JE, Wolinsky FD. Clinically relevant levels of depressive symptoms in community-dwelling middle-aged African Americans. *Journal of the American Geriatrics Society*. 2004; 52(5):741–748. [PubMed: 15086655]
- Morris JC, Heyman A, Mohs RC, Hughes JP, van Belle G, Fillenbaum G, Clark C. The consortium to establish a registry for Alzheimer's disease (CERAD). Part I. Clinical and neuropsychological assessment of Alzheimer's disease. *Neurology*. 1989; 39:1159–1165. [PubMed: 2771064]
- Nebes RD, Butters MA, Mulsant BH, Pollock BG, Zmuda MD, Houck PR, Reynolds CF III. Decreased working memory and processing speed mediate cognitive impairment in geriatric depression. *Psychological Medicine*. 2000; 30(3):679–691. [PubMed: 10883722]

- Norris JT, Gallagher D, Wilson A, Winograd CH. Assessment of depression in geriatric medical outpatients: The validity of two screening measures. *Journal of the American Geriatrics Society*. 1987; 35(11):989–995. [PubMed: 3668142]
- Ogunniyi A, Hall KS, Gureje O, Baiyewu O, Gao S, Unverzagt FW, Hendrie HC. Risk factors for incident Alzheimer's disease in African Americans and Yoruba. *Metabolic Brain Disease*. 2006; 21(2–3):235–240. [PubMed: 16850256]
- Ownby RL, Crocco E, Acevedo A, John V, Loewenstein D. Depression and risk for Alzheimer disease: Systematic review, meta-analysis, and metaregression analysis. *Archives of General Psychiatry*. 2006; 63(5):530–538. [PubMed: 16651510]
- Pedraza O, Dotson VM, Willis FB, Graff-Radford NR, Lucas JA. Internal consistency and test-retest stability of the geriatric depression scale-short form in African American older adults. *Journal of Psychopathology and Behavioral Assessment*. 2009; 31(4):412–416. [PubMed: 20161488]
- Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild cognitive impairment: Clinical characterization and outcome. *Archives of Neurology*. 1999; 56:303–308. [PubMed: 10190820]
- Pincus HA, Davis WW, McQueen LE. 'Subthreshold' mental disorders. A review and synthesis of studies on minor depression and other 'brand names.' *British Journal of Psychiatry*. 1999; 174:288–296. [PubMed: 10533546]
- Ravdin LD, Mattis PJ, Lachs MS. Assessment of cognition in primary care: Neuropsychological evaluation of the geriatric patient. *Geriatrics*. 2004; 59(2):37–40. 42, 44. [PubMed: 14989592]
- Reitan, R. Manual for administration of neuropsychological test batteries for adults and children. Tucson, AZ: Reitan Neuropsychological Laboratories Incorporated; 1978.
- Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, Turner MB. On behalf of the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2012 update: A report from the American Heart Association. *Circulation*. 2012; 225:e2–e220. [PubMed: 22179539]
- Roman GC, Tatemichi TK, Erkinjuntti T, Cummings JL, Masdeu JC, Garcia JH, Hofman A. Vascular dementia: Diagnostic criteria for research studies. Report of the NINDS-AIREN International Workshop. *Neurology*. 1993; 43(2):250–260. [PubMed: 8094895]
- Rovner BM, Casten RJ, Leiby BE. Cognitive and noncognitive determinants of everyday activities in a racially diverse population of older persons receiving health services. *The Journal of Nervous and Mental Disease*. 2012; 200(7):627–631. [PubMed: 22759942]
- Sachs-Ericsson N, Burns AB, Gordon KH, Eckel LA, Wonderlich SA, Crosby RD, Blazer DG. Body mass index and depressive symptoms in older adults: The moderating roles of race, sex, and socioeconomic status. *The American Journal of Geriatric Psychiatry*. 2007; 15(9):815–825. [PubMed: 17804833]
- Sims R, Madhere S, Callender C, Campbell A Jr. Patterns of relationships between cardiovascular disease risk factors and neurocognitive function in African Americans. *Ethnicity & Disease*. 2008; 18(4):471–476. [PubMed: 19157252]
- Steffens DC, Otey E, Alexopoulos GS, Butters MA, Cuthbert B, Ganguli M, Yesavage J. Perspectives on depression, mild cognitive impairment, and cognitive decline. *Archives of General Psychiatry*. 2006; 63(2):130–138. [PubMed: 16461855]
- Stoudemire A, Hill CD, Morris R, Martino-Saltzman D, Lewison B. Long-term affective and cognitive outcome in depressed older adults. *American Journal of Psychiatry*. 1993; 150:896–900. [PubMed: 8494065]
- Tang MX, Cross P, Andrews H, Jacobs DM, Small S, Bell K, Mayeux R. Incidence of AD in African-Americans, Caribbean Hispanics, and Caucasians in northern Manhattan. *Neurology*. 2001; 56(1): 49–56. [PubMed: 11148235]
- Unverzagt FW, Gao S, Baiyewu O, Ogunniyi AO, Gureje O, Perkins A, Hendrie HC. Prevalence of cognitive impairment: Data from the Indianapolis Study of Health and Aging. *Neurology*. 2001; 57(9):1655–1662. [PubMed: 11706107]
- U.S. Census Bureau. Summary Population and Housing Characteristics. Washington, DC: Bureau of the Census; 2001. Census of Population and Housing.

- Wechsler, D. WMS-III administration and scoring manual. San Antonio, TX: Psychological Corporation; 1997.
- Weintraub S, Salmon D, Mercaldo N, Ferris S, Graff-Radford NR, Chui H, Morris JC. The Alzheimer's Disease Centers' Uniform Data Set (UDS): The neuropsychological test battery. *Alzheimer's Disease and Associated Disorders*. 2009; 23(2):91–101.
- Wessels AM, Lane KA, Gao S, Hall KS, Unverzagt FW, Hendrie HC. Diabetes and cognitive decline in elderly African Americans: A 15-year follow-up study. *Alzheimers & Dementia*. 2011; 7(4): 418–424.
- Wilkinson, GS. Wide range achievement test: WRAT3. Wilmington, DE: Wide Range, Inc; 1993.
- Williams DR, Gonzalez HM, Neighbors H, Nesse R, Abelson JM, Sweetman J, Jackson JS. Prevalence and distribution of major depressive disorder in African Americans, Caribbean blacks, and non-Hispanic whites: Results from the National Survey of American Life. *Archives of General Psychiatry*. 2007; 64(3):305–315. [PubMed: 17339519]
- Wilson RS, Mendes De Leon CF, Bennett DA, Bienias JL, Evans DA. Depressive symptoms and cognitive decline in a community population of older persons. *J Neurol Neurosurg Psychiatry*. 2004; 75(1):126–129. [PubMed: 14707321]
- Wilson RS, Barnes LL, Mendes CF, Agarwal NT, Schneider JS, Bach J, Bennett DA. Depressive symptoms, cognitive decline, and risk of AD in older person. *Neurology*. 2002; 59:364–370. [PubMed: 12177369]
- Yaffe K, Blackwell T, Gore R, Sands L, Reus V, Browner WS. Depressive symptoms and cognitive decline in nondemented elderly women. *Archives of General Psychiatry*. 1999; 56:425–430. [PubMed: 10232297]
- Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, Leirer VO. Development and validation of a geriatric depression scale: A preliminary report. *Journal of Psychiatric Research*. 1983; 17:37–49. [PubMed: 7183759]
- Zahodne LB, Devanand DP, Stern Y. Coupled cognitive and functional change in Alzheimer's disease and the influence of depressive symptoms. *Journal of Alzheimers Disease*. 2013; 34(4):851–860.

Table 1

Demographic characteristics of non-demented participants by diagnostic group

Variable	Normal controls	CI	Total	Test statistic
	(<i>n</i> = 637)	(<i>n</i> = 307)	(<i>N</i> = 944)	
Sex (% women)	77.9	80.8	78.8	$\chi^2(1) = 1.05^{\text{ns}}$
Age, mean (SD)	68.62 (6.96)	71.03 (8.64)	69.40 (7.63)	$F(1, 942) = 21.01^{**}$
Years of education, mean (SD)	14.25 (2.80)	13.08 (3.21)	13.87 (2.99)	$F(1, 941) = 32.74^{**}$
GDS scores, mean (SD)	1.3 (1.92)	2.1 (2.61)	1.53 (2.2)	$F(1, 942) = 32.59^{**}$

Note. CI = participants with mild cognitive impairment with and without memory impairment and cognitively impaired, not MCI; ns = not significant.

**
p<.001.

Table 2

Unadjusted and adjusted regression model of GDS scores and cognitive performance

	Language			Executive function			Processing speed			Attention			Memory		
	B	SE(B)	β	B	SE(B)	β	B	SE(B)	β	B	SE(B)	β	B	SE(B)	β
Model 1															
GDS	-.048	.011	-.143**	.012	.014	.027	-.021	.008	.081*	-.026	.013	-.068*	-.030	.012	-.081*
Model 2															
GDS	-.037	.009	-.110**	-.032	.016	.069+	-.019	.008	.072*	-.007	.012	-.018	-.024	.012	-.062*
Age	-.027	.003	-.275	-.015	.005	-.110**	-.006	.002	.084*	-.007	.003	-.064*	-.030	.003	-.275
Sex	.029	.050	.016	-.070	.084	-.029	.178	.045	.131**	-.078	.062	-.039	.330	.062	.165
Education	.024	.008	.096	.040	.014	.110**	-.002	.007	.012	-.003	.010	-.011	.026	.010	.092
WRAT	.043	.004	.395**	.037	.006	-.069*	.013	.003	.150**	.059	.004	.472**	.015	.004	.121

Note. GDS = Geriatric Depression Scale; Edu = Education; WRAT = Wide Range Achievement Test; B = unstandardized beta; SE (B) = standard error of unstandardized beta. *Model 1*: Unadjusted model of main effects of GDS on cognitive performance. *Model 2*: Adjusted model controlling for age, sex, education, and WRAT.

* $p < .05$

** $p < .001$

+ $p = .05$.