Published in final edited form as:

Stat Sin. 2014 July ; 24(3): 1365–1387. doi:10.5705/ss.2012.316.

Variable Selection for Sparse High-Dimensional Nonlinear Regression Models by Combining Nonnegative Garrote and Sure Independence Screening

Shuang Wu^{1,†}, Hongqi Xue^{1,†}, Yichao Wu², and Hulin Wu¹

Shuang Wu: shuang_wu@urmc.rochester.edu; Hongqi Xue: hongqi_xue@urmc.rochester.edu; Yichao Wu: ywu11@ncsu.edu; Hulin Wu: hulin_wu@urmc.rochester.edu

¹Department of Biostatistics and Computational Biology, University of Rochester, Rochester, NY

²Department of Statistics, North Carolina State University, Raleigh, NC

Abstract

In many regression problems, the relations between the covariates and the response may be nonlinear. Motivated by the application of reconstructing a gene regulatory network, we consider a sparse high-dimensional additive model with the additive components being some known nonlinear functions with unknown parameters. To identify the subset of important covariates, we propose a new method for simultaneous variable selection and parameter estimation by iteratively combining a large-scale variable screening (the nonlinear independence screening, NLIS) and a moderate-scale model selection (the nonnegative garrote, NNG) for the nonlinear additive regressions. We have shown that the NLIS procedure possesses the sure screening property and it is able to handle problems with non-polynomial dimensionality; and for finite dimension problems, the NNG for the nonlinear additive regressions has selection consistency for the unimportant covariates and also estimation consistency for the parameter estimates of the important covariates. The proposed method is applied to simulated data and a real data example for identifying gene regulations to illustrate its numerical performance.

Keywords

Gene regulations; independence learning; nonlinear regressions; nonnegative garrote; sigmoid function; sure screening

1 Introduction

With the rapid advancement of modern technologies, high dimensional data now arise frequently in many areas of contemporary statistics, such as microarrays, RNA-seq, proteomics, biomedical imaging, signal processing and finance. In high dimensional statistical modeling, it is a fundamental problem to identify important explanatory variables and has drawn increasing interests. For linear regression models, many penalization methods have been proposed in the regularization framework to conduct variable selection and

Correspondence to: Shuang Wu, shuang_wu@urmc.rochester.edu.

[†]Joint first authors.

estimation. At the same time many efforts have been made to study their statistical properties in high dimensional settings. Existing methods include the bridge regression (Frank and Friedman, 1993; Huang et al., 2008), least absolute shrinkage and selection operator or Lasso (Tibshirani, 1996; Zhao and Yu, 2006; van de Geer, 2008; Zhang and Huang, 2008), the nonnegative garrote (Breiman, 1995; Yuan and Lin, 2007), the smoothly clipped absolute deviation (SCAD) penalty (Fan and Li, 2001; Fan and Peng, 2004; Kim et al., 2008), the adaptive Lasso (Zou, 2006) and the Dantzig selector (Candes and Tao, 2007) among others.

In many practical applications, there is little prior information to justify the assumption that the effects of covariates on the response take a linear form. For example, in economics and finance, a lot of continuous time finance models, such as bonding pricing models and diffusion processes describing yield curves, are highly nonlinear. Another example is the gene regulatory network (GRN), where the regulation and interaction between genes are known to be complex and inherently nonlinear. A widely adopted approach to address the nonlinearity is to extend the linear regression to the nonparametric additive model. The nonlinear effect of a covariate is modeled through a nonparametric function, which is usually approximated by a linear combination of some basis functions. So the selection of important variables corresponds to the selection of groups of basis functions. Methods for selecting grouped variables have been studied in Antoniadis and Fan (2001) and Yuan and Lin (2006). Lin and Zhang (2006) proposed the component selection and smoothing operator (COSSO) method for model selection in smoothing spline ANOVA with a fixed number of covariates. Recently, Ravikumar et al. (2009), Meier et al. (2009) and Huang et al. (2010) considered variable selection in high dimensional nonparametric additive models where the number of additive components is larger than the sample size.

The nonparametric additive models are flexible to account for a variety of nonlinear relationships. The model is essentially linear once the nonparametric components are expressed as linear combinations of basis functions. Consequently the computation is relatively easy. Although the nonparametric models enjoy many benefits of flexibility, robustness, and computational convenience, the alternative nonlinear models with known functional forms are necessary for some practical applications. In particular, the given nonlinear model is more interpretable and easy for predictions, and it also avoids the selection of nonparametric smoothing parameters such as the number of bases, whose impact on variable selection has not been well understood under nonparametric model settings. In this paper, we will focus on exploring the variable selection procedure for the nonlinear additive models as an alternative for the nonparametric additive models.

Our work is motivated by the application of reconstructing a gene regulatory network (GRN), in which the regulatory function of each gene is a sigmoid function. The usage of the sigmoid function model to describe a GRN has been discussed in literature, for example, Mestl et al. (1995) and Chen et al. (2004) among others. An example of the simple sigmoid function is

$$f(x) = \frac{1}{1 + e^{-\alpha x}}. \quad (1)$$

It is usually used to model nonlinear systems that exhibit “saturation” phenomena and the parameter $\alpha > 0$ indicates the transition rate. Since biological systems are seldom fully connected and most nodes are only directly connected to a small number of other nodes (Jeong et al., 2000), it is commonly assumed that the GRN is a sparse network. So the identification of a GRN requires to select the important regulators for each target gene using variable selection techniques.

We consider the following high-dimensional nonlinear additive model

$$Y = \beta_0 + \sum_{j=1}^p \beta_j f_j(X_j, \alpha_j) + \varepsilon, \quad (2)$$

where Y is the response variable, $\mathbf{X} = (X_1, \dots, X_p)^T$ is the vector of covariates, $f(\mathbf{X}, \boldsymbol{\alpha}) = \{f_1(X_1, \alpha_1), \dots, f_p(X_p, \alpha_p)\}^T$ are known nonlinear functions and ε is the random error term with mean 0 and variance $0 < \sigma^2 < \infty$. The parameters $\boldsymbol{\alpha} = (\alpha_1^T, \dots, \alpha_p^T)^T$ usually control the shapes of functions f . We assume that the dimension p of the covariates may increase with the sample size n and may be even larger than n , and the coefficient vector $\boldsymbol{\beta} = (\beta_1, \dots, \beta_p)^T$ is sparse in the sense that most of its elements are zeros. Given the i.i.d. observed data $\{(\mathbf{x}_i, y_i), i = 1, \dots, n\}$ with $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{ip})^T$, we are interested in identifying the subset of important covariates $\mathcal{M}_* = \{1 \leq j \leq p: \beta_j \neq 0\}$ and also estimating the parameters $(\beta_0, \beta_j, \alpha_j), j \in \mathcal{M}_*$.

Compared with the linear and generalized linear models, there exists very limited statistical research of parameter estimation, statistical inference, and especially variable selection for the nonlinear regression and generalized nonlinear models. Jennerich (1969), Malinvaud (1970) and Wu (1981) proposed the nonlinear least squares (NLS) estimation for nonlinear regression models and derived their asymptotic properties. A good review can be found in Seber and Wild (2003). Wei (1998), Kosmidis and Firth (2009) and Biedermann and Woods (2011) studied likelihood-based estimation and statistical inference for generalized nonlinear models. Xue et al. (2010) considered the NLS estimations for nonlinear ordinary differential equation models with measurement errors and established their large sample theories. Jiang et al. (2011) proposed a weighted composite quantile regression (WCQR) estimation and studied the model selection with the adaptive LASSO and SCAD penalties for nonlinear models with a diverging number of parameters. Their algorithm is tailored to the WCQR model and is not applicable to our model. Moreover, our model includes some unknown non-linear parameters α_j and they may have effects on the relationship between the response and the covariates, which makes the problem of selecting nonzero β_j 's even more complicated.

For given estimates $\hat{\alpha}_j$, model (2) can be treated as a linear regression model and variable selection methods developed for linear regressions can be then applied to estimate β_j . For

this purpose, we propose to use the nonnegative garrote (NNG), which requires a consistent initial estimate. Intuitively, the variable selection procedure works better if the estimates $\hat{\alpha}_j$ are closer to the true values. However, as the number of covariates p grows, it becomes quite difficult to get good estimates for α_j . To overcome the challenge of high dimensionality, we adopt the idea of independence screening introduced by Fan and Lv (2008). The sure independence screening (SIS) method performs efficient dimension reduction via marginal correlation learning for ultra-high dimensional feature selection. It has been shown to have a sure screening property; that is, with probability tending to 1, the independence screening technique retains all of the important features in the model. The SIS was later extended to high dimensional generalized linear models (Fan and Song, 2010) and nonparametric additive models (Fan et al., 2011). The idea of using marginal information to deal with high dimensionality was also adopted in other works. Hall et al. (2009) considered a different marginal utility, derived from an empirical likelihood point of view. Hall and Miller (2009) proposed a generalized correlation ranking, which allows nonlinear relationships. More recently, Wang (2012) studied a sure independence screening using a factor profiling approach; Xue and Zou (2011) considered sure independence screening for sparse signal recovery; Gorst-Rasmussen and Scheike (2012) investigated independent screening for single-index hazard rate models with right censored data; see Zhu et al. (2012) and Li et al. (2012) also for recent developments of model-free approaches for feature screening.

In this paper, we propose a nonlinear independence screening method based on the residual sum of squares of the marginal modeling and establish its sure screening property for model (2). We fit p marginal nonlinear regressions of the response Y against each covariate X_j separately and rank their importance to the joint model according to the residual sum of squares of the marginal models. The covariates whose residual sum of squares are smaller than a threshold are selected and then a more refined variable selection technique such as the nonnegative garrote can be applied to the nonlinear additive model conditional on these selected covariates. This nonlinear independence screening procedure can significantly reduce the dimension of the covariates, and more importantly, its sure screening property ensures that all the important covariates will be retained with probability tending to 1. Our method is not a trivial extension of Fan and Lv (2008) and Fan and Song (2010). For the nonlinear model (2), the minimum distinguishable signal of the marginal screening is closely related to the stochastic and numerical errors in the optimization of the nonlinear parameters. In addition, the objective function derived from the nonlinear model often has multiple local minima, making the estimation and marginal screening more challenging. For the theoretical development, we extend the sure independent screening approach from linear models to nonlinear models, where some global assumptions that are widely used in linear models, such as the convexity, are not appropriate for the nonlinear models and are replaced by some local assumptions, such as the local convexity. We will show in Section 3 that our nonlinear independence screening approach can handle problems with non-polynomial dimensionality or ultra-high dimensionality.

The remainder of the paper is organized as follows. Section 2 describes the nonnegative garrote for variable selection in nonlinear additive models with finite predictors and its asymptotic properties. We elaborate the independence screening for the nonlinear additive

models with ultra-high dimensional covariates and establish its sure independence screening property in Section 3. We present the results of simulation studies in Section 4 and provide an illustrative application in Section 5. Section 6 includes concluding remarks and some discussions. Proofs of the asymptotic results in Section 2 and Section 3 can be found in the web-appendix.

2 Nonnegative Garrote for Nonlinear Additive Models

2.1 Method

In this section, we consider the case of finite covariates, i.e., $p < n$. For the following multiple linear regression model

$$y_i = \sum_{j=1}^p \beta_j z_{ij} + \varepsilon_i, \quad i=1, \dots, n, \quad (3)$$

the original nonnegative garrote (Breiman, 1995) finds a set of nonnegative scaling factors c_j to minimize

$$\frac{1}{2} \sum_{i=1}^n \left[y_i - \sum_{j=1}^p c_j \hat{\beta}_j z_{ij} \right]^2 + n \lambda_n \sum_{j=1}^p c_j, \quad \text{subject to } c_j \geq 0, \quad (4)$$

with an initial estimates $\hat{\beta}_j$ for model (3). The final garrote estimates are then given by $\tilde{\beta}_j = \hat{c}_j \hat{\beta}_j$. An appropriately chosen λ_n can shrink some $\hat{c}_j$ to exact 0 and thus produces a sparse model. We omit the subscript n in λ_n in the following presentation if no confusion occurs. The selection consistency of nonnegative garrote was first proved by Zou (2006) and later Yuan and Lin (2007) showed that, as long as the initial estimate is consistent in terms of estimation, the nonnegative garrote estimate is consistent in terms of both estimation and model selection given that the tuning parameter λ is appropriately chosen. The ordinary least squares estimator is often chosen as the initial estimation.

We extend the idea of the nonnegative garrote (NNG) to the additive nonlinear regression (2) with finite p predictors. To simplify the presentation, the response variable Y and the nonlinear functions f_j are assumed to be centered with $E(Y) = 0$ and $E[f_j(X_j, \alpha_j)] = 0$ for all $j = 1, \dots, p$, so the intercept $\beta_0 = 0$. The proposed method can be easily adapted to include the intercept. Given initial estimates $\hat{\alpha}_j$, the nonlinear additive model (2) reduces to the linear regression model (3) with $z_{ij} = f_j(x_{ij}, \hat{\alpha}_j)$, $j = 1, \dots, p$, as the predictors. One can then proceed with the usual nonnegative garrote with the initial estimates $\hat{\beta}_j$ for variable selection and estimation. Suppose the nonnegative garrote selects covariates $X_j, j \in \hat{s} = \{1 \leq j \leq p: \hat{c}_j > 0\}$. The parameters α_j are then updated conditional on $\hat{\beta}_j, j \in \hat{s}$. These procedures iterate until some convergence criterion is met, for instance, the residual sum of squares of the fitted model does not change up to a tolerance. The initial estimates need to be carefully chosen, because the solution path consistency of the nonnegative garrote depends on the consistency of the initial estimator. We use the nonlinear least square estimates of (2) as the initial

estimates and when the model dimension of (2) is not very high, these estimates do have the consistency property.

The above iterative algorithm can be viewed as a solution of a separable nonlinear least squares problem (Ruhe and Wedin, 1980; Golub and Pereyra, 2003). The parameters are separated into two sets, where $\{\beta_j\}$ are linear parameters and $\{\alpha_j\}$ are nonlinear parameters. In each iteration, the optimization with respect to $\{\beta_j\}$ is performed first, and the correction to $\{\alpha_j\}$ follows after that. It has been shown that eliminating one set of parameters can result in faster convergence of the optimization problem. We refer to Ruhe and Wedin (1980) and Golub and Pereyra (2003) and references therein for detailed descriptions of separable nonlinear least squares problems and the convergence properties of related algorithms.

2.2 Asymptotic properties

To simplify our presentation, we introduce notations $\gamma = (\beta^T, \alpha^T)^T$ with $\beta = (\beta_1, \dots, \beta_p)^T$ and $\alpha = (\alpha_1^T, \dots, \alpha_p^T)^T$, and the true parameters are represented by $\gamma_0 = (\beta_0^T, \alpha_0^T)^T$. Denote $\ell(\gamma, X, Y) = [Y - F(X, \gamma)]^2$ with $F(X, \gamma) = \sum_{j=1}^p \beta_j f_j(X_j, \alpha_j)$ and $P_n \ell(\gamma, x, y) = n^{-1} \sum_{i=1}^n \ell(\gamma, x_i, y_i) = n^{-1} \sum_{i=1}^n [y_i - \sum_{j=1}^p \beta_j f_j(x_{ij}, \alpha_j)]^2$. In order to establish the asymptotic properties of the nonnegative garrote under the nonlinear additive model setting, the following conditions are needed:

- (A1) For all $j = 1, \dots, p$, $\beta_j \in \mathcal{B}_j \subset R$ and $\alpha_j \in \mathcal{H}_j \subset R^{d_j}$, where $\mathcal{B}_j$ and $\mathcal{H}_j$ are compact and convex with finite diameters A_1 and A_2 , respectively. Denote $\Gamma = \prod_{j=1}^p \mathcal{B}_j \times \mathcal{H}_j \subset R^{P + \sum_{j=1}^p d_j}$ and γ_0 is an interior of Γ .
- (A2) For any $\gamma \in \Gamma$, $E[F(X, \gamma) - F(X, \gamma_0)]^2 = 0$ if and only if $\beta_j = \beta_{0j}$ and $\alpha_j = \alpha_{0j}$ for all $j = 1, \dots, p$.
- (A3) Denote $\Omega_{1n} = \{X: \|X\|_\infty \leq K_n\}$ and $I_{1n}(X) = I(X \in \Omega_{1n})$ for some sufficiently large positive constants K_n , where $\|\cdot\|_\infty$ is the supremum norm. Assume that $F(X, \gamma)$, $\frac{\partial}{\partial \gamma} F(X, \gamma)$ and $\frac{\partial^2}{\partial \gamma \partial \gamma^T} F(X, \gamma)$ exist and are continuous and uniformly bounded for $X \in \Omega_{1n}$ and $\gamma \in \Gamma$ with bounds k_1 , k_2 and k_3 , respectively.
- (A4) The covariate vector X has a continuous density and there exist constants D_1 and D_2 such that the density function g_j of X_j satisfies $0 < D_1 \leq g_j(x) \leq D_2 < \infty$ on the support $[a, b]$ of X_j for every $1 \leq j \leq p$.
- (A5) There exists a sequence of random variables $\{\hat{\gamma}\}$ with $\frac{\partial P_n \ell(\gamma, x, y)}{\partial \gamma} \big|_{\gamma=\hat{\gamma}} = 0$ such that $P_n \ell(\gamma, x, y)$ is locally convex in some neighborhood of γ , say $\mathcal{A}(\delta) = \{\gamma \in \Gamma, \|\gamma - \hat{\gamma}\| \leq \delta\}$ for some $\delta > 0$ and $\gamma_0 \in \mathcal{A}(\delta)$.
- (A6) The Fisher information

$$I(\gamma) = E \left[\frac{\partial F(X, \gamma)}{\partial \gamma} \frac{\partial F(X, \gamma)}{\partial \gamma^T} \right]$$

is finite and positive definite at $\gamma = \gamma_0$.

(A7) Denote $\Omega_n\{(\mathbf{X}, Y): \|\mathbf{X}\|_\infty \leq K_n, |Y| \leq K_n^*\}$ and $I_n(\mathbf{X}, Y) = I\{(\mathbf{X}, Y) \in \Omega_n\}$ for some sufficiently large positive constants K_n^* . There exists a constant C_1 such that with $b_n = C_1 \max\{k_1^2, A_1 k_1 k_2\} V_1^{-1} (p/n)^{1/2}$ and V_1 given in (S.6) in the web-appendix,

$$\sup_{\gamma \in \Gamma, \|\gamma - \gamma_0\| \leq b_n} |E[\ell(\gamma, \mathbf{X}, Y) - \ell(\gamma_0, \mathbf{X}, Y)] (1 - I_n(\mathbf{X}, Y))| \leq o(p/n).$$

Conditions (A1), (A3), (A4) and (A7) are general bound requirements. Conditions (A2) and (A6) ensure the identifiability and the existence of the nonnegative garrote. We use the local convexity in (A5) to replace the global convexity in Condition C in Fan and Song (2010), because our model is nonlinear and the associated objective function may have multiple local minima. Note that the proposed nonnegative garrote for the nonlinear additive models requires an initial estimate and a natural choice is the NLS estimate. From the standard NLS estimation theories developed by Jennerich (1969), Malinvaud (1970) and Wu (1981), under (A1)–(A6), the NLS estimator $\hat{\gamma}$ defined in (A5) is consistent and asymptotically normal, so $\hat{\gamma}$ satisfies the consistency requirement of the nonnegative garrote initial estimates. Let $\hat{c}_j(\lambda)$ be the optimizer of minimizing (4). The final nonnegative garrote estimates are given by $\tilde{\beta}_j = \hat{c}_j(\lambda) \hat{\beta}_j$ and $\tilde{\alpha}_j = \hat{\alpha}_j$ for $j = 1, 2, \dots, p$. We have established the selection and estimation consistency of our nonnegative garrote estimates of the additive nonlinear linear model as follows.

Theorem 1

- i. Assume that $\text{var}\{f(\mathbf{X}, \alpha_0)\}$ is positive definite and the initial estimates $\hat{\gamma}$ are δ_n consistent, namely $\max_{1 \leq j \leq p} (|\hat{\beta}_j - \beta_{0j}| + \|\hat{\alpha}_j - \alpha_{0j}\|) = O_p(\delta_n)$ for some $\delta_n \rightarrow 0$. Under (A1)–(A3), for $\lambda \rightarrow 0$ satisfying $\delta_n = o(\lambda)$, we have $P\{\hat{c}_j(\lambda) = 0\} \rightarrow 1$ for any j such that $\beta_{0j} = 0$, and $\hat{c}_j(\lambda) = 1 + O_p(\lambda)$ for any j such that $\beta_{0j} \neq 0$.
- ii. Choose the nonlinear least squares estimate as the initial estimate $\hat{\gamma}$. Under (A1)–(A6) and $\lambda \rightarrow 0$ satisfying $n^{-1/2} = o(\lambda)$, we have selection consistency for unimportant predictors, namely $P\{\tilde{\beta}_j = 0\} \rightarrow 1$ for any j such that $\beta_{0j} = 0$, and also estimation consistency for the estimates corresponding to the important predictors almost surely.

We also establish an exponential bound for the tail probability of the NLS estimator $\hat{\gamma}$, which is parallel to Theorem 1 in Fan and Song (2010) and will be used in the next section. Since this result holds under general nonlinear conditions and has its own values from methodological and theoretical perspectives, we present it by the following theorem.

Theorem 2—For the NLS estimator $\hat{\gamma}$ defined in (A5), if (A1)–(A7) hold, then for any $t > 0$, we have

$$P(\sqrt{n}\|\hat{\gamma} - \gamma_0\| \geq 16k_n^* \max\{k_1, A_1 k_2\} (1+t)/V_1) \leq \exp(-2t^2) + nP(\Omega_n^c)$$

with $k_n^* = 2(k_1 + K_n^*)$.

3 Independence Screening for High Dimensional Nonlinear Regressions

As the dimension p of covariates in (2) grows with the sample size n , especially when $p \gg n$, the iterative algorithm described in Section 2 may not work well. This is partly because the nonlinear least squares for the models with a large number of parameters and redundant covariates may not converge properly to the local minimum. Moreover, the nonnegative garrote cannot be applied when the sample size is smaller than the number of covariates. To overcome the challenge of high dimensionality, we consider the idea of sure independence screening for linear or generalized linear regression models proposed by Fan and Lv (2008) and Fan and Song (2010), and extend it to the nonlinear additive regression models.

Now we rewrite p as p_n and fit p_n marginal nonlinear regressions of the response Y against each covariate X_j separately and rank their importance to the joint model according to a measure of the goodness of fit of their marginal models. In Fan and Lv (2008) and Fan and Song (2010), the covariates are standardized, i.e. $EX_j^2 = 1, j = 1, \dots, p_n$, so the magnitude of the coefficient estimate of the marginal model can preserve the nonsparsity of the joint model. To extend this idea to the nonlinear additive model (2), we need to standardize the

nonlinear functions $f_j(X_j, \alpha_j), j = 1, \dots, p_n$. Define $\hat{\beta}_j' = \hat{\beta}_j^M \sqrt{n^{-1} \sum_{i=1}^n f_j^2(x_{ij}, \hat{\alpha}_j^M)}$, where $\hat{\beta}_j^M$ and $\hat{\alpha}_j^M, j = 1, \dots, p_n$, are the NLS estimates of the marginal models. This $\hat{\beta}_j'$ can be considered as the marginal NLS estimates obtained by standardizing $f_j(X_j, \alpha_j)$ and we can use the magnitude of $\hat{\beta}_j'$ as the marginal utility for the independence screening. We refer to this strategy as the marginal least square estimate screening or MLSE screening, where we select a set of variables with the magnitude of $\hat{\beta}_j'$ greater than a prespecified threshold ζ_n :

$$\hat{\mathcal{N}} = \{1 \leq j \leq p_n : |\hat{\beta}_j'| \geq \zeta_n\}. \quad (5)$$

Note that the MLSE screening involves $\hat{\beta}_j^M$ and $\hat{\alpha}_j^M$, because both estimates contribute to determine the relationship between Y and X_j . Another natural way to incorporate the information of both $\hat{\beta}_j^M$ and $\hat{\alpha}_j^M$ is through the residual sum of squares (RSS) of the component wise nonlinear regressions. We propose to rank the covariates X_j 's according to the RSS of the marginal models and select a set of variables:

$$\hat{\mathcal{M}} = \{1 \leq j \leq p_n : \text{RSS}_j \leq \xi_n\}, \quad (6)$$

with $\text{RSS}_j = \min_{\beta_j, \alpha_j} \sum_{i=1}^n [y_i - \beta_j f_j(x_{ij}, \alpha_j)]^2$ as the residual sum of squares of the j th marginal fit and ξ_n as a prespecified threshold value. This strategy is analogous to the likelihood ratio screening proposed in Fan et al. (2009) and we refer to it as the RSS screening. We will show in the following section that it is asymptotically equivalent to the MLSE screening in preserving the nonsparsity of the joint model. In practical computations,

the sets $\mathcal{K}$ and $\hat{\mathcal{M}}$ may not be exactly the same because the nonlinear optimization problem may have several local minima. Moreover, the RSS screening is easier to implement as one does not need to standardize the nonlinear functions $f_j(X_j, \alpha_j)$. We found in our numerical analysis that the RSS screening is more robust than the MLSE screening, so the RSS screening is adopted in the simulation studies and the data application in this paper. To determine a data-driven threshold ξ_n for the RSS screening, one can use the random permutation to create null models as in Fan et al. (2011). An alternative thresholding scheme is to choose d covariates with the smallest marginal residual sum of squares. In case of large p_n , this thresholding approach can save a lot of computing effort by avoiding the random permutations.

The nonlinear independence screening (NLIS) procedure reduces the dimensionality of covariates from p_n to a possibly much smaller space with model size $|\hat{\mathcal{M}}|$. Following the variable screening, we can apply a more refined model selection technique to choose important covariates in the nonlinear additive model conditional on the selected variable set $|\hat{\mathcal{M}}|$, and for this purpose, we use the nonnegative garrote for nonlinear additive models described in Section 2. Based on the philosophy in Fan and Lv (2008), independence screening should be applied when p_n is very large; while for moderately high p_n , one should use penalized methods for variable selection. In our case, the penalized model selection via the nonnegative garrote for nonlinear additive models requires to obtain the initial parameter estimates through the NLS optimization and it is very difficult to get good initial estimates when p_n is large. So for variable selection in the nonlinear model (2), screening is necessary even when p_n is only relatively high.

The crucial part in our method is that the nonlinear independence screening procedure does not mistakenly miss any important covariates. We will show in the following section that our nonlinear independence screening procedure has a sure screening property, as defined by Fan and Lv (2008), so that all the important covariates will be retained with probability tending to 1. This is not a trivial extension of Fan and Lv (2008) and Fan and Song (2010), because in our case, the minimum distinguishable signal of selecting $\mathcal{M}$ is closely related to the stochastic and numerical errors in estimating the nonlinear parameters. In addition, fitting marginal models to a joint regression can be considered as a type of model misspecification (White, 1982) as most of the covariates are dropped out from the model fitting. The NLS objective function of a misspecified nonlinear model can be quite sensitive or unstable, making our implementation more difficult than that of linear models. For the theoretical development, since the objective function derived from our nonlinear additive model often has multiple local minima, some assumptions that have been widely used in linear regression models, such as global convexity, are not applicable in our case and should be replaced by assumptions that are appropriate for the nonlinear problem setting, such as local convexity.

3.1 Asymptotic properties

Let $\mathcal{M}_* = \{1 \leq j \leq p_n: \beta_{0j} \neq 0\}$ be the true sparse model with size $v_n = |\mathcal{M}_*|$, where $\beta_0 = (\beta_{01}, \dots, \beta_{0p_n})^T$ and $\alpha_0 = (\alpha_{01}^T, \dots, \alpha_{0p_n}^T)^T$ are the true parameter values. Denote $\gamma_j = (\beta_j, \alpha_j^T)^T$,

$F_j(X_j, \gamma_j) = \beta_j f_j(X_j, \alpha_j)$, $\ell(\gamma_j, X_j, Y) = [Y - F_j(X_j, \gamma_j)]^2$ and
 $P_n \ell(\gamma_j, \mathbf{x}_j, \mathbf{y}) = n^{-1} \sum_{i=1}^n [y_i - \beta_j f_j(x_{ij}, \alpha_j)]^2$. We also define

$$\gamma_j^M = \{\beta_j^M, (\alpha_j^M)^T\}^T = \arg \min_{\gamma_j} E \ell(\gamma_j, X_j, Y), \quad (7)$$

for $j = 1, \dots, p_n$, where E represents the expectation under the true model.

The following additional conditions are needed for proving the asymptotic properties of the nonlinear independence screening procedure.

(B1) There exist two positive constants k_4 and k_5 such that $k_5 \leq E f_j^2(X_j, \alpha_j) \leq k_4$ for all $j = 1, \dots, p_n$.

(B2) The marginal Fisher information:

$$I_j(\gamma_j) = E \left[\frac{\partial F_j(X_j, \gamma_j)}{\partial \gamma_j} \frac{\partial F_j(X_j, \gamma_j)}{\partial \gamma_j^T} \right]$$

is finite and positive definite at $\gamma = \gamma_j^M$ for $j = 1, \dots, p_n$.

(B3) There exists a constant $C_2 > 0$ such that for all $j = 1, \dots, p_n$,

$$\sup_{\gamma \in \Gamma, \|\gamma - \gamma_j^M\| \leq C_2} |E \beta_j^2 f_j^2(X_j, \alpha_j) I(|X_j| \geq K_n)| \leq o(n^{-1}).$$

(B4) There exist some positive constants m_0, m_1, s_0, s_1 and $a > 1$, such that for any real t ,

$$\begin{aligned} E \exp[t f_j(X_j, \alpha_j)] &\leq s_0 \exp(m_0 t^a), \quad j=1, \dots, p_n, \\ E \exp(t \varepsilon_i) &\leq s_1 \exp(m_1 t^a), \quad i=1, \dots, n. \end{aligned}$$

(B5) For $j = 1, \dots, p_n$, there exists a marginal least squares estimator (MLSE) $\hat{\gamma}_j^M$ with

$\frac{\partial P_n \ell(\gamma_j, \mathbf{x}_j, \mathbf{y})}{\partial \gamma_j} \Big|_{\gamma_j = \hat{\gamma}_j^M} = 0$ such that $P_n \ell(\gamma_j, \mathbf{x}_j, \mathbf{y})$ is locally convex in some neighborhood of $\hat{\gamma}_j^M$, say $\mathcal{A}_{1\delta} = \{\gamma_j \in \Gamma, \|\gamma_j - \hat{\gamma}_j^M\| \leq \delta\}$ for some $\delta > 0$ and $\gamma_j^M \in \mathcal{A}_{1\delta}$.

(B6) There exists a positive constant V_2 such that for all $\gamma_j \in \mathcal{A}_{1\delta}$, we have

$$E \ell(\gamma_j, X_j, Y) \geq V_2 \|\gamma_j - \gamma_j^M\|^2, \quad j = 1, \dots, p_n.$$

(B7) Denote $\Sigma_{\alpha} = \text{cov}[f(X, \alpha_0), f(X, \alpha^M)]$ and let $\lambda_{\max}(\Sigma_{\alpha})$ be its maximum eigenvalue. There exist two positive constants D_3 and D_4 such that

$$0 < D_3 \leq \beta_0^T \sum_{\alpha} \beta_0 \leq D_4 < \infty.$$

Condition (B1) is imposed to bound the nonlinear effects of the covariates so that their scales are of the same order. Condition (B4) is similar to Condition D in Fan and Song (2010) and it is satisfied if $f_j(X_j, \alpha_j)$ has a subgaussian distribution with a scale factor $\tau < \infty$, i.e., $P \exp[tf_j(X_j, \alpha_j)] \leq \exp(\tau^2 t^2/2)$ for all real t , or equivalently, $P\{|f_j(X_j, \alpha_j)| \geq t\} \leq c^* \exp\{-t^2/(2\tau^2)\}$ for some constant c^* . As we know, many types of distributions belong to the family of subgaussian distributions, such as the normal distribution and the bounded symmetric distribution. Condition (B5) is the marginal version of Condition (A5). The following Theorem 3 gives a uniform convergence result of the MLSEs defined in (B5) and the sure screening property of the MLSE screening, and Theorem 4 provides an asymptotic rate on the MLSE model size. These two theorems are extensions of Theorems 4 and 5 in Fan and Song (2010), respectively, to the nonlinear additive model setting (2). Condition (B7) is the parallel version with Condition F in Fan and Song (2010), where they required $\text{var}(\mathbf{X}^T \beta_0) = \beta_0^T \text{var}(\mathbf{X}) \beta_0$ to be bounded from above and below.

Theorem 3—Suppose that Conditions (A1), (A3), (A4) and (B1)–(B6) hold.

- i. If $n^{1-2\kappa}/(k_n^*)^2 \rightarrow \infty$ for some constant $0 < \kappa < 1/2$ given in Lemma 3, then for any $c_3 > 0$, there exists a positive constant c_4 such that

$$P \left(\max_{1 \leq j \leq p_n} \|\hat{\gamma}_j^M - \gamma_j^M\| \geq c_3 n^{-\kappa} \right) \leq p_n \{ \exp[-c_4 n^{1-2\kappa}/(k_n^*)^2] + n s_2 \exp[-m_2(K_n^*)^{\frac{a}{a-1}}] \},$$

$$\text{with } s_2 = s_0^{\nu_n} s_1.$$

- ii. We select a set of variables

$$\hat{\mathcal{N}}_{\zeta_n} = \{1 \leq j \leq p_n : |\hat{\beta}_j^M| \geq \zeta_n\},$$

where ζ_n is a predefined threshold value. If, in addition, conditions in Lemma 3 hold, then by taking $\zeta_n = c_5 n^{-\kappa}$ with $c_5 = c_2/2$ (c_2 is given in Lemma 3), we have

$$P(\mathcal{M}_* \subset \hat{\mathcal{N}}_{\zeta_n}) \geq 1 - \nu_n \{ \exp[-c_4 n^{1-2\kappa}/(k_n^*)^2] + n s_2 \exp[-m_2(K_n^*)^{\frac{a}{a-1}}] \}.$$

Theorem 4—Under Conditions (A1), (A3), (A4) and (B1)–(B7), for any $\zeta_n = c_5 n^{-2\kappa}$, we have

$$P[|\hat{\mathcal{N}}_{\zeta_n}| \leq O\{n^{2\kappa} \lambda \max(\sum_{\alpha})\}] \geq 1 - p_n \{ \exp[-c_4 n^{1-2\kappa}/(k_n^*)^2] + n s_2 \exp[-m_2(K_n^*)^{\frac{a}{a-1}}] \}.$$

Remark 1. Under Condition (B1), after some derivations, we can get that $\hat{\beta}_j^M$ and $\hat{\beta}_j'$ have the same order, with probability tending to 1 exponentially fast. Thus Theorem 3 and Theorem 4 also hold for $\hat{\beta}_j'$.

Remark 2. In order to obtain the optimal order of K_n^* , we balance the two terms in the upper bound of Theorem 3(i) and get $K_n^* = n^{(1-2\kappa)(a-1)/(3a-2)}$. It follows that

$$P\left(\max_{1 \leq j \leq p_n} \|\hat{\gamma}_j^M - \gamma_j^M\| \geq c_3 n^{-\kappa}\right) = O\{p_n \exp(-c_4 n^{(1-2\kappa)a/(3a-2)})\}.$$

Furthermore, if Y is bounded, then K_n^* can be taken as a finite constant. So

$$P\left(\max_{1 \leq j \leq p_n} \|\hat{\gamma}_j^M - \gamma_j^M\| \geq c_3 n^{-\kappa}\right) = O\{p_n \exp(-c_4 n^{1-2\kappa})\}.$$

In both of the above cases, the tail probability in Theorem 3 has the exponential rate. Thus, similar to Fan and Song (2010), we can handle the non-polynomial dimensionality with $\log p_n = o\{n^{(1-2\kappa)a/(3a-2)}\}$ or $\log p_n = o(n^{1-2\kappa})$.

Next we consider the sure screening property of the RSS screening (6). Let

$R^* = (R_1^*, \dots, R_{p_n}^*)^T$, where $R_j^* = E\ell(\gamma_0^M, X_j, Y) - E\ell(\gamma_j^M, X_j, Y)$ and $\gamma_0^M = \{0, (\alpha_j^M)^T\}^T$. The empirical values of R^* can be written as $R_n = (R_{1,n}, \dots, R_{p_n,n})^T$, where

$R_{j,n} = P_n \ell(\hat{\gamma}_0^M, x_j, y) - P_n \ell(\hat{\gamma}_j^M, x_j, y)$ and $\hat{\gamma}_0^M = \{0, (\hat{\alpha}_j^M)^T\}^T, j = 1, \dots, p_n$. Obviously, we

have $\ell(\gamma_0^M, X_j, Y) = Y^2 \equiv [Y - 0 \cdot f_j(X_j, \alpha_j^M)]^2$ and

$\ell(\hat{\gamma}_0^M, x_j, y) = y^2 \equiv [y - 0 \cdot f_j(x_j, \hat{\alpha}_j^M)]^2$. We sort the vector R_n in a descending order and select a set of variables:

$$\hat{\mathcal{M}}_{\xi_n} = \{1 \leq j \leq p_n : R_{j,n} \geq \xi_n\},$$

where ξ_n is a predefined threshold value. In the following two theorems, we show that this RSS screening is equivalent to the MLSE screening in the sense that they both possess the sure screening property and the numbers of selected variables of these two screening methods are of the same order.

Theorem 5—Suppose that Conditions (A1), (A3), (A4), (B1)–(B7) and conditions in Lemma 3 hold. By taking $\xi_n = c_7 n^{-2\kappa}$ for a sufficiently small $c_7 > 0$, there exists a c_8 such that

$$P(\mathcal{M}_* \subset \hat{\mathcal{M}}_{\xi_n}) \geq 1 - \nu_n \{\exp[-c_8 n^{1-2\kappa}/(k_n^*)^2] + n s_2 \exp[-m_2 (K_n^*)^{\frac{a}{a-1}}]\},$$

with $s_2 = s_0^{\nu_n} s_1$.

Theorem 6—Under the same conditions as those given in Theorem 5, we have

$$P[|\hat{\mathcal{M}}_{\xi_n}| \leq O\{n^{2\kappa} \lambda_{\max}(\sum_{\alpha})\}] \geq 1 - p_n \{\exp[-c_8 n^{1-2\kappa}/(k_n^*)^2] + n s_2 \exp[-m_2 (K_n^*)^{\frac{a}{a-1}}]\}.$$

3.2 Iterative NLIS-NNG Algorithm

Following Fan and Lv (2008) and Fan et al. (2009), we propose a strategy to iteratively combine a large-scale variable screening and a moderate-scale model selection for the nonlinear additive regressions to further enhance the performance of the method in terms of false selection errors. The complete algorithm works as follows:

1. For every $j \in \{1, \dots, p_n\}$, compute the marginal fit by solving

$$\min_{\beta_j, \alpha_j} \sum_{i=1}^n [y_i - \beta_j f_j(x_{ij}, \alpha_j)]^2. \quad (8)$$

Rank the covariates according to the marginal residual sum of squares

$$\text{RSS}_j = \sum_{i=1}^n [y_i - \hat{\beta}_j^M f_j(x_{ij}, \hat{\alpha}_j^M)]^2.$$

Select the top d covariates with the smallest RSS_j , or covariates with RSS_j smaller than a threshold ξ_n estimated from the random permutation. The set of selected covariates is denoted by $\mathcal{S}_1$.

2. Apply the NNG for the nonlinear additive model introduced in Section 2 on the set $\mathcal{S}_1$ to select a subset $\mathcal{M}_1$. The BIC score of the model with covariates in $\mathcal{M}_1$ is computed, denoted as $\text{BIC}(1)$.
3. For every $j \in \mathcal{M}_1^c = \{1, \dots, p_n\} \setminus \mathcal{M}_1$, minimize

$$\sum_{i=1}^n [y_i - \sum_{l \in \mathcal{M}_1} \beta_l f_l(x_{il}, \alpha_l) - \beta_j f_j(x_{ij}, \alpha_j)]^2, \quad (9)$$

with respect to $\beta_l, \alpha_l, l \in \mathcal{M}_1$ and $\beta_j, \alpha_j, j \in \mathcal{M}_1^c$. This regression reflects the additional contribution of the j th covariate conditioning on the existence of the variable set $\mathcal{M}_1$. After marginally screening similar as in Step 1 by ranking the RSS of model (9), we choose a set of covariates $\mathcal{S}_2 \in \mathcal{M}_1^c$. The NNG procedure is then applied on the set $\mathcal{M}_1 \cup \mathcal{S}_2$ to select a subset $\mathcal{M}_2$.

4. Repeat step 3 until $\mathcal{M}_k = \mathcal{M}_{k+1}$, or the size of $\mathcal{M}_k$ reaches a prespecified threshold d^* . The set of selected covariates is $\mathcal{M}_K$, where $K = \arg \min_k \text{BIC}(k)$.

In the marginal screening step, the data-driven threshold ξ_n estimated from the random permutation can be computed as follows. Suppose that covariates in $\mathcal{M}_k$ have been recruited after k iterations. We randomly permute the rows of $\{X_j\}$ to yield $\{X_j^*\}$, $j \in \mathcal{M}_k^c$ and then minimize the null model

$$\sum_{i=1}^n [y_i - \sum_{l \in \mathcal{M}_k} \beta_l f_l(x_{il}, \alpha_l) - \beta_j f_j(\tilde{x}_{ij}, \alpha_j)]^2, \quad (10)$$

with respect to β_l , α_l , $l \in \mathcal{M}_k$ and β_j , α_j , $j \in \mathcal{M}_k^c$. The threshold ξ_n is chosen to be the q -th quantile of the RSS of the null models (10). In the simulation examples, we use $q = 0$, so ξ_n is the minimum value of the permuted RSS. An alternative screening scheme is to choose d covariates with the smallest marginal RSS. This strategy is computationally more efficient than the permutation-based screening. In addition, we will show in Section 4 that this alternative thresholding scheme produces smaller numbers of false positives and also smaller prediction errors.

The proposed method can be extended to models with multiple possible parametric forms for covariates. For example, the additive components f_j in model (2) may be either linear or nonlinear. The marginal screening step in the above algorithm can be modified to include both the marginal nonlinear fit and the marginal linear fit for each covariate and the marginal RSS is then selected as the minimum value of the RSS of the linear and nonlinear marginal models. The top d covariates with the smallest RSS_j are selected, denoted as $\mathcal{S}_1$. We then apply the NNG for nonlinear additive models to the following model

$$y_i = \sum_{j \in \mathcal{S}_1} \beta_j^{(nlin)} f_j(x_{ij}, \alpha_j) + \sum_{j \in \mathcal{S}_1} \beta_j^{(lin)} x_{ij} + \varepsilon, \quad (11)$$

where $\beta_j^{(lin)}$ and $\beta_j^{(nlin)}$ are the coefficients for the linear and nonlinear components, respectively. In the practical implementation, the initial estimates of $(\beta_j^{(nlin)}, \alpha_j^T, \beta_j^{(lin)})$ need to be computed carefully, as the variable selection procedure can be very sensitive to these initial values. One may add some restrictions on the parameter searching space, such as $|\beta_j^{(nlin)} \beta_j^{(lin)}| = 0$, or use multiple initial estimates to improve the efficiency and accuracy of the nonlinear optimization. A simulation example in which the additive regression model contains both linear and nonlinear components is included in Section 4.

4 Simulation Studies

In this section, we investigate the numerical performance of our method by simulation studies. The simulation data are generated from the nonlinear additive model (2) without intercept, i.e. $\beta_0 = 0$. The covariates X_j are simulated according to the random effects model

$$X_j = 5 \frac{W_j + tU}{1+t} - 2.5, \quad j=1, \dots, p_n,$$

where $W_1, \dots, W_p$ and U are i.i.d. random variables from $\text{Unif}(0, 1)$. We use $t = 1$, so the covariates X_j are within the interval $(-2.5, 2.5)$ with pairwise correlation equal to 0.5. The sample size is set to be $n = 100$.

In the first example, all f_j 's are nonlinear functions, which are chosen to be the centered sigmoid function:

$$f_j(x, \alpha) = \frac{1}{1 + e^{-\alpha x}} - 0.5.$$

We consider the following simulation scenarios:

S1. $\beta_j = 3, 1 \leq j \leq 4$ and $\beta_j = 0, j > 4$; α_j 's are chosen equidistantly on $[1.5, 4.2]$, $1 \leq j \leq 4$; $\varepsilon_i \sim \mathcal{N}(0, 0.82^2)$;

S2. $p = 50$; $\beta_j = 3, 1 \leq j \leq 10$ and $\beta_j = 0, j > 10$; α_j 's are chosen equidistantly on $[1.5, 4.2]$, $1 \leq j \leq 10$; $\varepsilon_i \sim \mathcal{N}(0, 1.31^2)$;

The error term is chosen to give a signal-to-noise ratio (SNR) 3:1, which is defined as the square root of the ratio of the sum of nonzero components squared divided by the sum of residual squared. The total number of covariates p is set to be 50 or 200. For scenario S1, we also consider $p = 1000$ in order to illustrate the performance of our method under the ultra-high dimensional case.

We consider a second example in which the additive regression model contains both linear and nonlinear components:

$$Y = \sum_{j=1}^3 \beta_j \left(\frac{1}{1 + e^{-\alpha_j X_j}} - 0.5 \right) + \sum_{j=4}^6 \beta_j X_j + \varepsilon.$$

In this simulation, we not only need to choose the important covariates for predicting the response, but also need to determine the parametric form of each selected covariate, which in this case is either linear or sigmoid function. We set $p = 50$, $\beta_j = 3$, $\alpha_j = j + 1$, $1 \leq j \leq 3$, $\beta_j = 1$, $4 \leq j \leq 6$, $\beta_j = 0, j > 6$, and $\varepsilon_i \sim \mathcal{N}(0, 1)$. The SNR of this example is about 4:1.

We use cross-validation to select the tuning parameter λ in the nonnegative garrote and each of the above simulation settings is repeated 100 runs. The true positives (TP), false positives (FP) and prediction errors (PE) of the first example are summarized in Table 1. The prediction error is calculated on an independent test dataset of size $n/2$. The method c-NLIS-NNG in Table 1 refers to our proposed method with the direct cut-off screening scheme, where the top d covariates with the smallest marginal RSS are selected. Here we set $d = 1$ because it gives the smallest false selection rate. The method p-NLIS-NNG refers to the permutation-based screening scheme, which selects the covariates with RSS smaller than a data-driven threshold ξ_n estimated from the random permutation. We also compare the proposed method with the iterative nonparametric independence screening (INIS) method developed by Fan et al. (2011). The INIS method was designed for the nonparametric additive model. It can be applied to the variable selection in the nonlinear additive model (2) where the nonlinear functions f_j 's are assumed to be unknown and estimated nonparametrically. As suggested by Fan et al. (2011), we use the greedy modification of the INIS algorithm (g-INIS) where only one covariate is recruited in each screening step.

For scenario S1, all three methods have comparable performances in terms of selecting the true variables, but the numbers of false positives of both p-NLIS-NNG and g-INIS are larger than that of c-NLIS-NNG. For the more challenging scenario S2, c-NLIS-NNG and p-NLIS-NNG both perform well in selecting the true variables. However, g-INIS tends to miss about 2 important variables when $p = 50$ and miss about half of the important variables when $p = 200$. From the perspective of the prediction error, c-NLIS-NNG outperforms the other two methods in all cases. In Tables 2 and 3, we report the parameter estimations of the method c-NLIS-NNG conditioning on that the parameters are selected. We conclude from this simulation study that ignoring the information of knowing the nonlinear functions f_j and estimating them nonparametrically can lead to inefficient predictions and sometimes even seriously biased variable selection results. This simulation also indicates that the direct cut-off screening scheme works better than the permutation-based screening scheme in keeping down the false positives. So we will use the direct cut-off screening for the data application in Section 5.

For the second example with both linear and nonlinear components, we can see from Table 4(a) that the selected models by our method always contain all the nonzero components. The average numbers of true positives and false positives under the nonlinear and linear categories indicate both the selection accuracy as well as the accuracy of determining the corresponding parametric forms. Table 4(b) displays the parameter estimations conditioning on that the parameters are selected.

5 Application to Identify Gene Regulations

The regulation of gene expressions depends on the recognition of specific promoter sequences by transcriptional regulatory proteins and it often occurs through the coordinated action of multiple transcriptional regulators. Chen et al. (2004) used the nonlinear additive model (2) to model the transcription rate of a target gene as a combination of a set of regulatory functions from relevant regulators. Here the response Y is the transcription rate of the target gene; the covariate X_j is the gene expression of the j -th regulator; the intercept β_0 is the basal expression of the target gene; and the coefficient β_j is the regulatory capability from the j -th regulator. The regulatory function of the j -th regulator is a sigmoid function:

$$f_j\{X_j(t), \alpha_j\} = \frac{1}{1 + \exp\{-\alpha_j[X_j(t) - M_j]\}}, \quad (12)$$

where α_j is the transition rate and M_j is the mean expression level of regulator j . The goal is to select the important regulators from the potential set of regulators, i.e., regulators with $\beta_j > 0$.

We illustrate our proposed method using the dataset reported by Rabani et al. (2011). Using the short metabolic labeling of RNA with 4-thiouridine (4sU), the authors were able to distinguish recently transcribed RNA from the overall RNA and therefore obtained direct measurements of RNA transcription rate. The expression of RNA-total and RNA-4sU were measured for 254 representative signature genes during the response of mouse dendritic cells to lipopolysaccharide (LPS). There are 13 measurements for each gene, obtained at 15-min intervals over the first 3 hours after LPS stimulation. We focus on 44 genes that have

been identified to have varying degradation rates and are interested in finding genes that are related to these genes using model (2). The response is the RNA-4sU expression of a gene from these 44 genes and the predictors are the RNA-total expressions of 254 signature genes. The data are normalized by the mean of the 8 control genes (Ppp2r1a, Ndufs5, Psma7, Tomm7, Psmb4, Ndufa7, Eif4h, Capza1) and then standardized to have mean 0 and variance 1. So $M_j = 0$ in equation (12). We also center the sigmoid function (12) as in the simulation studies so there is no intercept in model (2).

We apply the proposed method to this dataset, where $n = 13$ and $p = 254$. The numbers of selected predictors for the 44 genes are between 1 and 3. Figure 1 displays the model fits for 6 randomly chosen genes. The selected regulators for these 6 genes and the corresponding parameter estimates and leave-one-out cross validation prediction errors are listed in Table 5. We found that some of our data-driven implications of gene regulations are consistent with previous biological findings. For example, Zfp36 is a known regulator of RNA stability that destabilizes its mRNA targets by binding AU-rich elements in their 3'UTR and it is known to autoregulate the stability of its own mRNA (Lai et al., 2006), which is in line with our finding that Zfp36 negatively regulates itself. Sadik et al. (2011) found in their studies that the expression of several pro-inflammatory mediators, including Il6 were decreased in Il17ra^{-/-} mice (mice lacking Il17ra) compared to wild-type mice, consistent with our result that Il17ra positively regulates Il6. We also identified some interesting genes, like Cebpb and Icosl, both of which are important regulators in immune and anti-inflammatory processes. We have shown in this example that our method can provide the biological investigators with a targeted list of genes, which could be very useful in subsequent studies.

6 Discussion

In many regression problems, the relations between the covariates and the response may be nonlinear. In this article, we propose a new method for simultaneous variable selection and parameter estimation in high-dimensional nonlinear additive models. To tackle the challenge of high dimensionality, we propose a nonlinear independence screening (NLIS) method based on the residual sum of squares of the marginal modeling, extending the idea of independence learning for the linear and generalized linear models (Fan and Lv, 2008; Fan and Song, 2010) to the nonlinear additive model setting. We have shown that our nonlinear independence screening procedure possesses the sure independence property and it is able to handle problems with non-polynomial dimensionality. After reducing the dimension of covariates to a moderate size through the NLIS procedure, we employ the nonnegative garrote (NNG) for nonlinear additive regressions to perform more refined variable selection for the submodel conditional on the covariates selected by the NLIS. We have shown that for model (2) with finite covariates, the NNG for nonlinear additive regressions has selection consistency for unimportant covariates and also estimation consistency for the parameter estimates of the important covariates. The proposed method is applied to simulated data and an real data example of identifying gene regulations to illustrate its numerical performance.

In the majority of statistical literature, the commonly adopted approach to address nonlinearity is through nonparametric modeling. However, the parametric approach has its own merit that should not be neglected. The parameters in the nonlinear regression model usually offer important biological implications. For example, the transition rate a_j in the sigmoid function (12) describes how fast the effect of a gene saturates. For small a_j , the regulatory effect is close to linear, while for large a_j , function (12) is close to a step function, indicating that a small deviation from the mean expression value of the gene leads to a large effect on the response, but this effect quickly stabilizes as the deviation increases. In practice, one can refine the parameter estimates using nonlinear least squares once the important covariates are determined, given the estimates from the nonnegative garrote as initial values. The inference of the selected model, such as the confidence intervals of the parameter estimates, can be carried out following the theories of nonlinear regression models (Jennerich, 1969; Malinvaud, 1970; Wu, 1981).

In our nonlinear regression model (2), the nonlinear functions f are assumed to be known, which requires some prior knowledge about the relations between the covariates and the response. Here we choose the sigmoid function, which is widely used in gene regulatory networks, to illustrate the proposed method. However, this is a general methodology that is also applicable to other nonlinear functions, such as the hill function and the gamma function. An important extension of our method is to include interactions of the covariates in the regression model. Another interesting problem is to relax the additive assumption in model (2) and allow more complicated nonlinear relationships. However, these topics are beyond the scope of this paper and will be studied in the future.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by the following grants: Center for Biodefense Immune Modeling, NIH/NIAID HHSN272201000055C; Center for AIDS Research, NIH/NIAID P30AI078498; Respiratory Pathogens Research Center, NIH/NIAID HHSN27220201200005C; Centers of Excellence for Influenza Research and Surveillance (CEIRS) at the University of Rochester, NIH/NIAID HHSN266200700008C; R01 AI087135; NSF grant DMS-1055210; NIH/NCI grant R01-CA149569; and two University of Rochester CTSI pilot awards (UL1RR024160) from the National Center For Research Resources.

References

- Antoniadis A, Fan J. Regularization of wavelet approximations. *Journal of the American Statistical Association*. 2001; 96:939–967.
- Biedermann S, Woods DC. Optimal designs for generalized non-linear models with application to second-harmonic generation experiments. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*. 2011; 60:281–299.
- Breiman L. Better subset regression using the nonnegative garrote. *Technometrics*. 1995; 37:373–384.
- Candes E, Tao T. The Dantzig selector: statistical estimation when p is much larger than n . *The Annals of Statistics*. 2007; 35:2313–2351.
- Chen H-C, Lee H-C, Lin T-Y, Li W-H, Chen B-S. Quantitative characterization of the transcriptional regulatory network in the yeast cell cycle. *Bioinformatics*. 2004; 20:1914–1927. [PubMed: 15044243]

- Fan J, Feng Y, Song R. Nonparametric independence screening in sparse ultra-high-dimensional additive models. *Journal of the American Statistical Association*. 2011; 106:544–557. [PubMed: 22279246]
- Fan J, Li R. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American Statistical Association*. 2001; 96:1348–1360.
- Fan J, Lv J. Sure independence screening for ultrahigh dimensional feature space. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 2008; 70:849–911. [PubMed: 19603084]
- Fan J, Peng H. Nonconcave penalized likelihood with a diverging number of parameters. *The Annals of Statistics*. 2004; 32:928–961.
- Fan J, Samworth R, Wu Y. Ultra-dimensional variable selection via independence learning: beyond the linear model. *Journal of Machine Learning Research*. 2009; 10:1829–1853.
- Fan J, Song R. Sure independence screening in generalized linear models with NP-dimensionality. *The Annals of Statistics*. 2010; 38:3567–3604.
- Frank I, Friedman J. A statistical view of some chemometrics regression tools. *Technometrics*. 1993; 35:109–148.
- Golub G, Pereyra V. Separable nonlinear least squares: the variable projection method and its applications. *Inverse Problems*. 2003; 19:R1–R26.
- Gorst-Rasmussen A, Scheike T. Independent screening for single-index hazard rate models with ultrahigh dimensional features. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 2012 In press.
- Hall P, Miller H. Using Generalized Correlation to Effect Variable Selection in Very High Dimensional Problems. *Journal of Computational and Graphical Statistics*. 2009; 18:533–550.
- Hall P, Titterton DM, Xue J-H. Tilting methods for assessing the influence of components in a classifier. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 2009; 71:783–803.
- Huang J, Horowitz JL, Ma S. Asymptotic properties of bridge estimators in sparse high-dimensional regression models. *The Annals of Statistics*. 2008; 36:587–613.
- Huang J, Horowitz JL, Wei F. Variable selection in nonparametric additive models. *The Annals of Statistics*. 2010; 38:2282–2313. [PubMed: 21127739]
- Jenrich RI. Asymptotic properties of non-linear least squares estimators. *Annals of Mathematical Statistics*. 1969; 40:633–643.
- Jeong H, Tombor B, Albert R, Oltvai Z, Barabási A-L. The large-scale organization of metabolic networks. *Nature*. 2000; 407:651. [PubMed: 11034217]
- Jiang X, Jiang J, Song X. Oracle model selection for nonlinear models based on weighted composite quantile regression. *Statistica Sinica*. 2011 In press.
- Kim Y, Choi H, Oh H-S. Smoothly clipped absolute deviation on high dimensions. *Journal of the American Statistical Association*. 2008; 103:1665–1673.
- Kosmidis I, Firth D. Bias reduction in exponential family nonlinear models. *Biometrika*. 2009; 96:793–804.
- Lai W, Parker J, Grissom S, Stumpo D, Blackshear P. Novel mRNA targets for tristetraprolin (TTP) identified by global analysis of stabilized transcripts in TTP-deficient fibroblasts. *Molecular and Cellular Biology*. 2006; 26:9196–9208. [PubMed: 17030620]
- Li R, Zhong W, Zhu L. Feature screening via distance correlation learning. *Journal of American Statistical Association*. 2012 to appear.
- Lin Y, Zhang H. Component selection and smoothing in multivariate non-parametric regression. *The Annals of Statistics*. 2006; 34:2272–2297.
- Malinvaud E. The consistency of nonlinear regressions. *Annals of Mathematical Statistics*. 1970; 41:956–969.
- Meier L, Geer V, Bühlmann P. High-dimensional additive modeling. *The Annals of Statistics*. 2009; 37:3779–3821.
- Mestl T, Plahte E, Omholt S. A mathematical framework for describing and analysing gene regulatory networks. *Journal of Theoretical Biology*. 1995; 176:291–300. [PubMed: 7475117]

- Rabani M, Levin J, Fan L, Adiconis X, Raychowdhury R, Garber M, Gnirke A, Nusbaum C, Hacohen N, Friedman N, Amit I, Regev A. Metabolic labeling of RNA uncovers principles of RNA production and degradation dynamics in mammalian cells. *Nature biotechnology*. 2011; 29:436–442.
- Ravikumar R, Liu H, Lafferty J, Wasserman L. Sparse additive models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 2009; 71:1009–1030.
- Ruhe A, Wedin P-A. Algorithm for separable nonlinear least squares problems. *SIAM Review*. 1980; 22:318–337.
- Sadik C, Kim N, Alekseeva E, Luster A. IL-17RA signaling amplifies antibody-induced arthritis. *PLoS Biology*. 2011; 6:e26342.
- Seber, GAF.; Wild, CJ. *Wiley Series in Probability and Statistics*. Hoboken, New Jersey: John Wiley & Sons, Inc; 2003. *Nonlinear Regression*.
- Tibshirani R. Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 1996; 58:267–288.
- van de Geer S. High-dimensional generalized linear models and the Lasso. *The Annals of Statistics*. 2008; 36:614–645.
- Wang H. Factor profiled sure independence screening. *Biometrika*. 2012; 99:15–28.
- Wei, BC. *Exponential family nonlinear models*. Singapore; Springer-Verlag: 1998.
- White H. Maximum likelihood estimation of misspecified models. *Econometrica*. 1982; 50:1–26.
- Wu C-F. Asymptotic theory of nonlinear least squares estimation. *The Annals of Statistics*. 1981; 9:501–513.
- Xue H, Miao H, Wu H. Sieve estimation of constant and time-varying coefficients in nonlinear ordinary differential equation models by considering both numerical error and measurement error. *Ann Stat*. 2010; 38:2351–2387. [PubMed: 21132064]
- Xue L, Zou H. Sure independence screening and compressed random sensing. *Biometrika*. 2011; 98:371–380.
- Yuan M, Lin Y. Model selection and estimation in regression with grouped variables. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 2006; 68:49–67.
- Yuan M, Lin Y. On the nonnegative garrote estimator. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 2007; 69:143–161.
- Zhang C-H, Huang J. The sparsity and bias of the lasso selection in high-dimensional linear regression. *The Annals of Statistics*. 2008; 36:1567–1594.
- Zhao P, Yu B. On model selection consistency of Lasso. *J Mach Learn Res*. 2006; 7:2541–2567.
- Zhu L-P, Li L, Li R, Zhu L-X. Model-Free Feature Screening for Ultrahigh-Dimensional Data. *Journal of the American Statistical Association*. 2012; 106:1464–1475. [PubMed: 22754050]
- Zou H. The adaptive Lasso and its oracle properties. *Journal of the American Statistical Association*. 2006; 101:1418–1429.

Figure 1. Standardized measurements (blue star) overlaid with the model fits (red) for 6 randomly chosen genes.

Table 1

Average values of the numbers of true positives (TP) and false positives (FP) and the medians of the prediction errors (PE) for the first example. c-NLIS-NNG and p-NLIS-NNG refer to our proposed method with the direct cut-off screening scheme and the permutation-based screening scheme, respectively and g-INIS refers to the iterative non-parametric independence screening method developed by Fan et al. (2011). Robust standard deviations are given in parentheses.

Simulation Setting	Method	TP	FP	PE
S1	c-NLIS-NNG	3.99 (0.02)	0.11 (0.19)	0.7070 (0.0848)
	p-NLIS-NNG	3.99 (0.02)	1.82 (1.11)	0.7443 (0.1017)
	g-INIS	4.00 (0.00)	0.93 (0.78)	0.9905 (0.1915)
	c-NLIS-NNG	3.95 (0.09)	0.27 (0.40)	0.7646 (0.1087)
	p-NLIS-NNG	3.97 (0.05)	1.91 (0.97)	0.7753 (0.0946)
	g-INIS	3.98 (0.04)	1.17 (0.97)	1.0648 (0.1722)
p = 1000	c-NLIS-NNG	3.90 (0.18)	0.52 (0.56)	0.7806 (0.1347)
	p-NLIS-NNG	3.89 (0.20)	2.67 (1.35)	0.8080 (0.1244)
	g-INIS	3.92 (0.15)	0.84 (0.85)	0.9906 (0.2211)
	c-NLIS-NNG	9.80 (0.35)	0.54 (0.64)	2.0932 (0.3186)
	p-NLIS-NNG	9.77 (0.37)	2.35 (1.31)	2.2965 (0.3428)
	g-INIS	7.90 (1.38)	1.23 (0.92)	4.0681 (1.0882)
d	c-NLIS-NNG	8.90 (1.26)	2.22 (1.42)	2.3630 (0.5701)
	p-NLIS-NNG	8.49 (1.39)	5.64 (2.30)	2.8248 (0.7689)
	g-INIS	4.58 (1.56)	2.81 (1.63)	9.0936 (2.5190)
	c-NLIS-NNG	9.77 (0.37)	2.35 (1.31)	2.2965 (0.3428)
	p-NLIS-NNG	9.77 (0.37)	2.35 (1.31)	2.2965 (0.3428)
	g-INIS	7.90 (1.38)	1.23 (0.92)	4.0681 (1.0882)

Table 2

Median of parameter estimates with median absolute deviation (MAD) in parentheses for simulation setting S1 using the method c-NLIS-NNG.

Parameters	$p = 50$	$p = 200$	$p = 1000$
$\beta_1 = 3$	3.1195 (0.5488)	2.7985 (0.4935)	3.0162 (0.5555)
$\beta_2 = 3$	3.0299 (0.3982)	3.0381 (0.3947)	2.8314 (0.3595)
$\beta_3 = 3$	3.0567 (0.3249)	2.9814 (0.2179)	2.9052 (0.2567)
$\beta_4 = 3$	3.0495 (0.2105)	3.0743 (0.2385)	2.9783 (0.2343)
$\alpha_1 = 1.5$	1.5039 (0.4945)	1.6768 (0.5351)	1.3106 (0.3106)
$\alpha_2 = 2.4$	2.3375 (0.6541)	2.2749 (0.4777)	2.5577 (0.5984)
$\alpha_3 = 3.3$	3.4061 (0.7139)	3.3341 (0.6510)	3.5423 (0.7391)
$\alpha_4 = 4.2$	4.3842 (0.6158)	4.0815 (0.6841)	4.1484 (0.8510)

Table 3

Median of parameter estimates with median absolute deviation (MAD) in parentheses for simulation setting S2 using the method c-NLIS-NNG.

Parameters	$p = 50$	$p = 200$	Parameters	$p = 50$	$p = 200$
$\beta_1 = 3$	3.1313 (0.7505)	2.8141 (0.7785)	$\alpha_1 = 1.5$	1.5785 (0.5785)	1.6189 (0.6189)
$\beta_2 = 3$	3.1759 (0.8522)	2.9269 (0.7809)	$\alpha_2 = 1.8$	1.7264 (0.7208)	2.0048 (0.9166)
$\beta_3 = 3$	3.1636 (0.5707)	2.8909 (0.6961)	$\alpha_3 = 2.1$	1.9644 (0.7856)	2.4203 (1.1869)
$\beta_4 = 3$	2.7876 (0.4590)	2.9831 (0.7133)	$\alpha_4 = 2.4$	2.483 (1.1055)	2.6863 (1.0092)
$\beta_5 = 3$	3.0042 (0.5454)	3.1175 (0.6674)	$\alpha_5 = 2.7$	2.7308 (0.9203)	2.8353 (1.1193)
$\beta_6 = 3$	3.004 (0.3352)	2.9959 (0.5895)	$\alpha_6 = 3.0$	3.1168 (0.9187)	3.3602 (1.3157)
$\beta_7 = 3$	3.0353 (0.4776)	3.1148 (0.6541)	$\alpha_7 = 3.3$	3.6671 (1.0256)	3.0564 (1.3226)
$\beta_8 = 3$	2.9735 (0.4068)	3.1066 (0.3664)	$\alpha_8 = 3.6$	3.6248 (1.0440)	3.5123 (1.3032)
$\beta_9 = 3$	3.0312 (0.4187)	3.112 (0.4006)	$\alpha_9 = 3.9$	3.8169 (1.1831)	3.9751 (1.0249)
$\beta_{10} = 3$	3.1164 (0.4113)	2.9493 (0.4248)	$\alpha_{10} = 4.2$	4.6199 (0.3801)	4.6018 (0.3982)

Table 4

(a) Average values of the numbers of true positives (TP) and false positives (FP) and the medians of the mean squared errors (MSE) for the second example with both linear and nonlinear components. Robust standard deviations are given in parentheses. (b) Median of parameter estimates with median absolute deviation (MAD) in parentheses for the second example with both linear and nonlinear components.

(a)			
	TP	FP	MSE
Combined	6.00 (0.00)	0.17 (0.40)	0.6564 (0.0994)
Nonlinear	2.37 (0.71)	0.76 (0.78)	
Linear	2.26 (0.77)	0.78 (0.79)	

(b)			
True	Estimate	True	Estimate
$\beta_1 = 3$	3.0989 (0.4574)	$\beta_4 = 1$	0.9952 (0.0981)
$\beta_2 = 3$	2.9050 (0.3134)	$\beta_5 = 1$	1.0009 (0.0607)
$\beta_3 = 3$	2.9623 (0.2646)	$\beta_6 = 1$	0.9709 (0.0706)
		$\alpha_1 = 2$	1.9741 (0.5687)
		$\alpha_2 = 3$	3.2921 (0.7273)
		$\alpha_3 = 4$	4.2079 (0.9409)

Table 5

Selected regulators and the corresponding parameter estimates and leave-one-out cross validation prediction errors (CVPE) for 6 randomly chosen genes.

Target genes	Regulators ($\hat{\beta}_j, \hat{\alpha}_j$)		CVPE
Il6	Il17ra (2.0400, 1.1731)	Dusp2 (4.1435, 0.7378)	0.3314
Il7r	Cebpb (-2.5906, 3.2332)	Hmga1 (2.1319, 2.1482)	1.3638
Zfp36	Zfp36 (-0.3415, 1.6778)	Btg2 (9.3860, 0.4893)	0.0410
Icosl	Icosl (6.8418, 1.8038)	tgif1 (-3.0583, 3.3271)	0.2570
Jun	Btg2 (3.3999, 1.3200)	Rilpl1 (-0.4657, 3.7681)	0.2509
Ifnb1	Nr4a1 (-2.5196, 2.4291)	Myd116 (8.2476, 1.0578)	0.0929