Published in final edited form as:

Chembiochem. 2014 September 5; 15(13): 1905–1910. doi:10.1002/cbic.201402255.

A Generalizable DNA-Catalyzed Approach to Peptide-Nucleic Acid Conjugation

Chih-Chi Chu^a, On Yi Wong^a, and Scott K. Silverman^{a,*}

^aDepartment of Chemistry, University of Illinois at Urbana-Champaign, 600 South Mathews Avenue, Urbana, IL 61801 (USA)

Abstract

We report DNA catalysts (deoxyribozymes) that join tyrosine-containing peptides to RNA and DNA in one step and without requiring protecting groups on either peptide or nucleic acid. Our previous efforts towards this goal required tethering the peptide to a DNA anchor oligonucleotide. Here, we establish direct in vitro selection for deoxyribozymes that use untethered, free peptide substrates. This approach enables imposition of selection pressure via reduced peptide concentration and leads to preparatively useful lower apparent K_m values of $\sim 100 \mu\text{M}$ peptide. Use of phosphorimidazolid (Imp) rather than triphosphate as the electrophile enables reactivity of either terminus (5' or 3') of both RNA and DNA. Our findings establish a generalizable means of joining unprotected peptide to nucleic acid in one step using DNA catalysts identified by in vitro selection.

Keywords

deoxyribozymes; DNA; in vitro selection; peptide-nucleic acid conjugate

Peptide-nucleic acid conjugates play key roles in many biological processes including recombination,^[1] viral propagation,^[2] topoisomerization,^[3] and activity of tumor suppressor proteins.^[4] Artificial nucleopeptides have also been used in various applications.^[5] To achieve site-specific side chain modification of biologically relevant peptide sequences, for which a fragment coupling approach is not feasible, chemical synthesis of these conjugates generally requires detailed and sophisticated approaches that involve extensive use of protecting groups.^[6] Therefore, alternative synthetic approaches are valuable when they enable direct one-step conjugation of unprotected nucleic acids to particular side chains of unprotected peptide substrates.

We have initiated efforts to identify deoxyribozymes (DNA enzymes)^[7] that catalyze various reactions of peptide substrates,^[8] including DNA-catalyzed formation of peptide-nucleic acid conjugates.^[9,10] To date, our in vitro selection experiments have used peptides that are covalently tethered to a DNA oligonucleotide anchor, which in turn is Watson-Crick base-paired to the invariant-sequence binding arm of the DNA pool (Fig. 1a). We have

scott@scs.illinois.edu, Homepage: <http://www.scs.illinois.edu/silverman>.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/cbic.20xxxxxxx>.

identified numerous deoxyribozymes that catalyze formation of peptide-RNA conjugates by reaction of a tyrosine (Tyr) or serine (Ser) hydroxyl group with the α -phosphate of a 5'-triphosphorylated RNA oligonucleotide (5'-pppRNA).^[9] Inclusion of the DNA anchor attached to the peptide enables a straightforward selection strategy based on polyacrylamide gel electrophoresis (PAGE) shift. Catalytically active DNA sequences grow substantially larger due to the DNA-catalyzed attachment of the RNA strand to the DNA-anchored peptide, noting that the RNA strand is already preattached through its remote 3'-end to the DNA pool via a catalytically dispensable loop. This size increase upon peptide-RNA conjugation enables PAGE-shift separation of the catalytically active DNA sequences.

Despite such success, the resulting deoxyribozymes have several practical disadvantages. The inclusion of the DNA anchor means that the peptide is not presented freely to candidate DNA sequences, and catalytic activity of the resulting deoxyribozymes is generally dependent on the Watson-Crick binding interaction of the anchor oligonucleotide with the pool binding arm. Also, selection pressure for the DNA catalysts to function at low peptide concentration cannot be imposed by reducing the peptide concentration during a series of iterated selection rounds, because doing so would forgo the anchoring effect; the deoxyribozymes would not bind their unanchored peptide substrates. As a consequence of these considerations, our previously identified deoxyribozymes often do not work at all when provided with the free peptide substrate that is unattached to its DNA anchor (even when the free oligonucleotide is additionally provided), and when they do function in this way, they require very high (>mM) free peptide concentration. Such failure of catalysis with the free peptide is likely observed because the deoxyribozymes were not required by the selection approach to interact sufficiently well with the peptide moiety itself, given the Watson-Crick binding energy separately provided by the DNA anchor.^[11]

In the present work, we sought to establish a different selection approach in which entirely free (not DNA-anchored) peptides are presented to the DNA pool during the key selection step of each round. The immediate difficulty in achieving this goal is that using a free peptide substrate during the selection step results in an insufficient PAGE shift to enable physical separation of the DNA sequences that just catalyzed the peptide-nucleic acid conjugation reaction. We therefore developed a new two-stage selection approach (Fig.1b). In each round, for the key selection step the DNA pool is presented with a free Tyr-containing peptide substrate that incorporates an azido (N_3) group connected to the peptide N-terminus. Catalytically active DNA sequences can attach the Tyr of this azido-peptide to the nucleic acid (e.g., 5'-pppRNA), albeit without yet adding sufficient mass to induce a useful PAGE shift. Nevertheless, the catalytically active DNA sequences become covalently linked to the azido-peptide, due to the presence of the remote RNA-DNA junction (i.e., the azido-peptide is now joined to the RNA, and the RNA was already connected to the DNA). Therefore, in a subsequent "capture" step, the DNA pool is provided with a 3'-alkyne-modified oligonucleotide and Cu^+ , such that copper-catalyzed azide-alkyne cycloaddition (CuAAC)^[12] increases the mass of *only* the active DNA sequences by the size of the alkyne-oligonucleotide and thereby enables PAGE-shift separation of these active sequences. Importantly, because the reaction step that involves DNA catalysis uses the entirely free peptide, the DNA sequences are directly challenged to interact well with this free peptide,

without depending in any way upon Watson-Crick binding energy contributed by an (absent) anchor oligonucleotide. Furthermore, selection pressure for activity at low peptide concentration can be straightforwardly imposed simply by decreasing the free peptide concentration in a series of selection rounds.^[13]

To implement the alternative selection approach of Fig. 1b, we first performed an initial set of selection experiments to establish viability. These experiments used an Ala-Tyr-Ala (AYA) tripeptide, modified on its N-terminus with an azido group on a short C₆ tether, along with a random N₄₀ DNA pool and 5'-pppRNA. Our previous experiments with DNA-catalyzed peptide-RNA conjugation used both Mg²⁺ and Mn²⁺ as catalytic cofactors,^[9] whereas several of our unrelated selections have revealed the utility of Zn²⁺ for DNA catalysis.^[8,14] Therefore, here we performed two selection experiments, using either all three of Mg²⁺/Mn²⁺/Zn²⁺ or only Zn²⁺ (for brevity, hereafter written respectively as Mg/Mn/Zn or Zn). During each key selection step, the incubation conditions were 70 mM HEPES, pH 7.5, each of 40 mM MgCl₂ and 20 mM MnCl₂ if included, 1 mM ZnCl₂, and 150 mM NaCl at 37 °C for 14 h. The free azido-AYA peptide substrate was provided at 1 mM in each round. The Mg/Mn/Zn and Zn selections had 11% yield at round 8 and 4% yield at round 10, respectively (see Fig. S1 for all selection progressions in this report).

Both selections were cloned, leading to individual deoxyribozymes whose catalytic activities were characterized (see Fig. S2 for all deoxyribozyme sequences in this report). In all, three deoxyribozymes were identified. The Mg/Mn/Zn selection provided a single deoxyribozyme, 8XJ105,^[15] that requires both Mn²⁺ and Zn²⁺ for optimal activity (even higher in the additional presence of Mg²⁺), whereas either Mg²⁺/Zn²⁺ or Zn²⁺ alone allows trace activity (Fig. 2). 8XJ105 had 45% yield using 1 mM peptide, with k_{obs} 0.41 h⁻¹ and apparent K_m of >1 mM peptide. The Zn selection led to two sequence-related deoxyribozymes that each require only Zn²⁺, with lower but reproducible yields of 10–15% with 1 mM peptide (data not shown). Two primary conclusions from this initial set of selection experiments are as follows. First, the alternative selection approach of Fig. 1b that uses the free peptide substrate is viable. Second, using different combinations of metal ions for this DNA-catalyzed reaction leads to functionally distinct deoxyribozymes. In particular, we have now found that Zn²⁺ can be a required catalytic cofactor for DNA-catalyzed synthesis of peptide-nucleic acid conjugates, noting that our previous experiments for this purpose used Mg²⁺ and Mn²⁺ but not Zn²⁺.^[9]

We also investigated the Mg/Mn/Zn ion combination with a different azido-peptide substrate, GPYSGN, which corresponds to the N-terminal sequence of the human rhinovirus 14 genome-linked (VPg) protein that is naturally connected at the Tyr to the 5'-end of viral ssRNA.^[16] After 11 rounds using 1 mM peptide, the pool yield was 16%. Cloning revealed a single deoxyribozyme sequence, 11EM103, with 60% yield, k_{obs} of 0.34 h⁻¹ at 1 mM peptide, and apparent K_m of 168 μM peptide (Fig. 2). 11EM103 requires all three of Mg²⁺, Mn²⁺, and Zn²⁺ for optimal activity; lesser but still strong activity is observed with Mn²⁺/Zn²⁺ and Mg²⁺/Zn²⁺. Although identified by selection at 1 mM Zn²⁺, 11EM103 was found to have optimum yield at 0.4 mM Zn²⁺ in the presence of Mg²⁺ and Mn²⁺ (Fig. S4). Substantial yield is retained by 11EM103 even at only 10 μM peptide, which is a practical concentration for preparative experiments (see below). Because the round 11 population had

already converged on a single sequence, further selection rounds with as little as 10 μ M peptide led only to the same sequence after cloning (data not shown). Assay of 11EM103 using GPFSGN and GPYAGN instead of GPYSGN showed equivalent reaction of GPYAGN but no reaction of GPFSGN (Fig. S6), consistent with nucleophilic reactivity of the tyrosine (but not the adjacent serine) with the pppRNA.

Until this point, all of the experiments — including those from our past efforts for this type of conjugation reaction^[9] — used solely 5'-pppRNA as the electrophile for reaction with the Tyr hydroxyl nucleophile. More generally, peptide-nucleic acid conjugates of both RNA and DNA are biologically relevant. Each of peptide-5'-RNA,^[2a,2b] peptide-5'-DNA,^[1,2c,2d,3a] and DNA-3'-peptide^[1,3b] junctions are formed naturally in various contexts. However, while 5'-pppRNA is readily prepared by in vitro transcription using a double-stranded DNA template and a phage RNA polymerase, there are strict limitations on the 5'-end sequence (purine-rich, and must begin with G or A).^[17] Several methods for solid-phase synthesis of 5'-pppRNA and 5'-pppDNA have been described, although their implementation requires access to a synthesizer.^[18] Preparation of 3'-triphosphorylated RNA or DNA is to our knowledge unreported. Therefore, deoxyribozymes that specifically require 5'-pppRNA as a substrate for peptide-nucleic acid conjugation have numerous limitations that would be relieved by using a different activated form of RNA or DNA.

Phosphorimidazolid (Imp; Fig. 3a) reacts analogously to triphosphate (ppp), but the leaving group upon nucleophilic attack at the α -phosphorus is imidazole for Imp rather than pyrophosphate for ppp. Imp is generally more reactive than ppp, by approximately 10^2 -fold. We evaluated the use of Imp as an alternative to ppp as the electrophile, noting that either terminus of either RNA or DNA can be activated in solution as Imp from a simple and readily introduced 5'- or 3'-phosphate.^[19] 5'-ImpDNA was prepared by reaction of 5'-phosphorylated DNA (5'-pDNA) with EDC and imidazole followed by a desalting column. Two selection experiments were then performed, using 5'-ImpDNA, 1 mM azido-AYA, and either Mg/Mn or Mg/Mn/Zn. The Mg/Mn selection led to 4% yield at round 13. The Mg/Mn/Zn selection led to 20% yield at round 11, and 8% yield at round 14 when the additional rounds were performed using 100 μ M peptide. We cloned the three rounds for which yields were stated, and individual deoxyribozymes were characterized.

The selection with Mg/Mn led to five unique deoxyribozymes, but each with ~10% yield at 60 h (data not shown); these DNA catalysts were not pursued further. The selection with Mg/Mn/Zn led to six and five unique deoxyribozymes at rounds 11 and 14, respectively; of the latter five, four were the same as a sequence found in round 11 and one was a new sequence. Each of the deoxyribozymes has optimum yield near 0.4 mM Zn^{2+} (Fig. S4). Yields at 10 h ranged from 12 to 45% (Fig. 3b); further incubation provided little further increase in yield. Apparent K_m values ranged from 93 μ M to >1 mM peptide (Fig. 3c). Notably, the sole deoxyribozyme that was uniquely identified in the round 14 population, 14EP125, has the lowest apparent K_m value of all of these deoxyribozymes (93 μ M). These findings demonstrate that Imp is a viable and more general alternative to ppp as the oligonucleotide electrophile for DNA-catalyzed peptide-nucleic acid conjugation. Moreover, the imposition of selection pressure via decreased peptide concentration successfully led to a

deoxyribozyme, 14EP125, that has a low apparent K_m value of $\sim 100 \mu\text{M}$ for its peptide substrate.

The relatively high reactivity of Imp correlates with modest instability, both during preparation and subsequent incubation. This instability is evident in the two closely spaced bands visible for the 5'-ImpDNA reactant in the PAGE images of Fig. 3b. Additional assays (Fig. S7) revealed that Imp is unstable to several treatments, including both annealing (heating/cooling) as commonly performed before deoxyribozyme assays and incubation under the assay conditions. When the annealing step was omitted, the final yield increased by $\sim 15\%$, limited by the instability of Imp under the assay conditions. These considerations should be noted for use of Imp substrates in such experiments.

The nine new deoxyribozymes of Fig. 2 and Fig. 3 were each evaluated with four other tyrosine-containing peptide substrates (Fig. S8). This was done to assess the catalytic activity of each deoxyribozyme with varying peptide sequences that were not used in the selection process itself. The results showed that 8XJ105 is highly tolerant of different amino acids near the reactive tyrosine, including those with potentially nucleophilic side chains (Ser, Thr, Lys, His, Arg, Asp, Glu). 11EP104 is moderately tolerant of changing the nearby amino acids, and the remaining seven deoxyribozymes are much less tolerant or intolerant. Therefore, the approach described here offers promise for sequence-general DNA catalysts for peptide-nucleic acid conjugation, although some of the deoxyribozymes are specific for particular tyrosine sequence contexts. Such specificity will be important for covalently modifying peptide and protein substrates that have more than one tyrosine residue, where discrimination on the basis of sequence context is necessary. We have not yet tested these deoxyribozymes with nucleic acid substrates of varying type (i.e., RNA versus DNA) and sequence.

Finally, the 11EM103 deoxyribozyme was evaluated for larger-scale synthesis of a peptide-nucleic acid conjugate using more realistic preparative concentrations and amounts of material. The GPYSGN peptide,^[20] the 11EM103 deoxyribozyme, and 5'-pppRNA were incubated in 1.0:1.5:2.0 mole ratio in two parallel reactions, using 1 nmol of peptide at either 10 or 20 μM . Both reactions led to substantial yield of the peptide-RNA conjugate after PAGE and extraction (Fig. 4). Based on the peptide as the limiting reagent, the yield of peptide-RNA conjugate was 22% and 33%, respectively, for 10 and 20 μM peptide, limited in part by $\sim 50\%$ gel extraction efficiency. 11EM103 has apparent K_m of 168 μM as described above; 10 or 20 μM peptide was used because higher concentrations are generally impractical for preparative experiments, especially with longer peptides.

In summary, through a series of in vitro selection experiments we have shown that free peptide substrates can be used directly to identify deoxyribozymes that create peptide-nucleic acid conjugates at Tyr residues. These deoxyribozymes can use as the conjugation partner the conventional 5'-triphosphate (ppp) electrophile, or they can use the more general phosphorimidazolid (Imp) electrophile. An Imp group can be placed readily at either the 5'-end or 3'-end of either RNA or DNA, and any of these activated oligonucleotide substrates can be incorporated readily into our selection approach.^[21] Therefore, on the basis of the CuAAC selection design and the use of the Imp electrophile, both of which are established

experimentally in this study, we anticipate that deoxyribozymes can be identified to form any desired Tyr-RNA or Tyr-DNA conjugate from the two corresponding unprotected peptide and nucleic acid substrates. Likely this approach can be expanded to include Ser (and Thr) as well,^[9b] and our success with deoxyribozymes that have phosphatase activity^[8] suggests that DNA-catalyzed nucleopeptide formation should be achievable with large proteins. In addition, the general ability to impose selection pressure by decreasing the peptide substrate concentration provides a rational means to seek DNA catalysts that function at low peptide concentrations. Our ongoing work aims to apply this class of deoxyribozymes to prepare and study biologically relevant peptide-nucleic acid conjugates.

Experimental Section

Peptides were prepared by solid-phase synthesis using Fmoc Rink amide MBHA resin as described.^[8] To introduce the azido group on the peptide N-terminus, the final coupling reaction was performed for 12 h using 5 equivalents (1 mmol) of 6-azidohexanoic acid (Chem-Impex), 5 equivalents (126 mg, 1 mmol) of diisopropyl carbodiimide (DIC; Acros), and 5 equivalents (135 mg, 1 mmol) of HOBt (Acros) in 5 mL of anhydrous dimethylformamide (DMF; Acros). At the end of the synthesis procedure, ethanedithiol was omitted in the step where the peptide was cleaved from the resin, because here no peptide included cysteine.

DNA oligonucleotides prepared by solid-phase synthesis were obtained from Integrated DNA Technologies (Coralville, IA) or synthesized on an ABI 394 instrument using reagents from Glen Research. 5'-pppRNA oligonucleotides were prepared by in vitro transcription using synthetic DNA templates and T7 RNA polymerase.^[17a] All oligonucleotides were purified by denaturing PAGE with running buffer 1× TBE (89 mM each Tris and boric acid and 2 mM EDTA, pH 8.3) as described previously.^[23] 5'-ImpDNA oligonucleotides were prepared by treatment of 5'-pDNA with EDC and imidazole followed by desalting as described in the Supporting Information.

The in vitro selection and cloning procedures were performed essentially as described previously in related efforts,^[9a,9c] with full procedures in the Supporting Information. Individual deoxyribozymes were prepared by solid-phase synthesis and assayed under single-turnover conditions using 3'-³²P-radiolabeled substrates. For 5'-pppRNA, radiolabeling was achieved starting from the unlabeled RNA using 5'-³²P-pCp and T4 RNA ligase.^[24] For 5'-ImpDNA, the unlabeled DNA was 3'-³²P-radiolabeled using α -³²P-dCTP and terminal deoxytransferase, followed by heat inactivation, 5'-phosphorylation using T4 polynucleotide kinase and ATP, PAGE purification, and finally formation of Imp by treatment with EDC and imidazole followed by desalting; see full procedure in the Supporting Information.

The general single-turnover assay procedure for each deoxyribozyme (Fig. 2 and Fig. 3) was as follows. A 10 μ L sample containing 0.4 pmol of 3'-³²P-radiolabeled 5'-pppRNA or 5'-ImpDNA substrate and 10 pmol of deoxyribozyme was annealed in 5 mM HEPES, pH 7.5, 15 mM NaCl, and 0.1 mM EDTA by heating at 95 °C for 1 min and cooling on ice for 5 min (for 8XJ105, 20 pmol of a short oligonucleotide complementary to the 3'-binding arm was

additionally included). The peptide-nucleic acid conjugation reaction was initiated by bringing the sample to 20 μL total volume containing 50 mM ($-\text{Zn}^{2+}$) or 70 mM ($+\text{Zn}^{2+}$) HEPES, pH 7.5, 1–5000 μM azido-peptide, $\text{Mg}^{2+}/\text{Mn}^{2+}/\text{Zn}^{2+}$ (chloride salts) as appropriate, and 150 mM NaCl. The sample was incubated at 37 $^{\circ}\text{C}$, and 2 μL aliquots were quenched at appropriate times with 5 μL of stop solution (80% formamide, 1 $\times$ TBE [89 mM each Tris and boric acid, 2 mM EDTA, pH 8.3], 50 mM EDTA, and 0.025% each bromophenol blue and xylene cyanol). For 8XJ105, 10 pmol of a decoy oligonucleotide (complementary to the deoxyribozyme's 40 nt catalytic region along with 10 nt of binding arms on either side) was added to each aliquot before PAGE, to displace the deoxyribozyme from the substrate and product. Samples were separated by 20% PAGE and quantified with a PhosphorImager. k_{obs} values were obtained by fitting the complete yield versus time data directly to first-order kinetics. Alternatively, for initial-rate kinetics as used for all apparent K_{m} plots, data for the initial linear portion (up to 1 or 2 h) was fit to a straight line.

The preparative-scale experiment using the 11EM103 deoxyribozyme (Fig. 4) was performed as follows. For 10 or 20 μM peptide, a 50 or 25 μL sample containing 1.5 nmol of 11EM103 and 2 nmol of 5'-pppRNA was annealed in 5 mM HEPES, pH 7.5, 15 mM NaCl, and 0.1 mM EDTA by heating at 95 $^{\circ}\text{C}$ for 1 min and cooling on ice for 5 min. The reaction was initiated by bringing the sample to 100 or 50 μL total volume containing 70 mM HEPES, pH 7.5, 1 nmol of azido-GPYSGN, 40 mM MgCl_2 , 20 mM MnCl_2 , 0.7 or 1 mM ZnCl_2 , and 150 mM NaCl. The sample was incubated at 37 $^{\circ}\text{C}$ for 60 h, precipitated with ethanol, and purified by 20% PAGE. The product and recovered substrate were quantified after gel extraction by UV absorbance (A_{260}). The product was analyzed by mass spectrometry on a Bruker UltrafleXtreME MALDI-TOF mass spectrometer with matrix 3-hydroxypicolinic acid in positive ion mode at the UIUC School of Chemical Sciences Mass Spectrometry Laboratory (instrument purchased with support from NIH S10RR027109A). $[\text{M}+\text{H}]^+$ calcd. 6268.7, found 6272.3 (10 μM peptide) and 6268.2 (20 μM peptide), = +0.06% and -0.008%.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by grants to S.K.S. by the National Institutes of Health (R01GM065966), the Defense Threat Reduction Agency (HDTRA1-09-1-0011), and the National Science Foundation (CHE0842534). We thank Benjamin Brandsen and Tania Velez for assistance with phosphorimidazolid preparation.

References

1. Grindley NDF, Whiteson KL, Rice PA. Annu. Rev. Biochem. 2006; 75:567–605. [PubMed: 16756503]
2. a) Ambros V, Baltimore D. J. Biol. Chem. 1978; 253:5263–5266. [PubMed: 209034] b) Rothberg PG, Harris TJ, Nomoto A, Wimmer E. Proc. Natl. Acad. Sci. USA. 1978; 75:4868–4872. [PubMed: 217003] c) Hermoso JM, Salas M. Proc. Natl. Acad. Sci. USA. 1980; 77:6425–6428. [PubMed: 6779279] d) Bamford DH, Mindich L. J. Virol. 1984; 50:309–315. [PubMed: 6368864] e) Rozovics JM, Virgen-Slane R, Semler BL. PLoS ONE. 2011; 6:e16559. [PubMed: 21408223]

3. a) Tse Y-C, Kirkegaard K, Wang JC. *J. Biol. Chem.* 1980; 255:5560–5565. [PubMed: 6155377] b) Wang JC. *Annu. Rev. Biochem.* 1985; 54:665–697. [PubMed: 2992360]
4. Samad A, Carroll RB. *Mol. Cell. Biol.* 1991; 11:1598–1606. [PubMed: 1705009]
5. a) Corey DR, Schultz PG. *Science*. 1987; 238:1401–1403. [PubMed: 3685986] b) Brenner S, Lerner RA. *Proc. Natl. Acad. Sci. USA*. 1992; 89:5381–538. [PubMed: 1608946] c) Vivès E, Schmidt J, Pèlerin A. *Biochim. Biophys. Acta*. 2008; 1786:126–138. [PubMed: 18440319] d) Said Hassane F, Saleh AF, Abes R, Gait MJ, Lebleu B. *Cell. Mol. Life Sci.* 2010; 67:715–726. [PubMed: 19898741] e) Juliano RL, Ming X, Nakagawa O. *Acc. Chem. Res.* 2012; 45:1067–1076. [PubMed: 22353142]
6. a) Flohr S, Jungmann V, Waldmann H. *Chem. Eur. J.* 1999; 5:669–681. b) Forget D, Boturyn D, Defrancq E, Lhomme J, Dumy P. *Chem. Eur. J.* 2001; 7:3976–3984. [PubMed: 11596939] c) Venkatesan N, Kim BH. *Chem. Rev.* 2006; 106:3712–3761. [PubMed: 16967918] d) Lu K, Duan Q-P, Ma L, Zhao D-X. *Bioconjug. Chem.* 2010; 21:187–202. [PubMed: 19856957]
7. a) Peracchi A. *Chem. Bio. Chem.* 2005; 6:1316–1322. b) Silverman SK. *Chem. Commun.* 2008:3467–3485. c) Schlosser K, Li Y. *Chem. Biol.* 2009; 16:311–322. [PubMed: 19318212] d) Silverman SK. *Angew. Chem. Int. Ed.* 2010; 49:7180–7201.
8. Chandrasekar J, Silverman SK. *Proc. Natl. Acad. Sci. USA*. 2013; 110:5315–5320. [PubMed: 23509279]
9. a) Pradeepkumar PI, Höbartner C, Baum DA, Silverman SK. *Angew. Chem. Int. Ed.* 2008; 47:1753–1757. b) Sachdeva A, Silverman SK. *Chem. Commun.* 2010; 46:2215–2217. c) Wong, O. Ph.D. thesis, University of Illinois at Urbana-Champaign; 2011. d) Velez TE, Singh J, Xiao Y, Allen EC, Wong O, Chandra M, Kwon SC, Silverman SK. *ACS Comb. Sci.* 2012; 14:680–687. [PubMed: 23088677]
10. Murakami H, Saito H, Suga H. *Chem. Biol.* 2003; 10:655–662. [PubMed: 12890539] Morimoto J, Hayashi Y, Iwasaki K, Suga H. *Acc. Chem. Res.* 2011; 44:1359–1368. [PubMed: 21711008] These “flexizyme” ribozymes developed by Suga and coworkers for tRNA 3'-aminoacylation are useful for preparing tRNA-amino acid conjugates by reaction at the carboxyl group of the amino acid, i.e., charging of tRNAs. However, these ribozymes do not enable side chain conjugation, and they do not function with polypeptide rather than single amino acid substrates.
11. Using a DNA “anchor” oligonucleotide that lacks base pairing to the pool binding arm, but still enables direct PAGE-shift selection, is not a viable strategy for two reasons. First, as a practical concern, the DNA-peptide conjugates are prepared on the nanomole scale, and prohibitive amounts of these conjugates would be required to perform selections that use mM concentrations; in contrast, with a base-paired anchor sequence, only tractable μ M concentrations are required. Second, as a conceptual consideration, ultimately we do not wish to require artificial attachment of a DNA oligonucleotide to the peptide substrate in order to achieve modification of that peptide's side chain.
12. a) Hong V, Presolski SI, Ma C, Finn MG. *Angew. Chem. Int. Ed.* 2009; 48:9879–9883. b) El-Sagheer AH, Brown T. *Chem. Soc. Rev.* 2010; 39:1388–1405. [PubMed: 20309492]
13. In a previous study, we attempted a selection approach analogous to that in Fig. 1b, except that the peptide had a free N-terminal amino group, and the capture reaction was reductive amination with the periodate-oxidized ribonucleotide 3'-terminus of a DNA strand. This effort was thwarted due to an unanticipated DNA-catalyzed reductive amination reaction during the capture step involving nucleophilic reactivity of a guanosine N²-amino group present on the pppRNA substrate. Wong O, Mulcrone AE, Silverman SK. *Angew. Chem. Int. Ed.* 2011; 50:11679–11684.
14. a) Hoadley KA, Purtha WE, Wolf AC, Flynn-Charlebois A, Silverman SK. *Biochemistry*. 2005; 44:9217–9231. [PubMed: 15966746] b) Kost DM, Gerdt JP, Pradeepkumar PI, Silverman SK. *Org. Biomol. Chem.* 2008; 6:4391–4398. [PubMed: 19005599]
15. Each deoxyribozyme in this study was named as, for example, 8XJ105, where 8 is the round number, XJ is the systematic alphanumeric designation for the particular selection, and 105 is the clone number.
16. a) Stanway G, Hughes PJ, Mountford RC, Minor PD, Almond JW. *Nucleic Acids Res.* 1984; 12:7859–7875. [PubMed: 6093056] b) Werner G, Rosenwirth B, Bauer E, Seifert JM, Werner FJ, Besemer J. *J. Virol.* 1986; 57:1084–1093. [PubMed: 3512851]

17. a) Milligan JF, Groebe DR, Witherell GW, Uhlenbeck OC. *Nucleic Acids Res.* 1987; 15:8783–8798. [PubMed: 3684574] b) Coleman TM, Wang G, Huang F. *Nucleic Acids Res.* 2004; 32:e14. [PubMed: 14744982]
18. a) Paul N, Springsteen G, Joyce GF. *Chem. Biol.* 2006; 13:329–338. [PubMed: 16638538] b) Zlatev I, Lavergne T, Debart F, Vasseur JJ, Manoharan M, Morvan F. *Org. Lett.* 2010; 12:2190–2193. [PubMed: 20420425] c) Zlatev I, Lackey JG, Zhang L, Dell A, McRae K, Shaikh S, Duncan RG, Rajeev KG, Manoharan M. *Bioorg. Med. Chem.* 2013; 21:722–732. [PubMed: 23260577] d) Goldeck M, Tuschl T, Hartmann G, Ludwig J. *Angew. Chem. Int. Ed.* 2014; 53 Early View.
19. a) Chu BC, Wahl GM, Orgel LE. *Nucleic Acids Res.* 1983; 11:6513–6529. [PubMed: 6622259] b) Wang F, Elbaz J, Willner I. *J. Am. Chem. Soc.* 2012; 134:5504–5507. [PubMed: 22404383]
20. The peptide included the azido group that was present during selection, because 11EM103 has ~2-fold reduced yield with the peptide that lacks the azido group.
21. For 3'-Imp substrates, the loop in Fig. 1 that enables the selection process can be switched from the right side to the left side, as we have already achieved in unrelated efforts.^[8,22]
22. a) Sachdeva A, Chandra M, Chandrasekar J, Silverman SK. *Chem. Bio. Chem.* 2012; 13:654–657. b) Walsh SM, Sachdeva A, Silverman SK. *J. Am. Chem. Soc.* 2013; 135:14928–14931. [PubMed: 24066831] c) Brandsen BM, Hesser AR, Castner MA, Chandra M, Silverman SK. *J. Am. Chem. Soc.* 2013; 135:16014–16017. [PubMed: 24127695]
23. a) Flynn-Charlebois A, Wang Y, Prior TK, Rashid I, Hoadley KA, Coppins RL, Wolf AC, Silverman SK. *J. Am. Chem. Soc.* 2003; 125:2444–2454. [PubMed: 12603132] b) Wang Y, Silverman SK. *Biochemistry.* 2003; 42:15252–15263. [PubMed: 14690435]
24. Wong O, Pradeepkumar PI, Silverman SK. *Biochemistry.* 2011; 50:4741–4749. [PubMed: 21510668]

Figure 1.

Selection strategies for DNA-catalyzed peptide-RNA conjugation. a) Previously used selection approach in which a DNA-anchored peptide is presented to the 5'-triphosphorylated RNA substrate (pppRNA), itself joined at its remote 3'-terminus to the DNA pool to enable PAGE-shift selection. This particular illustration shows reaction of the disulfide-linked CYA tripeptide.^[9c] b) Alternative selection approach established in this study, in which a free (non-anchored) peptide reacts with pppRNA. The peptide includes an N-terminal C₆-tethered azido group. After the DNA-catalyzed selection step, active DNA

sequences are captured by DNA-splinted reaction of the azide with a 3'-alkyne oligonucleotide (CuAAC), leading to a substantial PAGE shift that enables their separation. See Supporting Information for full procedures.

Figure 2.

Deoxyribozymes from the initial experiments with the AYA and GPYSGN peptide substrates, to establish the alternative selection approach of Fig. 1b. Assay conditions: 20 nM 3'-³²P-radiolabeled 5'-pppRNA, 0.5 μ M deoxyribozyme, 1 mM, 100 μ M, or 10 μ M azido-AYA peptide as indicated, 50 mM (–Zn²⁺) or 70 mM (+Zn²⁺) HEPES, pH 7.5, divalent metal ions as indicated (chloride salts; concentrations as stated for panel b), and 150 mM NaCl at 37 °C. (a) PAGE images for kinetic assays. Representative timepoints at $t = 30$ s, 2 h, 24 h. S = substrate; P = product. See Fig. S3 for quantification of these data. (b)

Kinetic plots. k_{obs} values for 8XJ105 at 1 mM and 100 μM peptide, respectively: 0.41 and 0.10 h^{-1} (40 mM Mg^{2+} , 20 mM Mn^{2+} , 1 mM Zn^{2+}). The yield for 8XJ105 was optimal at 0.4–1.0 mM Zn^{2+} (Fig. S4). k_{obs} values for 11EM103 at 1 mM, 100 μM , and 10 μM peptide, respectively: 0.34, 0.19, and 0.029 h^{-1} (40 mM Mg^{2+} , 20 mM Mn^{2+} , 0.4 mM Zn^{2+}). The yield for 11EM103 was optimal at 0.4 mM Zn^{2+} (Fig. S4). 11EM103 was clearly more active than 8XJ105 at lower peptide concentration. See Fig. S5 for apparent K_{m} plots.

Figure 3.

Alternative Imp electrophile, and deoxyribozymes for reaction of Tyr with 5'-ImpDNA. a) Structures of 5'-pppRNA and 5'-ImpDNA electrophiles, and their reactions with a Tyr nucleophile (X = RNA or DNA). b) Kinetic assays of deoxyribozymes for reaction of Tyr with 5'-ImpDNA. Assay conditions: 20 nM 3'-³²P-radiolabeled 5'-ImpDNA, 0.5 μM deoxyribozyme, 1 mM azido-AYA peptide, 70 mM HEPES, pH 7.5, 40 mM MgCl₂, 20 mM MnCl₂, 0.4 mM ZnCl₂, and 150 mM NaCl at 37 °C. *t* = 30 s, 2 h, 10 h for PAGE images (S = substrate, P = product). The 5'-ImpDNA substrate partially degrades to 5'-pDNA

throughout the reaction, partially limiting the yield (see Fig. S7 for details). Not shown are data for 11EP104 (very similar to 11EP101) and 11EP126 (very similar to 11EP103). k_{obs} values, top to bottom in figure legend: 0.53, 0.38, 0.39, 0.30, and 0.34 h^{-1} . c) Determination of peptide apparent K_{m} values using initial-rate kinetics. See Fig. S5 for quantified data with error bars.

Figure 4.

Preparative-scale reaction of the GPYSGN peptide with 5'-pppRNA, catalyzed by the 11EM103 deoxyribozyme. Reaction conditions: 10 or 20 μM azido-GPYSGN peptide, 15 or 30 μM 11EM103 deoxyribozyme, 20 or 40 μM pppRNA, 70 mM HEPES, pH 7.5, 40 mM MgCl_2 , 20 mM MnCl_2 , 0.7 mM or 1.0 mM ZnCl_2 , and 150 mM NaCl at 37 $^\circ\text{C}$ for 60 h (100 or 50 μL total volume; 1.0 nmol peptide, 1.5 nmol deoxyribozyme, 2.0 nmol pppRNA). Zn^{2+} concentrations were optimized for each peptide concentration using smaller-scale assays with 3'- ^{32}P -radiolabeled 5'-pppRNA, where extraction was unnecessary and yields were 60–70% (Fig. S9). The PAGE image was acquired by UV shadowing; the deoxyribozyme itself is clearly evident higher in both reaction lanes (not shown). The respective amounts of peptide-RNA conjugate product and recovered pppRNA were 219 and 734 pmol (10 μM peptide) and 326 and 584 pmol (20 μM peptide). From these values, the efficiency of gel extraction was only 46–48%, indicating that the 22% and 33% yields of peptide-RNA conjugate are about two-fold lower than the DNA-catalyzed chemical yields.