Published in final edited form as:

Biochem Biophys Res Commun. 2014 August 29; 451(3): 382–388. doi:10.1016/j.bbrc.2014.07.103.

Correlation of Glucocorticoid-mediated E4BP4 upregulation with altered expression of pro- and anti-apoptotic genes in CEM human lymphoblastic leukemia cells

Jessica A. Beach, Laura J. Nary, Rebeka Hovanessian, and Rheem D. Medh[§]

Department of Biology, California State University Northridge, Northridge, CA 91330-8303, USA

Jessica A. Beach: jessica.beach@cshs.org; Laura J. Nary: ljnary@gmail.com; Rebeka Hovanessian: rebeka888@gmail.com; Rheem D. Medh: rheem.medh@csun.edu

Abstract

In *C.elegans*, motoneuron apoptosis is regulated via a *ces-2 – ces-1 – egl-1* pathway. We tested whether human CEM lymphoblastic leukemia cells undergo apoptosis via an analogous pathway. We have previously shown that E4BP4, a *ces-2* ortholog, mediates glucocorticoid (GC)-dependent upregulation of *BIM*, an *egl-1* ortholog, in GC-sensitive CEM C7-14 cells and in CEM C1-15 mE#3 cells, which are sensitized to GCs by ectopic expression of E4BP4. In the present study, we demonstrate that the human *ces-1* orthologs, *SLUG* and *SNAIL*, are not significantly repressed in correlation with *E4BP4* expression. Expression of *E4BP4* homologs, the PAR family genes, especially *HLF*, encoding a known anti-apoptotic factor, was inverse to that of *E4BP4* and *BIM*. Expression of pro- and anti-apoptotic genes in CEM cells was analyzed via an apoptosis PCR Array. We identified *BIRC3* and *BIM* as genes whose expression paralleled that of *E4BP4*, while *FASLG*, *TRAF4*, *BCL2A1*, *BCL2L1*, *BCL2L2* and *CD40LG* as genes whose expression was opposite to that of *E4BP4*.

Keywords

glucocorticoids; E4BP4; BIM; leukemia; ces-2; CEM

Introduction

Glucocorticoids (GCs) are used widely in chemotherapy either as a primary cytotoxic drug targeting cancer cells (as in acute leukemia, lymphoma and multiple myeloma) or to reduce inflammation, prevent allergic reactions or reduce the side effects of chemotherapy [1]. The pharmacological actions of GC therapy include, among other effects, the suppression of the immune response and shrinkage of T-lymphocyte population. Loss of T-lymphocytes via apoptosis is a major cause of the immunosuppressive actions of GCs [1], however, the

© 2014 Elsevier Inc. All rights reserved.

[§]Corresponding author: Tel: 818-677-3338.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

mechanisms by which GCs evoke an apoptotic response is still not clear. Relapse of leukemia after GC therapy is often associated with development of resistant cells that no longer respond to GC-evoked apoptosis [2]. The primary action of GCs is mediated through their interactions with the glucocorticoid receptor (GR), a transcription factor from the nuclear receptor family, which modulates up or downregulation of genes containing GC Responsive Elements (GRE) or through interactions with other transcription factors, coactivators and corepressors [2]. We have demonstrated that GR transcriptional activity correlates with susceptibility of cells to GC-evoked apoptosis. Several laboratories have analyzed changes in gene expression profiles induced by GCs in an effort to identify candidate genes modulating GC-evoked apoptosis of leukemic lymphoid cells [3, 4].

We have established a crucial relationship between GC-dependent upregulation of *E4BP4* and sensitivity to GC-evoked apoptosis in the human leukemic CEM cell culture model [5]. *E4BP4* (or NFIL3) is a bZIP transcription factor with a role in anti-inflammatory response, circadian oscillation, apoptosis regulation, and immune cell development [6- 8]. *E4BP4* is an evolutionarily conserved homolog of the pro-apoptotic *C.elegans* death specification gene *ces-2*, which is known to downregulate the downstream survival gene *ces-1* to allow the upregulation of the pro-apoptotic gene *egl-1*, resulting in apoptosis in neurosecretory motor neuron cells. An analogous pathway in humans has been speculated (Figure 1) [9]. Human orthologs of *ces-1*, *SLUG* and *SNAIL*, have pro-survival functions. Expression of *SLUG* is abundant in hematopoietic stem cells (HSCs) and in undifferentiated progenitor cells but not in pro T and pro-B cells, or beyond [10]. Both Slug and Snail have been shown to promote tumor invasiveness, cell survival, protect from apoptosis, and induce chemoresistance [10-11]. The role of Slug and Snail in GC-evoked lymphoid cell apoptosis, or their regulation by *E4BP4*, has not yet been explored. The human ortholog of *egl-1*, Bim (a BH-3 only protein from the Bcl-2 family), an important regulator of the intrinsic pathway of apoptosis, has been previously shown to play a vital role in apoptosis in response to various stimuli [12, 13], including GCs [3, 14]. *BIM* is upregulated in GC sensitive human leukemic CEM C7-14 cells in conjunction with GC-evoked apoptosis. We have demonstrated that ectopic *E4BP4* expression restores sensitivity to GC-evoked apoptosis in a refractory CEM sister cell line, CEM C1-15, via restoration of *BIM* induction [15].

E4BP4 belongs to the PAR (Proline and Acid Rich) subfamily of bZIP transcription factors, based on its DNA-binding specificity, even though it lacks a PAR region (Figure 3A) [16]. Other key members of this subfamily include hepatic leukemia factor (HLF), thyrotrophic embryonic factor (TEF) and albumin D-site binding protein (DBP). *E4BP4* and PAR proteins regulate transcription by binding to an *E4BP4* response element (EBPRE), which has a consensus sequence of (G/A)T(G/T)A(C/T) GTAA (C/T) (21). Generally, *E4BP4* and other PAR proteins have temporally antiparallel expression patterns and antagonize each other's action, as has been extensively studied in the circadian regulation of clock genes [17]. In contrast to the apoptosis inducing action of *E4BP4*, TEF is reported to have antiapoptotic activity [18], while HLF is highly expressed in leukemic stem cells (LSCs) and has been implicated in chemoresistance [19]. While the PAR family transcription factors have been characterized as transcriptional activators, *E4BP4* is a repressor [16]. In several cases *E4BP4* has been shown to act as a transactivator, and a transactivation domain (TAD)

has been reported just upstream of the transrepressor domain (TRD). E4BP4 has also been recognized as a survival factor in a cell-type specific context. In pro-B lymphocytes, E4BP4 activates downstream genes and stimulates IL-3 (a cytokine) promoting cell survival [20]. Similarly, TEF and DBP have been shown to promote oxidative stress induced apoptosis via upregulation of BIK, a BH3-only proapoptotic member of the Bcl2 family [21].

To investigate the role of E4BP4 in GC-evoked apoptosis of human leukemic cells, we have previously created a cell line, CEM C1-15mE#3, that ectopically expresses mouse E4BP4 in the GC-resistant CEM C1-15 cells [15]. These cells are sensitized to GC-evoked apoptosis and *BIM* upregulation, analogous to the GC-sensitive sister cell line, CEM-C7-14. In the present study, we have further investigated GC-mediated regulation of *SLUG*, *SNAIL*, *HLF*, *TEF* and *DBP* in all three cell lines, in an effort to understand the role of these genes in GC-evoked apoptosis, and to evaluate the contribution of E4BP4 in their expression. Additionally, we have evaluated the GC-dependent regulation of transcripts of genes implicated in apoptosis using the RT² Profiler™ PCR Array from SA Biosciences.

Materials & Methods

Reagents

Dexamethasone (Dex) was purchased from EMD Biosciences (Madison, WI). RNA was extracted using TRIzol reagent (Cat #15596-026) from Invitrogen (La Jolla, CA). Reagents for reverse transcription (RT), including M-MLV reverse transcriptase, oligo(dT)₁₅ primer, RNasin® Ribonuclease inhibitor, dNTP mix were purchased from Promega Life Sciences (Madison, WI). SYBR® JumpStart™ Tag Ready Mix (Cat # 4438) for real-time qPCR was from Sigma-Aldrich (St. Louis, MO). RNeasy® Mini Kit (Cat #74104) and DNase Column (Cat #79254) were from Qiagen (Valencia, CA). RT² First Strand Kit (Cat #C-03), RT² RNA Quality Control (QC) PCR Array (Cat #PAHS-999) and RT² Profiler™ modified human apoptosis PCR Arrays (CAHP09369) were from SA Biosciences. Other reagent grade chemicals were purchased from Fisher Scientific (Pittsburgh, PA) or Sigma-Aldrich.

Cell culture

The CCRF-CEM derived human T-ALL cell lines CEM C7-14 and CEM C1-15 are sensitive and resistant to GCs, respectively, and are generous gifts from Dr. E. B. Thompson, University of Texas Medical Branch, Galveston. The CEM C1-15m#3 subclone, stably transfected with mouse E4BP4, has been characterized previously and is sensitive to GC-evoked apoptosis [15]. Cells were maintained in log phase at 37°C in a 5% CO₂ incubator in RPMI 1640 containing L-Glutamine (Catalog #50-020-PB) from Cellgro (Manassas, VA) supplemented with 5% heat-inactivated fetal bovine serum (FBS, Cat #S11150) from Atlanta Biologicals (Lawrenceville, GA).

Reverse Transcription and RT-qPCR (real-time-quantitative PCR) analysis

Cells were treated for 24 h at a density of 5×10^5 cells/ml with either absolute ethanol (control/vehicle) or 1 μ M Dex and total RNA was extracted in 1 ml of TRIzol reagent, according to instructions from the manufacturer. Reverse transcription was carried out by incubating 7 μ g of RNA for 3h at 42°C with 0.5 mg of oligo(dT)₁₅, 1 μ l (~200U) of M-MLV

reverse transcriptase, 0.5 mM dNTP mix, and 100 U of RNase inhibitor. For RT-qPCR, SYBR® Green JumpStart™ Taq ReadyMix was mixed with 250nM each of forward and reverse primers (Table 1) and 1 µl of reverse transcription product in a final volume of 25µl, and run on an ABI Prism 7300 Real Time PCR System.

To quantitate the relative expression levels, the cycle threshold (Ct) values for each sample were used to calculate fold inductions using the Pfaffl method formula:

$(E)^{C_{\text{Target}}(\text{control-sample})/(E)^{C_{\text{Reference}}(\text{control-sample})}}$, where E represents primer efficiency which was calculated for each sample reaction using LinRegPCR software. The reference gene was β-actin. Statistical significance was calculated using a paired Student's t-test, with a p = 0.05 denoting high significance and p = 0.1 denoting moderate significance.

RT² Profiler™ PCR Array

CEM-C7-14, CEM-C1-15, and CEM-C1-15m#3 cells were treated at a density of 5.0×10^5 cells/ml with ethanol or 1 µM Dex for a 24-hour period. Approximately 20 ml of cell suspension ($\sim 1.0 \times 10^7$ cells) were harvested using the Qiagen RNeasy® Mini Kit following the manufacture's protocol. Residual DNA was removed using a DNase column. First strand cDNA synthesis was completed with the RT² First Strand Kit, using 1.5 µg of total RNA. RNA quality was assessed on the ABI Prism 7300 using the RT² RNA QC PCR Array according to the manufacturer's protocol. Each RT sample that passed the quality control check was analyzed using a modified human apoptosis PCR array. Data were analyzed using C_t method through the SABioscience PCR Array Data Analysis Web Portal. Data presented are averages of two independent experiments.

Results and Discussion

We have previously demonstrated that E4BP4-dependent upregulation of Bim contributes to GC-evoked apoptosis of human leukemic CEM cells, using a cell line derived from the GC-resistant CEM C1-15 cells that ectopically expresses mouse E4BP4 (CEM C1-15mE#3) [15]. Here, we have extended our studies to evaluate the role of E4BP4 in GC-dependent regulation of genes whose protein products are known to crosstalk with E4BP4, and genes implicated in apoptosis.

Parallel regulation of *E4BP4* and *BIM* by Dex

As reported previously, significant Dex-mediated upregulation of *E4BP4* (8.1 and 7.2-fold) and *BIM* (6.3 and 6.5) transcripts was seen in CEM C7-14 and CEM C1-15mE#3 cells respectively, while CEM-C1-15 cells showed a distinctly blunted response for both *E4BP4* (2.2-fold) and *BIM* (2.3-fold) (Figure 2A). One thousand base pairs of the *BIM* promoter was scanned for the presence of EBPRES consensus sequences using transcription factor binding site prediction programs, AliBaba 2.1 and TRANSFAC, but none were detected (data not shown), suggesting that E4BP4 must regulate *BIM* expression either through other factors such as Slug/Snail or in conjunction with other transcription factors.

E4BP4 does not upregulate *BIM* via inhibition of *SLUG* and *SNAIL* expression

To test whether E4BP4-dependent apoptosis follows an *E4BP4*>*SLUG*/*SNAIL*>*BIM* pathway parallel to the evolutionarily conserved *ces-2*>*ces-1*>*egl-1* pathway in CEM cells (Figure 1), we tested whether expression of the survival genes *SLUG* and *SNAIL* had an inverse relationship with *E4BP4* and *BIM*. In contrast to this hypothesis, *SLUG* transcripts were upregulated by 1 μ M Dex in CEM C7-14 and CEM C1-15mE#3 cells by 3.4 and 5.9-fold respectively, and were marginally downregulated (0.84-fold) in CEM C1-15 cells. Data between individual cell lines were not statistically significant, as determined by a paired Student's *t*-test, except for CEM C1-15 and CEM C1-15mE#3, where *p* was 0.036 (Figure 2B). *SNAIL* expression was minimally repressed by 1 μ M Dex in all three cell lines (0.7-0.9-fold) (Figure 2B), and again the difference between pairs of cell lines was not significant, but the extent of repression (compared to ethanol treated controls) for CEM C7-14 and CEM C1-15mE#3 cells was marginally significant with a *p* value of 0.09 and 0.06 respectively. These data suggest that Dex-evoked *SNAIL* downregulation may facilitate *BIM* upregulation.

Inverse regulation of PAR-family transcripts in comparison to E4BP4

HLF, TEF and DBP share the same bZIP domain and DNA binding specificity as E4BP4, but lack a TRD (Figure 3A), and are known to have an opposite outcome on transcription of target genes compared to E4BP4. They have been shown to promote cell survival and chemoresistance, and their expression profile is opposite to that of E4BP4. To determine whether HLF, TEF and DBP have a role in sensitivity or resistance to GCs in the CEM cell model, we determined the effect of 1 μ M Dex on transcript levels of all three genes in GC-sensitive CEM C7-14 and CEM C1-15mE#3 cells and in GC-resistant CEM C1-15 cells (Figure 3B). All three transcripts were upregulated by Dex in the GC-resistant CEM C1-15 cells, and downregulated in GC-sensitive CEM C7-14 cells. In CEM C1-15mE#3 cells, both *HLF* and *DBP* were downregulated, while *TEF* was marginally upregulated. All data were significant for *HLF* with *p* = 0.05, and for *TEF* between CEM C7-14 and CEM C1-15 cells, with *p* = 0.10. While *E4BP4* is upregulated by Dex in all three cell lines, the extent of upregulation is 4-fold higher for the sensitive lines than for the resistant line (Figure 2A). Our results suggest a mutually antagonistic relationship between PAR proteins and E4BP4 in modulating cell survival and apoptosis, respectively (Figure 3C).

Regulation of genes implicated in apoptosis

In an effort to identify additional genes playing a role in GC-evoked apoptosis, and their dependence on E4BP4, gene expression profiles for CEM-C7-14, CEM-C1-15, and CEM-C1-15mE#3 cells were analyzed using the Human RT² Profiler™ Apoptosis PCR Array. Total RNA was extracted from cells treated for 24h with ethanol or 1 μ M Dex, and first-strand synthesis was performed and PCR Arrays were run as described in the Methods section. Two independent experiments, with six groups each (control and treated groups for each of three cell types) were performed. The software, RT² Profiler™ PCR Data Analysis v3.5, available through the SA Biosciences website, was used for data analysis. Briefly, control and Dex-treated threshold cycles for each gene were normalized to the average of three housekeeping genes, and the expression levels were calculated using the $\Delta\Delta$ Ct method. Only those genes that passed the software's quality check for all groups were analyzed.

Transcripts with a Ct value greater than 35 in any sample were excluded from the analysis. Additionally, any gene that showed a variation of greater than 3 standard deviations within the two replicates for any group was eliminated from the analysis.

For a gene to be considered as regulated by GCs in an E4BP4-dependent manner, it must be reproducibly up- or downregulated by GCs in the GC-sensitive CEM C7-14 and CEM C1-15mE#3 cells but not in the GC-resistant CEM C1-15 cells, suggesting that ectopic expression of E4BP4 restores GC-dependent transcriptional regulation in CEM C1-15mE#3 cells. Following these criteria, two genes, *BCL2L11* (*BIM*) and *BIRC3* showed a significantly greater upregulation by 1 μ M Dex in the sensitive cells compared to the resistant cells (Table 2A), although both these genes were upregulated >2-fold in CEM C1-15 cells. CEM C1-15 cells express functional GR, which has been shown to be transcriptionally active with a blunted response to GC stimulus, hence modest regulation of GC-dependent genes is to be expected (4). Bim is a well-known mediator of GC-evoked apoptosis of lymphoid leukemic cells [14], and we have shown that its expression correlates with *E4BP4* expression [15] (Figure 2A). *BIRC3* or IAP-1 (Inhibitor of Apoptosis-1) is known to bind to TRAFs (Tumor necrosis factor Receptor Associated Factors) and inhibit apoptosis by interfering with activation of caspases. *BIRC3* has been previously shown to be upregulated in correlation with GC-evoked apoptosis, and has been proposed as a rescue mechanism activated in cells trying to escape apoptosis [4]. TNFSF10, a gene encoding a member of the TNF family, that has been shown to induce apoptosis in tumor cells by activating caspases, was upregulated only in CEM C7-14 cells, suggesting that its upregulation by Dex is not mediated by E4BP4.

Six genes were upregulated by GCs in the resistant CEM C1-15 cells (Table 2B). *CD27*, was also upregulated in CEM C1-15mE#3 cells, hence may not have any correlation with E4BP4 expression. *CD27* is a member of the TNF receptor superfamily, and is thought to play a role in survival of activated T cells [22]. Three genes, *CFLAR*, *BAD* and *AKT1* are only upregulated in CEM C1-15 cells, and are not significantly regulated in the sensitive cells, and may contribute to resistance to GCs in CEM C1-15 cells. *CFLAR* is structurally similar to caspase-8, but lacks caspase activity and inhibits apoptosis [23]. *BAD* is a known proapoptotic member of the Bcl2 family [24], and CEM C1-15 cells resist apoptosis in spite of *BAD* upregulation. The proapoptotic activity of *BAD* is inhibited by its phosphorylation via *AKT* and MAP kinase [24]. It is possible that in CEM C1-15 cells, *BAD* is inactivated by phosphorylation. In support of this hypothesis, *AKT1*, encoding a serine/threonine-protein kinase survival factor known to phosphorylate *BAD* (31), is also upregulated only in the resistant cells. Two genes, *BCL2L2* and *CD40LG* are upregulated >2-fold in CEM C1-15 cells but downregulated >2-fold in CEM C1-15mE#3 cells, suggesting that E4BP4 may play a role in their repression. *BCL2L2* encodes the anti-apoptotic member of the Bcl2 family, Bcl-w, and has been shown to block GC-induced apoptosis [25]. *CD40LG* is a negative regulator of apoptosis, is expressed on T-cell surfaces, and engages CD40 on the B cell surface to facilitate immunoglobulin class switching. It has been shown to be negatively regulated by E4BP4 [26].

Of the three genes downregulated >2-fold in CEM-C7-14 cells (Table 2C), *FASLG* was also downregulated in the other two cell lines, but most strongly in CEM C1-15mE#3 cells. It is

a member of the TNF superfamily, known to antagonize with GCs in regulating T cell apoptosis [27]. *BCL2A1* and *TRAF4* were downregulated in CEM C7-14 cells >2-fold, in CEM C1-15mE#3 cells by about 1.8-fold and were not significantly altered in CEM C1-15 cells, thus E4BP4 may contribute to their repression. *BCL2A1* encodes an anti-apoptotic member of the Bcl2 family, Bfl1 [25], and its downregulation concurs with sensitivity of CEM cells. *TRAF4* is a TNF receptor associated factor and a negative regulator of NF κ B activation and apoptosis [28], thus its downregulation promotes apoptosis.

Six genes were downregulated significantly (>2-fold) only in CEM C1-15mE#3 cells, suggesting that ectopic E4BP4 expression may facilitate their repression. Notable among these was *SNAIL* encoding Snail, an ortholog of *C.elegans* *ces-1* gene which is a survival factor, and may be an intermediate in E4BP4-dependent upregulation of *BIM* (Figure 1). This is in agreement with our data in Figure 2B. Another notable gene in this group was *BCL2L1*, an anti-apoptotic member of the Bcl2 family, was upregulated 1.93-fold in CEM C1-15 cells, and its downregulation via E4BP4 may sensitize these cells to GC-evoked apoptosis.

Regulation of Bcl-2 family genes

The RT² ProfilerTM Apoptosis PCR Array included several pro- and anti-apoptotic genes belonging to the Bcl-2 family. Because of the critical role these genes are known to play in the execution of apoptosis, data corresponding to these genes was extracted and charted separately (Figure 4). Of the anti-apoptotic genes analyzed, all except *MCL1* were upregulated in the GC-resistant CEM C1-15 cells, although only *BCL2L1* and *BCL2L2* were upregulated approximately 2-fold (Figure 4A). The sensitive cells showed a general downregulation of anti-apoptotic genes, with a 2-fold repression of *BCL2L1* and *BCL2L2* in CEM C1-15mE#3 cells, suggesting that E4BP4 expression played a role in their repression. Among the pro-apoptotic genes analyzed, only *BIM* was markedly upregulated in the two sensitive cell lines compared to the resistant line (Figure 4B). Since these data are from duplicate arrays, they are not statistically significant, however the trends are informative.

Our studies demonstrate that E4BP4-mediated upregulation of *BIM* may be independent of *SLUG/SNAIL* repression, unlike the orthologous pathway in *C.elegans*. Expression patterns of the PAR family genes *TEF*, *HLF* and *DBP* are opposite to that of *E4BP4*, with *HLF* showing the most statistical significance. Ectopic expression of E4BP4 in the resistant CEM C1-15 cells restores GC-mediated regulation of key pro-and anti-apoptotic genes, depicting a complex network of gene regulatory signals that ultimately result in apoptosis.

Acknowledgments

Supported in part by NIH MBRS-SCORE (5SC3GM 081099) grant awarded to RDM, the CSUN Office of Graduate Studies, Research and International Programs, and the CSUN College of Science & Mathematics. We thank Dr. E. B. Thompson (UTMB, Galveston, TX) for kindly providing CEM-C7-14 and CEM-C1-15 cells. JAB and LJM performed some of this work as part of their MS thesis projects at CSUN.

References

- Schmidt S, Rainer J, Ploner C, et al. Glucocorticoid-induced apoptosis and glucocorticoid resistance: molecular mechanisms and clinical relevance. *Cell Death Differ.* 2004; 11(Suppl 1):S45–55. [PubMed: 15243581]
- Schaaf MJ, Cidlowski JA. Molecular mechanisms of glucocorticoid action and resistance. *JSMBB.* 2002; 83:37–48.
- Wang Z, Malone MH, He H, et al. Microarray analysis uncovers the induction of the proapoptotic BH3-only protein Bim in multiple models of glucocorticoid-induced apoptosis. *J Biol Chem.* 2003; 278:23861–23867. [PubMed: 12676946]
- Medh RD, Webb MS, Miller AL, et al. Gene expression profile of human lymphoid CEM cells sensitive and resistant to glucocorticoid-evoked apoptosis. *Genomics.* 2003; 81:543–555. [PubMed: 12782123]
- Priceman SJ, Kirzner JD, Nary LJ, et al. Calcium-dependent upregulation of E4BP4 expression correlates with glucocorticoid-evoked apoptosis of human leukemic CEM cells. *Biochem Biophys Res Commun.* 2006; 344:491–499. [PubMed: 16630563]
- Kamizono S, Duncan GS, Seidel MG, et al. Nfil3/E4bp4 is required for the development and maturation of NK cells in vivo. *J Exp Med.* 2009; 206:2977–2986. [PubMed: 19995955]
- Ohno T, Onishi Y, Ishida N. The negative transcription factor E4BP4 is associated with circadian clock protein PERIOD2. *Biochem Biophys Res Commun.* 2007; 354:1010–1015. [PubMed: 17274955]
- Zhang W, Zhang J, Kornuc M, et al. Molecular cloning and characterization of NF-IL3A, a transcriptional activator of the human interleukin-3 promoter. *Mol Cell Biol.* 1995; 15:6055–6063. [PubMed: 7565758]
- Seidel MG, Look AT. E2H-HLF usurps control of evolutionarily conserved survival pathways. *Oncogene.* 2001; 20:5718–5725. [PubMed: 11607821]
- Becker MW, Clarke MF. SLUGging away at cell death. *Cancer Cell.* 2002; 2:249–251. [PubMed: 12398886]
- Kurrey NK, Jalgaonkar SP, Joglekar AV, et al. Snail and Slug mediate radioresistance and chemoresistance by antagonizing p53-mediated apoptosis and acquiring a step-like phenotype in ovarian cancer cells. *Stem Cells.* 2009; 27:2059–2068. [PubMed: 19544473]
- Tan TT, Degenhardt K, Nelson DA, et al. Key roles of BIM-driven apoptosis in epithelial tumors and rational chemotherapy. *Cancer Cell.* 2005; 7:227–238. [PubMed: 15766661]
- Gordon PM, Fisher DE. Role for the proapoptotic factor BIM in mediating imatinib-induced apoptosis in a c-KIT-dependent gastrointestinal stromal tumor cell line. *J Biol Chem.* 2010; 285:14109–14114. [PubMed: 20231287]
- Lu J, Quearry B, Harada H. p38-MAP kinase activation followed by BIM induction is essential for glucocorticoid-induced apoptosis in lymphoblastic leukemia cells. *FEBS Lett.* 2006; 580:3539–3544. [PubMed: 16730715]
- Beach JA, Nary LJ, Hirakawa Y, et al. E4BP4 facilitates glucocorticoid-evoked apoptosis of human leukemic CEM cells via upregulation of Bim. *J Mol Signaling.* 2011; 6:13.
- Cowell IG. E4BP4/NFIL3, a PAR-related bZIP factor with many roles. *Bioessays.* 2002; 24:1023–1029. [PubMed: 12386933]
- Mitsui S, Yamaguchi S, Matsuo T, et al. Antagonistic role of E4BP4 and PAR proteins in the circadian oscillatory mechanism. *Genes & Development.* 2001; 15:995–1006. [PubMed: 11316793]
- Inukai T, Inaba T, Dang J, et al. TEF, an antiapoptotic bZIP transcription factor related to the oncogenic E2A-HLF chimera, inhibits cell growth by down-regulating expression of the common β chain of cytokine receptors. *Blood.* 2005; 105:4437–4444. [PubMed: 15665112]
- Dorn DC, Kou CA, Png KJ, et al. The effect of cantharidins on leukemic stem cells. *Int J cancer.* 2009; 124:2186–2199. [PubMed: 19123473]
- Ikushima S, Inukai T, Inaba T, et al. Pivotal role for the NFIL3/E4BP4 transcription factor in interleukin 3-mediated survival of pro-B lymphocytes. *Proc Natl Acad Sci USA.* 1997; 94:2609–2614. [PubMed: 9122243]

21. Ritchie A, Gutierrez O, Fernandez-Luna JL. PAR bZIP-bik is a novel transcriptional pathway that mediates oxidative stress-induced apoptosis in fibroblasts. *Cell Death and Differentiation*. 2009; 16:838–846. [PubMed: 19219069]
22. Hendricks J, Xiao Y, Borst J. CD27 promotes survival of activated T cells and complements CD28 in generation and establishment of the effector T cell pool. *J Exp Med*. 2003; 198:1369–1380. [PubMed: 14581610]
23. Fulda S. Targeting c-FLICE-like inhibitory protein (CFLAR) in cancer. *Expert Opinion on Therapeutic Targets*. 2013; 17:195–201. [PubMed: 23252466]
24. Datta SR, Dudek H, Tao X, et al. Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. *Cell*. 1997; 91:231–241. [PubMed: 9346240]
25. Tzifi F, Economopoulou C, Gourgoutis D, et al. The role of BCL2 family of apoptosis regulator proteins in acute and chronic leukemias. *Adv Hematol*. 2011; 2012:524308. [PubMed: 21941553]
26. Zhao M, Liu Q, Liang G, et al. E4BP4 overexpression: a protective mechanism in CD4+ T cells from SLE patients. *J Autoimmun*. 2013; 41:152–160. [PubMed: 23340290]
27. Novac N, Baus D, Dostert A, et al. Competition between glucocorticoid receptor and NF κ B for control of the human FasL promoter. *FASEB J*. 2006; 20:1074–1081. [PubMed: 16770006]
28. Xie P. TRAF molecules in cell signaling and in human diseases. *J Mol Signaling*. 2013; 8:7.

Highlights

- An apoptotic pathway akin to *C. elegans ces2-ces1-egl1* is evaluated in CEM cells.
- E4BP4 does not alter *BIM* expression via *SLUG/SNAIL* repression.
- PAR family gene expression profiles are inverse to that of *E4BP4*.
- An Apoptosis PCR Array identified a set of E4BP4-dependent genes.
- A number of Bcl2 family genes are regulated by E4BP4.

Figure 1. Conserved pathway for apoptosis

Analogy between the known pathway for apoptosis in neurosecretory motor neurons of *C. elegans* (left) and the hypothetical pathway in human leukemic T cells (right). In *C. elegans*, *ces-2* upregulates the proapoptotic *egl-1* via repression of the survival gene *ces-1*, to eventually activate CED-3 and trigger apoptosis. In human leukemic T cells, upregulation of *BIM* and subsequent inhibition of anti-apoptotic members of the Bcl-2 family leads to activation of caspases. Expression of the *ces-2* ortholog *E4BP4* has been shown to facilitate GC-mediated *BIM* upregulation, but the role of *SLUG/SNAIL* is not clear. Several other proteins including p27kip1, cyclin D3, c-Myc and c-Jun have been shown to facilitate apoptosis.

Figure 2. Dex-mediated upregulation of *BIM* and *E4BP4* in sensitive CEM cells does not correlate with downregulation of *SLUG*/*SNAIL*

CEM C7-14, CEM C1-15 and CEM C1-15mE#3 cells were treated with ethanol or 1 μ M Dex and real-time qPCR was performed using primers (Table 1) specific for *E4BP4*, *BIM* (Panel A), *SLUG* and *SNAIL* (Panel B). Fold change in expression of each transcript by 1 μ M Dex was calculated by the Pfaffl method using *ACTINB* as a reference. Data represent averages $\pm$ SD from three independent experiments. Statistical significance was calculated using a paired Student's t-test, shown next to the charts. Shaded boxes represent significant differences with p-values $\leq$ 0.05.

Figure 3. Inverse relationship between E4BP4 and other PAR-family homologs

Panel A: Representation of protein structural homology between *C. elegans* CES-2, and human PAR family proteins including E4BP4, HLF, TEF and DBP. TAD = transactivation domain; TRD = transrepression domain; PAR = proline & acid rich domain. **Panel B:** Real-time qPCR was performed on ethanol or 1 μ M Dex treated CEM cells and using primers (Table 1) specific for *TEF*, *HLF* and *DBP*. Fold change in expression was calculated by the Pfaffl method using *ACT1NB* as a reference. Data represent averages $\pm$ SD from three independent experiments. Statistical significance was calculated using a paired Student's t-test, shown below the chart. Dark shaded boxes represent significant differences with p-values ≤ 0.05 , light shaded boxes represent p-values ≤ 0.10 . **Panel C:** Proposed relationship between E4BP4 and other PAR genes and their effect on cell survival and apoptosis.

Figure 4. Correlation of sensitivity to GCs and expression of Bcl-2 family genes
Data for anti-apoptotic (Panel A) and pro-apoptotic (Panel B) members of the Bcl-2 family were extracted from the RT² Profiler™ Apoptosis PCR Array.

Table 1
PCR Primers

Transcript	Forward Primer	Reverse Primer	Product Size (bp)
E4BP4	5'ATGGGGAATTCTTTCTCTGG3'	5'CTTTGATCCGAGCTTGTGT3'	250
Bim	5'CAGATATGCGCCAGAGATA3'	5'ACCAGGCGGACAATGTAAC3'	163
Slug	5'CCCTGAAGATGCATATTCGGAC3'	5'CTTCTCCCCCGTGTGAGTTCTA3'	116
Snail	5'GCTGCAGGACTCTAATCCAGAGTT3'	5'GACAGAGTCCCAGATGAGCATTG3'	130
HLF	5'GCGTGCTCAGGTCCTGCTG3'	5'AGGCCCCAGGAATGCCGACT3'	143
TEF	5'GCTCTCCAGTGTGCCAGGCG3'	5'CGGGCATCCCGTGACCGTTT3'	189
DBP	5'TCCAGGTGCCGAGGAGCAG3'	5'GTTCTCCTTGAGCCGCCGGG3'	104
β-actin	5'AGTCCTCTCCCAAGTCCACA3'	5'CACGAAGGCTCATCATTCAA3'	130

Table 2
Genes Regulated Two-Fold or Higher by 1 μ M Dexamethasone

Symbol	Unigene	RefSeq	Description	Gene Name	CEM C7-14	CEM C1-15	C1-15 mE#3
A: Genes Upregulated in CEM C7-14 Cells							
BIRC3	Hs.127799	NM_001165	Baculovirus IAP repeat containing 3	API1/API2	9.80	3.28	13.83
BCL2L11 (BIM)	Hs.469658	NM_006538	BCL2-like 11 (apoptosis facilitator)	BAM/BIM	8.81	2.90	5.45
TNFSF10	Hs.478275	NM_003810	Tumor necrosis factor (ligand) superfamily, member 10	APO2L/Apo-2L	2.07	-1.06	-1.56
B: Genes Upregulated in CEM C1-15 Cells							
CD27	Hs.355307	NM_001242	CD27 molecule	SI52/T14	-1.07	3.15	2.08
CFLAR	Hs.390736	NM_003879	CASP8 and FADD-like apoptosis regulator	CASH/CASP8AP	1.16	2.41	1.47
BCL2L2	Hs.410026	NM_004050	BCL2-like 2	BCL-W/BCLW	-1.46	2.23	-2.01
BAD	Hs.370254	NM_004322	BCL2-associated agonist of cell death	BBC2/BCL2L8	1.12	2.17	1.40
CD40LG	Hs.592244	NM_000074	CD40 ligand	CD154/CD40L	-1.33	2.15	-2.14
AKT1	Hs.525662	NM_005163	V-akt murine thymoma viral oncogene homolog 1	AKT/PKB	-1.41	2.03	-1.01
C: Genes Downregulated in CEM C7-14 Cells							
FASLG	Hs.2007	NM_000639	Fas ligand (TNF superfamily, member 6)	APTILG1/CD178	-4.37	-3.29	-11.04
BCL2A1	Hs.227817	NM_004049	BCL2-related protein A1	ACC-1/ACC-2	-2.32	1.42	-1.85
TRAF4	Hs.8375	NM_004295	TNF receptor-associated factor 4	CART1/MLN62	-2.02	1.25	-1.82
D: Genes Downregulated in CEM C1-15m#3 Cells							
SNAIL (SNAIL)	Hs.48029	NM_005985	Snail homolog 1 (Drosophila)	SNAIL	-1.15	1.09	-5.05
CIDEA	Hs.249129	NM_001279	Cell death-inducing DFFA-like effector a	CIDE-A	-1.19	1.44	-4.13
TNFRSF25	Hs.462529	NM_003790	Tumor necrosis factor receptor superfamily, member 25	APO-3/DDR3	-1.27	1.96	-2.80
PYCARD	Hs.499094	NM_013258	PYD and CARD domain containing	ASC/CARD5	1.43	1.54	-2.65
NOL3	Hs.513667	NM_003946	Nuclear protein 3 (apoptosis repressor with CARD domain)	ARC/CARD2	-1.14	1.14	-2.02
BCL2L1	Hs.516966	NM_138578	BCL2-like 1	BCL0XL/S	-1.42	1.93	-2.01

Genes upregulated >2-fold are in red; genes downregulated >2-fold are in blue.