Published in final edited form as:

Brain Res Bull. 2014 August ; 0: 54–60. doi:10.1016/j.brainresbull.2014.07.003.

Eating habits modulate short term memory and epigenetical regulation of brain derived neurotrophic factor in hippocampus of low-and high running capacity rats

Ferenc Torma¹, Zoltan Bori¹, Erika Koltai¹, Klara Felszeghy¹, Gabriella Vacz¹, Lauren Koch², Steven Britton², Istvan Boldogh³, and Zsolt Radak¹

¹Research Institute of Sport Science, Semmelweis University, Budapest, Hungary

²Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, Michigan, USA

³Department of Microbiology and Immunology, University of Texas, Medical Branch at Galveston, Galveston, Texas 77555, USA

Abstract

Exercise capacity and dietary restriction (DR) are linked to improved quality of life, including enhanced brain function and neuro-protection. Brain derived neurotrophic factor (BDNF) is one of the key proteins involved in the beneficial effects of exercise on brain. Low capacity runner (LCR) and high capacity runner (HCR) rats were subjected to DR in order to investigate the regulation of BDNF. HCR-DR rats out-performed other groups in a passive avoidance test. BDNF content increased significantly in the hippocampus of HCR-DR groups compared to control groups ($p < 0.05$). The acetylation of H3 increased significantly only in the LCR-DR group. However, Chip-assay revealed that the specific binding between acetylated histone H3 and BDNF promoter was increased in both LCR-DR and HCR-DR groups. In spite of these increases in binding, at the transcriptional level only, the LCR-DR group showed an increase in BDNF mRNA content. Additionally, DR also induced the activity of cAMP response element-binding protein (CREB), while the content of SIRT1 was not altered. Peroxisome proliferator-activated receptor gamma coactivator-1 alpha (PGC-1 α) was elevated in HCR-DR groups. But, based on the levels of nuclear respiratory factor-1 and cytochrome c oxidase, it appears that DR did not cause mitochondrial biogenesis. The data suggest that DR-mediated induction of BDNF levels includes chromatin remodeling. Moreover, DR does not induce mitochondrial biogenesis in the hippocampus of LCR/HCR rats. DR results in different responses to a passive avoidance test, and BDNF regulation in LCR and HCR rats.

© 2014 Elsevier Inc. All rights reserved.

Correspondence: Zsolt Radak, Ph.D, Institute of Sport Science, Faculty of Physical Education and Sport Science, Semmelweis University, Alkotás u. 44.TF Budapest, Hungary, Tel: +36 1 3565764, Fax: +36 1 356 6337, radak@tf.hu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Brain-derived neurotrophic factor; histone; epigenetics; dietary restriction; exercise capacity

1. Introduction

It has been postulated that aerobic running could be important in the evolution of homo sapiens (Bramble and Lieberman, 2004; Dumas et al., 2007; Lieberman and Bramble, 2007). A life-style associated with physical activity provides excellent conditions for the regulation of brain derived neurotrophic factor (BDNF) (Mattson, 2012). Early in the evolution of humans it would have been difficult to have consistent, daily nutrition. Eating on alternate days during the hunting/gathering period would be considered normal. Therefore, genes apparently developed to store energy, based on irregular eating conditions (Chakravarthy and Booth, 2004). Alternate day eating has been used as a tool to study the effects of mild dietary restriction (DR) (Brown-Borg and Rakoczy, 2013; Pesic et al., 2010; Rodriguez-Bies et al., 2010; Terzibasi et al., 2009), and found to result in decreases in body mass and cardio protection (Garcia Ramos, 1980), and enhanced muscle performance, and increased lipid metabolism (Rodriguez-Bies et al., 2010). DR is well known to increase life expectancy (Sohal and Weindruch, 1996; Weindruch and May, 1995). Moreover, it also has been reported that DR can promote brain function (Qiu et al., 2012; Quintas et al., 2012; Singh et al., 2012) as it has been suggested that DR can mediate BDNF signaling and consequently play a role in memory neuro-protection, synaptic plasticity, and neurogenesis (Kumar et al., 2009; Radak et al., 2013b; Rich et al., 2010; Rothman et al., 2012).

The neuro-protective effects of physical exercise are well demonstrated (Barrientos et al., 2011; Radak et al., 2010; Radak et al., 2013a; Stranahan and Mattson, 2011) and include up-regulation of BDNF levels as well as improved brain function as measured by the Morris maze and passive avoidance tests (Griesbach et al., 2009; Radak et al., 2006; Sarga et al., 2013; Wu et al., 2013).

An experimental model was created for the running capacity of rats (Koch and Britton, 2001). Low capacity runners (LCR) and high capacity runners (HCR) were identified. LCR rats were found to have shorter life-spans and decreased resistance to oxidative stress (Hart et al., 2013; Koch et al., 2012; Koch et al., 2011b). LCR rats also readily develop cardiovascular disorders, and markers of metabolic syndrome, such as visceral adiposity, increased blood pressure, dyslipidemia and insulin resistance when compared to HCR rats (Koch et al., 2011a). Therefore, due to the shorter life span, adiposity, and impaired brain function of LCR rats, this appears to be an excellent model to study the effects of DR. HCR rats also have been shown to outperform LCR rats in cognitive tests (Wikgren et al., 2012).

Regular exercise and nutrition could have an effect of epigenetics (Barnes and Ozanne, 2011; Milagro et al., 2011; Radak et al., 2012; Vaquero and Reinberg, 2009). Reversible modification of lysine residues of histone proteins, such as acetylation and deacetylation, can readily alter the gene expression levels of proteins (Kaelin and McKnight, 2013; Sanders et al., 2013; Shankar et al., 2013), and these modifications can be inherited. It would be interesting to know whether DR can alter the level of histone acetylation at the promoter

regions of BDNF, and hence, have a long term effect on BDNF signaling. Therefore, the present investigation was carried out to study the effects of alternate day eating on the epigenetical modulation of BDNF in hippocampus of low and high running capacity rats.

2. Materials and methods

2.1. Animals

Selectively bred rat strains differing in intrinsic aerobic capacity – low capacity runners (LCR) and high capacity runners (HCR) – were used in this study (Koch and Britton, 2001). Endurance running capacity was assessed on a treadmill and the total distance run during a speed-ramped exercise test was used as a measure of maximal aerobic capacity. Rats with the highest running capacity were bred to produce the HCR strain and rats with poor running capacity were bred to generate LCR rats. A subgroup of male rats from generation 22 was phenotyped for intrinsic treadmill running capacity when 11 weeks old, at the University of Michigan (Ann Arbor, USA) and then shipped via air freight to Semmelweis University (Budapest, Hungary) for further study. Investigations were carried out according to the requirements of The Guiding Principles for Care and Use of Animals, EU, and approved by The Ethics Committee of Semmelweis University.

2.2. Animal setting and Dietary restriction (DR)

LCR and HCR male rats, aged 13 months, were assigned to control LCR (LCR-C), dietary restricted LCR (LCR-DR), control HCR (HCR-C) and dietary restricted HCR (HCR-DR) groups ($n =$ six rats per group). Dietary restriction was performed by feeding the animal every other day for 16 weeks (Fig. 1).

2.3. Tissue samples

After decapitation, brains were rapidly removed, hippocampi from both hemispheres were dissected and flash frozen in liquid nitrogen. All samples were kept at -80°C until homogenized. Samples thawed on ice were quickly pre-homogenized for 10 s at medium speed (IKA's Ultra-Turrax homogenizer) in two volume of PBS (pH 7.4) to gain a crude homogenate. Crude bilateral hippocampal homogenates were divided into four aliquots for ChIP, qRT-PCR, Western blot, and ELISA assays. Aliquots for ChIP assay were processed immediately, whereas the others were stored at -80°C awaiting further processing.

2.4. Passive avoidance test

The test was performed according to the step-through method as described earlier (Sarga et al., 2013). The apparatus consists of a two-compartment acrylic box with a lighted chamber connected to a darkened chamber by a guillotine door. As soon as the rats entered the dark chamber, they received an electrical shock (0.5 mA, 1 sec). The latency times for entering the dark chamber were measured in the training test, after 24h and ten days later in the retention test.

2.5. Assays

2.5.1. Ach3-ChIP—This assay was based on the Acetyl-Histone H3 Immunoprecipitation Assay Kit (Cat# 17-245Upstate/Millipore). In brief, DNA formaldehyde was added directly

to 100 μ l aliquots of total pre-homogenate (preH) representing about 30 mg of hippocampus in a final concentration of 1% and incubated for 15 minutes at 37°C. Formaldehyde was removed by washing of the pre-homogenate two times with ice cold PBS containing protease inhibitor (1mM phenylmethylsulfonyl fluoride (PMSF)), and pelleted for four minutes at 700 $\times$ g at 4°C.

Supernatants were diluted tenfold in ChIP Dilution Buffer (0.01% SDS, 1.1% Triton X-100, 1.2mM EDTA, 16.7mM Tris-HCl, pH 8.1, 167mM NaCl) supplemented with protease inhibitor PMSF. An aliquot of this chromatin solution was stored at -20°C to quantify the amount of DNA present in different samples before immunoprecipitation (Input).

Anti-acetyl histone H3 (AcH3) specific antibody (Catalog # 06-599, Upstate/Millipore) was added to 2 μ g of supernatant fraction (chromatin solution) and incubated overnight on ice with constant agitation. The other 1ml of chromatin solution was saved for a normal rabbit serum IgG (sIgG) (Cat# I5006, Sigma-Aldrich, St. Louis, MO, USA) no-antibody control. An immune complex of antigen-antibody-chromatin was collected by adding 60 μ l of salmon sperm DNA/Protein agarose slurry for one hour at 4°C with rotation. A beads complex was collected by centrifugation for 1 min at 700 $\times$ g at 4°C in a microcentrifuge.

Beads were washed for 3-5 minutes at 4°C with rotation with 1ml of each of the buffers: Low Salt Immune Complex Wash Buffer (0.1% SDS, 1% Triton X-100, 2mM EDTA, 20mM Tris-HCl, pH 8.1, 150mM NaCl), High Salt Immune Complex Wash Buffer (0.1% SDS, 1% Triton X-100, 2mM EDTA, 20mM Tris-HCl, pH 8.1, 500mM NaCl), LiCl Immune Complex Wash Buffer (0.25M LiCl, 1% NP40, 1% deoxycholate, 1mM EDTA, 10mM Tris-HCl, pH 8.1); and 2 $\times$ 1mL 1X TE (10mM Tris-HCl, 1mM EDTA, pH 8.0). Immune complexes were eluted by adding 250 μ l of freshly prepared elution buffer (1%SDS, 0.1M NaHCO₃) to pelleted beads. After vortexing, and brief centrifugation, the supernatant fraction was carefully transferred to another tube and elution was repeated, combining the two eluates. Protein-DNA crosslinks were reversed by adding 20 μ l of 5M NaCl to the combined eluates and incubated at 65°C for four hours. Proteins were digested by adding 10 μ l of 0.5M EDTA, 20 μ l 1M Tris-HCl (pH 6.5), and 2 μ l of 10mg/ml proteinase K to the eluate and incubated for one hour at 45°C. DNA was purified by phenol/chloroform extraction and ethanol precipitation. Before precipitation 20 μ g of glycogen were added to improve DNA yield. DNA pellets were washed with 70% ethanol, allowed to air dry, and resuspended in DNase/RNase-free water.

2.5.2. AcH3-ChIP – BDNF-qPCR—To quantify BDNF promoter-specific sequences, DNA samples obtained before and after immunoprecipitation, with AcH3 or sIgG, were subjected to quantitative real-time PCR (qPCR). PCR was performed using a primer pair specific for rat BDNF exon IV promoter (from -73 to +14 bp) forward: TCTATTTTCGAGGCAGAGGAGGTATC, reverse: AATGGGAAAGTGGGTGGGAG; as described by (Chen et al, 2003; Gomez-Pinilla et al, 2011)

PCR quantification: The threshold cycle (Ct) value was normalized by subtracting the negative control value for each corresponding sample.

2.5.3. RNA isolation—This assay was based on the Nucleospin® RNA II kit (Macherey-Nagel, Düren, Germany). Aliquots of crude bilateral hippocampal homogenates for mRNA quantification assay were stored at -80°C. Pre-homogenate samples were thawed quickly on ice and added to buffer R1, supplemented with β -mercaptoethanol and disrupted with Ultra Turrax® (IKA®-Werke, Staufen, Germany) for 60s at high speed. RNA isolation from the pre-homogenate was performed in accordance with the manufacturer's instructions. RNA was eluted by adding 60 μ l RNase-free water to columns. After brief centrifugation the flow-out fraction was applied to the eluate and placed onto the column for reelution.

RNA concentration, purity and integrity: The absorbance ratio of the RNA solution was measured at wavelengths 260nm/280nm ranging from 1.71 to 1.97 and at wavelengths 260nm/230nm ranging from 1.79 to 1.88. To check the integrity of RNA, aliquots were run on etidium bromide-stained agarose gel. RNA samples were stored at -80°C.

2.5.4. cDNA synthesis—cDNA was synthesized using a TetrocDNA Synthesis kit (Cat# BIO-65026 Bioline, Luckenwalde, Germany) in accordance with the manufactures' instructions. Briefly, the reaction conditions were as follows: 1 μ g of RNA, 1 μ l of random primers, 1 μ l of 10 mM dNTP, 1 μ l of RNase inhibitor, and 0.25 μ l of 200 μ g/mL reverse transcriptase in a final volume of 20 μ l, and the solution was incubated for ten minutes at 25°C, primer annealing, followed by 42°C for 60 minutes, primer elongation, and followed by 80°C for five min termination. cDNA samples were stored at -20°C.

2.5.5. Real time quantitative RT-PCR (qRT-PCR) of BDNF mRNA—Based on the principle of the Sybr Green detection method, EvaGreen® dye (Biotium, Hayward, CA, USA) was used to detect PCR products. The PCR was performed using a primer pair specific for rat BDNF mRNA, forward: CCATGAAAGAAGCAAACGT, reverse: CTCCAGCAGAAAGAGCAGA. PCR amplifications consisted of equal amounts of template DNA, 10 μ L of ImmoMix™ complete ready-to-use heat-activated 2 $\times$ reaction mix (Bioline GmbH, Luckenwalde, Germany), 1 μ L of 20 $\times$ EvaGreen (Biotium, Hayward, CA, U.S.A.), 2.5 μ L of 10 nmol/L forward and reverse primer (IBAGmbH, Göttingen, Germany) and water to a final volume of 20 μ L. Amplifications were performed in a Rotor-Gene 6000 thermal cycler (Corbett Life Science/Qiagen, London, UK) at 95°C for 10 min, followed by 40 cycles of 95°C for 10 sec, 60°C for 20 sec and 72°C for 30 sec. Melting analysis was performed as a final step after the PCR amplification process. The validity of the signal was evaluated by melting analysis and agarose gel electrophoresis. The melting analysis was accepted if an appropriate melting point value of the product was detected (neither other products nor primer dimer were detected). The gel analysis was accepted if only one band was visible with an appropriate size (neither other products nor primer dimer were visible). To eliminate the differences in signal between different PCR runs, three control cDNAs were selected to validate the samples of each run. Rat β -actin gene served as an endogenous control gene.

2.5.6. Western blot and ELISA analysis of BDNF—Aliquots of crude bilateral hippocampal homogenates for Western blot assay were stored at -80°C. The Western blot was carried out as described earlier (Radak et al, 2006). Samples were incubated in block

solution supplemented with primary antibody against BDNF, diluted 1:5000 (polyclonal rabbit anti-BDNF Santa Cruz SC-546) overnight at 4°C with constant agitation. To normalize the expression of BDNF, the blots were re-incubated with a mouse anti-GAPDH antibody. Antibody anti β -actin (monoclonal mouse Cat# sc-8035, Santa Cruz) was added to the incubation solution for endogenous control. The concentration of BDNF was determined from the hippocampal section of the brain, using the E-Max ImmunoAssay System (Promega, Madison, WI) as described previously (Radak et al, 2013b).

2.6. Statistics

The results were compared with a Kruskal-Wallis analysis of variance (ANOVA) using Mann-Whitney post-hoc analyses. Results were expressed as means $\pm$ SD. Significance level was set at $p < 0.05$.

3. Results

At the end of the study, the body masses of the DR animals were significantly lower than those of the LCR-C and HCR-C groups (LCR-C:510 $\pm$ 42g, HCR-C:475 $\pm$ 24g, LCR-DR: 430 $\pm$ 21g, HCR-DR:415 $\pm$ 17g).

The average VO₂max of the LCR animals was 40ml/kg/min while 65ml/kg/min, a significantly higher ($p < 0.05$) value, was measured in the HCR rats. The effects of DR were not significant for VO₂max.

The passive avoidance test is a fear-motivated test to assess memory of laboratory rodents. Our data revealed that in short term memory HCR-DR animals had better results than all of the other groups ($p < 0.05$, Fig.2). All of HCR-DR animal stayed the maximal time (5 min) in the lighted section of the chamber of the box.

DR has been used to increase the activity/content of histone deacetylase SIRT1, but differences amongst the groups were not detected at the protein levels of SIRT1 (Fig.3). Peroxisome proliferator-activated receptor gamma coactivator-1 alpha (PGC-1 α) is a downstream transcription factor of SIRT1, and increased levels of PGC-1 α were observed in the HCR-DR group (Fig. 4A). However, significant differences in the down-stream protein of PGC-1 α , nuclear respiratory factor 1 (NRF1) (Fig.4B) were not detected. COX4 content was measured to assess mitochondrial content. Neither the running capacity nor DR altered the mitochondrial content in the hippocampus (Fig. 4C).

The activity of cAMP response element-binding protein (CREB) transcription factor regulates the expression of BDNF, and DR increased the phosphorylation activity of CREB in both groups (Fig. 5A).

BDNF content was measured by ELISA and the content increased significantly in the hippocampus of HCR-DR groups compared to other experimental groups (Fig.5B). The protein levels of BDNF were also measured by Western blots (Fig 6A) and in order to know the epigenetical regulation of BDNF, the global acetylation level of histone H3 were measured and significantly increased in the LCR-DR group (Fig 6B). The specific binding between acetylated histone H3 and BDNF promoter IV region was detected and was

increased by alternate day eating in the LCR and HCR groups (Fig. 6C). However, the increases in these bindings did not result in enhanced transcriptional levels of mRNA of BDNF in the HCR group (Fig. 6D), indicating the complexity of BDNF regulation.

4. Discussion

Results revealed that alternate day eating can improve memory in rats with high capacity running, while the effects of DR are not significant in low capacity runners. The levels of BDNF changed in a similar fashion as memory. Gomez-Pinilla and co-workers (Gomez-Pinilla et al., 2011) have shown that voluntary exercise causes increased acetylation of H3 and remodels the chromatin at the promoter region of BDNF.

DR has been shown to alter H3 acetylation in aging liver (Kawakami et al., 2012), and it has been suggested that DR can cause epigenetical regulation of BDNF (Rosas-Vargas et al., 2011). BDNF is a neurotrophin and its gene contains a single coding exon with multiple promoter and alternative splicing transcripts (Bertaux et al., 2004). The binding between acetylated histone H3 and BDNF promoter IV was higher in the HC-DR group, although no changes in the mRNA levels were found. This seems to be controversial, but in the present study we focused on the BDNF promoter region IV, because it has been shown to be sensitive to exercise intervention (Gomez-Pinilla et al., 2011). The changes in mRNA or even protein levels could have originated from other promoter regions. BDNF is crucial for a variety of cellular functions in the nervous system, including proliferation, differentiation, synaptic activity, survival and neurotransmission (Gomez-Pinilla, 2008a, b; Lu et al., 2013). Moreover, BDNF is involved in the regulation of energy metabolism from appetite suppression, to glucose and lipid metabolism (Gomez-Pinilla, 2008a). BDNF can also be activated by the products of heme oxygenase-1 (HO-1), such as bilirubin and CO (Hung et al., 2010) and HO-1 can be activated by acute exercise (Niess et al., 2000). However, it must be noted that regular exercise down-regulates the level of HO-1. Thus, in the present experimental conditions, the possible role of HO-1 in the regulation of BDNF is uncertain.

It has been demonstrated that DR can elevate BDNF levels, in accordance with earlier observations (Duan et al., 2003; Lee et al., 2002) and the data from the present study revealed that the increases in DR-associated BDNF induction are dependent on the posttranslational modification of histones. Therefore, it cannot be ruled out that DR-dependent induction of BDNF is genetically controlled.

DR resulted in phosphorylation of the serine residue (S133) of the kinase-inducible domain of CREB, which initiates transcriptional activity, including the gene for BDNF (Tyagi et al., 2013). It is important to note that CREB can also regulate the transcription of SIRT1 in low nutrient conditions in skeletal muscle, liver and adipose tissue (Noriega et al., 2011). SIRT1 can also regulate the activation of CREB (Monteserin-Garcia et al., 2013). Under the present experimental conditions, SIRT1 protein content was not significantly altered in the hippocampus of rats by DR or running capacity. It has been suggested that energy deprivation, such as DR, can lead to increased AMP/ATP and NAD⁺/NADH ratios leading to AMPK mediated phosphorylation and SIRT1 mediated deacetylation of PGC-1 α (Scarpulla, 2011). PGC-1 α activation then can lead to increased mitochondrial biogenesis.

However, it has been questioned whether caloric restriction can cause mitochondrial biogenesis (Hancock et al., 2011). Moreover, there are conflicting data on whether or not SIRT1 mediated deacetylation can result in PGC-1 α induced mitochondrial biogenesis (Gurd, 2011; Higashida et al., 2013; Lagouge et al., 2006; Philp et al., 2011; Rodgers et al., 2005). The present investigation did not aim to study the interaction of PGC-1 α and SIRT1, but based on the COX4 and NRF1 data, it is suggested that DR does not result in increased mitochondrial biogenesis in the hippocampus of rats.

In summary, it is concluded that DR improves memory of rats with high capacity running, which is associated with enhanced BDNF content due to the acetylation of H3 at the promoter region of the BDNF gene. This observation suggests that, as a consequence of epigenetical regulation, DR induced BDNF levels can be inherited. Overall, our data confirm that the response of low and high running capacity runner rats to DR is different, emphasizing the importance of exercise and caloric control on brain health.

Acknowledgments

The LCR HCR rat model system was supported by grant R24 RR017718 from the National Center for Research Resources of the National Institutes of Health (to L.G.K. and S.L.B.), and National Institutes of Health grant RO1 DK077200 (to S.L.B.). The LCR and HCR model can be made available for collaborative study (contact: brittons@umich.edu or lgkoch@umich.edu).

References

- Barnes SK, Ozanne SE. Pathways linking the early environment to long-term health and lifespan. *Progress in biophysics and molecular biology*. 2011; 106:323–336. [PubMed: 21147148]
- Barrientos RM, Frank MG, Crysdale NY, Chapman TR, Ahrendsen JT, Day HE, Campeau S, Watkins LR, Patterson SL, Maier SF. Little exercise, big effects: reversing aging and infection-induced memory deficits, and underlying processes. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2011; 31:11578–11586. [PubMed: 21832188]
- Bertaux O, Toselli-Mollereau E, Auffray C, Devignes MD. Alternative usage of 5' exons in the chicken nerve growth factor gene: refined characterization of a weakly expressed gene. *Gene*. 2004; 334:83–97. [PubMed: 15256258]
- Bramble DM, Lieberman DE. Endurance running and the evolution of Homo. *Nature*. 2004; 432:345–352. [PubMed: 15549097]
- Brown-Borg HM, Rakoczy S. Metabolic adaptations to short-term every-other-day feeding in long-living Ames dwarf mice. *Experimental gerontology*. 2013; 48:905–919. [PubMed: 23832075]
- Chakravarthy MV, Booth FW. Eating, exercise, and “thrifty” genotypes: connecting the dots toward an evolutionary understanding of modern chronic diseases. *Journal of applied physiology (Bethesda, Md : 1985)*. 2004; 96:3–10.
- Chen WG, Chang Q, Lin Y, Meissner A, West AE, Griffith EC, Jaenisch R, Greenberg ME. Derepression of BDNF transcription involves calcium-dependent phosphorylation of MeCP2. *Science (New York, N Y)*. 2003; 302:885–889.
- Duan W, Guo Z, Jiang H, Ware M, Li XJ, Mattson MP. Dietary restriction normalizes glucose metabolism and BDNF levels, slows disease progression, and increases survival in huntingtin mutant mice. *Proceedings of the National Academy of Sciences of the United States of America*. 2003; 100:2911–2916. [PubMed: 12589027]
- Dumas L, Kim YH, Karimpour-Fard A, Cox M, Hopkins J, Pollack JR, Sikela JM. Gene copy number variation spanning 60 million years of human and primate evolution. *Genome research*. 2007; 17:1266–1277. [PubMed: 17666543]
- Garcia Ramos J. Physiology of the cortical dendrites. *Boletin de estudios medicos y biologicos*. 1980; 31:181–189. [PubMed: 6275865]

- Gomez-Pinilla F. Brain foods: the effects of nutrients on brain function. *Nature reviews. Neuroscience*. 2008a; 9:568–578. [PubMed: 18568016]
- Gomez-Pinilla F. The influences of diet and exercise on mental health through hormesis. *Ageing research reviews*. 2008b; 7:49–62. [PubMed: 17604236]
- Gomez-Pinilla F, Zhuang Y, Feng J, Ying Z, Fan G. Exercise impacts brain-derived neurotrophic factor plasticity by engaging mechanisms of epigenetic regulation. *The European journal of neuroscience*. 2011; 33:383–390. [PubMed: 21198979]
- Griesbach GS, Hovda DA, Gomez-Pinilla F. Exercise-induced improvement in cognitive performance after traumatic brain injury in rats is dependent on BDNF activation. *Brain research*. 2009; 1288:105–115. [PubMed: 19555673]
- Gurd BJ. Deacetylation of PGC-1alpha by SIRT1: importance for skeletal muscle function and exercise-induced mitochondrial biogenesis. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*. 2011; 36:589–597.
- Hancock CR, Han DH, Higashida K, Kim SH, Holloszy JO. Does calorie restriction induce mitochondrial biogenesis? A reevaluation. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2011; 25:785–791. [PubMed: 21048043]
- Hart N, Sarga L, Csende Z, Koltai E, Koch LG, Britton SL, Davies KJ, Kouretas D, Wessner B, Radak Z. Resveratrol enhances exercise training responses in rats selectively bred for high running performance. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*. 2013
- Higashida K, Kim SH, Jung SR, Asaka M, Holloszy JO, Han DH. Effects of resveratrol and SIRT1 on PGC-1alpha activity and mitochondrial biogenesis: a reevaluation. *PLoS biology*. 2013; 11:e1001603. [PubMed: 23874150]
- Hung SY, Liou HC, Fu WM. The mechanism of heme oxygenase-1 action involved in the enhancement of neurotrophic factor expression. *Neuropharmacology*. 2010; 58:321–329. [PubMed: 19925812]
- Kaelin WG Jr, McKnight SL. Influence of metabolism on epigenetics and disease. *Cell*. 2013; 153:56–69. [PubMed: 23540690]
- Kawakami K, Nakamura A, Goto S. Dietary restriction increases site-specific histone H3 acetylation in rat liver: possible modulation by sirtuins. *Biochemical and biophysical research communications*. 2012; 418:836–840. [PubMed: 22321393]
- Koch LG, Britton SL. Artificial selection for intrinsic aerobic endurance running capacity in rats. *Physiological genomics*. 2001; 5:45–52. [PubMed: 11161005]
- Koch LG, Britton SL, Wisloff U. A rat model system to study complex disease risks, fitness, aging, and longevity. *Trends in cardiovascular medicine*. 2012; 22:29–34. [PubMed: 22867966]
- Koch LG, Kemi OJ, Qi N, Leng SX, Bijma P, Gilligan LJ, Wilkinson JE, Wisloff H, Hoydal MA, Rolim N, Abadir PM, van Grevenhof EM, Smith GL, Burant CF, Ellingsen O, Britton SL, Wisloff U. Intrinsic aerobic capacity sets a divide for aging and longevity. *Circ Res*. 2011a; 109:1162–1172. [PubMed: 21921265]
- Koch LG, Kemi OJ, Qi N, Leng SX, Bijma P, Gilligan LJ, Wilkinson JE, Wisloff H, Hoydal MA, Rolim N, Abadir PM, van Grevenhof EM, Smith GL, Burant CF, Ellingsen O, Britton SL, Wisloff U. Intrinsic aerobic capacity sets a divide for aging and longevity. *Circulation research*. 2011b; 109:1162–1172. [PubMed: 21921265]
- Kumar S, Parkash J, Kataria H, Kaur G. Interactive effect of excitotoxic injury and dietary restriction on neurogenesis and neurotrophic factors in adult male rat brain. *Neuroscience research*. 2009; 65:367–374. [PubMed: 19732799]
- Lagouge M, Argmann C, Gerhart-Hines Z, Meziane H, Lerin C, Daussin F, Messadeq N, Milne J, Lambert P, Elliott P, Geny B, Laakso M, Puigserver P, Auwerx J. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1alpha. *Cell*. 2006; 127:1109–1122. [PubMed: 17112576]
- Lee J, Seroogy KB, Mattson MP. Dietary restriction enhances neurotrophin expression and neurogenesis in the hippocampus of adult mice. *Journal of neurochemistry*. 2002; 80:539–547. [PubMed: 11905999]

- Lieberman DE, Bramble DM. The evolution of marathon running : capabilities in humans. *Sports medicine* (Auckland, N Z). 2007; 37:288–290.
- Lu B, Nagappan G, Guan X, Nathan PJ, Wren P. BDNF-based synaptic repair as a disease-modifying strategy for neurodegenerative diseases. *Nature reviews. Neuroscience*. 2013; 14:401–416. [PubMed: 23674053]
- Mattson MP. Evolutionary aspects of human exercise--born to run purposefully. *Ageing research reviews*. 2012; 11:347–352. [PubMed: 22394472]
- Milagro FI, Campion J, Cordero P, Goyenechea E, Gomez-Uriz AM, Abete I, Zulet MA, Martinez JA. A dual epigenomic approach for the search of obesity biomarkers: DNA methylation in relation to diet-induced weight loss. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2011; 25:1378–1389. [PubMed: 21209057]
- Monteserin-Garcia J, Al-Massadi O, Seoane LM, Alvarez CV, Shan B, Stalla J, Paez-Pereda M, Casanueva FF, Stalla GK, Theodoropoulou M. Sirt1 inhibits the transcription factor CREB to regulate pituitary growth hormone synthesis. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2013; 27:1561–1571. [PubMed: 23292070]
- Niess AM, Sommer M, Schneider M, Angres C, Tschositsch K, Golly IC, Battenfeld N, Northoff H, Biesalski HK, Dickhuth HH, Fehrenbach E. Physical exercise-induced expression of inducible nitric oxide synthase and heme oxygenase-1 in human leukocytes: effects of RRR-alpha-tocopherol supplementation. *Antioxid Redox Signal*. 2000; 2:113–126. [PubMed: 11232592]
- Noriega LG, Feige JN, Canto C, Yamamoto H, Yu J, Herman MA, Matakaki C, Kahn BB, Auwerx J. CREB and ChREBP oppositely regulate SIRT1 expression in response to energy availability. *EMBO reports*. 2011; 12:1069–1076. [PubMed: 21836635]
- Pesic V, Marinkovic P, Janac B, Ignjatovic S, Popic J, Kanazir S, Ruzdijic S. Changes of behavioral parameters during long-term food restriction in middle-aged Wistar rats. *Physiology & behavior*. 2010; 101:672–678. [PubMed: 20713076]
- Philp A, Chen A, Lan D, Meyer GA, Murphy AN, Knapp AE, Olfert IM, McCurdy CE, Marcotte GR, Hogan MC, Baar K, Schenk S. Sirtuin 1 (SIRT1) deacetylase activity is not required for mitochondrial biogenesis or peroxisome proliferator-activated receptor-gamma coactivator-1alpha (PGC-1alpha) deacetylation following endurance exercise. *The Journal of biological chemistry*. 2011; 286:30561–30570. [PubMed: 21757760]
- Qiu G, Spangler EL, Wan R, Miller M, Mattson MP, So KF, de Cabo R, Zou S, Ingram DK. Neuroprotection provided by dietary restriction in rats is further enhanced by reducing glucocorticoids. *Neurobiology of aging*. 2012; 33:2398–2410. [PubMed: 22226488]
- Quintas A, de Solis AJ, Diez-Guerra FJ, Carrascosa JM, Bogonez E. Age-associated decrease of SIRT1 expression in rat hippocampus: prevention by late onset caloric restriction. *Experimental gerontology*. 2012; 47:198–201. [PubMed: 22143179]
- Radak Z, Hart N, Sarga L, Koltai E, Atalay M, Ohno H, Boldogh I. Exercise plays a preventive role against Alzheimer's disease. *Journal of Alzheimer's disease : JAD*. 2010; 20:777–783.
- Radak Z, Ihasz F, Koltai E, Goto S, Taylor AW, Boldogh I. The redox-associated adaptive response of brain to physical exercise. *Free radical research*. 2013a
- Radak Z, Silye G, Bartha C, Jakus J, Stefanovits-Banyai E, Atalay M, Marton O, Koltai E. The effects of cocoa supplementation, caloric restriction, and regular exercise, on oxidative stress markers of brain and memory in the rat model. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*. 2013b; 61:36–41. [PubMed: 23419390]
- Radak Z, Toldy A, Szabo Z, Siamilis S, Nyakas C, Silye G, Jakus J, Goto S. The effects of training and detraining on memory, neurotrophins and oxidative stress markers in rat brain. *Neurochemistry international*. 2006; 49:387–392. [PubMed: 16564605]
- Radak Z, Zhao Z, Koltai E, Ohno H, Atalay M. Oxygen Consumption and Usage During Physical Exercise: The Balance Between Oxidative Stress and ROS-Dependent Adaptive Signaling. *Antioxidants & redox signaling*. 2012

- Rich NJ, Van Landingham JW, Figueiroa S, Seth R, Corniola RS, Levenson CW. Chronic caloric restriction reduces tissue damage and improves spatial memory in a rat model of traumatic brain injury. *Journal of neuroscience research*. 2010; 88:2933–2939. [PubMed: 20544832]
- Rodgers JT, Lerin C, Haas W, Gygi SP, Spiegelman BM, Puigserver P. Nutrient control of glucose homeostasis through a complex of PGC-1 α and SIRT1. *Nature*. 2005; 434:113–118. [PubMed: 15744310]
- Rodriguez-Bies E, Santa-Cruz Calvo S, Fontan-Lozano A, Pena Amaro J, Berral de la Rosa FJ, Carrion AM, Navas P, Lopez-Lluch G. Muscle physiology changes induced by every other day feeding and endurance exercise in mice: effects on physical performance. *PloS one*. 2010; 5:e13900. [PubMed: 21085477]
- Rosas-Vargas H, Martinez-Ezquerro JD, Bienvenu T. Brain-derived neurotrophic factor, food intake regulation, and obesity. *Archives of medical research*. 2011; 42:482–494. [PubMed: 21945389]
- Rothman SM, Griffioen KJ, Wan R, Mattson MP. Brain-derived neurotrophic factor as a regulator of systemic and brain energy metabolism and cardiovascular health. *Annals of the New York Academy of Sciences*. 2012; 1264:49–63. [PubMed: 22548651]
- Sanders YY, Liu H, Zhang X, Hecker L, Bernard K, Desai L, Liu G, Thannickal VJ. Histone modifications in senescence-associated resistance to apoptosis by oxidative stress. *Redox biology*. 2013; 1:8–16. [PubMed: 24024133]
- Sarga L, Hart N, Koch LG, Britton SL, Hajas G, Boldogh I, Ba X, Radak Z. Aerobic endurance capacity affects spatial memory and SIRT1 is a potent modulator of 8-oxoguanine repair. *Neuroscience*. 2013; 252:326–336. [PubMed: 23973402]
- Scarpulla RC. Metabolic control of mitochondrial biogenesis through the PGC-1 family regulatory network. *Biochimica et biophysica acta*. 2011; 1813:1269–1278. [PubMed: 20933024]
- Shankar S, Kumar D, Srivastava RK. Epigenetic modifications by dietary phytochemicals: implications for personalized nutrition. *Pharmacology & therapeutics*. 2013; 138:1–17. [PubMed: 23159372]
- Singh R, Lakhanpal D, Kumar S, Sharma S, Kataria H, Kaur M, Kaur G. Late-onset intermittent fasting dietary restriction as a potential intervention to retard age-associated brain function impairments in male rats. *Age (Dordrecht, Netherlands)*. 2012; 34:917–933.
- Sohal RS, Weindruch R. Oxidative stress, caloric restriction, and aging. *Science (New York, N Y)*. 1996; 273:59–63.
- Stranahan AM, Mattson MP. Bidirectional metabolic regulation of neurocognitive function. *Neurobiology of learning and memory*. 2011; 96:507–516. [PubMed: 21236352]
- Terzibasi E, Lefrancois C, Domenici P, Hartmann N, Graf M, Cellerino A. Effects of dietary restriction on mortality and age-related phenotypes in the short-lived fish *Nothobranchius furzeri*. *Aging cell*. 2009; 8:88–99. [PubMed: 19302373]
- Tyagi E, Agrawal R, Zhuang Y, Abad C, Waschek JA, Gomez-Pinilla F. Vulnerability imposed by diet and brain trauma for anxiety-like phenotype: implications for post-traumatic stress disorders. *PloS one*. 2013; 8:e57945. [PubMed: 23483949]
- Vaquero A, Reinberg D. Calorie restriction and the exercise of chromatin. *Genes & development*. 2009; 23:1849–1869. [PubMed: 19608767]
- Weindruch R, May PC. Use of caloric restriction to investigate neuroendocrine involvement in aging. *Neurobiology of aging*. 1995; 16:845–847. discussion 855–846. [PubMed: 8532120]
- Wikgren J, Mertikas GG, Raussi P, Tirkkonen R, Ayravainen L, Peltö-Huikko M, Koch LG, Britton SL, Kainulainen H. Selective breeding for endurance running capacity affects cognitive but not motor learning in rats. *Physiology & behavior*. 2012; 106:95–100. [PubMed: 22285210]
- Wu A, Ying Z, Gomez-Pinilla F. Exercise facilitates the action of dietary DHA on functional recovery after brain trauma. *Neuroscience*. 2013; 248:655–663. [PubMed: 23811071]

Highlights

Dietary restriction improves memory of rats with high capacity running.

Dietary restriction shapes the chromatin at BDNF promoter region.

Different running capacity alters the regulation of BDNF.

Fig. 1. Experimental setting

Low capacity runner (LCR) and high capacity runner (HCR) male rats, aged 13 months, were assigned to control LCR (LCR-C), dietary restricted LCR (LCR-DR), control HCR (HCR-C) and dietary restricted HCR (HCR-DR) groups ($n =$ six rats per group). Dietary restriction was performed by feeding the animal every other day for 16 weeks

Fig. 2. Passive avoidance test

Fear-motivated memory was tested by a passive avoidance test, and significant differences were detected between high running capacity-dietary restricted (HCR-DR) and low running capacity-control (LCR-C), and low running capacity-dietary restricted (LCR-DR) and high running capacity-control (HCR-C) groups. Values are expressed as means $\pm$ SD for six animals per group (* $P < 0.05$).

Fig. 3. SIRT1 levels in rat hippocampus

The levels of SIRT1 content were evaluated by a SIRT1 antibody. Values are expressed as means $\pm$ SD for six animals per group (* $P < 0.05$).

Fig. 4. The levels of mitochondrial biogenesis- associated proteins

PGC1a, (A) NRF1 (B) and COXIV (C) levels show how dietary restriction (DR) affected these proteins in the hippocampus of LCR and HCR rats. Values are expressed as means $\pm$ SD for six animals per group (* $P < 0.05$).

Fig. 5. pCREB/CREB ratio and BDNF levels in hippocampus

cAMP response element-binding protein (CREB) activities are shown on the figure as a ratio of phosphorylated and total protein content of CREB (A). BDNF levels were evaluated by ELISA (B). Values are expressed as means $\pm$ SD for six animals per group (* P <0.05).

Fig. 6. BDNF protein, promoters, and mRNA levels and H3K14 contents in brain
BDNF (A), and H3K14 levels (B) were evaluated by Western blots, while the quantitative-DNA PCR for BDNF promoter IV region was performed on ACh3 histone immunoprecipitated ChIP samples (C). The mRNA levels of this samples are shown in panel D. Values are expressed as means $\pm$ SD for six animals per group (* $P < 0.05$).