Published in final edited form as:

Ann Behav Med. 2014 October ; 48(2): 142–144. doi:10.1007/s12160-014-9615-x.

Optimizing Approaches to Addressing Depression in Cardiac Patients: a Comment on O’Neil et al

Jesse C. Stewart, PhD¹ and Bruce L. Rollman, MD, MPH²

¹Department of Psychology, Indiana University-Purdue University Indianapolis, Indianapolis, IN

²Division of General Internal Medicine, Center for Research on Health Care, University of Pittsburgh School of Medicine, Pittsburgh, PA

Depression is frequently experienced by cardiac patients, with a prevalence ranging from 20% to 30% depending on the assessment method, clinical setting, disease severity, and patient gender, among other factors (1). Even the lower limit of this range is more than double the prevalence of this treatable condition in the general population (2). Moreover, a recent Scientific Statement from the American Heart Association (AHA) recommends that depression be considered a risk factor for adverse medical outcomes among patients with acute coronary syndrome (ACS), thus giving new urgency to the development and dissemination of effective and scalable interventions for these high-risk patients (3).

Strategies to treat depression in cardiac patients are of great interest because of their potential, in theory, to reduce morbidity. While the interventions of earlier trials had only modest effects on depressive symptoms (4), recent trials of collaborative care (5, 6) and more intensive treatments (7–9) have demonstrated moderate improvements in depressive symptoms and health-related quality of life. Although more intensive treatments often produce gains twice as large as collaborative care interventions (effect size: 0.6 vs. 0.3) (10), they can be difficult to provide at scale or to patients who have difficulty taking time off from work, live in rural areas, or have other transportation issues (11). Consequently, the report by O’Neil and colleagues (12) describing the efficacy and feasibility of a telehealth intervention for depressed ACS patients in this issue of the *Annals* is timely.

The MoodCare investigators screened for depression in patients hospitalized for ACS at one of six Australian hospitals. One hundred twenty-one patients with Patient Health Questionnaire-9 (PHQ-9) scores of 5–19 (mild to moderately severe symptoms) who were not already seeing a mental health specialist were enrolled in the trial. Following hospital discharge, these patients were randomized to either (a) their physicians' usual medical care or (b) a 6-month telehealth intervention that integrated 10 sessions of cognitive-behavioral therapy (CBT) for depression delivered by Master’s-level therapists into an existing cardiovascular disease risk reduction program. At baseline, the mean PHQ-9 score was 9.2. Half of the enrolled patients had a lifetime history of a depressive disorder, and 16% were

Correspondence and Reprint Requests: Jesse C. Stewart, Ph.D., Department of Psychology, Indiana University-Purdue University Indianapolis, 402 N. Blackford St., LD 100E, Indianapolis, IN 46202. Phone: (317) 274-6761. Fax: (317) 274-6756. jstew@iupui.edu.

Authors Statement of Conflict of Interests: Jesse C. Stewart and Bruce L. Rollman declare that they have no conflicts of interest.

taking an antidepressant or anxiolytic. At the 6-month follow-up, intervention patients exhibited significantly greater improvements in depressive symptoms relative to usual care patients (baseline-adjusted mean difference in post-treatment PHQ-9 score = -1.8 points, $p = .025$, effect size = 0.36) but did not display changes in health-related quality of life. The intervention was also found to be more efficacious for patients with a lifetime history of depression (mean PHQ-9 difference = -2.7 points, $p = .043$, effect size = 0.65). The investigators, however, did not report the impact of their intervention on cardiovascular morbidity and mortality.

Demonstrating that telephone-delivered CBT is feasible in ACS patients and yields an effect size similar to other depression interventions in cardiac patients (13) is an important finding. Because telephone-delivered CBT is likely more scalable, accessible, and cost effective than traditional CBT, using this approach could increase the number of cardiac patients receiving evidence-based depression care without a corresponding rise in required resources. However, the effect size of the MoodCare intervention in the total sample was approximately half that observed in trials delivering more intensive treatments (7–9). Possible explanations for this lower efficacy are: (1) patients with minimal or mild depressive symptoms (53% had a PHQ-9 score < 10) who have little room to improve from CBT were enrolled, (2) patients with PHQ-9 scores ≥ 20 who may benefit most from treatment were excluded, (3) depressive symptoms following hospitalization were not reevaluated to exclude those who experienced a transient elevation in symptoms rather than a true depressive episode, and (4) adherence to the intervention was lower (61% completed 5 or more of the 10 CBT sessions).

Given the recent AHA Scientific Statement (3) and earlier Scientific Advisory advocating routine screening and treatment of depression (14), there is a need for future research to identify ways to optimize approaches to addressing depression in cardiac patients. One potential approach that could expand treatment delivery at reasonable cost is to incorporate Internet-delivered computerized CBT (CCBT). At present, these programs are primarily used in Australia, the United Kingdom, and the Netherlands. When these programs are provided under the guidance of a non-therapist care manager, they produce depressive symptom improvements similar to more traditional CBT (effect size: 0.58) (15). Thus, CCBT could cost-effectively be offered as first-line treatment to all cardiac patients who screen positive for depression (with or without antidepressants), while reserving specialist-delivered CBT for nonresponders, those with complicated mental health histories, and others who lack Internet access.

Another potential and complementary approach is to adopt a prevention focus by aggressively screening and treating depression earlier in the natural history of cardiovascular disease, such as in primary care patients with cardiovascular risk factors but no overt disease. An advantage of earlier treatment with CBT is that it may prevent the onset of a depressive episode after the occurrence of a cardiac event, as patients treated to remission with CBT are half as likely to have a recurrence than those treated to remission with antidepressants (16). Earlier depression treatment may also yield cardiovascular benefits. In an 8-year follow-up of the Improving Mood-Promoting Access to Collaborative Treatment (IMPACT) trial, depressed primary care patients without baseline cardiovascular disease

who were randomized to the collaborative care intervention had a 48% lower risk of a myocardial infarction or stroke than patients randomized to usual care (17).

In sum, O'Neil et al.'s MoodCare Trial is a valuable step forward in the area of depression treatment for cardiac patients, as their scalable and accessible intervention could help meet the likely increasing demand for evidence-based depression care for cardiac patients (3, 14). At the same time, there remains a need for future research to identify ways to optimize approaches to addressing depression in this population, which could ultimately result in improved clinical outcomes for an even larger number of cardiac patients.

Acknowledgments

This work was supported by NIH grant R01 HL114016 (Rollman).

References

1. Thombs BD, Bass EB, Ford DE, et al. Prevalence of depression in survivors of acute myocardial infarction. *Journal of General Internal Medicine*. 2006; 21:30–38. [PubMed: 16423120]
2. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*. 2005; 62:617–627. [PubMed: 15939839]
3. Lichtman JH, Froelicher ES, Blumenthal JA, et al. Depression as a risk factor for poor prognosis among patients with acute coronary syndrome: systematic review and recommendations: a scientific statement from the American Heart Association. *Circulation*. 2014 (published online before print).
4. Thombs BD, de Jonge P, Coyne JC, et al. Depression screening and patient outcomes in cardiovascular care: a systematic review. *JAMA*. 2008; 300:2161–2171. [PubMed: 19001627]
5. Rollman BL, Belnap BH, LeMenager MS, et al. Telephone-delivered collaborative care for treating post-CABG depression: A randomized controlled trial. *JAMA*. 2009; 302:2095–2103. [PubMed: 19918088]
6. Huffman JC, Mastromauro CA, Sowden G, et al. Impact of a depression care management program for hospitalized cardiac patients. *Circulation: Cardiovascular Quality and Outcomes*. 2011; 4:198–205. [PubMed: 21386067]
7. Davidson KW, Bigger JT, Burg MM, et al. Centralized, stepped, patient preference-based treatment for patients with post-acute coronary syndrome depression: CODIACS Vanguard randomized controlled trial. *JAMA Intern Med*. 2013:1–8.
8. Davidson KW, Rieckmann N, Clemow L, et al. Enhanced depression care for patients with acute coronary syndrome and persistent depressive symptoms: coronary psychosocial evaluation studies randomized controlled trial. *Archives of Internal Medicine*. 2010; 170:600–608. [PubMed: 20386003]
9. Freedland KE, Skala JA, Carney RM, et al. Treatment of depression after coronary artery bypass surgery: a randomized controlled trial. *Archives of General Psychiatry*. 2009; 66:387–396. [PubMed: 19349308]
10. Archer J, Bower P, Gilbody S, et al. Collaborative care for depression and anxiety problems. *Cochrane Database Syst Rev*. 2012; 10 CD006525.
11. Berkman LF, Blumenthal J, Burg M, et al. Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICHD) Randomized Trial. *JAMA*. 2003; 289:3106–3116. [PubMed: 12813116]
12. O'Neil A, Taylor B, Sanderson K, et al. Efficacy and feasibility of a tele-health intervention for acute coronary syndrome patients with depression: Results of the "MoodCare" randomized controlled trial. *Annals of Behavioral Medicine*. 2014

13. Rutledge T, Redwine LS, Linke SE, Mills PJ. A meta-analysis of mental health treatments and cardiac rehabilitation for improving clinical outcomes and depression among patients with coronary heart disease. *Psychosomatic Medicine*. 2013; 75:335–349. [PubMed: 23630306]
14. Lichtman JH, Bigger JT Jr, Blumenthal JA, et al. Depression and coronary heart disease: recommendations for screening, referral, and treatment: a science advisory from the American Heart Association Prevention Committee of the Council on Cardiovascular Nursing, Council on Clinical Cardiology, Council on Epidemiology and Prevention, and Interdisciplinary Council on Quality of Care and Outcomes Research: endorsed by the American Psychiatric Association. *Circulation*. 2008; 118:1768–1775. [PubMed: 18824640]
15. Richards D, Richardson T. Computer-based psychological treatments for depression: a systematic review and meta-analysis. *Clinical Psychology Review*. 2012; 32:329–342. [PubMed: 22466510]
16. Gloaguen V, Cottraux J, Cucherat M, Blackburn IM. A meta-analysis of the effects of cognitive therapy in depressed patients. *Journal of Affective Disorders*. 1998; 49:59–72. [PubMed: 9574861]
17. Stewart JC, Perkins AJ, Callahan CM. Effect of collaborative care for depression on risk of cardiovascular events: data from the IMPACT randomized controlled trial. *Psychosomatic Medicine*. 2014; 76:29–37. [PubMed: 24367124]