Published in final edited form as:

J Reprod Immunol. 2014 October ; 0: 12–19. doi:10.1016/j.jri.2014.03.006.

The Microbiome, Parturition, and Timing of Birth: More questions than answers

Amanda L. Prince, PhD¹, Kathleen M. Antony, MD¹, Derrick M. Chu¹, and Kjersti M. Aagaard, MD PhD^{1,2,3}

¹Departments of Obstetrics and Gynecology, Division of Maternal-Fetal Medicine, Baylor College of Medicine, Houston, Texas

²Bioinformatics Research Lab, Baylor College of Medicine, Houston, Texas

³Molecular and Cell Biology, Baylor College of Medicine, Houston, Texas

Abstract

The causes of preterm birth are multifactorial, but its association with infection has been well-established. The predominant paradigm describes an ascending infection from the lower genital tract through the cervix and into the presumably sterile fetal membranes and placenta. Thus, an evaluation of the role of the vaginal microbiome in preterm birth is implicated. However, emerging fields of data described in this review suggest that the placenta might not be sterile, even in the absence of clinical infection. We thus propose an additional mechanism for placental colonization and infection: hematogenous spread. When considered in the context of decades of evidence demonstrating a strong risk of recurrence for preterm birth, studies on parturition are ideal for applying the rapidly expanding field of metagenomics and analytic pipelines. The translational implications toward identification of innovative treatments for the prevention of preterm birth are further discussed. In sum, exciting advances in understanding the role of both host and microbiota in parturition and preterm birth are on the horizon.

Introduction

In 2005, the World Health Organization estimated that 12.9 million births worldwide occurred preterm; up to 42% of these resulted in mortality (Beck et al., 2010). Preterm birth is the leading cause of neonatal morbidity and mortality, yet little is understood regarding the underlying etiology (Kilpatrick, 2013). It is traditionally thought that an ascending infection from the vagina causes preterm premature rupture of membranes (PPROM), which initiates preterm labor and ultimately birth. However, more recent studies have shown that

© 2014 Elsevier Ireland Ltd. All rights reserved.

Address for correspondence: Kjersti Aagaard, MD, PhD Department of Obstetrics and Gynecology Baylor College of Medicine Houston, TX 77030 (aagaardt@bcm.edu)..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

conflict_of_interest_statement

Dummy conflict of interest statement to allow submission of copyedited version.

bacteria from the oral cavity are most often found in the amniotic fluid of patients with preterm labor (Figure 1) (Madianos et al., 2013). Additionally, studies are now demonstrating that bacteria are naturally found in placental tissue, and our own lab has shown that the placenta harbors its own unique microbiome (Aagaard et al., 2014). Here we will discuss how the microbiome changes during pregnancy and how these changes may influence preterm birth (Table 1). Further research in this area will lead to a greater understanding of the etiologies of preterm birth and may result in innovative treatments to prevent preterm birth.

The microbiome during pregnancy

During a normal pregnancy, the gravidae undergoes a spectrum of anatomical, physiological, and biochemical changes. These functional alterations result from the influences of hormonal and physical fluctuations, and they affect every organ of the body. These are accompanied by concomitant changes in the microbiome, at least in the vagina and gut, which are the only sites that have been specifically examined in pregnancy to date (Aagaard et al., 2012; Koren et al., 2012; Romero et al., 2014).

During pregnancy, hormonal changes result in increased thickness of the vaginal mucosa, hypertrophy of the smooth muscle cells, and relaxation of the connective tissues. Recently, we cataloged the “normal” microbiota signature during pregnancy in a cross-sectional study sampling women at a variety of gestational ages (Aagaard et al., 2012). Using 454 pyrosequencing technology, we deep sequenced the V3–V5 region of 16S rRNA from samples obtained from the vaginal introitus, midvagina, and posterior fornix. Interestingly, we found that the vaginal microbial community differed by gestational age and proximity to the cervix (Aagaard et al., 2012). Furthermore, the microbial community structure resembled a non-pregnant state in late gestation, and we saw a decrease in alpha diversity, or within-sample diversity, with a corresponding increase in *Lactobacillus* species in gravid patients compared with nonpregnant subjects (Aagaard et al., 2012). Recently, Romero *et al.* took these studies further by examining the vaginal microbiome longitudinally during pregnancy at the posterior fornix (Romero et al., 2014). While the vaginal microbiome of gravid women could still be classified into distinct community state types, as previously described in nonpregnant women (Ravel et al., 2011), the vaginal microbiome became more stable and less diverse throughout pregnancy, as we previously described (Aagaard et al., 2012; Romero et al., 2014).

One such species that was discriminately and specifically enriched in our study was *L. johnsonii*. This species encodes enzymes and transporters that are essential for the release of bile salt hydrolase and is primarily found in the upper gastrointestinal tract (Pridmore et al., 2004). *L. johnsonii* also produces Lactacin F, which limits other lactobacillus and *Enterococcus* species in the gastrointestinal tract (Abee et al., 1994). Thus, the increase in *L. johnsonii* may be important for the inoculation of neonates in order to promote the digestion of breast milk postpartum.

While these alterations in the microbiome may serve to inoculate the neonatal gut, they may also contribute to pregnancy maintenance. In addition to the aforementioned enrichment in

L. johnsonii, there was enrichment of *L. jensenii* and *L. crispatus*. *L. jensenii* anaerobically metabolizes glycogen, and the increased estrogen levels in pregnancy lead to increased glycogen, which, through *L. jensenii* metabolism, contributes to the acidic vaginal environment. This acidic environment likely suppresses the growth of bacteria associated with bacterial vaginosis, which may be a natural mechanism for preventing infections associated with preterm birth (O'Hanlon et al., 2011). The finding that the vaginal microbiome becomes more similar to the nonpregnant microbiome in late pregnancy may also support the microbiome's role in pregnancy maintenance, as a return to a nonpregnant state may herald changes associated with parturition.

With regard to the enteric microbiome, alpha diversity is decreased as the pregnancy progresses from the first to third trimester; however, beta diversity increases during this time (Koren et al., 2012; Romero et al., 2014). Further, as gravidae progress through their pregnancy, the ratio of *Firmicutes* and *Bacteroidetes* changes similar to that seen between lean and obese subjects in various studies (Koren et al., 2012; Turnbaugh et al., 2009, 2006). In fact, when the enteric microbiome from gravidae in their first or third trimester is transplanted into germ-free mice, mice inoculated with microbiota from third-trimester patients are more prone to an increase in adipose tissue and the inflammatory cytokines, IL-1 β , IL-5, IL-6, and GM-CSF (Koren et al., 2012). These data indicate that as pregnancy progresses, the body undergoes changes to increase the capacity for energy-harvesting microbes.

The microbiome, parturition, and preterm birth

As we begin to understand changes in the microbiome that occur during pregnancy, a natural question to ponder is the role of the microbiome in the regulation of the timing of parturition and notably preterm birth. While the dogma currently states that preterm birth is initiated by an ascending infection from the vaginal cavity to the placenta, recent studies have challenged this idea (Gonçalves et al., 2002; Madianos et al., 2013). In fact, the colonization of the placenta with bacteria may not be exclusive to patients with evidence of infection, such as preterm labor. For example, Stout *et al.* recently demonstrated that bacteria could be found in the basal plate of term and preterm placentas following spontaneous deliveries (Stout et al., 2013). Additionally, our own lab has found that the placenta harbors a unique microbiome, and that this microbiome differs in term and preterm subjects (Figure 1) (Aagaard et al., 2014). Supporting our observations, a bacterium found in the amniotic fluid of preterm patients was most likely to be found in the oral cavity as well, rather than the vaginal cavity (Bearfield, 2002; Douvier et al., 1999; Han et al., 2006) and both term and preterm neonates appear to be colonized early with bacteria normally found in the oral cavity (Figure 1) (Human Microbiome Project Consortium, 2012; Jost et al., 2012; Milisavljevic et al., 2013; Palmer et al., 2007). When *Fusobacterium nucleatum* is isolated from preterm patients and injected intravenously into pregnant mice, the placenta becomes selectively colonized with bacteria while other organs, such as the liver or spleen, do not (Han et al., 2004). This study also documented that while *F. nucleatum* infection did not result in preterm birth in mice, it resulted in stillbirth (Han et al., 2004). This is akin to the “failed escape hypothesis” of stillbirth, which postulates that stillbirth occurs if a fetus fails

to initiate a fetal inflammatory response with subsequent preterm birth (Blackwell et al., 2003; Romero et al., 1998).

An independent study isolated bacteria from the oral cavity of a variety of individuals and subsequently used this mixture to infect gravid mice. Similar results were seen in this study where the placenta was colonized by the oral bacteria (Fardini et al., 2010). Arce *et al.* took these types of studies one step further and inoculated mice by feeding them chow containing either the gram-negative bacteria *Campylobacter rectus* alone or a combination of *C. rectus* and *Porphyromonas gingivalis*. They found that infected mice had decreased fecundity and increased resorption rates (Arce et al., 2009). Additionally, infected mice that were impregnated had inflammation in the placenta, along with an increase in the expression of Toll-like receptor-4 (TLR-4), which recognizes lipopolysaccharide (LPS) on gram-negative bacteria (Arce et al., 2009). Taken together, these studies demonstrate that hematogenous infections have the ability to colonize and infect the placenta, which provides support to the notion that preterm birth might be caused by infections not originating in the vaginal cavity.

Despite the recent explosion of metagenomic studies involving the microbiome, there are relatively few studies using metagenomics to examine the role of the microbiome in preterm birth. Most recently, Hyman *et al.* examined the posterior fornix of term and preterm patients using 16S sequencing. When examining 37 Caucasian gravidae, with only seven patients resulting in a preterm birth, Hyman *et al.* found that there was a decrease in the alpha diversity of the overall microflora of preterm patients (Hyman et al., 2014). Interestingly, while *Lactobacillus* has been found to be the dominating genus of the vaginal cavity, there were no differences in this genus between term and preterm patients, and there appeared to be no bacterium in particular dominating the posterior fornix in preterm patients (Hyman et al., 2014). For example, in one preterm patient *Bifidobacterium* dominated the vaginal cavity while in another the vaginal cavity was dominated by *Ureaplasma* (Hyman et al., 2014). These results are interesting because when infants with early-onset necrotizing colitis (EONS) are examined by 16S sequencing, Wang *et al.* determined that the majority of patients had a mono-microbial infection (Wang et al., 2013). Notably, the study by Hyman *et al.* neglected to examine the oral cavity of term and preterm patients, which the recent studies above implicate as playing an important role in preterm birth.

Preterm birth and inflammation

Given the early and emerging evidence pertaining to the potential role of oral and hematogenous infections (of oral microbes) in preterm birth, one wonders why the placenta might harbor largely commensal microbes. This may be due to the unique immune environment of the placenta during pregnancy which ultimately prevents rejection of the fetus. While increased levels of IL-6 have been found in the amniotic fluid of preterm gravidae, the immunosuppressive/immunomodulatory cytokine, IL-10, has also been found (Greig et al., 1995; Han et al., 2009). Further, T cells are found at the maternal–fetal interface, and specialized NK cells have also been characterized in the placenta (Christiansen, 2013; Zenclussen, 2013). The effect of these changes may not only prevent rejection of the fetus, but may alter the composition of the mother's microbiome in order to

promote energy harvest required for pregnancy as well as to inoculate the neonate with the proper microbiome required for normal digestion and development.

However, these changes in the gravid immune system may functionally alter susceptibility of the placenta to hematogenous colonization, which in turn may trigger an inflammatory response (Figure 2). While the microbial species themselves may not be pathogenic, the relatively altered community structure (*e.g.*, dysbiosis) would thereby render the risk of localized inflammation. For example, infusion of LPS into placental explants *in vitro* has been shown to trigger production of pro-inflammatory cytokines, such as IL-1 β , IL-6, and IL-8 (Fortunato et al., 1996; Holcberg et al., 2008). Furthermore, when IL-1 β is infused intra-amniotically into pregnant rhesus macaques, amniotic levels of TNF and prostaglandins are raised and uterine contractions are induced (Sadowsky et al., 2003). These findings imply a pro-inflammatory etiology, but it is not clear whether this can be attributed to a single taxa or a dysbiotic placental microbiome community profile.

These changes in the gravid immune system within the placental microenvironment must strike a delicate balance between the prevention of fetal rejection and susceptibility to dysbiosis and ensuing inflammation. The observation that preterm birth tends to recur in individual mothers suggests that human host genetic variation may disrupt this balance and render the placenta more susceptible to infection in the face of a microbial challenge. In support of this hypothesis, significant efforts have been made to associate preterm birth with small nucleotide polymorphisms (SNPs) within key cytokines altered in the gravid immune system. For example, in a study of 62 Caucasian women who delivered preterm, mothers, with a homozygous C to T polymorphism within the promoter region of IL-1 β at position -31 (IL-1 β -31C>T), were found to have a higher rate of preterm birth (Odds ratio = 6.36) (Hollegaard et al., 2008). The strength of this SNP association appears to be dependent on both the fetal and maternal genotypes (Yilmaz et al., 2012). However, most polymorphisms of IL-1 β do not increase total serum levels, but the TT genotype at position -31 increases local monocyte IL-1 β secretion in response to LPS (Chen et al., 2006; Cullup et al., 2004). Additionally, similar studies have examined polymorphisms of the IL-6 promoter. As mentioned previously, IL-6 is a pro-inflammatory cytokine increased in the amniotic fluid of preterm mothers (Han et al., 2009). Individuals homozygous for a G to C polymorphism at position -174 within the IL-6 promoter (IL-6 -175G>C) have significantly decreased plasma levels of IL-6 (Fishman et al., 1998). In concordance with this result, Simhan *et al.* found that the CC genotype at this locus was associated with fewer preterm births (Simhan, 2003). In further stratifications, Macones *et al.* discovered that women with a polymorphism in the promoter region of TNF α (-308) and diagnosed with bacterial vaginosis had a higher rate of preterm birth (adjusted odds ratio = 6.0) (Macones et al., 2004). Altogether, these results suggest a host gene–microbiome interaction that influences a mother's risk of preterm birth. Hematogenous microbial colonization of the placenta (albeit in low abundance) may cause local inflammation that induces premature labor. However, certain polymorphisms of key cytokines may heighten the sensitivity of the gravid immune system to bacterial stimuli, leading to an inappropriate immune response to an otherwise normal placental microbiome.

Timing of placental colonization and inflammation

Another intriguing idea is that bacterial infiltration into placental tissue and the resulting inflammation may occur in term labors as a method of inducing labor. Recently, Lappas examined the expression of cytokines and transcription factors in the myometrium of gravidae undergoing an elective or laboring cesarean. This group found that the forkhead box O1 (FOXO1) was upregulated in patients undergoing a labored cesarean (Lappas, 2013). Further, when FOXO1 expression was diminished in myometrial cell lines using small interfering RNAs (siRNAs), this group demonstrated that IL-6 and IL-8 expression was diminished in the absence of FOXO1 upon stimulation with IL-1 β (Lappas, 2013). This study, combined with the finding that bacteria were found in the basal plate of the placenta of term and preterm gravidae (Stout et al., 2013), indicates that microbes may permeate the placenta to induce labor. However, questions pondering why this process occurs early in preterm patients and how to prevent the early onset of labor still remain. The evolving knowledge base of human genomics, the microbiome, and microbial metabolic pathways will hopefully elucidate answers to these questions. Further work in these areas is desperately needed in order to answer these questions definitively.

Conclusions

Preterm birth is one of the leading causes of neonatal mortality in the United States, yet its definitive attributable etiology remains to be clarified. Its known association with genitourinary infections and periodontal disease has prompted investigations into the microbiome of these sites. Until recently, the paradigm was that the majority of intrauterine infections originated in the lower genital tract with microbiota ascending into an otherwise sterile environment resulting in infection of the placenta, fetal membranes (chorioamnionitis), umbilical cord (funisitis), and the fetus (sepsis) (Gonçalves et al., 2002). However, mounting evidence from human and animal models points to an alternative hematogenous source of infection, namely from the oral cavity (Madianos et al., 2013). The discovery of associations between the oral and placental microbiome, particularly in cases of periodontal disease and preterm birth, would have a particularly high impact, and the current innovations in metagenomic research serve as the perfect arena in which to apply microbiome principles to research and prevention of preterm birth. Utilizing current and developing state-of-the-science technology and analysis tools, science is at the point in which the role of the microbiome in preterm birth can, and should, be prospectively investigated. Only this will lead to the understanding necessary to propose causative etiologies and diagnostics of preterm birth. Once understood, innovative treatments, including simple dietary changes, may be developed to tackle this pressing perinatal epidemic.

Whether we are obstetrician gynecologists, reproductive biologists, or physician scientists, we are called upon to challenge long-standing assumptions that have failed to lead to major breakthroughs in peripartum care for women and their children. Beyond the introduction of progesterone-based prophylaxis and cerclage for the prevention of recurrent preterm birth, we have yet to make a meaningful dent in the rate of preterm births. Both nationally and globally our rates of prematurity have only incrementally fluctuated over the past decades.

Concomitant with our emerging understanding that our microbiome largely comprises microbiota that are more friend than foe is an appreciation that longitudinal studies with deep phenotypic and clinical metadata are going to be instrumental in discerning association from causation. As more whole genome shotgun data are combined with metatranscriptomics analysis (thereby telling us not only who is present, but what they are actually doing), we will begin to acquire necessary snapshots of biological flow and disease progression over time. However, these more complex datasets will necessitate tools and personnel with the capacity to convert big data to applicable knowledge. We anticipate that future studies that are capable of making a significant and lasting impact will be those that marry big data metagenomic science with inherent clinical knowledge from larger population-based cohorts with longitudinal samples. These studies will ultimately enable inferred causation analysis, which is critical to deciphering the role of the microbiome in modulating the risk of preterm birth, and the impact of known and novel interventions on this process. Such approaches will yield not only potentially novel biomarkers, but the capacity for knowledge-based interventions.

REFERENCES

- Aagaard K, Riehle K, Ma J, Segata N, Mistretta T-A, Coarfa C, Raza S, Rosenbaum S, Van den Veyver I, Milosavljevic A, Gevers D, Huttenhower C, Petrosino J, Versalovic J. A metagenomic approach to characterization of the vaginal microbiome signature in pregnancy. *PLoS One*. 2012; 7:e36466. doi:10.1371/journal.pone.0036466. [PubMed: 22719832]
- Aagaard K, Ma J, Antony KM, Anders A, Ganu R, Petrosino J, Versalovic J. The Placenta Harbors a Unique Microbiome. *Sci. Transl. Med.* 2014 In Press.
- Abee T, Klaenhammer TR, Letellier L. Kinetic studies of the action of lactacin F, a bacteriocin produced by *Lactobacillus johnsonii* that forms poration complexes in the cytoplasmic membrane. *Appl. Environ. Microbiol.* 1994; 60:1006–13. [PubMed: 8161167]
- Arce RM, Barros SP, Wacker B, Peters B, Moss K, Offenbacher S. Increased TLR4 expression in murine placentas after oral infection with periodontal pathogens. *Placenta*. 2009; 30:156–62. doi: 10.1016/j.placenta.2008.11.017. [PubMed: 19101032]
- Bearfield C. Possible association between amniotic fluid micro - organism infection and microflora in the mouth. *J. Obstet.* 2002; 109:527–33.
- Beck S, Wojdyla D, Say L, Betran AP, Merialdi M, Requejo JH, Rubens C, Menon R, Van Look P.F. a. The worldwide incidence of preterm birth: a systematic review of maternal mortality and morbidity. *Bull. World Health Organ.* 2010; 88:31–8. doi:10.2471/BLT.08.062554. [PubMed: 20428351]
- Blackwell S, Romero R, Chaiworapongsa T, Kim YM, Bujold E, Espinoza J, Camacho N, Hassan S, Yoon BH, Refuerzo JS. Maternal and fetal inflammatory responses in unexplained fetal death. *J. Matern. Fetal. Neonatal Med.* 2003; 14:151–7. doi:10.1080/jmf.14.3.151.157. [PubMed: 14694969]
- Chen H, Wilkins LM, Aziz N, Cannings C, Wyllie DH, Bingle C, Rogus J, Beck JD, Offenbacher S, Cork MJ, Rafie-Kolpin M, Hsieh C-M, Kornman KS, Duff GW. Single nucleotide polymorphisms in the human interleukin-1B gene affect transcription according to haplotype context. *Hum. Mol. Genet.* 2006; 15:519–29. doi:10.1093/hmg/ddi469. [PubMed: 16399797]
- Christiansen OB. Reproductive immunology. *Mol. Immunol.* 2013; 55:8–15. doi:10.1016/j.molimm.2012.08.025. [PubMed: 23062611]
- Cullup H, Middleton PG, Duggan G, Conn JS, Dickinson A M. Environmental factors and not genotype influence the plasma level of interleukin-1 receptor antagonist in normal individuals. *Clin. Exp. Immunol.* 2004; 137:351–8. doi:10.1111/j.1365-2249.2004.02531.x. [PubMed: 15270852]

- Douvier S, Neuwirth C, Filipuzzi L, Kisterman JP. Chorioamnionitis with intact membranes caused by *Capnocytophaga sputigena*. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 1999; 83:109–12. [PubMed: 10221619]
- Fardini Y, Chung P, Dumm R, Joshi N, Han YW. Transmission of diverse oral bacteria to murine placenta: evidence for the oral microbiome as a potential source of intrauterine infection. *Infect. Immun.* 2010; 78:1789–96. doi:10.1128/IAI.01395-09. [PubMed: 20123706]
- Fishman D, Faulds G, Jeffery R, Mohamed-Ali V, Yudkin JS, Humphries S, Woo P. The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis. *J. Clin. Invest.* 1998; 102:1369–76. doi:10.1172/JCI2629. [PubMed: 9769329]
- Fortunato SJ, Menon RP, Swan KF, Menon R. Inflammatory cytokine (interleukins 1, 6 and 8 and tumor necrosis factor- α) release from cultured human fetal membranes in response to endotoxic lipopolysaccharide mirrors amniotic fluid concentrations. *Am. J. Obstet. Gynecol.* 1996; 174:1855–61. discussion 1861–2. [PubMed: 8678151]
- Gonçalves LF, Chaiworapongsa T, Romero R. Intrauterine infection and prematurity. *Ment. Retard. Dev. Disabil. Res. Rev.* 2002; 8:3–13. doi:10.1002/mrdd.10008. [PubMed: 11921380]
- Greig PC, Herbert WN, Robinette BL, Teot L. a. Amniotic fluid interleukin-10 concentrations increase through pregnancy and are elevated in patients with preterm labor associated with intrauterine infection. *Am. J. Obstet. Gynecol.* 1995; 173:1223–7. [PubMed: 7485325]
- Han Y, Redline R, Li M, Yin L. *Fusobacterium nucleatum* induces premature and term stillbirths in pregnant mice: implication of oral bacteria in preterm birth. *Infect.* 2004; 72:2272–9. doi:10.1128/IAI.72.4.2272.
- Han Y, Ikegami A, Bissada N. Transmission of an uncultivated *Bergeyella* strain from the oral cavity to amniotic fluid in a case of preterm birth. *J. Clin.* 2006; 44:1475–83. doi:10.1128/JCM.44.4.1475-1483.2006.
- Han YW, Shen T, Chung P, Buhimschi IA, Buhimschi CS. Uncultivated bacteria as etiologic agents of intra-amniotic inflammation leading to preterm birth. *J. Clin. Microbiol.* 2009; 47:38–47. doi:10.1128/JCM.01206-08. [PubMed: 18971361]
- Holcberg G, Amash A, Sapir O, Sheiner E, Levy S, Myatt L, Huleihel M. Perfusion with lipopolysaccharide differently affects the secretion of interleukin-1 beta and interleukin-1 receptor antagonist by term and preterm human placentae. *Placenta.* 2008; 29:593–601. doi:10.1016/j.placenta.2008.03.008. [PubMed: 18471873]
- Hollegaard MV, Grove J, Thorsen P, Wang X, Mandrup S, Christiansen M, Norgaard-Pedersen B, Wojdemann KR, Tabor A, Attermann J, Hougaard DM. Polymorphisms in the tumor necrosis factor alpha and interleukin-1 beta promoters with possible gene regulatory functions increase the risk of preterm birth. *Acta Obstet. Gynecol. Scand.* 2008; 87:1285–90. doi:10.1080/00016340802468340. [PubMed: 18951205]
- Human Microbiome Project Consortium. Structure, function and diversity of the healthy human microbiome. *Nature.* 2012; 486:207–14. doi:10.1038/nature11234. [PubMed: 22699609]
- Hyman RW, Fukushima M, Jiang H, Fung E, Rand L, Johnson B, Vo KC, Caughey AB, Hilton JF, Davis RW, Giudice LC. Diversity of the vaginal microbiome correlates with preterm birth. *Reprod. Sci.* 2014; 21:32–40. doi:10.1177/1933719113488838. [PubMed: 23715799]
- Jost T, Lacroix C, Braegger CP, Chassard C. New insights in gut microbiota establishment in healthy breast fed neonates. *PLoS One.* 2012; 7:e44595. doi:10.1371/journal.pone.0044595. [PubMed: 22957008]
- Kilpatrick, SJ. Prediction and Prevention of Preterm Birth: What have we learned since 2001?. 2013. [WWW Document]. <http://contemporaryobgyn.modernmedicine.com/sites/default/files/images/Logos/coglogo.jpg>. URL <http://contemporaryobgyn.modernmedicine.com/contemporary-obgyn/news/modernmedicine/modern-medicine-feature-articles/prediction-and-prevention-pr>
- Koren O, Goodrich JK, Cullender TC, Spor A, Laitinen K, Bäckhed HK, Gonzalez A, Werner JJ, Angenent LT, Knight R, Bäckhed F, Isolauri E, Salminen S, Ley RE. Host remodeling of the gut microbiome and metabolic changes during pregnancy. *Cell.* 2012; 150:470–80. doi:10.1016/j.cell.2012.07.008. [PubMed: 22863002]

- Lappas M. Forkhead box O1 (FOXO1) in pregnant human myometrial cells: a role as a pro-inflammatory mediator in human parturition. *J. Reprod. Immunol.* 2013; 99:24–32. doi:10.1016/j.jri.2013.04.005. [PubMed: 23778262]
- Macones GA, Parry S, Elkousy M, Clothier B, Ural SH, Strauss JF. A polymorphism in the promoter region of TNF and bacterial vaginosis: preliminary evidence of gene-environment interaction in the etiology of spontaneous preterm birth. *Am. J. Obstet. Gynecol.* 2004; 190:1504–8. discussion 3A. doi:10.1016/j.ajog.2004.01.001. [PubMed: 15284722]
- Madianos PN, Bobetsis Y. a, Offenbacher S. Adverse pregnancy outcomes (APOs) and periodontal disease: pathogenic mechanisms. *J. Periodontol.* 2013; 84:S170–80. doi:10.1902/jop.2013.1340015. [PubMed: 23631577]
- Milisavljevic V, Garg M, Vuletic I, Miller JF, Kim L, Cunningham TD, Schröder I. Prospective assessment of the gastroesophageal microbiome in VLBW neonates. *BMC Pediatr.* 2013; 13:49. doi:10.1186/1471-2431-13-49. [PubMed: 23560555]
- O'Hanlon DE, Moench TR, Cone RA. In vaginal fluid, bacteria associated with bacterial vaginosis can be suppressed with lactic acid but not hydrogen peroxide. *BMC Infect. Dis.* 2011; 11:200. doi: 10.1186/1471-2334-11-200. [PubMed: 21771337]
- Palmer C, Bik EM, DiGiulio DB, Relman DA, Brown PO. Development of the human infant intestinal microbiota. *PLoS Biol.* 2007; 5:e177. doi:10.1371/journal.pbio.0050177. [PubMed: 17594176]
- Pridmore RD, Berger B, Desiere F, Vilanova D, Barretto C, Pittet A-C, Zwahlen M-C, Rouvet M, Altermann E, Barrangou R, Mollet B, Mercenier A, Klaenhammer T, Arigoni F, Schell MA. The genome sequence of the probiotic intestinal bacterium *Lactobacillus johnsonii* NCC 533. *Proc. Natl. Acad. Sci. U. S. A.* 2004; 101:2512–7. [PubMed: 14983040]
- Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SSK, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. Vaginal microbiome of reproductive-age women. *Proc. Natl. Acad. Sci. U. S. A.* 108 Suppl. 2011:4680–7. doi:10.1073/pnas.1002611107.
- Romero R, Gomez R, Ghezzi F, Yoon BH, Mazor M, Edwin SS, Berry SM. A fetal systemic inflammatory response is followed by the spontaneous onset of preterm parturition. *Am. J. Obstet. Gynecol.* 1998; 179:186–93. [PubMed: 9704786]
- Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosch DW, Nikita L, Galuppi M, Lamont RF, Chaemsathong P, Miranda J, Chaiworapongsa T, Ravel J. The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome.* 2014; 2:4. doi:10.1186/2049-2618-2-4. [PubMed: 24484853]
- Sadowsky DW, Novy MJ, Witkin SS, Gravett MG. Dexamethasone or interleukin-10 blocks interleukin-1 β -induced uterine contractions in pregnant rhesus monkeys. *Am. J. Obstet. Gynecol.* 2003; 188:252–263. doi:10.1067/mob.2003.70. [PubMed: 12548226]
- Simhan H. Interleukin-6 promoter –174 polymorphism and spontaneous preterm birth. *Am. J. Obstet. Gynecol.* 2003; 189:915–918. doi:10.1067/S0002-9378(03)00843-3. [PubMed: 14586325]
- Stout MJ, Conlon B, Landeau M, Lee I, Bower C, Zhao Q, Roehl KA, Nelson DM, Macones GA, Mysorekar IU. Identification of intracellular bacteria in the basal plate of the human placenta in term and preterm gestations. *Am. J. Obstet. Gynecol.* 2013; 208:226, e1–7. doi:10.1016/j.ajog.2013.01.018. [PubMed: 23333552]
- Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature.* 2006; 444:1027–31. doi:10.1038/nature05414. [PubMed: 17183312]
- Turnbaugh PJ, Hamady M, Yatsunenkov T, Cantarel BL, Duncan A, Ley RE, Sogin ML, Jones WJ, Roe BA, Affourtit JP, Egholm M, Henrissat B, Heath AC, Knight R, Gordon JI. A core gut microbiome in obese and lean twins. *Nature.* 2009; 457:480–4. doi:10.1038/nature07540. [PubMed: 19043404]
- Wang X, Buhimschi CS, Temoin S, Bhandari V, Han YW, Buhimschi IA. Comparative microbial analysis of paired amniotic fluid and cord blood from pregnancies complicated by preterm birth and early-onset neonatal sepsis. *PLoS One.* 2013; 8:e56131. doi:10.1371/journal.pone.0056131. [PubMed: 23437088]

- Yılmaz Y, Verdi H, Taneri A, Yazıcı AC, Ecevit AN, Karaka NM, Tarcın A, Haberal A, Ozbek N, Atac FB. Maternal-fetal proinflammatory cytokine gene polymorphism and preterm birth. *DNA Cell Biol.* 2012; 31:92–7. doi:10.1089/dna.2010.1169. [PubMed: 21682548]
- Zenclussen AC. Adaptive immune responses during pregnancy. *Am. J. Reprod. Immunol.* 2013; 69:291–303. doi:10.1111/aji.12097. [PubMed: 23418773]

Figure 1. The fetal and neonatal microbiome does not replicate the vaginal microbiome in pregnancy, but more closely resembles the oral and placental microbiome

The oral microbiome has been previously characterized in the non-gravid population by the Human Microbiome Project Consortium (2012). Oral bacteria associations with preterm birth have been described in a very limited number of studies, and primarily with targeted 16S-based methodologies (Bearfield, 2002; Douvier et al., 1999; Fardini et al., 2010; Han et al., 2006, 2004). However, recent studies have characterized the vaginal microbiome during pregnancy, and demonstrated that the species richness decreases during pregnancy with a corresponding dominance in *Lactobacillus* spp (Aagaard et al., 2012; Romero et al., 2014).

While one recent study reported an association between the vaginal microbiome richness and diversity and preterm birth (Hyman et al., 2014), others suggest that the vaginal microbiome normally varies across the gestational age spectrum (Aagaard et al., 2012; Romero et al., 2014). What is striking is the lack of shared taxonomy in the microbiome profiles of term or VLBW or preterm infant microbiomes in the first week of life with the maternal vaginal microbiome.

So what might colonize the infant at birth? We recently described the placental microbiome, including comparisons between term and preterm subjects, and found differences in the overall beta diversity with distinct clustering by virtue of preterm birth (Stout et al., 2013; Aagaard et al., 2014). Of interest, we similarly observed that the placental microbiome (in aggregate) most closely resembles the oral microbiome (Aagaard et al., 2014). While the examination of the cord blood and amniotic fluid microbiome is still needed in term neonates, recent studies in preterm neonates have revealed a prevalence of bacteria normally residing in the maternal oral cavity (Wang et al., 2013). Within the first week of life, the development of the microbiome has been studied in term and preterm neonates (Jost et al., 2012; Milisavljevic et al., 2013; Palmer et al., 2007). While Firmicutes has been described to be a major phylum colonizing neonates, these are not predominantly *Lactobacillus*. At a coarse taxonomic level, these observations collectively lead one to speculate that it is neither ascending bacteria from the vaginal tract (nor contact with bacteria from the vaginal tract during delivery) that are the sole or even majority colonizers of the infant microbiome. Alternatively, it appears that the placenta and maternal oral flora are most closely mirrored in the infant during the first week of life.

Figure 2. Dysbiosis of the placental microbiome and risk of preterm birth

Recent studies have suggested that the placenta harbors a low-abundance community of microbes (Stout et al., 2013; Aagaard et al., 2014). On the left, we schematically depict a term labor placenta harboring commensal flora with few associations with periodontal disease, inflammation, or innate immune receptors. On the right, we depict the alternative state of dysbiotic flora, which would be associated with an induction or propagation of inflammation. The resulting imbalance favoring perpetuation of inflammation would lead to a weakening of MMP and other structural proteins, resulting in preterm premature rupture of membranes (PPROM) and induction of preterm labor.

Table 1

Microbiome Studies of Non-gravid and Gravid Populations

Authors	Site	Technique(s)	Study Design	Findings
<i>Non-Gravid</i>				
NIH HMP Consortium (Human and Project, 2012)	Skin, nares, oral, vagina	Sequencing	Longitudinal	Characterized healthy reference population
Ravel <i>et al.</i> (2011)	Mid-vagina (self-collected)	Sequencing	Cross-sectional	Characterized healthy, non-gravid vaginal microbiome
<i>Gravid</i>				
Aagaard <i>et al.</i> (2012)	Vaginal introitus, posterior fornix, and midvagina	Sequencing	Cross-sectional	Characterized healthy, gravid vaginal microbiome
Romero <i>et al.</i> (2014)	Posterior fornix	Sequencing	Longitudinal	Characterized healthy, gravid vaginal microbiome throughout pregnancy
Koren <i>et al.</i> (2012)	Stool	Sequencing	Longitudinal	Characterized 1 st and 3 rd trimester stool
Douvier <i>et al.</i> (1999)	Amniotic fluid, placenta, maternal endocervix	Bacterial culture	Case study	Placenta colonized with oral bacteria, <i>Capnocytophaga sputigena</i>
Han <i>et al.</i> (2006)	Amniotic fluid	PCR	Cross-sectional	Amniotic fluid was colonized with oral bacteria of the <i>Bergeyella</i> spp
Bearfield (2002)	Amniotic fluid, vagina, and dental plaque	Bacterial culture and PCR	Cross-sectional	<i>Streptococcus</i> spp or <i>F. nucleatum</i> mostly found in the oral cavity of gravaidae testing positive in the amniotic fluid.
Stout <i>et al.</i> (2013)	Placenta	Histopathology	Cross-sectional	Bacteria were found in the basal plate of the placenta
Han <i>et al.</i> (2004)	Placenta, liver, spleen, amniotic fluid, fetus	Bacterial culture	Murine study	<i>F. nucleatum</i> isolated from preterm births colonized placenta of gravid mice
Fardini <i>et al.</i> (2010)	Placenta	Sequencing	Murine study	Oral bacteria colonize murine placenta
Aagaard <i>et al.</i> (2014)	Placenta	Sequencing	Population-based, cross-sectional	The placenta harbors a unique microbiome profile, most akin to the oral microbiome and varies by virtue of

Authors	Site	Technique(s)	Study Design	Findings
				preterm birth and a remote history of antenatal infection
<i>Neonatal</i>				
Palmer <i>et al.</i> (2007)	Stool	Sequencing, microarray, PCR	Longitudinal	Characterized healthy neonatal microbiome
Jost <i>et al.</i> (2012)	Stool	Sequencing	Longitudinal	Characterized healthy neonatal microbiome
Wang <i>et al.</i> (2013)	Cord blood, amniotic fluid	Bacterial culture and Sequencing	Cross-sectional	Neonates with necrotizing colitis had predominantly one species of bacteria dominating
Milisavljevic <i>et al.</i> (2013)	Gastroesophageal	Sequencing	Longitudinal	Characterized the microbiome in VLBW infants

PCR polymer chain reaction, VLBW very low birth weight