Published in final edited form as:

Urol Oncol. 2014 January ; 32(1): 24.e7–24.11. doi:10.1016/j.urolonc.2012.08.014.

High resolution ultrasound features of prostatic rib metastasis: A prospective feasibility study with implication in the high-risk prostate cancer patient

Kenneth S. Lee, M.D.^{a,*}, Arthur A. De Smet, M.D.^a, Glenn Liu, M.D.^b, and Mary Jane Staab, R.N.^b

^aUniversity of Wisconsin School of Medicine and Public Health, Department of Radiology, 600 Highland Avenue CSC E3/311, Madison, WI 53792-3252

^bUniversity of Wisconsin Carbone Cancer Center, Division of Hematology/Oncology

Abstract

Objective—In a prior study, high resolution ultrasound (US) was shown to be accurate for evaluating rib metastasis detected on bone scan. However, that study did not address the specific US appearance typical of osteoblastic rib metastasis. Our objective was to determine the specific US imaging appearance of osteoblastic prostate carcinoma rib metastasis using osteolytic renal cell carcinoma rib metastasis as a comparison group.

Materials and methods—The Institutional Review Board approval and informed consent were obtained for this prospective feasibility study. We performed high resolution US of 16 rib metastases in 4 patients with prostate carcinoma metastases and compared them to 8 rib metastases in 3 male patients with renal cell carcinoma. All patients had rib metastases proven by radiographs and computed tomography (CT). High resolution US scanning was performed by a musculoskeletal radiologist using a 12–5 MHz linear-array transducer. Transverse and longitudinal scans were obtained of each rib metastasis.

Results—All 16 prostate carcinoma metastases demonstrated mild cortical irregularity of the superficial surface of the rib without associated soft tissue mass, cortical disruption, or bone destruction. 7 of 8 (88%) renal cell carcinoma rib metastases demonstrated cortical disruption or extensive bone destruction without soft tissue mass. One of 8 (12%) renal cell carcinoma rib metastases demonstrated only minimal superficial cortical irregularity at the site of a healed metastasis.

Conclusion—Osteoblastic prostate carcinoma rib metastases have a distinctive appearance on US. Our success in visualizing these lesions suggests that US may be a useful tool to characterize isolated rib abnormalities seen on a bone scan in high-risk prostate cancer patients who are being evaluated for curative surgery or radiation treatment.

Keywords

Musculoskeletal ultrasound; Rib metastasis; Fracture; Osteoblastic; Osteolytic

1. Introduction

Prostate cancer is the most common cancer in men, and the second most frequent cause of cancer-specific death in the United States [1]. Although controversial, the advent of prostate cancer screening has led to an increased diagnosis of the cancer at earlier stages resulting in high cure rates for those who have organ-confined disease. For men whose prostate specific antigen is >20 ng/ml, Gleason 8, or have clinical T3 or T4 disease (high risk), a staging whole body bone scan is recommended to evaluate for osseous metastasis [2,3].

Unfortunately, given the high sensitivity of bone scintigraphy, areas of increased technetium-99 m-MDP uptake are often found and may be related to trauma, degenerative disease, infection, or healing bone which complicates the staging for osseous metastasis [4]. Therefore, the decision whether or not to subject the patient to morbidities of surgery or radiation therapy can be greatly affected by the bone scan interpretation, particularly in high-risk prostate cancer patients.

One such area that often results in confusion is the ribs [5]. Given that prostate cancer affects older men, who have various degrees of osteoporosis and less soft tissue (fat and muscle) padding the lateral ribs from daily trauma, it is not uncommon to see positive uptake in ribs on a staging bone scan. When no obvious history of trauma can be ascertained, one often attempts to further characterize these lesions with rib x-rays, computed tomography (CT), or magnetic resonance imaging, each with its limitations in evaluating this anatomic area.

Musculoskeletal ultrasound (US) is growing in popularity in the United States and may play a significant role in evaluating rib lesions, especially in the high-risk prostate cancer patient population. Well-known advantages of sonography include its accessibility, use of non-ionizing radiation, the ability to evaluate the costochondral junction [6], and not being affected by respiratory motion. One study showed that US is significantly more sensitive for detecting rib fractures compared to radiography [6].

In patients with a known primary malignancy, only 17% of rib lesions found on bone scintigraphy were malignant [7], and 6% to 8% of malignant bone lesions appeared as a single focus of radiotracer activity involving the ribs on bone scintigraphy [4]. Additional imaging is often needed to determine if a scintigraphic rib lesion represents a metastasis or a fracture. In a previous study, high resolution US was shown to be accurate in differentiating a rib fracture from a rib metastasis [5]. However, the study did not address the specific US appearance of an osteoblastic metastasis, which is the most common appearance for prostate cancer osseous metastasis. Therefore, our objective was to determine the specific US appearance of osteoblastic prostate carcinoma rib metastasis using osteolytic renal cell carcinoma rib metastasis as a comparison group.

2. Materials and methods

2.1. Patient selection

Institutional Review Board approval and informed consent were obtained for this prospective case-controlled study. We enrolled 7 consecutive patients through the University of Wisconsin Comprehensive Cancer Center Oncology Clinic from January 2006 to December 2008. The inclusion criteria included: (1) proven primary prostate or renal cell carcinoma with definite skeletal metastasis by imaging and clinical verification; (2) positive evidence on radiograph, CT, and bone scan of rib metastases by consensus review; and (3) agreement to participate in this study. We selected patients with these 2 primary malignancies for study as almost all prostate carcinoma bony metastases are osteoblastic while almost all renal cell carcinoma bony metastases are osteolytic. All patients were undergoing chemotherapy for their malignancy but had not received radiation therapy to their chest or ribs.

2.2. Sonographic technique

High resolution US scanning was performed and interpreted by 2 experienced musculoskeletal radiologists with a combined 15 years of musculoskeletal US experience; both were blinded to the type of primary malignancy. US examination was performed using a Philips iU22 ultrasound system (Philips Ultrasound, Bothell, WA) and a variable high frequency, high resolution 12–5 MHz linear-array transducer. Each subject was scanned in both the prone and lateral decubitus positions after determining the location of each rib lesion seen on the bone scan. Localization of rib level was performed using US and by counting the number of ribs in the paraspinal line just lateral to the costovertebral junction, locating the most inferior rib as the reference standard (Fig. 1). Once each rib metastasis was localized, transverse and longitudinal sonographic gray scale and Doppler images were obtained and stored on a radiology workstation (Horizon Rad Station, version 11.0 service package 6, McKesson) and DICOM viewer (eFilm Lite software, Merge Healthcare).

We did not evaluate lesions in the lateral arc of the first to fourth ribs because of shadowing from the scapula and clavicle, nor the costovertebral junction which was obscured by the orientation of the rib at this location. These limitations of US in assessing the ribs were noted in the study by Paik et al. [5].

2.3. Retrospective clinical review

The medical records of each subject were reviewed to determine the date of last follow-up at our hospital, the site of primary cancer, and the history for recent trauma or surgery.

2.4. Data analysis

The findings of all available radiographs, CT, and bone scintigraphy of each abnormal rib lesion site were reviewed along with the sonographic images, blinded to the diagnosis. Radiograph and CT findings of each scintigraphic rib lesion were classified into osteoblastic or osteolytic lesions, rib fracture, or normal appearance and then correlated with US findings.

The sonographic images were reviewed to characterize osteoblastic or osteolytic metastasis according to a modification of the criteria used by Paik et al. [5]: cortical disruption described as focal disruption or angulation of the normally smooth echogenic cortical line of the rib surface, cortical surface irregularity as loss of the normal smooth contour of the echogenic line, bone destruction as large area of complete loss of the echogenic line with or without vascularity, and the presence of an exophytic soft tissue mass.

3. Results

We studied 24 rib lesions in 7 consecutive patients (all men; 46–84 years old; mean 67.3 years old) with 16 rib metastases in 4 patients with prostate carcinoma and 8 rib metastases in 3 male patients with renal cell carcinoma. All patients had rib metastases proven by radiographs, CT, or bone scan.

All 16 prostate carcinoma metastases demonstrated mild cortical irregularity of the superficial surface of the rib without associated soft tissue mass, cortical disruption, or bone destruction. 7 of 8 (88%) renal cell carcinoma rib metastases demonstrated cortical disruption or extensive bone destruction without soft tissue mass. 1 of 8 (12%) renal cell carcinoma rib metastases demonstrated only minimal superficial cortical irregularity. This lesion was clinically considered to be a regressing rib metastasis according to follow-up imaging. Of the metastatic rib lesions, 13 were located posteriorly, 6 were located posterolaterally, and 5 were located anteriorly.

4. Discussion

Rib lesions found on bone scintigraphy can complicate an already difficult treatment approach in high-risk prostate cancer patients. Confirmed metastatic disease precludes a surgical option. However, bone scintigraphic findings in the ribs can be non-specific, and eventually a clinical decision needs to be made based on non-specific results. Developing tools to better characterize these incidental findings is of high importance as the clinical decision can either result in an attempt at curative therapy or not.

The use of musculoskeletal US may offer a quick non-invasive approach to diagnosis. US application in the bony thorax has been limited by structures difficult to evaluate due to shadowing from closely positioned ribs of the thoracic cage and air from the subjacent lungs [8]. One known application of sonography is its use in detecting occult rib fractures [6]. Sonography can detect 10 times more fractures than in radiographs, and in 6 times as many patients [6]. Only 1 prior study expanded the use of high resolution sonography by showing that US can differentiate rib fracture from rib metastasis [5]. However, we know of no study that characterizes the specific sonographic imaging appearance of osteoblastic prostate rib metastasis, which may be important in high-risk patients with a non-specific rib finding on bone scintigraphy.

A feature that suggested metastatic rib involvement in a prior study was if bone scintigraphy showed a hot spot while sonographic evaluation in this region was negative [5,9]. This study suggested that the only role of US was to exclude a traumatic fracture. However, a subsequent study has shown that metastatic rib involvement has distinct US features, which

include mass formation and irregular bone destruction, but did not categorize typical US findings with the type of skeletal metastases [5]. We found that osteoblastic rib lesions have different sonographic appearances (Table 1), which could be contrasted with osteolytic rib lesions from renal cell carcinoma.

In our study, osteoblastic prostate carcinoma metastasis demonstrated mild cortical irregularity of the superficial surface of the rib without associated soft tissue mass, cortical disruption, or bone destruction (Fig. 2). In contradistinction, the sonographic features of osteolytic renal cell carcinoma rib metastasis were more aggressive and distinctly different from osteoblastic prostatic lesions. The majority of osteolytic renal cell rib lesions demonstrated frank cortical disruption with extensive bone destruction (Fig. 3), but interestingly, no soft tissue mass or increased vascularity (Fig. 4). Our finding of no associated soft tissue mass in osteolytic lesions was different compared to Lin et al. who found that renal cell metastatic involvement to be a highly vascular soft tissue replacement process of bone [10]. The lack of associated vascularity and soft tissue mass may be the sequela of chemotherapy, similar to our finding of a regressing rib metastasis in the same patient with renal cell carcinoma.

A previous study showed that 12% of solitary rib hot spots were caused by metastases [9]. Solitary rib hot spots on bone scintigraphy can be a diagnostic dilemma for clinicians compared to multiple hot spots in those with known primary carcinoma. Cancer patients may be fragile and cachexic and predisposed to rib fracture even after simple movements or coughing. The ability to confirm a rib metastasis is important for clinical management, especially in the high-risk prostate cancer patient where surgery may be contemplated. Confirmation of a rib metastatic lesion can help direct therapy, or more importantly, avoid an unnecessary biopsy that delays treatment. Our study further characterizes rib metastatic lesions that may prove helpful in patients that present with a solitary hot spot on bone scintigraphy.

The strength of our study is its prospective design, but there were several limitations. Although a feasibility study was done, we found distinctive differences in osteoblastic prostatic rib lesions compared to osteolytic rib lesions which are consistent with the known histopathologic features of these types of metastatic lesions. Thus, we feel that these sonographic appearances will be consistent between prostate carcinoma and renal cell carcinoma rib metastases in a larger study. Another limitation was that subjects were not controlled for the duration of chemotherapy, which may introduce variability of the US appearance at imaging presentation. This may help to explain our finding of no associated soft tissue mass in renal cell carcinoma rib lesions. Benign rib fractures were not included in this study, which could be a limitation. However, rib metastatic lesions have already been well differentiated from rib fracture in a prior study by Paik et al. [5], which our findings support. Our initial findings in this feasibility study support a larger scale study that would include other types of cancers and comparison across additional modalities such as radiographs, CT, and magnetic resonance imaging. We suspect that US would be both highly sensitive and specific for its high resolution, and the modality of choice due to its accessibility and low cost.

In conclusion, osteoblastic prostate metastatic rib lesions have distinctive US features. With this knowledge, high resolution US evaluation of the rib may be a quick and non-invasive method for not only differentiating rib fracture from metastasis, but also confirming osteoblastic prostatic rib metastasis. However, larger studies need to be done.

References

1. Siegel R, Naishadham D, Jemal A. Cancer statistics. *CA Cancer J Clin*. 2012; 62(1):10–29. [PubMed: 22237781]
2. Gnanapragasam VJ, Mason MD, Shaw GL, Neal DE. The role of surgery in high-risk localised prostate cancer. *BJU Int*. 2012; 109(5):648–58. [PubMed: 21951841]
3. Schmitges J, Trinh QD, Walz J, Graefen M. Surgery for high-risk localized prostate cancer. *Ther Adv Urol*. 2011; 3(4):173–82. [PubMed: 21969847]
4. Corcoran RJ, Thrall JH, Kyle RW, Kaminski RJ, Johnson MC. Solitary abnormalities in bone scans of patients with extraosseous malignancies. *Radiology*. 1976; 121(3 Pt 1):663–7. [PubMed: 981663]
5. Paik SH, Chung MJ, Park JS, Goo JM, Im JG. High-resolution sonography of the rib: can fracture and metastasis be differentiated? *AJR Am J Roentgenol*. 2005; 184(3):969–74. [PubMed: 15728626]
6. Griffith JF, Rainer TH, Ching AS, Law KL, Cocks RA, Metreweli C. Sonography compared with radiography in revealing acute rib fracture. *AJR Am J Roentgenol*. 1999; 173(6):1603–9. [PubMed: 10584808]
7. McNeil BJ. Value of bone scanning in neoplastic disease. *Semin Nucl Med* Oct. 1984; 14(4):277–86.
8. Mathis G. Thoraxsonography—part I: chest wall and pleura. *Ultrasound Med Biol*. 1997; 23(8): 1131–9. [PubMed: 9372561]
9. Wu PS, Chiu NT, Lee BF, Yao WJ, WCHH. Clinical significance of solitary rib hot spots on bone scans in patients with extraskkeletal cancer: correlation with other clinical manifestations. *Clin Nucl Med*. 2002; 27(8):567–71. [PubMed: 12170001]
10. Lin PP, Mirza AN, Lewis VO, et al. Patient survival after surgery for osseous metastases from renal cell carcinoma. *J Bone Joint Surg Am*. 2007; 89(8):1794–801. [PubMed: 17671020]

Fig. 1.

Localizing method to evaluate a posterior seventh rib lesion in a 61-year-old male with renal cell carcinoma while in prone position. Extended field of view with US probe oriented longitudinally in the posterior paraspinal line demonstrating the typical appearance of the lower rib cage used for localizing rib lesions. The inferior most rib is the reference point used to locate the seventh rib-containing lesion (arrow).

Fig. 2.

Sonographic appearance of an osteoblastic prostate rib metastasis in a 70-year-old male with prostate carcinoma. (A) Longitudinal and (B) transverse sonographic images of the posterolateral eighth rib demonstrate mild cortical irregularity of the superficial surface of the rib (arrows) without associated soft tissue mass, cortical disruption or bone destruction.

Fig. 3.

Sonographic appearance of an osteolytic renal cell rib metastasis in a 46-year-old male with renal cell carcinoma. (A) Axial computed tomography image of the chest shows an expansile lytic lesion of the posterolateral rib (arrow). (B) Corresponding longitudinal sonographic image of the same rib lesion demonstrates cortical disruption with underlying bone expansion and destruction (arrows).

Fig. 4. Sonographic appearance of another osteolytic renal cell rib metastasis in a 70-year-old male with renal cell carcinoma. Longitudinal Doppler sonographic image of a posterior rib lesion shows extensive rib destruction with complete loss of the normal cortical echogenic line (arrows), but without soft tissue mass or vascularity.

Table 1

Differentiating sonographic features of osteoblastic prostate and osteolytic renal rib metastasis

	Osteoblastic	Osteolytic
Cortical irregularity	+	–
Cortical disruption	–	+
Cortical destruction	–	+
Soft tissue mass	–	–