Published in final edited form as:

Fertil Steril. 2010 July ; 94(2): e44–e440. doi:10.1016/j.fertnstert.2010.04.069.

Reply of the Authors: Cellular senescence in usual type uterine leiomyoma

Jian-Jun Wei, M.D.

Department of Pathology, Robert H. Lurie Comprehensive Cancer Center, Northwestern University, Chicago, Illinois

Reply of the Authors

We thank Drs. Baird, Davis, and Peddada for sharing their findings of the natural history for fibroid growth in association with age and ethnicity (1). Since racial disparity for fibroid growth is a very interesting topic, it deserves another look for racial differences in cellular senescence.

We examined a total of 86 fibroids (36 from black women, 28 from white women, and 22 from women of other races) for the study (Table 1) (2). The average percentage of senescence was 20.1 ± 2.84 from white women and 19.1 ± 3.05 from black women. No statistical significance was found in this small cohort. However, the experiment was not designed for the study of racial differences. In particular, the sample size was small, and age and clinical settings were not matched between black and white women. Apparently it requires a well-designed study to address the issue of age-related cellular senescence for fibroid growth in different races.

References

1. Peddada SD, Laughlin SK, Miner K, Guyon JP, Haneke K, Vahdat HL, et al. Growth of uterine leiomyomata among premenopausal black and white women. *Proc Natl Acad Sci USA.* 2008; 105:19887–92. [PubMed: 19047643]
2. Laser J, Lee P, Wei J-J. Cellular senescence in usual type uterine leiomyoma. *Fertil Steril.* 2010; 93:2020–6. [PubMed: 19217096]