Published in final edited form as:

Endocr Relat Cancer. 2014 October ; 21(5): T301–T313. doi:10.1530/ERC-14-0166.

Molecular Approaches to Thyroid Cancer Diagnosis

Susan J. Hsiao and Yuri E. Nikiforov

Division of Molecular and Genomic Pathology, Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh PA

Abstract

Thyroid nodules are common, and the accurate diagnosis of cancer or benign disease is important for the effective clinical management of these patients. Molecular markers are a helpful diagnostic tool, particularly for cytologically indeterminate thyroid nodules. In the past few years, significant progress has been made in developing molecular markers for clinical use in fine needle aspiration (FNA) specimens, including gene mutation panels and gene expression classifiers. With the availability of next generation sequencing technology, gene mutation panels can be expanded to interrogate multiple genes simultaneously and to provide yet more accurate diagnostic information. In addition, recently several new molecular markers in thyroid cancer have been identified that offer diagnostic, prognostic, and therapeutic information that could potentially be of value in guiding individualized management of patients with thyroid nodules.

Keywords

thyroid cancer; thyroid nodule; molecular marker; prognostic marker

Introduction

Among endocrine tumors, thyroid cancer is the most common, with an estimated incidence of 12.2 per 100,000 per year in the United States (Howlader, et al. 2013). The incidence of thyroid cancer, both in the United States and world-wide, has been increasing over the last four decades (Burgess and Tucker 2006; Davies and Welch 2006; Albores-Saavedra, et al. 2007). The increased incidence of thyroid cancer diagnoses has been attributed, in part, to improved detection of small or subclinical thyroid nodules on thyroid ultrasonography and by other imaging techniques; however, increased incidence of thyroid tumors of all sizes has also been reported (Albores-Saavedra et al. 2007; Jung, et al. 2013). The increased number of cases of papillary thyroid cancer is predominantly of follicular variant, *RAS* mutation positive tumors, suggesting a potential role for environmental (chemical/dietary) factors (Jung et al. 2013).

In addition to environmental factors, genetic factors are involved in thyroid cancer predisposition. Aside from the well-characterized familial forms of medullary thyroid

Corresponding Author: Yuri E. Nikiforov MD PhD, Department of Pathology, University of Pittsburgh, 3477 Euler Way, Room 8031, Pittsburgh, PA 15213, nikiforovye@upmc.edu.

Declaration of Interest: Y.E.N. serves as consultant for Quest Diagnostics. S.J.H. has no potential conflicts of interest.

cancer, non-medullary thyroid cancer in a first-degree relative increases the risk 4-10 fold higher than in the general population (Frich, et al. 2001; Hemminki, et al. 2005). Familial non-medullary thyroid cancer is characterized by autosomal dominant inheritance with reduced penetrance, and has been estimated to account for approximately 5-10% of all thyroid cancers (Charkes 2006; Malchoff and Malchoff 2006; Moses, et al. 2011; Mazeh and Sippel 2013). Genetic linkage studies have mapped susceptibility loci to several regions including 1q21, 2q21, 8p23, 8q24, 9q22, 14q31, and 19p13 (Bignell, et al. 1997; Canzian, et al. 1998; Malchoff, et al. 2000; McKay, et al. 2001; Cavaco, et al. 2008; He, et al. 2009; Tomaz, et al. 2012). Definitive germline genetic mutations underlying thyroid cancer predisposition remain yet to be identified within candidate genes in these regions. Thyroid tumor development likely involves a complex interplay between genetic predisposition and environmental risk factors.

Thyroid cancer typically presents as a thyroid nodule. However, thyroid nodules are commonly found incidentally and may be seen in up to 50% of patients older than 60 years (Mazzaferri 1992, 1993; Guth, et al. 2009). Only 5% of thyroid nodules are malignant (Brito, et al. 2012). Most thyroid cancers are well-differentiated papillary carcinomas or follicular carcinomas and are associated with a low mortality rate, particularly in patients with stage I or II disease (survival rate >98%). However, a subset of these patients will have recurrent disease (Mazzaferri and Jhiang 1994). In addition, patients who present with higher stage disease or distant metastases, and patients with poorly differentiated or anaplastic thyroid cancer have higher mortality rates (Volante, et al. 2004; Tanaka, et al. 2011). Accurate identification of subsets of patients with risk factors for aggressive disease and higher mortality rates can help guide treatment and management and as well as prevent overtreatment of patients with low-risk disease.

Thyroid Cancer Diagnosis in FNA and the Need for Molecular Markers

The diagnosis of thyroid cancer is typically obtained through ultrasound examination and fine-needle aspiration (FNA) biopsy of suspicious nodules. Cytologic examination of cells collected by FNA biopsy is the most reliable diagnostic method for evaluating thyroid nodules, and is able to definitely classify thyroid nodules as benign or malignant in the majority of cases (Cooper, et al. 2009; Gharib, et al. 2010). The Bethesda reporting system for classifying thyroid cytology was proposed in 2007 by the National Cancer Institute and provides diagnostic categories with accompanying risk stratification and recommended clinical management (Baloch, et al. 2008; Ali and Cibas 2010). Thyroid FNA specimens in the benign category have a low risk of malignancy (~0-3%), and thyroid FNAs in the malignant category have a 97-99% risk of malignancy (Ali and Cibas 2010). However, 20-30% of thyroid FNA specimens are indeterminate and fall into one of the following categories: atypia of undetermined significance/follicular lesion of undetermined significance (AUS/FLUS), follicular or oncocytic (Hürthle cell) neoplasm/suspicious for a follicular or oncocytic (Hürthle cell) neoplasm (FN/SFN), and suspicious for malignant cells (SUSP) (Baloch et al. 2008; Otori and Schoedel 2011). The AUS/FLUS diagnostic category is associated with a 5-15% risk of malignancy and the recommended management is repeat FNA; for the FN/SFN category, the risk of malignancy is 15-30% and the recommended management is surgical lobectomy; and for the SUSP category, the risk of malignancy is

60-75% and the recommended management is near-total thyroidectomy or surgical lobectomy (Ali and Cibas 2010).

Of the indeterminate thyroid nodules that are surgically resected, 10-40% are confirmed to be malignant (Mazzaferri 1993; Baloch, et al. 2002; Baloch et al. 2008). As a result, most diagnostic surgeries are performed for benign thyroid nodules. On the other hand, for patients who have undergone a surgical lobectomy and found to have a tumor larger than 1 cm, a second surgery is usually performed to remove the remaining thyroid lobe. Therefore, additional diagnostic markers are needed to guide the management of patients with indeterminate thyroid nodules in order to reduce the frequency of unnecessary diagnostic lobectomies and two-step surgeries.

Several types of ancillary approaches have been used to improve the diagnostic yield of fine needle aspiration biopsies in indeterminate thyroid nodules. These include immunohistochemical stains, microRNAs, gene mutations/rearrangements, and gene expression panels (Bartolazzi, et al. 2008; Nikiforov, et al. 2011; Alexander, et al. 2012; Keutgen, et al. 2012). While many of these ancillary tools have not yet reached clinical practice, two such approaches, gene mutation/rearrangement panels and gene expression classifier, are currently being used for clinical management (Nikiforov et al. 2011; Alexander et al. 2012). These panels are based on the wealth of knowledge on thyroid cancer genetics accumulated over the last three decades. In addition, gene mutation/rearrangement panels have the added benefit of providing information that could be potentially useful in prognostication and targeted therapy.

The utility of molecular markers in indeterminate thyroid nodules can be evaluated based on the sensitivity, specificity, negative predictive value, and positive predictive value of each test. Sensitivity is a measure of the proportion of actual positives which are correctly identified as such, and specificity is the proportion of negatives which are correctly identified as such. Negative predictive value (NPV) is the percentage of patients with a negative test result who do not have the disease, and positive predictive value (PPV) is the percentage of patients with a positive test result who have the disease. A negative result in a molecular test with high sensitivity and negative predictive value, would suggest that the patient with an indeterminate thyroid nodule has a higher likelihood of having a benign nodule and could be managed with active surveillance, whereas a positive result in a molecular test with high specificity and positive predictive value suggests that the patient with an indeterminate thyroid nodule likely has thyroid cancer and would be an indication for surgery. Importantly, although specificity and sensitivity depend only on test performance, NPV and PPV depend on the prevalence of disease in the tested population. Therefore, institutional differences in malignancy rates conferred by each cytologic diagnosis may result in significant variation of NPV and PPV, which should be taken into account when any molecular test is used clinically.

Molecular Genetics of Thyroid Cancer

From the 1990s, when pathogenesis of only approximately 25% of thyroid cancers was understood, to the present, when genes involved in the pathogenesis of greater than 90% of

thyroid cancers have been described, much progress has been made in elucidating the molecular mechanisms underlying thyroid cancer (Fig. 1). This progress provides the basis upon which new diagnostic and prognostic markers, as well as new targeted therapies, have been developed.

The molecular pathogenesis of the majority of thyroid cancer involves dysregulation of the mitogen-activated protein kinase (MAPK) and phosphatidylinositol-3 kinase (PI3K)/AKT signaling pathways (Fig. 2). The MAPK pathway is frequently activated in thyroid cancer through point mutations of the *BRAF* and *RAS* genes and *RET/PTC* and *TRK* rearrangements (Kimura, et al. 2003; Soares, et al. 2003; Frattini, et al. 2004; Adeniran, et al. 2006). Point mutation in *BRAF* is found in approximately 45% of papillary thyroid cancers (Cohen, et al. 2003; Kimura et al. 2003). *BRAF* is a serine-threonine kinase which, upon activation by *RAS*, activates *MEK* and leads to activation of downstream effectors of the MAPK pathway. In nearly all cases (98-99% of cases) activating point mutations of *BRAF* involve codon 600 and result in the V600E mutation, and in 1-2% of cases other *BRAF* mutations such as the K601E mutation, small in-frame insertions or deletions, or *BRAF* rearrangement can occur (Soares et al. 2003; Ciampi, et al. 2005; Ciampi and Nikiforov 2005; Hou, et al. 2007b; Chiosea, et al. 2009).

RAS genes (*HRAS*, *KRAS*, and *NRAS*) are G proteins that signal to both the MAPK and PI3K/AKT pathways. Point mutations in the *RAS* genes typically occur in codons 12, 13, and 61, and are found in 40-50% of follicular carcinomas and in 10-20% of papillary thyroid carcinomas (Lemoine, et al. 1989; Namba, et al. 1990; Suarez, et al. 1990). *RAS*-mutated papillary thyroid carcinomas typically are of the follicular variant (Zhu, et al. 2003; Adeniran et al. 2006). *RAS* mutations are also seen in 20-40% of follicular adenomas (Lemoine et al. 1989; Namba et al. 1990; Suarez et al. 1990; Motoi, et al. 2000). Whereas *NRAS*, *HRAS* and *KRAS* mutations are found in follicular-cell derived thyroid tumors, mutations in *HRAS* and *KRAS* also occur in medullary thyroid cancers (Agrawal, et al. 2013).

The *RET* gene is a receptor tyrosine kinase that is expressed in thyroid C cells, but not in follicular cells. The *RET* gene can be activated by fusion with various partners that drive the expression of the 3' portion of the *RET* gene coding for the tyrosine kinase domain of the receptor, and provide the dimerization motif to lead to the constitutive activation of RET kinase. The most common rearrangement types are *RET/PTC1* (formed by fusion of *RET* with the *CCDC6* gene) and *RET/PTC3* (formed by fusion of *RET* with the *NCOA4* gene) (Grieco, et al. 1990; Santoro, et al. 1994). The *RET/PTC1* and *RET/PTC3* rearrangements are found in 10-20% of papillary thyroid carcinomas (Nikiforov 2006; Zhu, et al. 2006), and their incidence is progressively decreasing (Jung et al. 2013). These rearrangements are found at higher frequencies in children/young adults and in patients with a history of radiation exposure (Nikiforov, et al. 1997; Fenton, et al. 2000; Rabes, et al. 2000). In addition to rearrangements involving *RET* which are found in papillary thyroid tumors, the *RET* gene is commonly found to be mutated in medullary thyroid carcinomas, in both familial and sporadic cases (de Groot, et al. 2006; Kloos, et al. 2009).

The *PAX8/PPAR γ* rearrangement, a fusion between a paired domain transcription factor and the peroxisome proliferator-activated receptor genes, is found in 30-40% of follicular carcinomas (Dwight, et al. 2003; French, et al. 2003; Nikiforova, et al. 2003). The *PAX8/PPAR γ* rearrangement can also be seen, at lower prevalence, in the follicular variant of papillary thyroid carcinoma and in follicular adenomas (Marques, et al. 2002; Nikiforova, et al. 2002; Dwight et al. 2003; French et al. 2003; Nikiforova et al. 2003).

The importance of the PI3K/AKT pathway in thyroid tumorigenesis has been increasingly recognized in the last decade. The PI3K/AKT pathway can be activated by activating mutations in *PIK3CA* and *AKT1* as well as by inactivation of *PTEN*, which negatively regulates this pathway (Fig. 2). Somatic mutations of *PTEN* have been reported in follicular thyroid tumors and anaplastic thyroid carcinoma, and germline mutations of *PTEN* can result in follicular thyroid tumors arising in patients with Cowden syndrome (Dahia, et al. 1997; Gustafson, et al. 2007; Hou, et al. 2007a; Nikiforova, et al. 2013). Activating mutations in *PIK3CA* typically occur at hotspots within exon 9 and exon 20 and have been reported in follicular thyroid carcinomas, poorly differentiated thyroid carcinomas, and anaplastic thyroid carcinomas (Garcia-Rostan, et al. 2005; Hou et al. 2007a; Ricarte-Filho, et al. 2009). *AKT1* mutations have been reported in metastatic thyroid cancer (Ricarte-Filho et al. 2009).

Additional genes mutated in thyroid cancer include *TP53* and *CTNNB1* (beta-catenin). *TP53* is a tumor suppressor that plays important roles in cell cycle regulation and DNA repair and *CTNNB1* is involved in Wnt signaling. These genes tend to be mutated in more aggressive and advanced thyroid tumors (Dobashi, et al. 1994; Garcia-Rostan, et al. 2001). In addition to these well-characterized mutations, mutations in thyroid stimulating hormone receptor (*TSHR*) and *GNAS* have also been shown to play a role in thyroid tumorigenesis. Also, novel markers have been recently identified, including *ETV6/NTRK3*, *STRN/ALK*, and *TERT*. These markers will be discussed in further detail below.

Utility of Mutational Molecular Markers in Preoperative FNA Samples

BRAF and *RAS* point mutations and *RET/PTC* and *PAX8/PPAR γ* rearrangements are the most common genetic alterations found in thyroid cancer and have been used for cancer detection in thyroid nodules with indeterminate FNA cytology. A seven (or eight) gene mutation panel which includes *BRAF*, *KRAS*, *HRAS*, *NRAS*, and *RET/PTC1*, *RET/PTC3*, *PAX8/PPAR γ* (and *TRK*) rearrangements is the best characterized molecular panel. The presence of any mutation has high specificity and high positive predictive value for malignancy, as demonstrated in three prospective studies, one of which involved two institutions and the rest were based on a single institution (Nikiforov, et al. 2009; Cantara, et al. 2010; Nikiforov et al. 2011). All three studies have demonstrated a high specificity and positive predictive value of the positive test for cancer detection in all categories of indeterminate cytology (Nikiforov et al. 2009; Cantara et al. 2010; Nikiforov et al. 2011). These results provide strong evidence that the presence of any of these mutations, with the possible exception of *RAS* (discussed below), is an indication for surgery, and specifically for total thyroidectomy. The strategy of using mutational markers combined with cytologic diagnosis to guide the extent of surgery was recently shown to be effective in reducing the

need for two-step surgery, i.e. initial lobectomy followed by completion thyroidectomy (Yip, et al. 2014). In a series of 471 FNA biopsies that had indeterminate cytology (AUS/FLUS or FN/SFN), patients who did not have mutational marker testing were 2.5-fold more likely to require a two-stage surgery (Yip et al. 2014).

RAS mutations, compared to *BRAF* mutation or *RET/PTC* and *PAX8/PPAR γ* rearrangements, have a lower positive predictive value for cancer of 74% to 87% (Nikiforov et al. 2009; Cantara et al. 2010; Nikiforov et al. 2011). The benign nodules positive for *RAS* mutation were found, on surgical resection, to be follicular adenomas. Although follicular adenomas are benign, evidence exists that follicular adenomas represent a pre-cancerous change that can progress to malignancy (Burns, et al. 1992; Fagin 2002; Zhu et al. 2003; Nikiforov and Ohori 2012). Thus such patients would benefit from removal of these nodules before possible progression.

Negative predictive value of mutational panels has been best characterized for the 7-gene panel. In mutation-negative nodules with AUS/FLUS cytology, the residual risk of malignancy in a large series of cases (with the disease prevalence of 14%) was 6%, with a 2.3% risk of invasive cancer (Nikiforov et al. 2011). In these AUS/FLUS nodules, the negative predictive value is high at 94% (Nikiforov et al. 2011). Because the residual risk of cancer approaches that of the risk in nodules with benign cytology, and because most missed cancers are intra-thyroidal (0.5% risk of extra-thyroidal spread), low grade tumors, this test can be used to eliminate the need for surgery in AUS/FLUS nodules, as long as the disease prevalence in the AUS/FLUS category does not exceed 14-15%.

In mutation-negative nodules with FN/SFN cytology, the 7-gene panel decreases the risk of malignancy from 27% to 14% and in mutation-negative nodules with SUSP cytology from 54% to 28% (Nikiforov et al. 2011). As the risk of malignancy is significantly decreased, these patients could be offered a diagnostic lobectomy rather than a total thyroidectomy. However, an improved negative predictive value would still be desired to reduce the risk further.

This molecular mutational panel may also play a valuable role in pediatric patients with thyroid nodules, in which thyroid FNA biopsy may result in an indeterminate diagnosis in as many as 38% of cases (Monaco, et al. 2012). In a small series of cases, preoperative molecular testing was able to guide surgical management and prevent the need for a second surgery in 60% of cases (Buryk, et al. 2013).

Gene Expression Markers for FNA Diagnosis of Indeterminate Nodules

A panel of gene expression markers was identified by examining mRNA expression profiles in thyroid nodules and utilizing that data to train a molecular classifier (Chudova, et al. 2010). This gene expression classifier uses the expression of 142 genes to classify thyroid nodules into a benign or suspicious category (Chudova et al. 2010). The reported genes used in the gene expression panel are involved in many processes, including energy metabolism and cell differentiation/development (Alexander et al. 2012).

The molecular test based on the gene expression classifier is commercially offered as Afirma. The test was validated in one study, which was a multi-institutional prospective double blind study which included 265 nodules with indeterminate cytology from 49 clinical sites (Alexander et al. 2012). The study limitations include a relatively small sample size (129 AUS/FLUS, 81 FN/SFN, and 55 SUSP samples) and a high rate of post-unblind exclusion of samples. The study demonstrated that, with a disease prevalence of 24%, 25%, and 62% in AUS/FLUS, FN/SFN, and SUSP cytology groups, respectively, the negative predictive value was 95% in AUS/FLUS, 94% in FN/SFN, and 85% in SUSP cytology nodules. The positive predictive value was significantly lower and varied by cytologic category (AUS/FLUS – 38%, FN/SFN – 37%, and SUSP – 76%) (Alexander et al. 2012). Therefore, this test is marketed as a “rule-out” test, i.e. the test to identify nodules likely to be benign and thus avoid unnecessary surgery (Duick, et al. 2012; Alexander, et al. 2014). Importantly, the NPV for this test also depends on the disease prevalence in each category of indeterminate cytology. A recent study showed a lower NPV for the Afirma gene expression classifier test, which was found to be 89.6% for AUS/FLUS and FN/SFN cytology nodules (Harrell and Bimston 2013). This is likely due to a higher disease prevalence in patients with indeterminate cytology nodules, which was 33% in this study. Therefore, additional independent, not industry supported studies are required to establish the performance of this test in different patient populations.

Expanded Mutational Marker Panels for FNA Specimens

The continuing discovery of genes involved in thyroid carcinogenesis together with the availability of new high-throughput technologies has led to a rapid expansion of mutational panels that can be used in FNA samples. The expanded panels can detect more mutations and with higher sensitivity, which is expected to increase significantly the sensitivity and NPV of mutational panels. Next generation sequencing (NGS) technologies allow the high-throughput, massively parallel sequencing of nucleic acid sequences and offer a cost-effective way to analyze a large number of genetic alterations in small samples. NGS-based approaches generate an increased amount of complex information with specialized requirements for analysis and reporting that can be effectively managed with bioinformatics applications (e.g. SeqReporter) (Roy, et al. 2014). A 15-gene mutational panel expanded to include other clinically significant genes that have been found to be mutated in thyroid cancer, such as *PIK3CA*, *TP53*, *TSHR*, *PTEN*, *GNAS*, *CTNNB1*, *AKT1*, and *RET*, has recently been reported (Nikiforova et al. 2013).

Mutations in *PIK3CA* and *AKT1* have been reported in thyroid tumors, and occur more frequently in advanced and dedifferentiating tumors (Garcia-Rostan et al. 2005; Hou et al. 2007a; Ricarte-Filho et al. 2009). Mutations in *PIK3CA* were detected, in a limited series of this expanded mutational panel, in papillary thyroid carcinoma, poorly differentiated thyroid carcinoma, and anaplastic thyroid carcinoma, and were found both in the presence and absence of other mutations (Nikiforova et al. 2013). *PTEN* mutations, consistent with previous reports, were detected in follicular carcinoma, as well as in benign follicular adenomas (Dahia et al. 1997; Hou et al. 2007a; Nikiforova et al. 2013).

Additional mutations known to occur in more aggressive types of well differentiated thyroid cancer as well as in poorly differentiated and anaplastic carcinomas involve *TP53* and *CTNNB1* (Dobashi et al. 1994; Garcia-Rostan et al. 2001). Interestingly, in one study, *TP53* mutations were found not only anaplastic carcinomas, but also in 22% of oncocytic follicular carcinomas (Nikiforova et al. 2013).

Somatic mutations of the *TSHR* gene are known to frequently occur in autonomously functioning thyroid nodules (Garcia-Jimenez and Santisteban 2007; Nishihara, et al. 2009). However, *TSHR* mutations at specific hotspots were also found in thyroid carcinomas (Nikiforova et al. 2013). Similar to *TSHR* mutations, mutations in *GNAS*, a gene which encodes an alpha subunit of heterotrimeric G protein complexes, occur predominantly in benign hyperfunctioning nodules. In a limited series, all nodules found to carry an isolated *GNAS* mutation were found to be benign after surgery (Nikiforova et al. 2013). Therefore, *GNAS* mutations may uniquely function as markers of benign nodules.

The *RET* gene is commonly found to be mutated in medullary thyroid carcinomas, in both familial and sporadic cases. Among sporadic medullary carcinomas, the *RET* M918T mutation is the most common, accounting for more than 75% of all somatic *RET* mutations found in these tumors (de Groot et al. 2006; Kloos et al. 2009). The *RET* M918T mutation is also the most common mutation in MEN2B, whereas in MEN2A and familial medullary thyroid carcinoma, mutations in *RET* typically occur in one of five cysteine codons within the cysteine-rich extracellular domain (Mulligan, et al. 1995; Hansford and Mulligan 2000). Detection of specific mutations of *RET* in medullary thyroid carcinoma (or mutation of *KRAS* or *HRAS*, which can occur in sporadic cases), thus can not only inform diagnosis, but can also facilitate genetic analysis of germline mutations (Agrawal et al. 2013).

An expanded 15-gene NGS-based panel has been validated in a study of 228 samples from thyroid nodules including 51 FNA samples and showed accurate detection of multiple mutations with a sensitivity of 3-5% of mutant allele (Nikiforova et al. 2013). This analysis was able to identify mutations in 27 tumors which would not have been detected by mutational analysis of *BRAF* and *RAS* genes, indicating that it is expected to increase the sensitivity and NPV of cancer detection in thyroid nodules (Nikiforova et al. 2013). In addition, although most tumors in this study were positive for a single mutation, in 9 tumors, 2-3 mutations were identified (Nikiforova et al. 2013). These results suggest that in clinical practice, a targeted NGS panel will further increase the accuracy of cancer risk assessment in thyroid nodules and will provide information about the presence of multiple mutations, which has prognostic implications.

Emerging Novel Molecular Markers

In addition to the *RET/PTC* and *PAX8/PPAR γ* rearrangements which are found in approximately 15% of thyroid tumors, many other gene fusions have been described such as rearrangements involving *NTRK* or *BRAF*. Fusion of *BRAF* with A kinase anchor protein 9 (*AKAP9*) is a rearrangement rarely found in sporadic thyroid cancer, although it also occurs at higher frequencies (up to 11%) in patients with a history of radiation exposure (Ciampi et al. 2005). *NTRK1* is a receptor tyrosine kinase and when rearranged with one of 3 potential

fusion partners, activates MAPK pathway signaling (Bongarzone, et al. 1998; Musholt, et al. 2000). *NTRK1* rearrangements are found in approximately 1-5% of papillary thyroid carcinomas and in higher frequencies in patients with radiation exposure (Ciampi et al. 2005; Leeman-Neill, et al. 2013a). More recently, whole-transcriptome (RNA-Seq) analyses led to the discovery of novel gene fusions in thyroid cancer. RNA-Seq analysis of radiation-associated thyroid cancer identified a novel *ETV6-NTRK3* chromosomal rearrangement, which occurs in 2% of sporadic papillary thyroid cancers and 14.5% of radiation-associated tumors (Leeman-Neill, et al. 2013b). Another interesting gene fusion with therapeutic implications was recently identified by RNA-Seq analysis of aggressive forms of thyroid cancer. The fusion of the striatin (*STRN*) gene and anaplastic lymphoma kinase (*ALK*) gene were found in 9% of poorly differentiated thyroid cancers, 4% of anaplastic thyroid cancers, and 1.2% of well-differentiated papillary thyroid cancer (Kelly, et al. 2014). Whereas in the past testing for rare fusions and other mutational events was difficult to apply to clinical FNA samples due to high cost and need for large amount of DNA and RNA for testing using individual gene assays, at the present time the availability of targeted NGS offers a convenient and cost-effective way for detection of multiple point mutations and gene rearrangements in clinical FNA samples.

In addition to these markers, another recently discovered molecular marker is mutation of the telomerase reverse transcriptase (*TERT*) promoter (Landa, et al. 2013; Liu, et al. 2013b; Melo, et al. 2014). *TERT* mutations have not been found in benign thyroid nodules, and therefore are expected to be useful diagnostically, but are also likely to play an important role in disease prognostication, and will be discussed in the next section. In our preliminary validation study, the use of an extended panel of mutational markers that includes point mutations and gene fusions in >60 genes results in a negative predictive value of >95% for thyroid nodules with AUS/FLUS and FN/SFN cytology that were found to be negative for these mutations (Nikiforov et al. Manuscript in preparation).

Molecular Markers for Cancer Prognostication

The utility of molecular markers in FNA biopsies may extend beyond diagnostic information to have a role in preoperatively identifying subsets of tumors with more aggressive biological behavior. Such patients may potentially benefit from more extensive initial surgery to include central compartment lymph node dissection to prevent tumor recurrence. Among prognostic markers, one of the best studied is the *BRAF* V600E mutation. The presence of the *BRAF* V600E mutation in papillary thyroid cancer was found to be associated with poor prognostic factors such as extrathyroidal invasion, lymph node metastases, and recurrence [reviewed in (Xing 2007)]. In thyroid FNA, preoperative testing for *BRAF* V600E mutation was found to be useful in predicting disease persistence and recurrence (Xing, et al. 2009). However, not all studies have shown such associations (Kim, et al. 2005; Liu, et al. 2005; Ito, et al. 2009). A recent meta-analysis of 14 studies including a total of 2,470 patients demonstrated that *BRAF* V600E mutation was significantly associated with tumor recurrence or persistent disease, which was found in 25% of *BRAF* V600E-positive tumors vs. 13% of *BRAF* mutation negative tumors (Tufano, et al. 2012). In addition, a large, multicenter study of 1849 patients found the presence of the *BRAF* V600E mutation to be significantly associated with increased mortality from papillary thyroid

cancer (Xing, et al. 2013). The overall mortality was 5% in patients with *BRAF* V600E mutation and 1% in *BRAF* mutation negative patients. These data suggest that overall *BRAF* V600E-positive tumors have a higher chance to have more aggressive disease features at presentation and as a group to have increased risk of recurrence and overall mortality. However, it is important to note that the majority of patients with *BRAF* V600E mutation do not have recurrent disease and overall survival remains very high in both groups of patients (Fig. 3). This suggests that *BRAF* V600E taken in isolation is a relatively sensitive, but not specific marker of tumor recurrence and tumor-related mortality.

Recently, additional and more specific markers of more aggressive tumor behavior have emerged. One of such markers is the presence of multiple driver mutations in thyroid cancer. Co-existing mutations in the early driver genes such as *BRAF* or *RAS* with mutations in *PIK3CA*, *AKT1*, or *TP53* in the same tumor has been shown to occur in poorly differentiated and anaplastic tumors (Garcia-Rostan et al. 2005; Hou et al. 2007a; Liu, et al. 2008). More recently, an NGS-based mutational analysis revealed that ~4% of well differentiated papillary cancers have more than one mutation, and these tumors are distinctively aggressive and typically present with distant metastases (Nikiforova et al. 2013).

TP53 mutations are known to occur with high frequency in poorly differentiated thyroid cancer (25%) and anaplastic thyroid cancer (70-80%), and is a well characterized genetic event governing thyroid tumor dedifferentiation (Donghi, et al. 1993; Fagin, et al. 1993). However, *TP53* mutation has also been found in some well-differentiated cancers such as papillary thyroid carcinoma and oncocytic follicular carcinoma (Nikiforova et al. 2013). It is likely that well differentiated cancers carrying *TP53* mutation have potential for tumor dedifferentiation and more aggressive clinical course, which should be addressed in further studies.

Another very promising prognostic molecular marker is mutation of the telomerase reverse transcriptase (*TERT*) promoter. Telomerase is a reverse transcriptase that utilizes an RNA template to add telomeric repeats to the ends of chromosomes. Telomerase is not expressed in most normal tissues, but is frequently activated in tumor cells (Shay and Bacchetti 1997). Maintenance of telomere length, either through telomerase activation or a recombination-based mechanism known as alternative lengthening of telomeres (ALT), is required for immortalization of cancer cells. Recently two mutations in the promoter of *TERT* (chr5:1295228C>T, termed C228T, and chr5:1295250C>T, termed C250T) were discovered in melanoma and were found to result in increased transcriptional activity of the promoter (Horn, et al. 2013; Huang, et al. 2013). The C228T and C250T *TERT* promoter mutations were also detected in follicular cell-derived thyroid cancers but were absent in benign lesions and in medullary thyroid cancers (Landa et al. 2013; Liu et al. 2013a; Liu et al. 2013b; Melo et al. 2014). The C228T and C250T mutations have a significantly higher prevalence in aggressive thyroid tumors including widely invasive oncocytic carcinoma and anaplastic thyroid carcinoma (Landa et al. 2013; Liu et al. 2013a; Liu et al. 2013b; Melo et al. 2014). Interestingly, *TERT* mutations in some studies were found to be more common in tumors with *BRAF* V600E mutation, which may suggest a possible synergistic interplay between MAPK pathway activation and telomerase activation to promote aggressive tumor behavior (Landa et al. 2013; Liu et al. 2013b). In a recent large study of 469 patients with a

mean follow-up of 8 years, *TERT* promoter mutations were found to be an independent risk factor for persistent disease, distant metastases, and disease-specific mortality for well differentiated thyroid cancer (Melo et al. 2014).

Summary and future directions

Over the last several years, significant progress has been made in understanding the genetic mechanisms of thyroid cancer and in the development of molecular tests for cancer diagnosis in thyroid nodules. Work from multiple research labs as well as genomic sequencing data from papillary thyroid carcinomas from The Cancer Genome Atlas (TCGA) has led to the identification of mutations and other driver genetic alterations in greater than 90% of thyroid cancers, making it one of the best characterized human cancers from a genetic standpoint. Moreover, next generation sequencing (NGS) technology offers the reliable detection of most of these genetic alterations in limited cell samples obtained by FNA biopsy, offering significant improvement in the accuracy of cancer detection in thyroid nodules as compared to currently available clinical tests. As a result, it is likely that in the near future NGS-based molecular tests will be able to predict the risk of cancer in thyroid nodules with very high accuracy, which will eliminate the uncertainty of indeterminate FNA cytology. Furthermore, as the cost of next generation sequencing continues to plummet and analytical tools become more efficient, the cost of molecular testing will decrease, making routine use of NGS-based molecular tests even more feasible and cost-effective for management of patients with cytologically indeterminate thyroid nodules.

Moreover, molecular markers are expected to have a significant impact on cancer prognostication. While *BRAF* V600E mutation can be considered a relatively sensitive prognostic marker for papillary cancer, it is not specific and can not be used in isolation for tumor prognostication. Recent results obtained by broad tumor genotyping show that several specific molecular signatures (such as the presence of several driver mutations, *TP53* mutations, or *TERT* promoter mutations) are found in a small fraction of well differentiated papillary and follicular cancers and are associated with more aggressive tumor behavior. It is expected that these molecular signatures will be confirmed and perhaps further improved in additional studies and will offer more specific detection of well differentiated thyroid cancers that have higher risk for tumor recurrence and cancer-related mortality. Future studies will be needed to define the optimal surgical and post-surgical management of patients based on these molecular signatures. With these advances, determination of a personalized cancer genome will become feasible in thyroid nodules in the near future, and will offer truly individualized medicine for patients with thyroid nodules and cancer.

Acknowledgments

This was supported in part by the NIH grant R01 CA88041 to Y.E.N.

References

- American Cancer Society. Cancer Facts & Figures 2012. Atlanta: American Cancer Society; 2012.
- Adeniran AJ, Zhu Z, Gandhi M, Steward DL, Fidler JP, Giordano TJ, Biddinger PW, Nikiforov YE. Correlation between genetic alterations and microscopic features, clinical manifestations, and

prognostic characteristics of thyroid papillary carcinomas. *Am J Surg Pathol*. 2006; 30:216–222. [PubMed: 16434896]

Agrawal N, Jiao Y, Sausen M, Leary R, Bettegowda C, Roberts NJ, Bhan S, Ho AS, Khan Z, Bishop J, et al. Exomic sequencing of medullary thyroid cancer reveals dominant and mutually exclusive oncogenic mutations in RET and RAS. *J Clin Endocrinol Metab*. 2013; 98:E364–369. [PubMed: 23264394]

Albores-Saavedra J, Henson DE, Glazer E, Schwartz AM. Changing patterns in the incidence and survival of thyroid cancer with follicular phenotype—papillary, follicular, and anaplastic: a morphological and epidemiological study. *Endocr Pathol*. 2007; 18:1–7. [PubMed: 17652794]

Alexander EK, Kennedy GC, Baloch ZW, Cibas ES, Chudova D, Diggans J, Friedman L, Kloos RT, LiVolsi VA, Mandel SJ, et al. Preoperative diagnosis of benign thyroid nodules with indeterminate cytology. *N Engl J Med*. 2012; 367:705–715. [PubMed: 22731672]

Alexander EK, Schorr M, Kloppner J, Kim C, Sipos J, Nabhan F, Parker C, Steward DL, Mandel SJ, Haugen BR. Multicenter clinical experience with the Afirma gene expression classifier. *J Clin Endocrinol Metab*. 2014; 99:119–125. [PubMed: 24152684]

Ali, SZ.; Cibas, ES. *The Bethesda System for Reporting Thyroid Cytopathology*. New York: Springer; 2010.

Baloch ZW, Fleisher S, LiVolsi VA, Gupta PK. Diagnosis of “follicular neoplasm”: a gray zone in thyroid fine-needle aspiration cytology. *Diagn Cytopathol*. 2002; 26:41–44. [PubMed: 11782086]

Baloch ZW, LiVolsi VA, Asa SL, Rosai J, Merino MJ, Randolph G, Vielh P, DeMay RM, Sidawy MK, Frable WJ. Diagnostic terminology and morphologic criteria for cytologic diagnosis of thyroid lesions: a synopsis of the National Cancer Institute thyroid Fine-Needle Aspiration State of the Science Conference. *Diagn Cytopathol*. 2008; 36:425–437. [PubMed: 18478609]

Bartolazzi A, Orlandi F, Saggiorato E, Volante M, Arecco F, Rossetto R, Palestini N, Ghigo E, Papotti M, Bussolati G, et al. Galectin-3-expression analysis in the surgical selection of follicular thyroid nodules with indeterminate fine-needle aspiration cytology: a prospective multicentre study. *Lancet Oncol*. 2008; 9:543–549. [PubMed: 18495537]

Bignell GR, Canzian F, Shayeghi M, Stark M, Shugart YY, Biggs P, Mangion J, Hamoudi R, Rosenblatt J, Buu P, et al. Familial nontoxic multinodular thyroid goiter locus maps to chromosome 14q but does not account for familial nonmedullary thyroid cancer. *Am J Hum Genet*. 1997; 61:1123–1130. [PubMed: 9345104]

Bongarzone I, Vigneri P, Mariani L, Collini P, Pilotti S, Pierotti MA. RET/NTRK1 rearrangements in thyroid gland tumors of the papillary carcinoma family: correlation with clinicopathological features. *Clin Cancer Res*. 1998; 4:223–228. [PubMed: 9516975]

Brito JP, Yarur AJ, Prokop LJ, McIver B, Murad MH, Montori V. Prevalence of Thyroid Cancer in Multinodular Goiter vs. Single Nodule: A Systematic Review and Meta-analysis. *Thyroid*. 2012

Burgess JR, Tucker P. Incidence trends for papillary thyroid carcinoma and their correlation with thyroid surgery and thyroid fine-needle aspirate cytology. *Thyroid*. 2006; 16:47–53. [PubMed: 16487013]

Burns JS, Blaydes JP, Wright PA, Lemoine L, Bond JA, Williams ED, Wynford-Thomas D. Stepwise transformation of primary thyroid epithelial cells by a mutant Ha-ras oncogene: an in vitro model of tumor progression. *Mol Carcinog*. 1992; 6:129–139. [PubMed: 1388684]

Buryk MA, Monaco SE, Witchel SF, Mehta DK, Gurtunca N, Nikiforov YE, Simons JP. Preoperative cytology with molecular analysis to help guide surgery for pediatric thyroid nodules. *Int J Pediatr Otorhinolaryngol*. 2013; 77:1697–1700. [PubMed: 23993207]

Cantara S, Capezzone M, Marchisotta S, Capuano S, Busonero G, Toti P, Di Santo A, Caruso G, Carli AF, Brilli L, et al. Impact of proto-oncogene mutation detection in cytological specimens from thyroid nodules improves the diagnostic accuracy of cytology. *J Clin Endocrinol Metab*. 2010; 95:1365–1369. [PubMed: 20130073]

Canzian F, Amati P, Harach HR, Kraimps JL, Lesueur F, Barbier J, Levillain P, Romeo G, Bonneau D. A gene predisposing to familial thyroid tumors with cell oxyphilia maps to chromosome 19p13.2. *Am J Hum Genet*. 1998; 63:1743–1748. [PubMed: 9837827]

Cavaco BM, Batista PF, Martins C, Banito A, do Rosario F, Limbert E, Sobrinho LG, Leite V. Familial non-medullary thyroid carcinoma (FNMTc): analysis of fPTC/PRN, NMTC1, MNG1

- and TCO susceptibility loci and identification of somatic BRAF and RAS mutations. *Endocr Relat Cancer*. 2008; 15:207–215. [PubMed: 18310288]
- Charkes ND. On the prevalence of familial nonmedullary thyroid cancer in multiply affected kindreds. *Thyroid*. 2006; 16:181–186. [PubMed: 16613533]
- Chiosea S, Nikiforova M, Zuo H, Ogilvie J, Gandhi M, Seethala RR, Ohori NP, Nikiforov Y. A novel complex BRAF mutation detected in a solid variant of papillary thyroid carcinoma. *Endocr Pathol*. 2009; 20:122–126. [PubMed: 19370421]
- Chudova D, Wilde JJ, Wang ET, Wang H, Rabbee N, Egidio CM, Reynolds J, Tom E, Pagan M, Rigl C Ted, et al. Molecular Classification of Thyroid Nodules Using High-Dimensionality Genomic Data. *J Clin Endocrinol Metab*. 2010
- Ciampi R, Knauf JA, Kerler R, Gandhi M, Zhu Z, Nikiforova MN, Rabes HM, Fagin JA, Nikiforov YE. Oncogenic AKAP9-BRAF fusion is a novel mechanism of MAPK pathway activation in thyroid cancer. *J Clin Invest*. 2005; 115:94–101. [PubMed: 15630448]
- Ciampi R, Nikiforov YE. Alterations of the BRAF gene in thyroid tumors. *Endocr Pathol*. 2005; 16:163–172. [PubMed: 16299399]
- Cohen Y, Xing M, Mambo E, Guo Z, Wu G, Trink B, Beller U, Westra WH, Ladenson PW, Sidransky D. BRAF mutation in papillary thyroid carcinoma. *J Natl Cancer Inst*. 2003; 95:625–627. [PubMed: 12697856]
- Cooper DS, Doherty GM, Haugen BR, Kloos RT, Lee SL, Mandel SJ, Mazzaferri EL, McIver B, Pacini F, Schlumberger M, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2009; 19:1167–1214. [PubMed: 19860577]
- Dahia PL, Marsh DJ, Zheng Z, Zedenius J, Komminoth P, Frisk T, Wallin G, Parsons R, Longy M, Larsson C, et al. Somatic deletions and mutations in the Cowden disease gene, PTEN, in sporadic thyroid tumors. *Cancer Res*. 1997; 57:4710–4713. [PubMed: 9354427]
- Davies L, Welch HG. Increasing incidence of thyroid cancer in the United States, 1973–2002. *Jama*. 2006; 295:2164–2167. [PubMed: 16684987]
- de Groot JW, Links TP, Plukker JT, Lips CJ, Hofstra RM. RET as a diagnostic and therapeutic target in sporadic and hereditary endocrine tumors. *Endocr Rev*. 2006; 27:535–560. [PubMed: 16849421]
- Dobashi Y, Sugimura H, Sakamoto A, Mernyei M, Mori M, Oyama T, Machinami R. Stepwise participation of p53 gene mutation during dedifferentiation of human thyroid carcinomas. *Diagn Mol Pathol*. 1994; 3:9–14. [PubMed: 8162258]
- Donghi R, Longoni A, Pilotti S, Michieli P, Della Porta G, Pierotti MA. Gene p53 mutations are restricted to poorly differentiated and undifferentiated carcinomas of the thyroid gland. *J Clin Invest*. 1993; 91:1753–1760. [PubMed: 8473515]
- Duick DS, Klopper JP, Diggans JC, Friedman L, Kennedy GC, Lanman RB, McIver B. The impact of benign gene expression classifier test results on the endocrinologist-patient decision to operate on patients with thyroid nodules with indeterminate fine-needle aspiration cytopathology. *Thyroid*. 2012; 22:996–1001. [PubMed: 22873825]
- Dwight T, Thoppe SR, Foukakis T, Lui WO, Wallin G, Hoog A, Frisk T, Larsson C, Zedenius J. Involvement of the PAX8/peroxisome proliferator-activated receptor gamma rearrangement in follicular thyroid tumors. *J Clin Endocrinol Metab*. 2003; 88:4440–4445. [PubMed: 12970322]
- Fagin JA. Minireview: branded from the start-distinct oncogenic initiating events may determine tumor fate in the thyroid. *Mol Endocrinol*. 2002; 16:903–911. [PubMed: 11981026]
- Fagin JA, Matsuo K, Karmakar A, Chen DL, Tang SH, Koeffler HP. High prevalence of mutations of the p53 gene in poorly differentiated human thyroid carcinomas. *J Clin Invest*. 1993; 91:179–184. [PubMed: 8423216]
- Fenton CL, Lukes Y, Nicholson D, Dinauer CA, Francis GL, Tuttle RM. The ret/PTC mutations are common in sporadic papillary thyroid carcinoma of children and young adults. *J Clin Endocrinol Metab*. 2000; 85:1170–1175. [PubMed: 10720057]
- Frattini M, Ferrario C, Bressan P, Balestra D, De Cecco L, Mondellini P, Bongarzone I, Collini P, Gariboldi M, Pilotti S, et al. Alternative mutations of BRAF, RET and NTRK1 are associated with

- similar but distinct gene expression patterns in papillary thyroid cancer. *Oncogene*. 2004; 23:7436–7440. [PubMed: 15273715]
- French CA, Alexander EK, Cibas ES, Nose V, Laguette J, Faquin W, Garber J, Moore F Jr, Fletcher JA, Larsen PR, et al. Genetic and biological subgroups of low-stage follicular thyroid cancer. *Am J Pathol*. 2003; 162:1053–1060. [PubMed: 12651598]
- Frich L, Glatte E, Akslen LA. Familial occurrence of nonmedullary thyroid cancer: a population-based study of 5673 first-degree relatives of thyroid cancer patients from Norway. *Cancer Epidemiol Biomarkers Prev*. 2001; 10:113–117. [PubMed: 11219767]
- Garcia-Jimenez C, Santisteban P. TSH signalling and cancer. *Arq Bras Endocrinol Metabol*. 2007; 51:654–671. [PubMed: 17891229]
- Garcia-Rostan G, Camp RL, Herrero A, Carcangiu ML, Rimm DL, Tallini G. Beta-catenin dysregulation in thyroid neoplasms: down-regulation, aberrant nuclear expression, and CTNNB1 exon 3 mutations are markers for aggressive tumor phenotypes and poor prognosis. *Am J Pathol*. 2001; 158:987–996. [PubMed: 11238046]
- Garcia-Rostan G, Costa AM, Pereira-Castro I, Salvatore G, Hernandez R, Hermsem MJ, Herrero A, Fusco A, Cameselle-Teijeiro J, Santoro M. Mutation of the PIK3CA gene in anaplastic thyroid cancer. *Cancer Res*. 2005; 65:10199–10207. [PubMed: 16288007]
- Gharib H, Papini E, Paschke R, Duick DS, Valcavi R, Hegedus L, Vitti P. American Association of Clinical Endocrinologists, Associazione Medici Endocrinologi, and European Thyroid Association medical guidelines for clinical practice for the diagnosis and management of thyroid nodules: executive summary of recommendations. *J Endocrinol Invest*. 2010; 33:51–56. [PubMed: 20543551]
- Grieco M, Santoro M, Berlingieri MT, Melillo RM, Donghi R, Bongarzone I, Pierotti MA, Della Porta G, Fusco A, Vecchio G. PTC is a novel rearranged form of the ret proto-oncogene and is frequently detected in vivo in human thyroid papillary carcinomas. *Cell*. 1990; 60:557–563. [PubMed: 2406025]
- Gustafson S, Zbuk KM, Scacheri C, Eng C. Cowden syndrome. *Semin Oncol*. 2007; 34:428–434. [PubMed: 17920899]
- Guth S, Theune U, Aberle J, Galach A, Bamberger CM. Very high prevalence of thyroid nodules detected by high frequency (13 MHz) ultrasound examination. *Eur J Clin Invest*. 2009; 39:699–706. [PubMed: 19601965]
- Hansford JR, Mulligan LM. Multiple endocrine neoplasia type 2 and RET: from neoplasia to neurogenesis. *J Med Genet*. 2000; 37:817–827. [PubMed: 11073534]
- Harrell RM, Bimston DN. Surgical Utility of Afirma: Effects of High Cancer Prevalence and Oncocytic Cell Types in Patients with Indeterminate Thyroid Cytology. *Endocr Pract*. 2013;1–16.
- He H, Nagy R, Liyanarachchi S, Jiao H, Li W, Suster S, Kere J, de la Chapelle A. A susceptibility locus for papillary thyroid carcinoma on chromosome 8q24. *Cancer Res*. 2009; 69:625–631. [PubMed: 19147577]
- Hemminki K, Eng C, Chen B. Familial risks for nonmedullary thyroid cancer. *J Clin Endocrinol Metab*. 2005; 90:5747–5753. [PubMed: 16030170]
- Horn S, Figl A, Rachakonda PS, Fischer C, Sucker A, Gast A, Kadel S, Moll I, Nagore E, Hemminki K, et al. TERT promoter mutations in familial and sporadic melanoma. *Science*. 2013; 339:959–961. [PubMed: 23348503]
- Hou P, Liu D, Shan Y, Hu S, Studeman K, Condouris S, Wang Y, Trink A, El-Naggar AK, Tallini G, et al. Genetic alterations and their relationship in the phosphatidylinositol 3-kinase/Akt pathway in thyroid cancer. *Clin Cancer Res*. 2007a; 13:1161–1170. [PubMed: 17317825]
- Hou P, Liu D, Xing M. Functional characterization of the T1799-1801del and A1799-1816ins BRAF mutations in papillary thyroid cancer. *Cell Cycle*. 2007b; 6:377–379. [PubMed: 17297294]
- Howlander, N.; Noone, AM.; Krapcho, M.; Garshell, J.; Neyman, N.; Altekruse, SF.; Kosary, CL.; Yu, M.; Ruhl, J.; Tatalovich, Z., et al. SEER Cancer Statistics Review, 1975-2010. National Cancer Institute; Bethesda, MD: 2013. http://seer.cancer.gov/csr/1975_2010/, based on November 2012 SEER data submission, posted to the SEER web site, April 2013
- Huang FW, Hodis E, Xu MJ, Kryukov GV, Chin L, Garraway LA. Highly recurrent TERT promoter mutations in human melanoma. *Science*. 2013; 339:957–959. [PubMed: 23348506]

- Ito Y, Yoshida H, Maruo R, Morita S, Takano T, Hirokawa M, Yabuta T, Fukushima M, Inoue H, Tomoda C, et al. BRAF mutation in papillary thyroid carcinoma in a Japanese population: its lack of correlation with high-risk clinicopathological features and disease-free survival of patients. *Endocr J*. 2009; 56:89–97. [PubMed: 18840924]
- Jung CK, Little MP, Lubin JH, Brenner AV, Wells SA Jr, Sigurdson AJ, Nikiforov YE. The increase in thyroid cancer incidence during the last four decades is accompanied by a high frequency of BRAF mutations and a sharp increase in RAS mutations. *J Clin Endocrinol Metab*. 2013 jc20132503.
- Kelly LM, Barila G, Liu P, Evdokimova VN, Trivedi S, Panebianco F, Gandhi M, Carty SE, Hodak SP, Luo J, et al. Identification of the transforming *STRN-ALK* fusion as a potential therapeutic target in the aggressive forms of thyroid cancer. *PNAS*. 2014 Published online before print on February 3, 2014. doi: 2010.1073/pnas.1321937111.
- Keutgen XM, Filicori F, Crowley MJ, Wang Y, Scognamiglio T, Hoda R, Buitrago D, Cooper D, Zeiger MA, Zarnegar R, et al. A panel of four miRNAs accurately differentiates malignant from benign indeterminate thyroid lesions on fine needle aspiration. *Clin Cancer Res*. 2012; 18:2032–2038. [PubMed: 22351693]
- Kim TY, Kim WB, Song JY, Rhee YS, Gong G, Cho YM, Kim SY, Kim SC, Hong SJ, Shong YK. The BRAF mutation is not associated with poor prognostic factors in Korean patients with conventional papillary thyroid microcarcinoma. *Clin Endocrinol (Oxf)*. 2005; 63:588–593. [PubMed: 16268813]
- Kimura ET, Nikiforova MN, Zhu Z, Knauf JA, Nikiforov YE, Fagin JA. High prevalence of BRAF mutations in thyroid cancer: genetic evidence for constitutive activation of the RET/PTC-RAS-BRAF signaling pathway in papillary thyroid carcinoma. *Cancer Res*. 2003; 63:1454–1457. [PubMed: 12670889]
- Kloos RT, Eng C, Evans DB, Francis GL, Gagel RF, Gharib H, Moley JF, Pacini F, Ringel MD, Schlumberger M, et al. Medullary thyroid cancer: management guidelines of the American Thyroid Association. *Thyroid*. 2009; 19:565–612. [PubMed: 19469690]
- Landa I, Ganly I, Chan TA, Mitsutake N, Matsuse M, Ibrahimasic T, Ghossein RA, Fagin JA. Frequent somatic TERT promoter mutations in thyroid cancer: higher prevalence in advanced forms of the disease. *J Clin Endocrinol Metab*. 2013; 98:E1562–1566. [PubMed: 23833040]
- Leeman-Neill RJ, Brenner AV, Little MP, Bogdanova TI, Hatch M, Zurnadzy LY, Mabuchi K, Tronko MD, Nikiforov YE. RET/PTC and PAX8/PPARGgamma chromosomal rearrangements in post-Chernobyl thyroid cancer and their association with iodine-131 radiation dose and other characteristics. *Cancer*. 2013a; 119:1792–1799. [PubMed: 23436219]
- Leeman-Neill RJ, Kelly LM, Liu P, Brenner AV, Little MP, Bogdanova TI, Evdokimova VN, Hatch M, Zurnadzy LY, Nikiforova MN, et al. ETV6-NTRK3 is a common chromosomal rearrangement in radiation-associated thyroid cancer. *Cancer*. 2013b
- Lemoine NR, Mayall ES, Wyllie FS, Williams ED, Goyns M, Stringer B, Wynford-Thomas D. High frequency of ras oncogene activation in all stages of human thyroid tumorigenesis. *Oncogene*. 1989; 4:159–164. [PubMed: 2648253]
- Liu RT, Chen YJ, Chou FF, Li CL, Wu WL, Tsai PC, Huang CC, Cheng JT. No correlation between BRAFV600E mutation and clinicopathological features of papillary thyroid carcinomas in Taiwan. *Clin Endocrinol (Oxf)*. 2005; 63:461–466. [PubMed: 16181240]
- Liu T, Wang N, Cao J, Sofiadis A, Dinets A, Zedenius J, Larsson C, Xu D. The age- and shorter telomere-dependent TERT promoter mutation in follicular thyroid cell-derived carcinomas. *Oncogene*. 2013a
- Liu X, Bishop J, Shan Y, Pai S, Liu D, Murugan AK, Sun H, El-Naggar AK, Xing M. Highly prevalent TERT promoter mutations in aggressive thyroid cancers. *Endocr Relat Cancer*. 2013b; 20:603–610. [PubMed: 23766237]
- Liu Z, Hou P, Ji M, Guan H, Studeman K, Jensen K, Vasko V, El-Naggar AK, Xing M. Highly prevalent genetic alterations in receptor tyrosine kinases and phosphatidylinositol 3-kinase/akt and mitogen-activated protein kinase pathways in anaplastic and follicular thyroid cancers. *J Clin Endocrinol Metab*. 2008; 93:3106–3116. [PubMed: 18492751]
- Malchoff CD, Malchoff DM. Familial nonmedullary thyroid carcinoma. *Cancer Control*. 2006; 13:106–110. [PubMed: 16735984]

- Malchoff CD, Sarfarazi M, Tendler B, Forouhar F, Whalen G, Joshi V, Arnold A, Malchoff DM. Papillary thyroid carcinoma associated with papillary renal neoplasia: genetic linkage analysis of a distinct heritable tumor syndrome. *J Clin Endocrinol Metab.* 2000; 85:1758–1764. [PubMed: 10843148]
- Marques AR, Espadinha C, Catarino AL, Moniz S, Pereira T, Sobrinho LG, Leite V. Expression of PAX8-PPAR gamma 1 rearrangements in both follicular thyroid carcinomas and adenomas. *J Clin Endocrinol Metab.* 2002; 87:3947–3952. [PubMed: 12161538]
- Mazeh H, Sippel RS. Familial nonmedullary thyroid carcinoma. *Thyroid.* 2013; 23:1049–1056. [PubMed: 23734600]
- Mazzaferri EL. Thyroid cancer in thyroid nodules: finding a needle in the haystack. *Am J Med.* 1992; 93:359–362. [PubMed: 1415298]
- Mazzaferri EL. Management of a solitary thyroid nodule. *N Engl J Med.* 1993; 328:553–559. [PubMed: 8426623]
- Mazzaferri EL, Jhiang SM. Long-term impact of initial surgical and medical therapy on papillary and follicular thyroid cancer. *Am J Med.* 1994; 97:418–428. [PubMed: 7977430]
- McKay JD, Lesueur F, Jonard L, Pastore A, Williamson J, Hoffman L, Burgess J, Duffield A, Papotti M, Stark M, et al. Localization of a susceptibility gene for familial nonmedullary thyroid carcinoma to chromosome 2q21. *Am J Hum Genet.* 2001; 69:440–446. [PubMed: 11438887]
- Melo M, Rocha AG, Vinagre J, Batista R, Peixoto J, Tavares C, Celestino R, Almeida A, Salgado C, Eloy C, et al. TERT promoter mutations are a major indicator of poor outcome in differentiated thyroid carcinomas. *J Clin Endocrinol Metab.* 2014 jc20133734.
- Monaco SE, Pantanowitz L, Khalbuss WE, Benkovich VA, Ozolek J, Nikiforova MN, Simons JP, Nikiforov YE. Cytomorphological and molecular genetic findings in pediatric thyroid fine-needle aspiration. *Cancer Cytopathol.* 2012; 120:342–350. [PubMed: 22605559]
- Moses W, Weng J, Kebebew E. Prevalence, clinicopathologic features, and somatic genetic mutation profile in familial versus sporadic nonmedullary thyroid cancer. *Thyroid.* 2011; 21:367–371. [PubMed: 21190444]
- Motoi N, Sakamoto A, Yamochi T, Horiuchi H, Motoi T, Machinami R. Role of ras mutation in the progression of thyroid carcinoma of follicular epithelial origin. *Pathol Res Pract.* 2000; 196:1–7. [PubMed: 10674266]
- Mulligan LM, Marsh DJ, Robinson BG, Schuffenecker I, Zedenius J, Lips CJ, Gagel RF, Takai SI, Noll WW, Fink M, et al. Genotype-phenotype correlation in multiple endocrine neoplasia type 2: report of the International RET Mutation Consortium. *J Intern Med.* 1995; 238:343–346. [PubMed: 7595170]
- Musholt TJ, Musholt PB, Khaladj N, Schulz D, Scheumann GF, Klempnauer J. Prognostic significance of RET and NTRK1 rearrangements in sporadic papillary thyroid carcinoma. *Surgery.* 2000; 128:984–993. [PubMed: 11114633]
- Namba H, Rubin SA, Fagin JA. Point mutations of ras oncogenes are an early event in thyroid tumorigenesis. *Mol Endocrinol.* 1990; 4:1474–1479. [PubMed: 2283998]
- Nikiforov YE. RET/PTC Rearrangement—a link between Hashimoto's thyroiditis and thyroid cancer... or not. *J Clin Endocrinol Metab.* 2006; 91:2040–2042. [PubMed: 16757533]
- Nikiforov, YE.; Otori, NP. Follicular carcinoma. In: Nikiforov, YE.; Biddinger, PW.; Thompson, LDR., editors. *Diagnostic pathology and molecular genetics of the thyroid.* 2nd. Philadelphia: Lippincott Williams & Wilkins; 2012. p. 152-182.
- Nikiforov YE, Otori NP, Hodak SP, Carty SE, LeBeau SO, Ferris RL, Yip L, Seethala RR, Tublin ME, Stang MT, et al. Impact of mutational testing on the diagnosis and management of patients with cytologically indeterminate thyroid nodules: a prospective analysis of 1056 FNA samples. *J Clin Endocrinol Metab.* 2011; 96:3390–3397. [PubMed: 21880806]
- Nikiforov YE, Rowland JM, Bove KE, Monforte-Munoz H, Fagin JA. Distinct pattern of ret oncogene rearrangements in morphological variants of radiation-induced and sporadic thyroid papillary carcinomas in children. *Cancer Res.* 1997; 57:1690–1694. [PubMed: 9135009]
- Nikiforov YE, Steward DL, Robinson-Smith TM, Haugen BR, Klopper JP, Zhu Z, Fagin JA, Falciglia M, Weber K, Nikiforova MN. Molecular testing for mutations in improving the fine-needle

- aspiration diagnosis of thyroid nodules. *J Clin Endocrinol Metab.* 2009; 94:2092–2098. [PubMed: 19318445]
- Nikiforova MN, Biddinger PW, Caudill CM, Kroll TG, Nikiforov YE. PAX8-PPARgamma rearrangement in thyroid tumors: RT-PCR and immunohistochemical analyses. *Am J Surg Pathol.* 2002; 26:1016–1023. [PubMed: 12170088]
- Nikiforova MN, Lynch RA, Biddinger PW, Alexander EK, Dorn GW 2nd, Tallini G, Kroll TG, Nikiforov YE. RAS point mutations and PAX8-PPAR gamma rearrangement in thyroid tumors: evidence for distinct molecular pathways in thyroid follicular carcinoma. *J Clin Endocrinol Metab.* 2003; 88:2318–2326. [PubMed: 12727991]
- Nikiforova MN, Wald AI, Roy S, Durso MB, Nikiforov YE. Targeted next-generation sequencing panel (ThyroSeq) for detection of mutations in thyroid cancer. *J Clin Endocrinol Metab.* 2013; 98:E1852–1860. [PubMed: 23979959]
- Nishihara E, Amino N, Maekawa K, Yoshida H, Ito M, Kubota S, Fukata S, Miyauchi A. Prevalence of TSH receptor and Gsalpha mutations in 45 autonomously functioning thyroid nodules in Japan. *Endocr J.* 2009; 56:791–798. [PubMed: 19550078]
- Othori NP, Schoedel KE. Variability in the atypia of undetermined significance/follicular lesion of undetermined significance diagnosis in the Bethesda system for reporting thyroid cytopathology: sources and recommendations. *Acta Cytol.* 2011; 55:492–498. [PubMed: 22156456]
- Rabes HM, Demidchik EP, Sidorow JD, Lengfelder E, Beimfohr C, Hoelzel D, Klugbauer S. Pattern of radiation-induced RET and NTRK1 rearrangements in 191 post-chernobyl papillary thyroid carcinomas: biological, phenotypic, and clinical implications. *Clin Cancer Res.* 2000; 6:1093–1103. [PubMed: 10741739]
- Ricarte-Filho JC, Ryder M, Chitale DA, Rivera M, Heguy A, Ladanyi M, Janakiraman M, Solit D, Knauf JA, Tuttle RM, et al. Mutational profile of advanced primary and metastatic radioactive iodine-refractory thyroid cancers reveals distinct pathogenetic roles for BRAF, PIK3CA, and AKT1. *Cancer Res.* 2009; 69:4885–4893. [PubMed: 19487299]
- Roy S, Durso MB, Wald A, Nikiforov YE, Nikiforova MN. SeqReporter: automating next-generation sequencing result interpretation and reporting workflow in a clinical laboratory. *J Mol Diagn.* 2014; 16:11–22. [PubMed: 24220144]
- Santoro M, Dathan NA, Berlingieri MT, Bongarzone I, Paulin C, Grieco M, Pierotti MA, Vecchio G, Fusco A. Molecular characterization of RET/PTC3; a novel rearranged version of the RET proto-oncogene in a human thyroid papillary carcinoma. *Oncogene.* 1994; 9:509–516. [PubMed: 8290261]
- Shay JW, Bacchetti S. A survey of telomerase activity in human cancer. *Eur J Cancer.* 1997; 33:787–791. [PubMed: 9282118]
- Soares P, Trovisco V, Rocha AS, Lima J, Castro P, Preto A, Maximo V, Botelho T, Seruca R, Sobrinho-Simoes M. BRAF mutations and RET/PTC rearrangements are alternative events in the etiopathogenesis of PTC. *Oncogene.* 2003; 22:4578–4580. [PubMed: 12881714]
- Suarez HG, du Villard JA, Severino M, Caillou B, Schlumberger M, Tubiana M, Parmentier C, Monier R. Presence of mutations in all three ras genes in human thyroid tumors. *Oncogene.* 1990; 5:565–570. [PubMed: 2183158]
- Tanaka K, Sonoo H, Saito W, Ohta Y, Shimo T, Sohda M, Yamamoto Y, Kurebayashi J. Analysis of clinical outcome of patients with poorly differentiated thyroid carcinoma. *ISRN Endocrinol.* 2011; 2011:308029. [PubMed: 22363874]
- Tomaz RA, Sousa I, Silva JG, Santos C, Teixeira MR, Leite V, Cavaco BM. FOXE1 polymorphisms are associated with familial and sporadic nonmedullary thyroid cancer susceptibility. *Clin Endocrinol (Oxf).* 2012; 77:926–933. [PubMed: 22882326]
- Tufano RP, Teixeira GV, Bishop J, Carson KA, Xing M. BRAF mutation in papillary thyroid cancer and its value in tailoring initial treatment: a systematic review and meta-analysis. *Medicine (Baltimore).* 2012; 91:274–286. [PubMed: 22932786]
- Volante M, Landolfi S, Chiusa L, Palestini N, Motta M, Codegone A, Torchio B, Papotti MG. Poorly differentiated carcinomas of the thyroid with trabecular, insular, and solid patterns: a clinicopathologic study of 183 patients. *Cancer.* 2004; 100:950–957. [PubMed: 14983490]

- Xing M. BRAF mutation in papillary thyroid cancer: pathogenic role, molecular bases, and clinical implications. *Endocr Rev.* 2007; 28:742–762. [PubMed: 17940185]
- Xing M, Alzahrani AS, Carson KA, Viola D, Elisei R, Bendlova B, Yip L, Mian C, Vianello F, Tuttle RM, et al. Association between BRAF V600E mutation and mortality in patients with papillary thyroid cancer. *Jama.* 2013; 309:1493–1501. [PubMed: 23571588]
- Xing M, Clark D, Guan H, Ji M, Dackiw A, Carson KA, Kim M, Tufaro A, Ladenson P, Zeiger M, et al. BRAF mutation testing of thyroid fine-needle aspiration biopsy specimens for preoperative risk stratification in papillary thyroid cancer. *J Clin Oncol.* 2009; 27:2977–2982. [PubMed: 19414674]
- Yip L, Wharry L, Armstrong M, Silbermann A, McCoy KL, Stang MT, Ohori NP, LeBeau SO, Coyne C, Nikiforova MN, et al. A Clinical Algorithm for Fine-Needle Aspiration Molecular Testing Effectively Guides the Appropriate Extent of Initial Thyroidectomy. *Ann Surg.* 2014 Published online before print on October 10, 2013. doi:10.1097/SLA.0000000000000215.
- Zhu Z, Ciampi R, Nikiforova MN, Gandhi M, Nikiforov YE. Prevalence of RET/PTC rearrangements in thyroid papillary carcinomas: effects of the detection methods and genetic heterogeneity. *J Clin Endocrinol Metab.* 2006; 91:3603–3610. [PubMed: 16772343]
- Zhu Z, Gandhi M, Nikiforova MN, Fischer AH, Nikiforov YE. Molecular profile and clinical-pathologic features of the follicular variant of papillary thyroid carcinoma. An unusually high prevalence of ras mutations. *Am J Clin Pathol.* 2003; 120:71–77. [PubMed: 12866375]

Figure 1.
Progress in identifying mutational markers in thyroid cancer.

Figure 2. Molecular pathways in thyroid cancer

The molecular pathogenesis of thyroid cancer involves dysregulation of the mitogen-activated protein kinase (MAPK), phosphatidylinositol-3 kinase (PI3K)/AKT, and the TSHR cAMP signaling pathways. The MAPK pathway is frequently activated in thyroid cancer through point mutations of the *BRAF* and *RAS* genes and *RET/PTC* and *TRK* rearrangements, and the PI3K pathway is frequently activated through point mutations of *PIK3CA* and mutation/deletion of *PTEN*.

Figure 3. Associations between *BRAF* V600E and tumor recurrence and mortality

Based on the results reported by Tufano et al. (Tufano et al. 2012) and Xing et al. (Xing et al. 2013).