Published in final edited form as:

J Biomol NMR. 2014 September ; 60(1): 23–27. doi:10.1007/s10858-014-9854-y.

Effective strategy to assign ^1H - ^{15}N heteronuclear correlation NMR signals from lysine side-chain NH_3^+ groups of proteins at low temperature

Alexandre Esadze, Levani Zandarashvili, and Junji Iwahara*

Department of Biochemistry and Molecular Biology, Sealy Center for Structural Biology and Molecular Biophysics, University of Texas Medical Branch, Galveston, TX 77555-1068

Abstract

Recent studies have shown that lysine side-chain NH_3^+ groups are excellent probes for NMR investigations of dynamics involving hydrogen bonds and ion pairs relevant to protein function. However, due to rapid hydrogen exchange, observation of ^1H - ^{15}N NMR cross peaks from lysine NH_3^+ groups often requires use of a relatively low temperature, which renders difficulty in resonance assignment. Here we present an effective strategy to assign ^1H and ^{15}N resonances of NH_3^+ groups at low temperatures. This strategy involves two new $^1\text{H}/^{13}\text{C}/^{15}\text{N}$ triple-resonance experiments for lysine side chains. Application to a protein-DNA complex is demonstrated.

Keywords

Proteins; side chains; Lysine; NH_3^+ groups; $^1\text{H}/^{13}\text{C}/^{15}\text{N}$ resonances

Hydrogen bonds and ion pairs involving protein side chains are of fundamental importance in protein functions such as molecular recognition and catalysis. Recently it was demonstrated that lysine (Lys) side-chain NH_3^+ groups are very useful probes for NMR studies of protein dynamics involving hydrogen bonds and ion pairs (Anderson et al. 2013; Esadze et al. 2011; Zandarashvili et al. 2013; Zandarashvili et al. 2011). Heteronuclear ^1H - ^{15}N cross peaks from Lys NH_3^+ groups are also useful for identifying molecular interfaces between protein and ligand (Blaum et al. 2010; Poon et al. 2006). Resonance assignment for Lys NH_3^+ groups typically requires not only standard experiments such as (H)C(CO)NH, H(CCO)NH, and HCCH-TOCSY, which correlate side-chain and backbone resonances, but also Lys-selective NMR experiments (Andre et al. 2007; Iwahara et al. 2007). This assignment process is not trivial because rapid hydrogen exchange renders difficulty in ^1H detection of NH_3^+ groups. To mitigate this problem, observation of ^1H signals from Lys NH_3^+ groups often requires the use of relatively low pH and temperature. However, the use of a low temperature can substantially reduce sensitivity of most triple-resonance experiments due to a longer molecular rotational correlation time, and thereby hamper resonance assignment.

*To whom correspondence should be addressed. j.iwahara@utmb.edu.

Resonance assignment for Lys side-chain NH_3^+ groups requires long-range correlation spectra because these amino groups are located at the fifth positions (i.e., ζ position) from $\text{C}\alpha$ atoms. Short-range correlations between NH_3^+ and $^1\text{H}\epsilon/^{13}\text{C}\epsilon$ resonances are typically insufficient for unambiguous assignment due to poor dispersion of their chemical shifts. While long-range correlation NMR methods such as the 3D (H)CCENH3 (Iwahara et al. 2007) may in principle allow for assignment of Lys NH_3^+ resonances, this approach can be impractical at low temperatures due to rapid ^{13}C transverse relaxation. In fact, it is well known that rapid ^{13}C transverse relaxation can diminish the efficiency of ^{13}C - ^{13}C Hartmann-Hahn cross polarization if a molecular rotational correlation time τ_r is relatively long (Fischer et al. 1996).

In this paper, we present two distinct approaches to assign Lys side-chain NH_3^+ groups at a low temperature. One of the approaches is to obtain long-range ^{15}N - ^{13}C correlations by taking advantage of very slow ^{15}N transverse relaxation of NH_3^+ groups. Due to bond rotations along the symmetry axis and cross correlation between three ^{15}N - ^1H dipole-dipole interactions, NMR relaxation of the in-phase single-quantum term N_x for NH_3^+ groups is extremely slow and occurs in a bi-exponential manner (Esadze et al. 2011; Zandarashvili et al. 2013). In our previous studies, initial transverse relaxation rates $R_{2,\text{ini}}$ for NH_3^+ groups were in the range between 0.9 and 4.5 s^{-1} for the systems with $\tau_r = 8.5$ and 10.8 ns (Anderson et al. 2013; Esadze et al. 2011); the average $R_{2,\text{ini}}$ for the Lys side-chain NH_3^+ groups that directly interact with DNA phosphate groups in the HoxD9 homeodomain-DNA complex was 1.7 s^{-1} (Anderson et al. 2013), which corresponds to a transverse relaxation time of ~ 580 ms. Therefore, even a time over one hundred milliseconds can readily be used for ^{15}N - ^{13}C coherence transfer via J -evolution between N_x and $2N_yC_z$ terms for a relatively small scalar coupling. This coherence transfer is substantially more efficient than coherence transfer via J -evolution from C_x to $2C_yN_z$, because the ^{15}N transverse relaxation of NH_3^+ groups is far slower than relevant ^{13}C transverse relaxation of CH_2 groups.

From this perspective, we have developed the 3D H3NCG experiment (Figure 1), which provides a long-range correlation between $^1\text{H}/^{15}\text{N}$ resonances of the NH_3^+ groups and $^{13}\text{C}\gamma$ resonance for Lys side chains. This 3D experiment relies on relatively small scalar coupling constants $^3J_{\text{NC}}$ between the $^{15}\text{N}\zeta$ and $^{13}\text{C}\gamma$ nuclei. Absolute values of the $^3J_{\text{NC}}$ constants are in a range of 1 – 3 Hz, and depend on Lys χ_4 torsion angles, which can undergo dynamic transitions between gauche and trans conformers (Huang and MacKerell 2013; Zandarashvili et al. 2011). Owing to the slow ^{15}N transverse relaxation of NH_3^+ groups during the coherence transfers between the N_x and $2N_yC_z$ terms, the magnetizations of interest survive a total of 200 ms for the evolution of the small $^3J_{\text{NC}}$ couplings between the $^{15}\text{N}\zeta$ and $^{13}\text{C}\gamma$ nuclei in the 3D H3NCG experiment. Because ^{15}N chemical shifts of lysine side-chain NH_3^+ groups are unique (~ 33 ppm), ^{15}N r-SNOB pulses (Kupce et al. 1995) selective to $^{15}\text{N}\zeta$ are used to suppress potentially aliased signals from arginine and backbone ^{15}N nuclei. In this experiment, undesirable coherence transfer between $^{15}\text{N}\zeta$ and $^{13}\text{C}\epsilon$ nuclei is avoided by using ^{13}C I-BURP2 pulses (Geen and Freeman 1991) that invert $^{13}\text{C}\gamma$ (~ 25 ppm) but do not affect $^{13}\text{C}\epsilon$ (~ 42 ppm) nuclei. Although these I-BURP2 pulses also invert $^{13}\text{C}\delta$ (~ 28 ppm), $^2J_{\text{NC}\delta}$ coupling constants are undetectably small in lysine side chains (Zandarashvili et al. 2011). Thus, these ^{15}N and ^{13}C shaped pulses permit selective

observation of $^1\text{H}\zeta/^{15}\text{N}\zeta/^{13}\text{C}\gamma$ cross peaks from lysine side chains. Using the 3D H3NCECD (Iwahara et al. 2007) and 3D H3NCG spectra, three resonances of $^{13}\text{C}\gamma$, $^{13}\text{C}\delta$, and $^{13}\text{C}\epsilon$ nuclei can be obtained for each lysine side-chain NH_3^+ group. Because the ^{13}C - ^{15}N coherence transfer in the H3NCG and H3NCECD experiments is not exposed to rapid ^{13}C transverse relaxation, these experiments are more sensitive than the (H)CCENH3 experiment that involves coherence transfer via J -evolution from C_x to $2C_yN_z$. It should also be noted that the H3NCG and H3NCECD experiments with ^2H decoupling can be readily performed for perdeuterated proteins. Although deuteration should substantially slow ^{13}C transverse relaxation and therefore improve ^{13}C - ^{13}C coherence transfer (Yamazaki et al. 1994), the lack of side-chain ^1H nuclei in perdeuterated proteins makes the (H)CCENH3 experiment impractical.

Using a 750-MHz spectrometer, we recorded 3D H3NCG and 3D H3NCECD spectra for the Egr-1 – DNA complex at 10 °C. This complex contains the Egr-1 zinc-finger protein (90 amino acids) and 12-bp DNA with Egr-1's target sequence. Our group uses the complex to study dynamics in DNA scanning and recognition by Egr-1 (Esadze et al. 2014; Zandarashvili et al. 2012). The Egr-1 zinc-finger protein and 12-bp DNA were purified as previously described (Esadze and Iwahara 2014; Takayama et al. 2010). A 280- μl solution of 0.8 mM complex in a buffer of 20 mM potassium succinate (pH 5.8), 2 mM ZnCl_2 , and 20 mM KCl was sealed into an inner tube of co-axial NMR tube. To avoid deuterated species of NH_3^+ groups, D_2O for NMR lock was separately sealed in a thin outer layer of the co-axial NMR tube (Iwahara et al. 2007). The relatively low temperature, 10 °C, was required to observe ^1H - ^{15}N HISQC (Iwahara et al. 2007) cross peaks from all 6 Lys side-chain NH_3^+ groups of the Egr-1 – DNA complex under these solution conditions. 3D (H)CCENH3 spectrum (Iwahara et al. 2007) recorded at this temperature did not show signals from any Lys residues other than Lys89, which is located at the disordered C-terminal tail (data not shown). Because the (H)CCENH3 experiment involves the ^{13}C - ^{15}N coherence transfer schemes via J -evolution from C_x to $2C_yN_z$ terms and the ^{13}C - ^{13}C cross polarization scheme, rapid ^{13}C transverse relaxation can severely reduce sensitivity at low temperatures. In contrast, the 3D H3NCECD and 3D H3NCG spectra clearly show signals from all Lys side-chain NH_3^+ groups (Figure 2). As described above, higher sensitivity of these experiments can be attributed largely to efficient ^{13}C - ^{15}N coherence transfer via J -evolution from N_x to $2N_yC_z$ terms.

Our second approach to assign lysine side-chain NH_3^+ resonances at low temperatures is to observe $^{15}\text{N}\zeta$ resonances through ^1H detection of lysine side-chain CH_2 groups at higher temperatures. In this approach, we use a new 2D heteronuclear long-range correlation spectrum, (H2C)N(CC)H-TOCSY, which shows cross peaks of lysine side-chain $^{15}\text{N}\zeta$ and CH_2 proton resonances. Figure 3 illustrates the pulse sequence for this experiment. Owing to the detection of carbon-attached ^1H nuclei, this experiment is not hindered by rapid hydrogen exchange of NH_3^+ groups, and therefore can be performed using a high temperature at which ^{13}C - ^{13}C Hartmann-Hahn cross polarization can occur efficiently. As described previously (Iwahara et al. 2007; Andre et al. 2007), ^{15}N transverse magnetization is maintained to be in-phase with respect to ^1H via ^1H -WALTZ-16 decoupling so that scalar relaxation due to hydrogen exchange is avoided. Although this experiment can be

implemented as a 3D experiment with a ^{13}C dimension, we used the 2D version to achieve a higher resolution in the ^{15}N dimension as well as a higher sensitivity. Figure 4 shows the 2D (H2C)N(CC)H-TOCSY spectrum recorded for the Egr-1 – DNA complex. While rapid hydrogen exchange makes it impossible to detect ^1H signals from the Lys NH_3^+ groups of the Egr-1 – DNA complex at 35 °C, the (H2C)N(CC)H-TOCSY spectrum for the same sample at this temperature exhibited many ^1H - ^{15}N cross peaks from Lys side chains (Figure 4C). In this spectrum, there was significant variation in signal intensity, presumably because individual Lys side chains exhibit different degrees of relaxation loss, depending on fast internal motions or slow conformational exchange (Foster et al. 1997). Because multiple side-chain ^1H resonances are observed at high resolution along the ^1H direct-detection dimension with excellent separation by sharp ^{15}N resonances, the (H2C)N(CC)H-TOCSY spectrum greatly facilitates ^{15}N resonances assignment for lysine NH_3^+ groups. The temperature dependence of the H2(C)N spectra (Andre et al. 2007) allows us to track the change of ^{15}N resonances as a function of temperature (Figure 4B). These data together with the 3D H3NCG and H3NCECD spectra allowed us to unambiguously assign ^1H - ^{15}N HISCQ signals from the Lys NH_3^+ groups of the Egr-1 – DNA complex at 10 °C (Figure 4A).

In conclusion, we have presented an effective strategy for resonance assignment of lysine NH_3^+ groups that can be observed only at relatively low temperatures. Our new approaches can help expand applicability of the NMR methods for characterizing Lys side-chain NH_3^+ groups (Anderson et al. 2013; Andre et al. 2007; Esadze et al. 2011; Iwahara et al. 2007; Segawa et al. 2008; Tomlinson et al. 2009; Williamson et al. 2013; Zandarashvili et al. 2013; Zandarashvili et al. 2011). As a consequence, dynamics of hydrogen bonds and ion pairs involving Lys side chains may become more assessable for a larger number of proteins and protein complexes.

Acknowledgments

We thank Tianzhi Wang for maintenance of the spectrometers at UTMB NMR facility and Dan Nguyen for language editing. This work was supported by Grant R01GM105931 from the National Institutes of Health (to J.I.).

References

- Anderson KM, Esadze A, Manoharan M, Bruschweiler R, Gorenstein DG, Iwahara J. Direct Observation of the Ion-Pair Dynamics at a Protein-DNA Interface by NMR Spectroscopy. *J Am Chem Soc.* 2013; 135:3613–3619. [PubMed: 23406569]
- Andre I, Linse S, Mulder FA. Residue-specific pKa determination of lysine and arginine side chains by indirect ^{15}N and ^{13}C NMR spectroscopy: application to apo calmodulin. *J Am Chem Soc.* 2007; 129:15805–15813. [PubMed: 18044888]
- Blaum BS, Deakin JA, Johansson CM, Herbert AP, Barlow PN, Lyon M, Uhrin D. Lysine and arginine side chains in glycosaminoglycan-protein complexes investigated by NMR, cross-linking, and mass spectrometry: a case study of the factor H-heparin interaction. *J Am Chem Soc.* 2010; 132:6374–81. [PubMed: 20394361]
- Esadze A, Iwahara J. Stopped-flow fluorescence kinetic study of protein sliding and intersegment transfer in the target DNA search process. *J Mol Biol.* 2014; 426:230–244. [PubMed: 24076422]
- Esadze A, Kemme CA, Kolomeisky AB, Iwahara J. Positive and negative impacts of nonspecific sites during target location by a sequence-specific DNA-binding protein: origin of the optimal search at physiological ionic strength. *Nucleic Acids Res.* 2014; 42:7039–46. [PubMed: 24838572]

- Esadze A, Li DW, Wang T, Brüschweiler R, Iwahara J. Dynamics of lysine side-chain amino groups in a protein studied by heteronuclear ^1H - ^{15}N NMR spectroscopy. *J Am Chem Soc.* 2011; 133:909–919. [PubMed: 21186799]
- Fischer MWF, Zeng L, Zuiderweg ERP. Use of ^{13}C - ^{13}C NOE for the assignment of NMR lines of larger labeled proteins at larger magnetic fields. *J Am Chem Soc.* 1996; 118:12457–12458.
- Foster MP, Wuttke DS, Radhakrishnan I, Case DA, Gottesfeld JM, Wright PE. Domain packing and dynamics in the DNA complex of the N-terminal zinc fingers of TFIIIA. *Nat Struct Biol.* 1997; 4:605–8. [PubMed: 9253405]
- Geen H, Freeman R. Band-selective radiofrequency pulses. *J Magn Reson.* 1991; 93:93–141.
- Huang J, MacKerell AD Jr. CHARMM36 all-atom additive protein force field: validation based on comparison to NMR data. *J Comput Chem.* 2013; 34:2135–2145. [PubMed: 23832629]
- Iwahara J, Jung YS, Clore GM. Heteronuclear NMR spectroscopy for lysine NH_3 groups in proteins: unique effect of water exchange on ^{15}N transverse relaxation. *J Am Chem Soc.* 2007; 129:2971–2980. [PubMed: 17300195]
- Kay LE, Xu GY, Singer AU, Muhandiram DR, Formankay JD. A Gradient-Enhanced HCCH-TOCSY Experiment for Recording Side-Chain ^1H and ^{13}C Correlations in H_2O Samples of Proteins. *J Magn Reson Ser B.* 1993; 101:333–337.
- Kup e E, Boyd J, Campbell ID. Short selective pulses for biochemical applications. *J Magn Reson Ser B.* 1995; 106:300–303. [PubMed: 7719630]
- Marion D, Ikura M, Tschudin R, Bax A. Rapid Recording of 2d Nmr-Spectra without Phase Cycling - Application to the Study of Hydrogen-Exchange in Proteins. *J Magn Reson.* 1989; 85:393–399.
- Piotto M, Saudek V, Sklenár V. Gradient-tailored excitation for single-quantum NMR spectroscopy of aqueous solutions. *J Biomol NMR.* 1992; 2:661–5. [PubMed: 1490109]
- Poon DKY, Schubert M, Au J, Okon M, Withers SG, McIntosh LP. Unambiguous determination of the ionization state of a glycoside hydrolase active site lysine by ^1H - ^{15}N Heteronuclear Correlation Spectroscopy. *J Am Chem Soc.* 2006; 128:15388–9. [PubMed: 17132001]
- Segawa T, Kateb F, Duma L, Bodenhausen G, Pelulessy P. Exchange rate constants of invisible protons in proteins determined by NMR spectroscopy. *Chembiochem.* 2008; 9:537–542. [PubMed: 18247446]
- Shaka AJ, Keeler J, Freeman R. Evaluation of a New Broad-Band Decoupling Sequence - Waltz-16. *J Magn Reson.* 1983; 53:313–340.
- Shaka AJ, Lee CJ, Pines A. Iterative Schemes for Bilinear Operators - Application to Spin Decoupling. *J Magn Reson.* 1988; 77:274–293.
- Takayama Y, Sahu D, Iwahara J. NMR studies of translocation of the Zif268 protein between its target DNA Sites. *Biochemistry.* 2010; 49:7998–8005. [PubMed: 20718505]
- Tomlinson JH, Ullah S, Hansen PE, Williamson MP. Characterization of salt bridges to lysines in the protein G B1 domain. *J Am Chem Soc.* 2009; 131:4674–4684. [PubMed: 19281232]
- Williamson MP, Hounslow AM, Ford J, Fowler K, Hebditch M, Hansen PE. Detection of salt bridges to lysines in solution in barnase. *Chem Commun.* 2013; 49:9824–9826.
- Yamazaki T, Lee W, Arrowsmith CH, Muhandiram DR, Kay LE. A suite of triple resonance NMR experiments for the backbone assignment of ^{15}N , ^{13}C , ^2H labeled proteins with high sensitivity. *J Am Chem Soc.* 1994; 116:11655–66.
- Zandarashvili L, Esadze A, Iwahara J. NMR studies on the dynamics of hydrogen bonds and ion pairs involving lysine side chains of proteins. *Adv Protein Chem Struct Biol.* 2013; 93:37–80. [PubMed: 24018322]
- Zandarashvili L, Li DW, Wang T, Brüschweiler R, Iwahara J. Signature of mobile hydrogen bonding of lysine side chains from long-range ^{15}N - ^{13}C scalar J-couplings and computation. *J Am Chem Soc.* 2011; 133:9192–9195. [PubMed: 21591797]
- Zandarashvili L, Vuzman D, Esadze A, Takayama Y, Sahu D, Levy Y, Iwahara J. Asymmetrical roles of zinc fingers in dynamic DNA-scanning process by the inducible transcription factor Egr-1. *Proc Natl Acad Sci U S A.* 2012; 109:E1724–1732. [PubMed: 22675124]

Figure 1.

Pulse sequences for the 3D H3NCG experiment for Lys side-chain NH_3^+ resonance assignment. Thin and bold bars in black represent hard rectangular 90° and 180° pulses, respectively. Unless indicated otherwise, pulse phases are along x . Carrier positions: ^1H , the position of the water resonance; ^{15}N , 33 ppm; and ^{13}C , 20 ppm. Short-bold bars represent water-selective soft-rectangular ^1H 90° pulses (1.2 ms). The 3-9-19 pulse train-based WATERGATE scheme (Piotto et al. 1992) was used to suppress the water signal. The $^{13}\text{C}_\gamma$ -selective pulses should not affect $^{13}\text{C}_\epsilon$ nuclei ($\sim 42\text{ppm}$). RF strengths for ^1H and ^{15}N WALTZ-16 composite pulses (Shaka et al. 1983) were 3.3 kHz and 1.0 kHz, respectively. Shaped pulses: ^1H half-Gaussian 90° pulse (2.1 ms); ^{13}C I-BURP2 180° pulse (1.2 ms); and ^{15}N r-SNOB 180° pulse (1.03 ms). Delays: $\tau_a = 2.7$ ms; $\delta = 2.6$ ms; and $T = 50$ ms. Phase cycles: $\phi_1 = [2x, 2(-x)]$, $\phi_2 = [x, -x]$, $\phi_3 = [4x, 4y]$, and receiver = $[x, -x, -x, x, -x, x, x, -x]$. Quadrature detections for indirect ^{13}C and ^{15}N dimensions were achieved using States-TPPI (Marion et al. 1989) for ϕ_1 and ϕ_2 , respectively.

Figure 2.

Strips of the 3D H3NCG and H3CECD spectra for all Lys side-chain NH_3^+ groups of the Egr-1 – DNA complex at 10 °C. Negative contours are shown in green. The ^1H and ^{15}N spectral widths for were 18.0 ppm and 4.7 ppm, respectively, in both experiments. The ^{13}C spectral widths were 20.7 ppm in H3NCG and 26.0 ppm in H3NCECD. In each experiment, 32 scans were accumulated per FID, and numbers of complex points for ^1H , ^{13}C , and ^{15}N dimensions were 810, 32, and 32, respectively. The total experimental time was 60 hours each. These spectra were recorded at the ^1H frequency of 750 MHz using a Bruker Avance III spectrometer equipped with a non-cryogenic TXI probe.

Figure 3.

The pulse sequence of the 2D (H₂C)N(CC)H-TOCSY experiment. This experiment provides correlation between Lys side-chain ¹⁵N_ζ and ¹H resonances, and corresponds to a hybrid of the H₂CN (Andre et al. 2007) and HCCH-TOCSY (Kay et al. 1993) experiments for H₂O samples. ¹³C shaped pulses are Lys ¹³C_ε-selective r-SNOB pulses (980 us). Length of the ¹³C- spin lock (SL) was 1 ms. ¹³C carrier position was 43 ppm. The rf strength of the DIPSI-3 scheme (the total length, 16.9 ms) was 9.6 kHz and that of hard ¹³C pulses was 20.8 kHz. ¹⁵N shaped pulses are Lys ¹⁵N_ζ-selective r-SNOB pulses (1.0 ms). ¹⁵N carrier position was 32 ppm. Delays: $\tau_1 = 1.6$ ms; $\tau_2 = 1.1$ ms; $\delta = 1.7$ ms; and $T_c = 20$ ms. Phase cycles: $\phi_1 = [x, -x]$; $\phi_2 = [2x, 2(-x)]$, $\phi_3 = [4x, 4y]$; $\phi_4 = [4x, 4(-x)]$; and receiver = $[x, -x, -x, x, -y, y, y, -y]$. Quadrature detections for indirect ¹⁵N dimensions was achieved using States-TPPI for ϕ_1 .

Figure 4.

Use of 2D (H₂C)N(CC)H-TOCSY spectrum and temperature dependence of 2D H₂(C)N spectra for assignment of Lys NH₃⁺ groups. The spectra were recorded for 0.8 mM ¹³C/¹⁵N-labeled Egr-1 – DNA complex at pH 5.8. (A) The lysine NH₃⁺-selective HISQC (Iwahara et al. 2007) spectrum recorded at 10 °C. (B) Lysine side-chain-specific H₂(C)N (Andre et al. 2007) spectra recorded at 10 °C (black), 18 °C (blue), 27 °C (magenta), and 35 °C (red). (C) (H₂C)N(CC)H-TOCSY spectrum recorded at 35 °C. The spectra widths and numbers of complex points: 13.4 ppm and 400 points for ¹H; and 2.2 ppm and 50 points for ¹⁵N. The mixing time of the ¹³C DIPSI-3 scheme (Shaka et al. 1988) was 16.9 ms. 1024 scans were accumulated per FID. The total time to record the 2D (H₂C)N(CC)H-TOCSY spectrum was 56 hours. All spectra shown in this figure were recorded with a Bruker Avance III spectrometer equipped with a QCI cryogenic probe operated at the ¹H frequency of 600 MHz.