Published in final edited form as:

Org Biomol Chem. 2014 October 7; 12(37): 7345–7354. doi:10.1039/c4ob00953c.

Cooperative Hybridization of γ PNA Miniprobos to a Repeating Sequence Motif and Application to Telomere Analysis

Ha H. Pham^{#†§}, Connor T. Murphy^{#†§}, Gopalsamy Sureshkumar^{†§}, Danith H. Ly^{†§}, Patricia L. Opresko^{†§}, and Bruce A. Armitage^{†§*}

[†] Department of Chemistry, Carnegie Mellon University, 4400 Fifth Avenue, Pittsburgh, PA 15213

[§] Center for Nucleic Acids Science and Technology, Carnegie Mellon University, 4400 Fifth Avenue, Pittsburgh, PA 15213

[‡] Department of Environmental and Occupational Health, University of Pittsburgh, Graduate School of Public Health, Pittsburgh, PA 15219

[#] These authors contributed equally to this work.

Abstract

GammaPNA oligomers having one or two repeats of the sequence AATCCC were designed to hybridize to DNA having one or more repeats of the complementary TTAGGG sequence found in the human telomere. UV melting curves and surface plasmon resonance experiments demonstrate high affinity and cooperativity for hybridization of these miniprobos to DNA having multiple complementary repeats. Fluorescence spectroscopy for Cy3-labeled miniprobos demonstrate increases in fluorescence intensity for assembling multiple short probes on a DNA target compared with fewer longer probes. The fluorescent γ PNA miniprobos were then used to stain telomeres in metaphase chromosomes derived from U2OS cells possessing heterogeneous long telomeres and Jurkat cells harboring homogenous short telomeres. The miniprobos yielded comparable fluorescence intensity to a commercially available PNA 18mer probe in U2OS cells, but significantly brighter fluorescence was observed for telomeres in Jurkat cells. These results suggest that γ PNA miniprobos can be effective telomere-staining reagents with applications toward analysis of critically short telomeres, which have been implicated in a range of human diseases.

Hybridization probes are central to many biotechnological applications involving nucleic acids, including microarrays, PCR amplification and diagnostics, fluorescence *in situ* hybridization (FISH), affinity purification and antisense regulation of gene expression. Modified backbones such as phosphorothioate, morpholino and locked nucleic acids (LNA) have established footholds in various applications due to superior performance relative to unmodified DNA oligonucleotides. For example, the nuclease stability of morpholino oligomers has led to their common use as antisense agents to block translation of specific

*To whom correspondence should be addressed. Tel 412-268-4196, Fax 412-268-1061, army@cmu.edu.

Supporting Information. UV melting curves, SPR sensorgrams, microscope images and MALDI-TOF spectra and HPLC chromatograms for γ PNA oligomers. This material is available free-of-charge via the Internet at <http://pubs.rsc.org>.

mRNAs in microinjected zebrafish embryos^{1, 2}, while LNA is gaining increasing use due in large part to its exceptionally high affinity for complementary DNA and RNA³.

Peptide nucleic acid (PNA, Chart 1) is another synthetic nucleic acid analogue that has found applications, primarily in FISH analyses. Considerable interest in PNA has arisen from its high affinity for complementary DNA and RNA in spite of its radically altered backbone structure relative to DNA.⁴ (The enhanced affinity is due, at least in part, to the lack of a negative charge on the PNA backbone.⁵) A fluorescent PNA 18mer complementary to three repeats of the human telomere sequence is widely used for telomere staining and quantitative length determination.⁶ PNA FISH probes targeted to ribosomal RNA are also commonly used in clinical pathogen identification assays.⁷

An important advance in the development of PNA came with the reports of γ PNA, which exhibits further enhancement of affinity,⁸ comparable to LNA,⁹ but also improved water solubility with appropriate substituents at the gamma carbon¹⁰ (Chart 1). The high affinity of γ PNA has been attributed to its right-handed helical structure, which pre-organizes γ PNA for hybridization to complementary DNA and RNA.^{8, 11} The added affinity extends the strand invasion capability first reported for unmodified PNA^{12, 13} to virtually any sequence of double-stranded DNA.^{14, 15}

The high affinity of γ PNA also provides an opportunity to address an ongoing concern with FISH probes, namely brightness. In order to sensitively detect genomic DNA or low abundance RNA targets *in situ* without signal amplification, bright probes are needed. cDNA probes prepared by enzymatic methods can contain multiple fluorescent dyes by virtue of using labeled dNTPs, although such long, polyanionic probes typically require lengthy incubation steps due to slow hybridization kinetics. Alternatively, relatively short (20mer), synthetic 2'-O-methoxy probes, each bearing a fluorescent dye and targeted to a different subsequence within a transcript, have been used to stain mRNA.¹⁶ Multimer probes consisting of up to four copies of a biotinylated probe assembled via a streptavidin core have also been used to achieve single molecule imaging of endogenous mRNAs.¹⁷

In the context of nucleic acid hybridization, higher affinity can be translated into shorter probe length to realize comparable equilibrium constant and melting temperature as lower affinity probes. These “miniprobos” offer a new opportunity to increase the brightness of a labeled DNA or RNA target: more probes and, therefore, more dyes can be delivered to a target sequence of a given length. This strategy can be particularly effective for staining repeating sequence targets, such as telomeric DNA. Herein, we describe the DNA hybridization properties of γ PNA miniprobos and demonstrate effective staining of telomeres in two human cell lines.

RESULTS AND DISCUSSION

Design

γ PNA miniprobos were designed to be complementary to the repeating hexamer sequence found in human telomeric DNA: 5'-TTAGGG-3'. Three oligomers were synthesized (Table 1): a 12mer complementary to two telomere repeats, a 6mer complementary to a single

repeat, and a 6mer in which two of the cytosine bases were replaced by guanidino G-clamp, a synthetic nucleobase that forms more stable base pairs with G than does C, by virtue of additional hydrogen bonding and pi stacking ability.^{18, 19} Ly and coworkers reported that incorporation of guanidino G-clamp residues into γ PNA significantly enhances strand invasion into duplex DNA.²⁰ As shown in Scheme 1, hybridization of the 12mer or 6mer γ PNA miniprobcs to a complementary DNA of a given length will result in 1.5 or 3 times as many dyes being delivered to the target nucleic acid as the conventional PNA probe.

γ PNAs were synthesized by standard solid-phase methods,^{21, 22} end-labeled with Cy3 prior to cleavage from the polystyrene support, purified by reversed phase HPLC and characterized by MALDI-ToF mass spectrometry (supporting information).

UV Melting Analysis

The DNA hybridization properties of the γ PNAs were assessed using UV melting curve analysis. We first compared formation of 1:1 duplexes between each of the γ PNAs and a complementary DNA oligonucleotide consisting of the corresponding number of telomeric repeats. For example γ **P₁₂** was hybridized to **Telo-2**, consisting of two telomeric repeats, while γ **P₆** and γ **P_{6-clamp}** were hybridized to **Telo-1**. For comparison, **P₁₈** was hybridized to **Telo-3**.

Figure 1 shows UV melting curves recorded at 260 nm for samples containing PNA or γ PNA hybridized to the complementary DNA in a Tris buffer containing 100 mM KCl. Under these conditions, the traditional PNA probe **P₁₈** exhibits a melting temperature $T_m = 66$ °C with the complementary **Telo-3** probe. The 12mer γ PNA miniprobe γ **P₁₂** forms a more stable duplex with its complementary DNA **Telo-2** ($T_m = 77$ °C) in spite of having 6 fewer base pairs, illustrating the exceptionally high affinity of the gamma-modified backbone. In contrast, the γ **P₆**/**Telo-1** duplex has a much lower stability ($T_m = 24$ °C), although substituting two G-clamp bases into the γ PNA significantly increases the stability to the point where the 6mer γ PNA-DNA duplex is almost as stable as the 18mer PNA-DNA duplex ($T_m = 62$ °C versus 66 °C).

We next characterized the ability of multiple γ PNA miniprobcs to hybridize to a single DNA oligonucleotide having multiple repeating complementary target sites. The experiments included hybridization of γ **P₁₂** to either **Telo-2** (1:1 duplex) or **Telo-4** (2:1 duplex), and of γ **P₆** or γ **P_{6-clamp}** to **Telo-1**/**Telo-4** (1:1 up to 4:1 duplexes). The experiments were designed to maintain the same total base pair concentration (24 μ M) by adjusting the concentration of the DNA complement. For example, γ **P₆** and **Telo-1** were mixed in 4 μ M concentration, whereas **Telo-4** was used at 1 μ M concentration.

Melting curves are shown in Supplementary Figures 1 and 2 and melting temperatures are reported in Table 2. It is evident that the stabilities of the duplexes formed by γ **P₁₂** and γ **P_{6-clamp}** increase with the length of the DNA complement, in spite of the entropic penalty inherent to formation of 2:1, 3:1 or 4:1 duplexes compared with 1:1 duplexes. These results indicate considerable cooperativity in hybridization of γ PNA miniprobcs to adjacent sites on the DNA complement, with end-to-end stacking of the adjacent probes likely responsible for the added stability for the longer DNA complements. A similar phenomenon was proposed

to account for the higher thermal stability of a PNA₂-DNA triplex formed on a polymeric deoxyadenosine template relative to a dA₈ single strand.²³ Cooperative hybridization has also been proposed as a possible explanation for the ability of antisense oligonucleotides to inhibit translation of *huntingtin* mRNA bearing greater numbers of triplet CAG repeats relative to alleles with fewer repeats²⁴. It is also worth noting that **Telo-4** folds into a G-quadruplex structure at room temperature, but this is far less stable ($T_m = 40^\circ\text{C}$, Supplementary Figure 3) than the corresponding duplexes formed with γP_{12} and $\gamma\text{P}_{6\text{-clamp}}$. (γP_6 initially shows a substantial increase in T_m for **Telo-2** versus **Telo-1**, but this is reduced at longer DNA lengths, perhaps due in part to competitive G-quadruplex formation, at least for **Telo-4**.) Taken together, the UV melting experiments reveal that the γ PNA miniprobcs can hybridize to complementary DNA oligonucleotides to form heteroduplexes stabilized by cooperative end-stacking of adjacent miniprobcs.

Fluorescence Spectroscopy

The advantage of miniprobcs for fluorescent labeling is the ability to deliver more dyes per target nucleic acid (Scheme 1). To demonstrate this effect, we compared fluorescence spectra recorded for **P₁₈**, γP_{12} and $\gamma\text{P}_{6\text{-clamp}}$ after hybridization to a 36mer DNA bearing 6 repeats of the AATCCC complementary sequence. Thus, the three probes should hybridize in 2:1, 3:1 and 6:1 stoichiometries, respectively (Scheme 1). Figure 2 shows the spectra, normalized to **P₁₈**. The fluorescence intensity increases by factors of 1.7 and 3.4 for the 12mer and 6mer γ PNA miniprobcs, respectively, relative to the PNA 18mer, slightly exceeding the theoretical 1.5- and 3-fold increases based on the stoichiometries. These results demonstrate that fluorescently labeled γ PNA miniprobcs as short as 6mers can be hybridized directly adjacent to one another along a DNA target with no apparent loss of quantum yield for the fluorophore.

SPR Analysis

We also used surface plasmon resonance experiments to compare the kinetics of hybridization and dissociation for the various PNA and γ PNA oligomers to immobilized DNA targets. For this experiment, approximately the same mass of biotinylated **Telo-1**, **Telo-2** and **Telo-4** was immobilized on separate flow cells of a streptavidin-coated SPR chip. Since the molecular weight of the three DNAs approximately doubles with each increase in length, this insures that the same number of binding sites is present on each flow cell, although the surface density of the DNA varies. Each of the γ PNA miniprobcs was flowed over the surface of the chip at a concentration of 50 nM for a period of 420 sec, then the flow solution was changed to γ PNA-free buffer to promote dissociation of the bound γ PNA. Sensorgrams representing the average of three trials are shown in Figures 3 and 4 and initial on- and off-rates are shown in Tables 3 and 4.

Figure 3 (left) shows SPR sensorgrams for the hexamer miniprobe γP_6 . No binding is observed with **Telo-1**, consistent with the low melting temperature recorded for this duplex, even at much higher concentration ($T_m = 24^\circ\text{C}$, Table 2). Extending the DNA target to include two complementary sites (**Telo-2**) leads to observable binding of γP_6 , although the γ PNA also dissociates relatively quickly from the immobilized DNA. Binding of the γ PNA probe to **Telo-4** is significantly slower than to **Telo-2** under these conditions, presumably

due to folding of **Telo-4** into a G-quadruplex structure on the chip, as we observed previously for a related DNA oligonucleotide.²⁵ This hypothesis was supported by the observation that replacing KCl with LiCl, where the quadruplex would be destabilized,²⁶ resulted in a large increase in γ PNA hybridization (Supplementary Figure 4).

The same experiment was performed using the higher affinity miniprobe γ **P_{6-clamp}** (Figure 3, center). In contrast to γ **P₆**, γ **P_{6-clamp}** binds rapidly to **Telo-1**, followed by slower dissociation. Binding to **Telo-2** appears to occur in two phases, a faster phase that slightly exceeds that of **Telo-1**, followed by a slower phase. Since there are two binding sites for the γ PNA on **Telo-2**, one possible explanation for these observations is that the kinetics reflect differences in rates for binding to the two sites. Within this model, the fast rate could correspond to binding to the distal site, i.e. the site furthest from the chip surface, while the slower rate could be due to lower accessibility of the site proximal to the streptavidin-coated surface. Regardless of the order of hybridization to the two sites, once bound, the γ PNA molecules strongly resist dissociation. Although the slower off-rate from **Telo-2** relative to **Telo-1** could reflect the presence of two binding sites per DNA strand (which would promote re-capture of a dissociated γ PNA), another contribution likely comes from the stabilizing effect of end-stacking between adjacent γ PNA molecules hybridized to the same strand. Notably, while association of γ **P_{6-clamp}** is slower for **Telo-4**, those γ PNA molecules that bind to the immobilized DNA are even slower to dissociate than for **Telo-2**. In addition, comparing the data for γ **P_{6-clamp}** and γ **P₆** with **Telo-2**, these results indicate that the guanidine G-clamp nucleobase modifications exert their affinity-enhancing effect primarily through a slower dissociation rate.

We next performed SPR analysis of the 12mer miniprobe γ **P₁₂**. Figure 3 (right) compares binding and dissociation from **Telo-2** and **Telo-4**. Even the 1:1 complex between γ **P₁₂** and **Telo-2** shows very slow dissociation, as we reported previously for γ PNA-DNA duplexes.¹⁰ Interestingly, four-fold less γ **P₁₂** binds to the quadruplex target **Telo-4** versus the unfolded **Telo-2**, whereas this difference for γ **P_{6-clamp}** is only two-fold (compare the central and right panels of Figure 3), perhaps indicating a kinetic advantage of shorter probes for invading structured targets.

The data in Figure 3 were normalized to the highest binding signal within each set of data. To compare different γ PNA miniprobos for binding to a single target, we renormalized the sensorgrams for binding to **Telo-2** (Figure 4). These data show the slightly faster binding of the γ **P_{6-clamp}** versus γ **P₁₂** but slower dissociation of the longer probe, where twice as many base pairs need to be disrupted in order to allow the probe to dissociate.

Table 3 shows the initial on- and off-rates determined from the SPR sensorgrams. Several conclusions can be drawn from the data, which reinforce the qualitative observations made from the sensorgrams. First, the data for **Telo-2** illustrate similar on-rates for the three γ PNAs but off-rates that vary by a factor of 33 from the fastest (γ **P₆**) to the slowest (γ **P₁₂**). These results illustrate the helix-stabilizing effects of the guanidine G-clamp nucleobase for γ **P_{6-clamp}** and the longer probe length for γ **P₁₂**. Next, comparing data across a row, e.g. γ **P_{6-clamp}** with **Telo-1**, **-2** and **-4**, The increasing kinetic stability with target length is evident as the off-rates get progressively slower as more probes can hybridize adjacent to

one another. In particular, comparing **Telo-1** with **Telo-2**, the on-rate is only 29% greater for the longer target, whereas the off-rate is nearly 7-fold slower, connecting the cooperativity of hybridization to greater resistance to dissociation. In addition, the much slower on-rates due to the inherent quadruplex structure for **Telo-4** are also evident in comparing the absolute values with those of **Telo-2**.

Telomere Analysis

Since the initial report by Lansdorp and coworkers⁶, the 18mer Cy3-labeled PNA probe **P₁₈** has become widely used for telomere imaging and length analysis. However, growing epidemiological evidence indicating that critically short telomeres are implicated in a wide range of aging-related diseases^{27, 28} motivates the development of brighter telomere probes, since shorter telomeres will bind fewer probes and therefore give lower fluorescence intensity relative to longer telomeres. Our hypothesis was that the higher affinity of γ PNA would permit 12mer or perhaps even 6mer versions of the original 18mer PNA probe to be used for telomere analysis since this would allow a greater number of probes to bind to a telomere of a given length (Scheme 1).

To test the ability of the fluorescent γ PNA miniprobcs to stain telomeres using quantitative fluorescence *in situ* hybridization (qFISH), two human cell lines, U2OS osteosarcoma and Jurkat T-lymphocytes, were stained using standard telomere-FISH (teloFISH) protocols.²⁹ These cell lines were chosen for their divergent telomere maintenance mechanisms. Approximately 85% of human cancers maintain stable telomere lengths through the activation of the holoenzyme telomerase.³⁰ Conversely, approximately 15% of cancers maintain their telomeres in the absence of telomerase through a recombination-based pathway known as alternative lengthening of telomeres (ALT).³⁰ This includes U2OS cells, which have relatively long mean telomere lengths (~21kb) compared to normal human cell lines.³¹⁻³³ ALT-active cells exhibit highly variable telomere lengths, leading to a notable number of signal free ends (SFEs) in U2OS cells despite their long mean length. In contrast, Jurkat cells are telomerase positive and have shorter average telomere lengths than U2OS cells.³¹⁻³³

Cells were stalled in metaphase by the addition of colcemid and metaphase chromosome spreads were prepared and fixed as previously described.³⁴ Chromosomes were hybridized for two hours with **P₁₈**, γ **P₁₂** or γ **P_{6-clamp}** and imaged for telomere staining. The fluorescence intensities of the Cy3 spots were determined using TFL-Telo quantitation software. Results for U2OS cells are shown in Figures 5 and 6. The yellow fluorescent spots observed at the ends of the chromosomes (counterstained with DAPI and colored blue) indicate successful telomere staining by all three probes (Figure 5). Quantitative analysis (Figure 6) revealed similar performance for the three probes in the low intensity region of the histogram, including equal number of unstained termini (SFEs), which result from loss of a telomere or telomeres that are too short to bind sufficient probe for detection. However, at higher fluorescence intensities, slightly higher percentages of telomeres were stained by the γ PNA probes, particularly γ **P_{6-clamp}** (Figure 6, inset).

The similarity in SFEs detected using all three probes (18, 12 and 6-mer), despite the improved fluorescence intensity of the γ -modified probes, implies that the shortest telomeres

in U2OS cells may be below the threshold for detection even for the shorter probes. Alternatively, these SFEs may represent chromatid ends that lack telomeric DNA or contain degenerative repeats such as those in subtelomeric DNA, since telomere lengthening is not templated by telomerase in these cells (due to the absence of telomerase).³⁵

Similar experiments were performed with Jurkat cells (Figure 7A and Supplementary Figure 5), which rely on telomerase for telomere maintenance and exhibit stable lengths that average 6 kb.³² Considerable non-telomeric staining of chromosomes was observed for γ **P_{6-clamp}** (data not shown) so the analysis is restricted to **P₁₈** and γ **P₁₂**. In contrast to U2OS cells, staining Jurkat cells with γ **P₁₂** resulted in approximately 5-fold fewer SFEs than staining with **P₁₈**, consistent with improved staining of critically short telomeres. The γ **P₁₂** staining also revealed a significantly greater percentage of telomeres with brighter signals compared to **P₁₈**. The increase in the number of signal free ends and brighter overall fluorescence in this case is likely due to hybridization of more γ **P₁₂** probes relative to **P₁₈** and suggests that analyzing the shortest telomeres in Jurkat cells is improved by using a more sensitive probe.

The standard qFISH protocol requires denaturation of the chromosomal DNA prior to hybridization of the probe to the telomere, which consists of a relatively short single-stranded region extending into a much longer double-stranded region, both of which contain TTAGGG repeats complementary to the probe. However, previous studies have shown that γ PNA affinity for complementary DNA is sufficiently high to allow direct invasion of double-stranded DNA in order to access target sites.¹⁵ Therefore, we hypothesized that the denaturing step could be omitted when using the γ **P₁₂** probe to stain the metaphase chromosome preparations. To test this, we compared unmodified **P₁₈** and γ **P₁₂** under non-denaturing hybridization conditions, in which formamide was omitted from the hybridization buffer and heating and RNase A treatment steps were deleted from the protocol. Under these conditions, ca. 75% of the telomeres were unstained (i.e. SFEs) by the unmodified **P₁₈** probe whereas this number was reduced to 40% for the higher affinity γ **P₁₂** (Figure 7B and Supplementary Figure 6). For those telomeres that were stained, γ **P₁₂** gave considerably brighter signals than **P₁₈**. Thus, while overall staining efficiency was reduced for both probes under non-denaturing conditions, the γ **P₁₂** probe yielded a greater percentage of telomeres with brighter signals compared to the unmodified **P₁₈** probe. Overall, these results demonstrate that, in metaphase chromosome spreads prepared from Jurkat cells, the shorter γ **P₁₂** probe exhibited significantly improved telomere staining under both denaturing and non-denaturing conditions. This finding opens potential new applications of qFISH using nondenaturing conditions, such as flow cytometry coupled with FISH. This technique is useful for examining telomere lengths in specific cell subpopulations in human blood including granulocytes, T cells and or B cells, which are identified with specific antibodies against antigens that would not withstand harsh denaturing conditions needed for hybridization of lower affinity probes.³⁶

Another potential application of γ PNA miniprobcs is in studies of telomere dysfunction, which rely on observation of co-localization of DNA damage response (DDR) proteins with telomeres, leading to inappropriate recognition as a DNA double strand break due to uncapping. These DDR markers include 53BP1 protein and the phosphorylation of histone

H2AX at the telomeres which are detected with antibodies or GFP-tagging of exogenous protein.^{37, 38} Current protocols require that cells or chromosome spreads are immunostained with DDR-specific antibodies and then fixed prior to subjecting the samples to denaturing FISH conditions for PNA staining of the telomeric DNA. γ PNA hybridization under milder non-denaturing conditions should allow more reliable detection of fluorescent protein tags or antibody-conjugated fluorophores or substrates.

Conclusion—The results described above demonstrate that γ PNA miniprobos as short as 6 nucleobases assemble cooperatively on repeating sequence DNA targets to form stable γ PNA/DNA heteroduplexes. The miniprobos successfully stained telomeric DNA in metaphase chromosome spreads. FISH results for Jurkat cells stained with the 12mer probe γ P₁₂ are particularly promising, with the observation of significantly fewer signal free ends (SFEs) and a greater number of telomeres exhibiting brighter signals, compared with a conventional 18mer PNA probe. Our findings support the hypothesis that shorter probes can produce more intense fluorescence signal and allow better sensitivity and detection of short telomeres by allowing more probes, and thus more fluorophores, to bind to the same number of DNA bases. The imaging and quantitation data show that this is a viable application of fluorescent miniprobos, although the extent of improvement depends on the experimental conditions and cell line. Future work will involve testing the miniprobos in additional cell lines to see if the performance in qFISH assays depends on the telomere maintenance mechanism (ALT versus telomerase).

Any hybridization probe requires optimization in order to maximally capture or detect its target. Short probes have the benefit of being more sensitive to single mismatches (i.e. a mismatch has a greater destabilizing effect than in a long probe), but are more likely to find perfect-match sequences at unintended sites. This is a possible reason for the less optimal performance of the 6mer miniprobos: without the G-clamp modification, the probe did not have sufficient affinity to stain telomeres under our washing conditions, whereas with two G-clamp bases, the affinity was so high that staining of other sites in the genome is likely to occur, particularly under denaturing conditions. It is also worthwhile to recognize the considerably higher amount of effort required to synthesize the G-clamp γ PNA monomer for a miniprobe that would be, at best, only twofold brighter than the 12mer version which does not need any modified bases. On the other hand, long probes are less likely to find multiple perfect-match sites but are more likely to hybridize to single-mismatch sites. Ongoing work is directed toward optimizing the length and affinity of γ PNA miniprobos to obtain maximum telomere brightness while minimizing background staining of nontelomeric genomic regions. (It is also possible that, in the absence of an RNase treatment, telomeric RNA hybridizes to PNA and γ PNA probes.)

Finally, we note that γ PNA miniprobos should be useful for chromosome or mRNA “painting”, in which sets of fluorescent probes targeted to different regions of the same chromosome³⁹ or mRNA¹⁶ are applied in combination, resulting in brighter and/or multicolor fluorescence. The short length and lack of a negative charge on the γ PNA backbone should permit faster hybridization relative to oligonucleotide or cDNA FISH probes.

METHODS

Materials

DNA oligonucleotides were purchased from Integrated DNA Technologies and used after standard desalting by gel filtration chromatography. **P₁₈** was obtained from Panagene. DNAs contained different numbers of telomeric repeats **Telo-n** (5'-TTAGGG-3')_n, where n = 1-4 or 6. γ -miniPEG-t-Boc-protected PNA monomers and oligomers were synthesized according to published procedures,^{10, 21} purified using reverse-phase high performance liquid chromatography (HPLC), and characterized by matrix-assisted laser desorption/ionization time of flight (MALDI-ToF) mass spectrometry (Applied Biosystems, Voyager DE sSTR) using α -cyano-4-hydroxycinnamic acid as the matrix. Cy3 was a generous gift from B. Schmidt of the Molecular Biosensor and Imaging Center at Carnegie Mellon University. For Cy3 labeling, a carboxylic acid derivative of Cy3 was coupled to the γ PNA prior to cleavage using similar procedures to all of the other monomers. Samples were run with linear detection and positive ionization. γ PNA characterization data are shown in Figures S7-S9.

All PNA, γ PNA and DNA concentrations were determined by measuring the absorbance at 260 nm at 90°C on a Cary 3 Bio spectrophotometer. At high temperature, all the bases are assumed to be unstacked, and the extinction coefficient can be calculated as the sum of the individual bases. Extinction coefficients for PNA monomers were obtained from Applied Biosystems whereas DNA extinction coefficients were used as reported in the literature. Calculated extinction coefficients (ϵ_{260} , M⁻¹cm⁻¹) are **P₁₈** = 167,400; γ **P₆** = 55,800; Quantitation of FISH results = 65,400 and γ **P₁₂** = 111,600.

UV Melting Experiments

UV-vis measurements were conducted on a Varian Cary 300 Bio UV-vis spectrophotometer equipped with thermoelectrically controlled multicell holders. Samples were prepared in a buffer containing 10mM Tris-HCl (pH 7) and 0.1 mM Na₂EDTA. For LiCl experiments, a stock solution of 10 mM Li₂EDTA was used. Thermal experiments were conducted in the presence of either 100 mM KCl or 100 mM LiCl. The solutions were heated to 95 °C followed by cooling to 15 °C at a rate of 1 °C min⁻¹. The samples were allowed to equilibrate for 5 minutes at 15 °C and then heated back to 95 °C at the same rate. The absorbance at 260 nm for complementary binding or 295nm for G-quadruplex formation was recorded at an interval of 0.5 °C. All samples contained concentrations of 4 μ M PNA and DNA repeat sequences, for example 2 μ M γ **P₁₂** (2 repeat sequences) + 2 μ M **Telo-2** (2 repeat sequences) or 1 μ M **Telo-4** (4 repeat sequence); 4 μ M γ **P₆** or Quantitation of FISH results (1 repeat sequence) + 4 μ M **Telo-1**, 2 μ M **Telo-2**, 1.3 μ M **Telo-3** or 1 μ M **Telo-4**. All experiments were run in triplicate and the averaged data were plotted as the fraction of melted PNA-DNA duplex as a function of temperature. Melting transition temperature (T_m), the temperature at which PNA-DNA hybrid was half-melted, was determined as described by Marky and Breslauer.⁴⁰

Fluorescence Spectroscopy

Fluorescence emission spectra were collected on Cary Eclipse Fluorescence Spectrophotometer at 25°C. Spectra for samples containing a final concentration of 4 μ M (PNA + DNA repeats) buffered in 10 mM Tris-HCl (pH=7), 0.1 mM EDTA and 100 mM KCl were recorded with excitation at 530 nm after annealing of the samples from 95-5°C.

Surface Plasmon Resonance Experiments

SPR experiments were performed on a Biacore T100 system with a four-channel sensor chip coated with carboxymethylated dextran on the gold surface. Approximately 10000 response units (RU) of streptavidin was immobilized on the chip's surface using standard N-hydroxysuccinimide/1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (NHS/EDC) coupling. Then 120 RU of 5'-biotinylated DNAs (25 nM) were attached to separate flow cells via non-covalent capture to streptavidin-coated surface. Before sample injection, sensor chips were conditioned with five consecutive 1 min-injections of 1M NaCl, 50 mM NaOH followed by multiple washing flow with LiCl/HBS-EP buffer (0.01 M HEPES, 0.15 M LiCl, 3mM EDTA, and 0.005% Surfactant P-20, pH 7.4). Biotinylated DNA in LiCl/HBS-EP buffer was then injected at a flow rate of 2 μ L/min to achieve long contact times with the surface until a sufficient amount of DNA was immobilized. Synthesized PNAs were dissolved in water and then diluted to a working stock solution of 5 μ M. Samples of varying PNA concentration were then prepared in filtered and degassed running buffers by serial dilutions (10-50 nM in 10 nM increments) from stock solutions. Samples were run in triplicate at a flow rate of 50 L/min with either LiCl/HBS-EP or KCl/HBS-EP running buffers. The sensor surface was then regenerated with two injections of 10 mM NaOH/1 M NaCl. Buffer injections were conducted to remove remaining regeneration solution before the next sample was injected. Prior to data analysis, we corrected the raw sensorgrams by performing "double referencing". Briefly, we subtracted control flow cell 1 from all sample flow cells to remove changes in RU contributed by bulk shift. Within the same flow cell, we subtracted an average of the two buffer injections from PNA injections to account for any nonspecific binding or baseline shifts due to experimental artifacts.

Cell Culture

Jurkat cells and U2OS cells were cultured in standard cell culture conditions. Cells were grown in a temperature controlled incubator at 37°C and 5% CO₂ in Dulbecco's Modified Eagle's Medium (DMEM) with 10% Fetal Bovine Serum (FBS), 50U Penicillin/mL and 50 μ g Streptomycin/mL.

Quantitative Fluorescence In-situ Hybridization (qFISH)

For staining and imaging of telomeres in metaphase chromosome spreads the cells were treated with 0.05 μ g/mL colcemid for 10 hours. 1×10^6 cells (1×10^5 cell/mL) were subsequently harvested and treated with 75 mM KCl hypotonic buffer for 12 min at 37°C and then fixed and stored at -20 °C in methanol/acetic acid fixative (3:1). Lysed cells were dropped onto water-coated glass microscope slides and dried at 42 °C for 3 minutes to spread the chromosomes and were then incubated overnight at room temperature. The prepared microscope slides were fixed in 4% formaldehyde and then treated with 0.1%

Pepsin in 0.01N HCl for 10 minutes at 37 °C. Fixation and washing were repeated. Slides were then dehydrated in successive ethanol solutions of 70%, 90%, and 100% for 5 minutes each. Slides were then treated with 500µg/mL of RNase A for 10 minutes at 37 °C followed by an additional dehydration series. Samples were denatured for 3 min at 80 °C, in a hybridization mixture containing 70% deionized formamide, unless otherwise indicated, 10% NEN blocking reagent [Roche], 0.1 M Tris-HCl [pH=7.4], MgCl₂ buffer [82 mM NaH₂PO₄, 9 mM citric acid, 20 mM MgCl₂], and 0.5 µg/ml of the indicated PNA or γPNA. Non-denaturing conditions were identical except formamide was omitted and samples were incubated at room temperature rather than 80°C. After 2h hybridization at room temperature, the samples were washed twice with wash solution (70% deionized formamide, 0.1% BSA, and 10 mM Tris-HCl [pH=7.4]), and three times with a second wash solution (100 mM Tris-HCl [pH=7.4], 150 mM NaCl, 0.1% Tween 20). Samples were finally counterstained with DAPI and mounted with coverslips.

Image Analysis

Stained metaphase spreads were imaged using a Nikon Ti90 epifluorescence microscope (Nikon Inc., NY) equipped with PlanApo 60×/1.40 oil immersion objective and run by Nikon Elements. A series of nine Z-stack 0.25 µm cross sections were captured with 80ms exposure for DAPI and a 300ms exposure time for Cy3 unless otherwise indicated. Images were converted to 8-bit .TIFF files using Image J, and optimal Z-stacks were analyzed for telomere brightness by employing TFL-Telo software.⁴¹ The telomere threshold was set to three for U2OS cells and two for Jurkat cells. The chromosome threshold was set to 9. Telomere spots and chromosomes were inspected and adjusted for accuracy as needed. Telomere fluorescence was binned by brightness, and calculated as a ratio of telomere objects per bin to total telomere objects in the metaphase using Excel (Microsoft). Error was calculated as standard error of the means from three repeats of each experiment (two repeats for nondenaturing conditions). Column graphs were generated using Origin software.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

The authors gratefully acknowledge generous support for this research from the U.S. National Institutes of Health (R43GM108187 to BAA and PLO), the National Science Foundation (CHE 1012467 to DHL) and from the DSF Charitable Foundation. SPR instrumentation was purchased with support from NSF Major Research Instrumentation Award CHE 0821296.

References

1. Bedell VM, Westcot SE, Ekker SC. Briefings Func. Genom. 2011; 10:181–188.
2. Huang P, Zhu Z, Lin S, Zhang B. J. Genet. Genom. 2012; 39:421–433.
3. Lundin KE, Hoejland T, Hansen BR, Persson R, Bramsen JB, Kjems J, Koch T, Wengel J, Smith CIE. Adv. Genet. 2013; 82:47–107. [PubMed: 23721720]
4. Egholm M, Buchardt O, Christensen L, Behrens C, Freier SM, Driver DA, Berg RH, Kim SK, Nørdén B, Nielsen PE. Nature. 1993; 365:566–568. [PubMed: 7692304]

5. Tomac S, Sarkar M, Ratilainen T, Wittung P, Nielsen PE, Nordén B, Gräslund A. *J. Am. Chem. Soc.* 1996; 118:5544–5552.
6. Lansdorp PM, Verwoerd NP, van de Rijke FM, Dragowska V, Little M-T, Dirks RW, Raap AK, Tanke HJ. *Hum. Mol. Gen.* 1996; 5:685–691. [PubMed: 8733138]
7. Stender H, Fiandaca M, Hyldig-Nielsen JJ, Coull J. *J. Microbiol. Methods.* 2002; 48:1–17. [PubMed: 11733079]
8. Dragulescu-Andrasi A, Rapireddy S, Frezza BM, Gayathri C, Gil RR, Ly DH. *J. Am. Chem. Soc.* 2006; 128:10258–10267. [PubMed: 16881656]
9. Wengel J. *Acc. Chem. Res.* 1999; 32:301–310.
10. Sahu B, Sacui I, Rapireddy S, Zanotti KJ, Bahal R, Armitage BA, Ly DH. *J. Org. Chem.* 2011; 76:5614–5627. [PubMed: 21619025]
11. Yeh JI, Shivachev B, Rapireddy S, Crawford MJ, Gil RR, Du S, Madrid M, Ly DH. *J. Am. Chem. Soc.* 2010; 132:10717–10727. [PubMed: 20681704]
12. Nielsen PE, Egholm M, Berg RH, Buchardt O. *Science.* 1991; 254:1498–1500.
13. Lohse J, Dahl O, Nielsen PE. *Proc. Natl. Acad. Sci. USA.* 1999; 96:11804–11808. [PubMed: 10518531]
14. He G, Rapireddy S, Bahal R, Sahu B, Ly DH. *J. Am. Chem. Soc.* 2009; 131:12088–12090. [PubMed: 19663424]
15. Bahal R, Sahu B, Rapireddy S, Lee CM, Ly DH. *ChemBioChem.* 2012; 13:56–60. [PubMed: 22135012]
16. Raj A, van den Bogaard P, Rifkin SA, van Oudenaarden A, Tyagi S. *Nature Methods.* 2008; 5:877–879. [PubMed: 18806792]
17. Lifland AW, Zurla C, Santangelo PJ. *Bioconjugate Chem.* 2010; 21:483–488.
18. Wilds CJ, Maier MA, Tereshko V, Manoharan M, Egli M. *Angew. Chem. Int. Ed.* 2002; 41:115–117.
19. Ortega J-A, Blas JR, Orozco M, Grandas A, Pedroso E, Robles J. *Org. Lett.* 2007; 9:4503–4506. [PubMed: 17915883]
20. Rapireddy S, Bahal R, Ly DH. *Biochemistry.* 2011; 50:3913–3918. [PubMed: 21476606]
21. Christensen L, Fitzpatrick R, Gildea B, Petersen KH, Hansen HF, Koch T, Egholm M, Buchardt O, Nielsen PE, Coull J, Berg RH. *J. Pept. Sci.* 1995; 3:175–183. [PubMed: 9222994]
22. Koch T, Nielsen PE. *Peptide Nucleic Acids: Protocols and Applications.* Horizon Bioscience, Norfolk (2nd edn.). 2004:37–60.
23. Kim SK, Nielsen PE, Egholm M, Buchardt O, Berg RH, Nordén B. *J. Am. Chem. Soc.* 1993; 115:6477–6481.
24. Gagnon KT, Watts JK, Pendergraff HM, Montmaillier C, Thai D, Potier P, Corey DR. *J. Am. Chem. Soc.* 2011; 133:8404–8407. [PubMed: 21539318]
25. Roy S, Zanotti KJ, Murphy CT, Tanious FA, Wilson WD, Ly DH, Armitage BA. *Chem. Commun.* 2011; 47:8524–8526.
26. Williamson JR, Raghuraman MK, Cech TR. *Cell.* 1989; 59:871–880. [PubMed: 2590943]
27. Sanders JL, Newman AB. *Epidemiol. Rev.* 2013; 35:112–131. [PubMed: 23302541]
28. Aviv A. *J. Gerontol. A Biol. Sci. Med. Sci.* 2008; 63:979–983. [PubMed: 18840804]
29. Poon SS, Lansdorp PM. *Curr. Protoc. Cell Biol.* 2001 Chapter 18, Unit 18 14.
30. Cesare AJ, Reddel RR. *Nat. Rev. Genet.* 2010; 11:319–330. [PubMed: 20351727]
31. Bryan TM, Englezou A, Dalla-Pozza L, Dunham MA, Reddel RR. *Nature Med.* 1997; 3:1271–1274. [PubMed: 9359704]
32. Sarzotti-Kelsoe M, Daniell XG, Whitesides JF, Buckley RH. *Immunol. Res.* 2011; 49:44–48. [PubMed: 21120634]
33. Wang F, Pan X, Kalmbach K, Seth-Smith ML, Ye X, Antunes DMF, Yin Y, Liu L, Keefe DL, Weissman SM. *Proc. Natl. Acad. Sci. USA.* 2013; 110:E1906–E1912. [PubMed: 23661059]
34. Liu F-J, Barchowsky A, Opresko PL. *PLoS One.* 2010:5.
35. Coleman J, Baird DM, Royle NJ. *Hum. Mol. Gen.* 1999; 8:1637–1646. [PubMed: 10441326]

36. Baerlocher GM, Vulto I, de Jong G, Lansdorp PM. *Nature Protocols*. 2006; 1:2365–2376. [PubMed: 17406480]
37. Cesare AJ, Kaul Z, Cohen SB, Napier CE, Pickett HA, Neumann AA, Reddel RR. *Nature Struct. Mol. Biol.* 2009; 16:1244–1251. [PubMed: 19935685]
38. Takai H, Smogorzewska A, de Lange T. *Curr. Biol.* 2003; 13:1549–1556. [PubMed: 12956959]
39. Ried T, Schröck E, Ning Y, Wienberg J. *Hum. Mol. Gen.* 1998; 7:1619–1626. [PubMed: 9735383]
40. Marky LA, Breslauer KJ. *Biopolymers*. 1987; 26:1601–1620. [PubMed: 3663875]
41. Poon, SS.; Lansdorp, PM. *Methods in Cell Biology: Flow Cytometry*. 64 edn.. Darzynkiewicz, Z.; Crissman, HA.; Robinson, JP., editors. Vol. 64. Academic Press; San Diego, CA: 2001. p. 69-96.

Chart 1.
Chemical structures of PNA, γ PNA and the guanidino G-clamp nucleobase.

Scheme 1.

Fig. 1. UV melting curves recorded for 1:1 duplexes between PNA or γ PNA and the complementary DNA strands **Telo-1**, **Telo-2** or **Telo-3**. Samples contained 4 μ M each of PNA/ γ PNA and DNA repeat sequences, as explained in the Experimental section.

Fig. 2. Fluorescence emission spectra recorded for samples containing 20 nM **Telo-6** DNA and 40 nM **P₁₈**, 60 nM γP_{12} or 120 nM $\gamma P_{6\text{-clamp}}$.

Fig. 3. SPR sensorgrams recorded for association and dissociation of γ P₆, γ P₆-clamp or γ P₁₂ to immobilized DNA targets. 10 nM γ PNA was flowed over the SPR chip during the association phase (400 s) before switching to γ PNA-free buffer to promote dissociation.

Fig. 4. SPR sensorgrams for association and dissociation of 50 nM γP_6 , $\gamma P_{6-clamp}$ and γP_{12} to immobilized. **Telo-2**

Fig. 5.

Telomere staining of U2OS metaphase chromosome spreads using Cy3-labeled PNA and γ PNA probes. Staining by fluorescent *in situ* hybridization under denaturing conditions was achieved using **P₁₈** (left), **γ P₁₂** (center), and **γ P_{6-clamp}** (right).

Fig. 6.

Quantitation of FISH results for U2OS cells. Percent values represent the average of three independent experiments. Error bars represent standard error of the mean. Bars represent fluorescence values quantified by bins of 30,000 arbitrary fluorescence units (dashed box indicates inset).

Fig. 7.

Quantitative analysis of telomere staining in Jurkat cells by fluorescent PNA and γ PNA probes under (A) denaturing and (B) non-denaturing conditions. SFE = signal-free ends. Percent values represent the average of three independent experiments. Error bars represent standard error of the mean. Bars represent fluorescence values quantified by bins of 10,000 arbitrary fluorescence units (dashed box indicates inset).

Table 1Sequences of PNA, γ PNA and DNA oligomers used in this study.

Name	Sequence [†]
P₁₈	H ₂ N-Lys-(AATCCC) ₃ -Cy3
γP₁₂	H ₂ N-Lys-(AATCCC) ₂ -Cy3
γP₆	H ₂ N-Lys-AATCCC-Cy3
γP_{6-clamp} [§]	H ₂ N- Lys-AAT <u>X</u> C <u>X</u> -Cy3
Telo-<i>n</i> (DNA)	5'-(TTAGGG) _{<i>n</i>} -3'

[†] PNA and γ PNA sequences are written in the C-to-N direction.[§] X = G-clamp nucleobases

Table 2

UV melting temperature (T_m , °C) values recorded for PNA- and γ PNA-DNA duplexes.

	P_{18}	γP_{12}	γP_6	$\gamma P_{6\text{-clamp}}$
Telo-1	<i>nd</i>	<i>nd</i>	24.3 ± 1.7	62.3 ± 1.1
Telo-2	<i>nd</i>	77.3 ± 3.2	42.8 ± 1.9	72.4 ± 0.4
Telo-3	66.0	<i>nd</i>	38.4 ± 0.7	78.4 ± 0.3
Telo-4	<i>nd</i>	82.5 ± 0.3	38.2 ± 1.3	81.3 ± 0.7

nd = not determined

Table 3

Initial on-rates and off-rates (RU/s) of duplexes formed in 150 mM KCl. (Values in parentheses are normalized to the γP_{12} data for each column.)

	Telo-1		Telo-2		Telo-4	
	<i>On-rate</i>	<i>Off-rate</i>	<i>On-rate</i>	<i>Off-rate</i>	<i>On-rate</i>	<i>Off-rate</i>
γP_6	0	0	0.768 (0.8)	0.427 (33)	0.117 (0.7)	0.68 (34)
$\gamma P_{6\text{-clamp}}$	0.789	0.250	1.016 (1.0)	0.036 (2.7)	0.276 (1.8)	0.011 (5.5)
γP_{12}			1.015 (1.0)	0.013 (1.0)	0.155 (1.0)	0.002 (1.0)