Published in final edited form as:

Physiol Behav. 2014 September ; 136: 145–154. doi:10.1016/j.physbeh.2014.03.002.

Is fat taste ready for primetime?

Nicholas V. DiPatrizio¹

¹Department of Anatomy and Neurobiology, University of California, Irvine, School of Medicine

Abstract

Mounting evidence suggests that gustation is important for the orosensory detection of dietary fats, and might contribute to preferences that humans, rodents, and possibly other mammals exhibit for fat-rich foods. In contrast to sweet, sour, salty, bitter, and umami, fat is not widely recognized as a primary taste quality. Recent investigations, however, provide a wealth of information that is helping to elucidate the specific molecular, cellular, and neural mechanisms required for fat detection in mammals. The latest evidence supporting a fat taste will be explored in this review, with a particular focus on recent studies that suggest a surprising role for gut-brain endocannabinoid signaling in controlling intake and preference for fats based on their proposed taste properties.

Keywords

fat; taste; endocannabinoid; food reward; periphery

1. Introduction

Flavor is comprised of several sensory modalities, which include gustatory (i.e., taste), somatosensory (e.g., tactile), and olfactory components [1]. Primary taste qualities are widely agreed to include sweet, sour, salty, bitter, and recently, umami [2]. The classification of fat as a primary taste, however, is slow to gain acceptance, largely due to disagreement on what constitutes a “fatty” taste (see for review [3, 4]). In possibly reductionist terms, primary tastes are generally defined by A) dedicated receptor mechanisms and cells (i.e., taste receptor cells) capable of transducing a discrete chemical signal into an electrical signal, B) distinct sensations unique to a particular stimuli, C) specific neural pathways that transmit electrical signals from the oral cavity to the brainstem, and recently extended by Richard Mattes to include several other elements [4] (see Table 1). Based on the evidence discussed in this review, fat is proposed to meet these basic criteria for classification as a primary taste quality.

¹Corresponding author: 3216 Gillespie Neuroscience Research Facility, Irvine, California, 92697-1275, Phone: 949-824-7080, ndipatri@uci.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Gustatory signals from food are detected by taste cells localized to the chemosensory organs [5], the taste buds, within distinct papillae on the tongue [3], and recent studies suggest several candidate fat taste receptors in taste cells [6–10]. Taste buds are also located in the pharynx and larynx [11]. Taste cells are further categorized [5] into three major types: type 1 cells, which, in addition to possibly functioning as taste receptor cells (see [12]), might act to maintain the local environment—similar to glial cells—by degrading neurotransmitters released from other cells; type 2 cells that contain G-protein coupled receptors for sweet, bitter and umami, and release ATP, which is likely a major source of activation of associated gustatory afferent nerves via interactions with P2X2/3 receptors located on these nerve fibers [13]; type 3 cells, which are presynaptic, respond to sour stimuli, possess conventional synapses, and release the neurotransmitters, norepinephrine, serotonin, and GABA. Fungiform and foliate papillae are located on the anterior two-thirds and lateral regions of the tongue, respectively [5]. Taste buds in fungiform papillae are innervated by the chorda tympani branch of the facial nerve (VII), while taste buds in foliate papillae are primarily innervated by the lingual branch of the glossopharyngeal nerve (IX). Circumvallate papillae located on the posterior one-third of the tongue are also innervated by the lingual branch of IX. Branches of the vagus nerve (X) carry afferent taste signals from the laryngeal epithelium to the brainstem. Gustatory neurotransmission from the oral cavity is carried by cranial nerves VII, IX, and X to the nucleus of the solitary tract in the caudal brainstem, which in turn, sends fibers to the parabrachial nucleus in the pons (in rodents) and communicates bi-directionally with numerous forebrain regions and peripheral organs [14–16]. Somatosensory (e.g., fat texture) signals from food are transmitted to the brainstem by trigeminal (V) neurons and IX, [17, 18]. Importantly, increasing evidence suggests that gustatory neural pathways are critical for preferences that rodents display for fatty foods: surgical disruption of afferent neurotransmission from the tongue to the brainstem, carried by the chorda tympani or glossopharyngeal nerves, reduces intake and preference for fats [19–22]. Furthermore, gustatory signals from dietary fats and sugars are suggested to be important for maintaining food reward [15], which shares neurobiological commonalities with drug addiction [23–26], and might contribute to obesity. Indeed, obese humans, when compared to normal weight subjects, significantly prefer, crave, and consume foods containing high levels of fat [27–30].

Endocannabinoids are a class of lipid-derived endogenous signaling molecules that are synthesized in most mammalian tissues, and are proposed to drive the seeking and sensing of energy-dense foods, and the accumulation of their energy content for future utilization (see [15]). Recent studies reveal that the endocannabinoids act in concert with taste pathways to control the intake of palatable foods [15, 31–33]. An increasing body of work suggests a significant role in feeding for post-ingestive cues derived from dietary fats (see for review [34–36]); however, this review will focus on preabsorptive cephalic mechanisms and evidence for fat as a primary taste quality. Moreover, new studies are discussed that support an important function for gut-brain endocannabinoid signaling in the gustatory control of dietary fat intake, preference, and possibly obesity.

2. Cephalic components of fat intake

2.1. Orosensory fat exposure and dopamine signaling

It is well established that mammals display robust preferences for fat-rich diets [37]. Several laboratories demonstrated that rodents maintain ingestion of fatty diets when post-ingestive influences are eliminated by sham feeding, which isolates the orosensory component of feeding from its post-ingestive consequences [37, 38] (see Figure 1 for details of the sham feeding paradigm in rat). Sham feeding rats corn oil (100%) increased levels of dopamine within the nucleus accumbens [39], a brain region that is essential for processing food reward [40, 41]. This effect is not isolated to the ventral striatum: recent work revealed that tasting dietary fat drives dopamine release in the basolateral amygdala (BLA)[42]. For these studies, rats were trained to develop a preference for solutions containing corn oil. After a preference was established, the rodents were presented with corn oil (100%), a low concentration of linoleic acid (18:2 FA; 1%), or non-nutritive mineral oil (100%; has similar textural properties to that of nutritive oils, but does not contain any fatty acids), and dopamine levels were measured in the nucleus accumbens and BLA via microdialysis. Oral exposure to corn oil or linoleic acid increased DA levels in both brain regions within 10 minutes after access to the test diets, whereas mineral oil failed to elicit a similar response. The short latency for these effects after access to the nutritive oils suggests that oral cues from dietary fats are primarily responsible for enhanced dopamine outflow, and possibly preference. Supporting this notion, earlier studies in mice confirmed that post-oral administration of corn oil into the duodenum elicited a robust place preference only when paired with concurrent oral exposure to oils [43].

2.2. Cephalic-phase feeding and peripheral cholinergic signaling

Studies by Gerard P. Smith and colleagues suggest that a peripheral mechanism is responsible for maintaining intake based on the orosensory qualities of foods [44]. Treatment with the peripherally-restricted muscarinic acetylcholine receptor antagonist, atropine methyl nitrate, inhibited sham feeding of a palatable liquid diet. Earlier studies found that atropine methyl nitrate inhibited normal food intake [45]; however, anticholinergic drugs were thought to possibly influence food intake indirectly by disrupting saliva secretion and swallowing. Smith and colleagues treated rats with atropine methyl nitrate prior to the sham feeding of water and found no changes in its sham intake [44]. Together, the results suggest that inhibiting peripheral muscarinic acetylcholine receptors selectively reduces intake of foods with nutritional value by a mechanism that does not include a general disruption in swallowing. Specific peripheral targets for muscarinic acetylcholine receptor antagonism and downstream mechanisms that influence intake remain to be determined. Recent evidence, however, suggests that endocannabinoid signals in the proximal small intestine, driven by vagal cholinergic activity, might contribute to fat sham feeding based on the orosensory qualities of fats [38, 46].

2.3. Orosensory fat exposure and peripheral endocannabinoid signaling

Increasing evidence reveals a significant role for the endocannabinoid system in controlling the intake of palatable foods [15, 47], and some of our recent studies suggest that gut-brain endocannabinoid signaling controls the intake of dietary fats based on their orosensory

properties [38]. Individual groups of rats were sham fed separate macronutrients, followed by collection of brain and other organs for analysis of endocannabinoid content by high-performance liquid chromatography/mass spectrometry [38]. Rats were sham fed for 30 minutes at a fixed concentration of 10mL in order to ensure that all animals had equal oral exposure to the test diets. Sham feeding either a corn oil emulsion at a concentration that is readily sham fed by rats (25% vol/vol [48]), or the nutritionally-complete liquid test diet, Ensure (vanilla; 25% vol/vol), robustly enhanced levels of the two main endocannabinoids, 2-arachidonoyl-*sn*-glycerol and anandamide, in the jejunum. These effects were tissue-specific, because tasting corn oil failed to significantly affect endocannabinoid levels in other peripheral organs included in our evaluation (i.e., tongue, stomach, ileum, liver, pancreas) or throughout the brain (i.e., parabrachial nucleus, dorsal striatum, ventral striatum, medial aspects of the hypothalamus, lateral aspects of the hypothalamus, cerebellum). Importantly, cephalic-phase gut endocannabinoid signaling was macronutrient specific: sham feeding carbohydrate or protein at concentrations (8% w/v) that are readily sham fed by rats in other studies [49] did not affect endocannabinoid levels in the jejunum [38]. Furthermore, full subdiaphragmatic vagotomy completely blocked endocannabinoid signaling in the jejunum after tasting corn oil, which suggests that critical oral signals from fat are transmitted to the gut by cholinergic neurotransmission carried by efferent vagal fibers.

We next evaluated the physiological relevance for this macronutrient- and organ-specific signaling event. Twenty minutes before presentation of the corn oil emulsion, rats were infused with low concentrations of the cannabinoid CB₁ receptor antagonist/inverse agonist, rimonabant (0.3–1.0 mg/kg), directly into the small intestine through an intraduodenal catheter. Rimonabant dose-dependently blocked the 30 min sham intake of corn oil. A similar reduction in sham fat intake was observed following pretreatment with the non-brain penetrant neutral CB₁ receptor antagonist, URB 447, which supports a peripheral mechanism in the response. In contrast to corn oil, a higher dose of rimonabant (1mg/kg) was required to affect the intake of standard rodent chow. This result is likely due to the brain-penetrant rimonabant, while administered locally into the intestine, gaining access at higher concentrations to central CB₁Rs that control feeding, and thus, reducing the intake of standard chow. For this reason, peripheralized neutral CB₁R antagonists were used in subsequent studies. Collectively, the results suggest that gustatory exposure to dietary fats, but not protein or carbohydrate, initiates endocannabinoid signaling in the small intestine, but not other organs, and this local signaling event might provide positive feedback to the brain to maintain the intake of palatable fatty foods.

Cholinergic signals transmitted by the efferent vagus nerve are necessary for induction of endocannabinoid signaling in the gut; however, specific constituents of the cholinergic system (e.g., muscarinic acetylcholine receptors) that are responsible for initiating production of the endocannabinoids are unknown and currently under investigation. Furthermore, CB₁R activation in the gut might drive feeding by modifying satiation and/or hunger signaling communicated to the brain by vagal afferent fibers [50, 51]. Recent evidence from rodents supports this hypothesis: CB₁R mRNA was found in enteroendocrine I cells of the proximal small intestine, which produce and secrete the satiation hormone,

cholecystokinin [52], and CB₁Rs in the stomach control the release of the hunger signal, ghrelin [53].

3. Orosensory qualities of fats, and the endocannabinoids

3.1. Olfaction

Despite the role for olfaction in some aspects of feeding, evidence from a variety of labs suggests that olfactory cues are not necessary for the orosensory detection of dietary fats in humans and rodents. Humans reliably detect a wide range of short, medium, and long-chain free fatty acids when olfaction is eliminated through use of nose clips [54]. Rodents rendered anosmic by either zinc sulfate perfusion through the nasal cavity, or surgical ablation of olfactory signaling, resulted in only mild reductions in preference for nutritive oils at low concentrations in two-bottle choice tests [55–57]. Furthermore, zinc sulfate failed to affect conditioned place preferences that mice exhibit for corn oil [57]. Together, the data suggest that olfactory cues, while likely important, are dispensable for the detection of dietary fats.

3.2. Gustation

Increasing evidence from rodents and humans suggests that gustatory signals derived from free fatty acids are important for the detection of dietary fats in the oral cavity (see for review [36, 58–62]). When given the option of consuming emulsions that contain nutritive oils (e.g., corn oil or 18:2 FA), or those containing a non-nutritive mineral oil, rodents reliably prefer those containing nutritive oils [46, 48, 63, 64]. Free fatty acids, however, do not comprise a significant portion of dietary lipids from foods. Instead, fats are typically found in the form of triacylglycerols and phospholipids. These lipids release a portion of their fatty acid component after hydrolysis by lingual lipases secreted from glands (e.g., von Ebner's glands) located in the tongue and oral cavity in rodents [65, 66] and humans [67, 68]. Prevention of hydrolysis with an inhibitor of lingual lipase activity, tetrahydrolipstatin, blocked preferences for triacylglycerols, but not free fatty acids, in a rodent two-bottle choice test [69], and substantially increased detection thresholds in humans [70]. The results strongly suggest that free fatty acids are detected in the oral cavity, and their liberation by hydrolysis might contribute to fat detection. Nonetheless, low levels of free fatty acids are found in some foods (e.g., nuts) and might serve as gustatory cues [71, 72]. Indeed, a recent study in humans revealed that when palatable fatty foods (e.g., walnuts, olive oil) were masticated for one minute then expectorated, free fatty acid levels in saliva reached μ molar concentrations, which are likely adequate to drive taste receptor signaling [71]. The source of the free fatty acids, however, was not determined and could have been derived from the fat source directly, or released from the triacylglycerols contained in the food after hydrolysis by lingual lipase. Lingual lipase in humans is reported to have low activity [73] when compared to rodents [69]. Thus, these tests should be repeated with the addition of a lingual lipase inhibitor in order to more definitively ascertain the degree to which the hydrolysis of triacylglycerols is responsible for the free fatty acids found in human saliva after mastication of fatty foods.

In contrast to rodents, which readily consume and prefer emulsions containing free fatty acids, humans are reported to find free fatty acids unpleasant [4] despite a clear attraction to fat-rich foods [3, 74]. Furthermore, it has been suggested that human preferences for fatty foods are largely supported by the textural qualities of fats (e.g., creaminess attributed to triacylglycerols), and that free fatty acids might serve as an alarm signal for the presence of oxidized unsaturated free fatty acids (e.g., rancid meat), which are associated with several pathologies when consumed [4]. Thus, potential species differences should be noted when considering the generalization of studies in rodents to humans; however, given the evidence [71], it is plausible—but not definitive—that free fatty acids contribute to the robust preferences that humans display for fatty foods.

3.3. Fat texture, free fatty acids, and peripheral endocannabinoid signaling

Tactile qualities of fats are considered important for the perception of a fat flavor in humans [75, 76] and likely other mammals. We explored the role of texture and the fatty acid content of dietary fats in driving gut endocannabinoid signaling in rat [46]. Confirming our previous studies [38], sham feeding a small fixed quantity of a corn oil emulsion for 30 min (25% vol/vol, 10mL) increased jejunal endocannabinoid levels; however, sham feeding mineral oil (25% vol/vol, 10mL) failed to elicit the same response [46]. This result suggests that tactile qualities of oils alone are not sufficient for induction of gut endocannabinoid signaling. We next asked what specific fatty acids are responsible for cephalic-phase gut endocannabinoid signaling. We sham fed rats a variety of free fatty acids with a fixed carbon chain length of 18 carbons, and increasing levels of unsaturation. In order to mask tactile cues from the fatty acids, we maintained a fixed concentration of mineral oil (20% vol/vol) and added specific free fatty acids (5% vol/vol) to this emulsion (10mL). Sham feeding the monounsaturated fatty acid, oleic acid (18:1 FA), or the diunsaturated fatty acid, linoleic acid (18:2 FA), increased endocannabinoid levels in the jejunum by greater than two-fold. In contrast to 18:1 FA and 18:2 FA, sham feeding the saturated fatty acid, stearic acid (18:0 FA), or the polyunsaturated fatty acid, linolenic fatty acid (18:3 FA), failed to increase jejunal endocannabinoid levels. A failure for 18:3 FA to drive endocannabinoid signaling in the jejunum was surprising and warrants further study given that this unsaturated fat drives pancreatobiliary secretion [6] and inhibits delayed rectifying K⁺ channels (DRK) in rat taste cells [77]. Nonetheless, the specificity for certain fatty acids to drive endocannabinoid signaling in the jejunum provides additional support for a receptor-mediated event in the oral cavity that mediates fat taste. In order to control for possible post-ingestive influences for 18:2 FA on jejunal endocannabinoid production during sham feeding (i.e., leakage of a small amount of emulsion into intestine), animals were fitted with intraduodenal catheters and infused with the same amount of 18:2 FA emulsion that drove intestinal endocannabinoid signaling during sham feeding. In contrast to sham feeding, post-ingestive exposure to 18:2 FA failed to increase endocannabinoid levels. This result strongly supports a cephalic mechanism that drives jejunal endocannabinoid signaling during sham feeding of 18:2 FA, rather than post-ingestive consequences.

An increasing source of cheap food energy in the Western diet throughout the 20th and into the 21st century includes corn and soybean based diets, which contain mostly 18:2 FA (50–60% total content; 18:1 FA, 20–30% total content) [78, 79], and are readily preferred by

rodents [80]. Thus, it is plausible that—similar to rodents [38, 46]—attraction to foods rich in 18:2 FA (e.g., french fries or potato chips) is also driven by gut-brain endocannabinoid signaling in humans, and might contribute to obesity. Indeed, the concentration of 18:2 FA in the Western diet has increased over the 20th century from 1% of total energy consumption to 8%, and this increase positively correlates with obesity rates throughout this time period [81]. Furthermore, polyunsaturated fatty acids (e.g., 18:2 FA) are preferentially oxidized, and when consumed, have been suggested to lead to several other pathologies, including atherosclerosis and some cancers (see for review [4]). Experiments in rodents support possible causation for consuming diets high in 18:2 FA and obesity. Mice maintained for 14 weeks on a diet in which total energy from 18:2 FA was increased from 1% to 8% exhibited significantly increased food intake and adiposity, when compared to mice that were maintained on the control diet [81]. The experimental diet was formulated so that increases in 18:2 FA from 1% to 8% were met with reciprocal decreases in saturated fat content in order to keep the diets equicaloric. Therefore, the results suggest that the 18:2 FA content of the diet, but not absolute levels of fats, led to the obese phenotype. Early studies in humans also suggest a similar obesogenic response to high levels of 18:2 FA [82]. In contrast to elderly male control subjects maintained on a standard cafeteria diet (40% total energy from fats), subjects chronically fed an equicaloric diet enriched with 18:2 FA (fat energy mostly from the addition of vegetable oil), at the expense of saturated fats, displayed increases in the levels of 18:2 FA in adipose tissue that positively correlated with a modest body weight gain. Together, the studies suggest that diets high in 18:2 FA, but not necessarily saturated fats, might drive obesity. Other investigations in humans, however, reveal a greater obesogenic response to saturated versus unsaturated fats (see for review [83]). Furthermore, disruptions in energy balance (e.g., increased energy intake versus expenditure), rather than the specific source of energy, is widely regarded to be primarily responsible for changes in body weight (see for review [84]). Thus, further work will be critical to delineate the biological underpinnings that govern body weight, including the role for endocannabinoid signaling in these functions.

3.4. Dietary fat preference and peripheral endocannabinoids

Rodents reliably exhibit robust preferences for nutritive oils, including corn oil or 18:2 FA, over mineral oil when given the choice in two-bottle choice tests [48, 63, 64]. In light of our results that revealed an enhancement in jejunal endocannabinoid signaling after tasting 18:2 FA, but not mineral oil [46], we next tested the hypothesis that endocannabinoid signaling at CB₁Rs in the gut contributes to the preferences that rodents exhibit for fat-rich diets containing high levels of 18:2 FA (e.g., corn oil). We confirmed that rats strongly prefer emulsions containing 18:2 FA (5% + 20% vol/vol mineral oil) to those containing only mineral oil (25% vol/vol) in the 30 min sham two-bottle choice test [46]. We then pretreated separate groups of sham feeding rats with the peripherally restricted CB₁R antagonists, AM6545 [85–87], or URB447 [38, 88], and presented 18:2 FA and mineral oil emulsions in the sham two-bottle choice test. Blocking peripheral CB₁Rs inhibited the robust preferences displayed for 18:2 FA versus mineral oil alone. We next gave rats the choice between sham feeding two test diets that failed to increase endocannabinoid content in the jejunum, 18:0 FA (5% w/vol + 25% vol/vol mineral oil) and mineral oil (25% vol/vol) [46]. Surprisingly, rats preferred mineral oil to 18:0 FA, and this preference for mineral oil was not affected by

pretreatment with AM6545. Collectively, the results support the hypothesis that tasting 18:2 FA drives endocannabinoid signaling in the gut and this signal is critical for the preferences that rats exhibit for fat-rich foods. Furthermore, the lack of effect for AM6545 on preferences for mineral oil versus 18:0 FA provides additional support for specific fatty acid signals in the oral cavity that drive gut endocannabinoid-mediated preference.

Several lines of investigation reveal that long chain unsaturated fatty acids (LCUFAs) activate taste receptor cells (see for review [4, 89]). Evidence for a similar response in taste cells to saturated fats, on the other hand, is lacking. Saturated fats failed to, both, elicit endocannabinoid signaling in the gut in sham feeding rats [46], and drive pancreatobiliary secretions following application to the tongue in rats with esophageal ligation [6]. In addition, saturated fatty acids applied to isolated rat fungiform papillae failed to induce a prolonged depolarization by the inhibition of DRK channels that occurs in the presence of most LCUFAs [90]. Nonetheless, humans are reported to detect saturated fats [91, 92] and only female rats, when compared to male, were able to reliably avoid the saturated free fatty acid, lauric acid (12:0 FA), in a conditioned taste aversion paradigm [93]. The saturated fatty acid, myristic acid (14:0 FA), but not caproic acid (6:0 FA), increased intracellular Ca^{2+} levels in cultured rat trigeminal neurons [17]. Moreover, the saturated fats, caprylic acid (8:0 FA) or capric acid (10:0 FA), induced signaling within HEK 293T cells transfected with human GPR40 or GPR120, albeit mostly to a lesser magnitude than for LCUFAs [94]. Further studies will be critical to delineate potential differential responses to free fatty acids with differing levels of unsaturation on signaling in specific taste cell subtypes. In addition, due to discrepant results between humans and rodent studies, and male and female rodents, it will be important to address potential species and sex differences for the detection of specific fatty acids.

3.5. Endocannabinoids and the tongue

Evidence from the Ninomiya group suggests that endocannabinoid mechanisms in the oral cavity control sensory neurotransmission from sweet taste [95]. CB_1R protein was found in fungiform and circumvallate papillae on the mouse tongue, where almost two-thirds of type 2 cells containing CB_1Rs also co-localized with the sweet taste receptor component, T1r3. [95]. Peripheral administration of endocannabinoids enhanced chorda tympani neural responses to sweeteners, but not sour, bitter, salty, or umami. This effect was also found *in vitro* for isolated taste cells in the presence of endocannabinoids, and was absent in mice lacking cannabinoid CB_1Rs , which implicates endocannabinoid signaling at CB_1Rs in these responses. Together, the evidence reveals that endocannabinoids act at CB_1Rs on taste cells and might mediate the perception of sweet taste by controlling its associated neurotransmission into the brainstem.

A direct test for CB_1R activation in the tongue and neural responses to fatty acids has not been reported to date. Sham feeding of corn oil, however, failed to significantly increase levels of the endocannabinoids in the whole tongue in rat [38]. The tongue contains several subtypes of taste cells [3, 5]. Accordingly, it is possible that tasting fat might trigger endocannabinoid signaling at CB_1Rs in distinct subpopulations of taste cells on the tongue, which might in turn, control afferent neurotransmission associated with fat taste. A recent

study found elevations in endocannabinoid levels in the saliva of obese fasting humans when compared to normal weight subjects, and endocannabinoid levels were positively correlated with waist circumference and body mass index (BMI) [96]. While these results are preliminary, it is plausible that alterations in endocannabinoid signaling in the oral cavity under conditions of obesity might contribute to the dysregulation of oral sensitivities to fatty acids observed in obese rodents and humans [73, 97–100]. Further investigations, however, will be necessary to identify a direct role for endocannabinoids in the oral cavity in taste signaling and their role in obesity.

4. Proposed fatty acid receptors

Several proteins located on taste receptor cells are important for dietary fat detection in rodent, including the membrane glycoprotein protein CD 36 [6–8, 101], the G-protein coupled receptors GPR40 and GPR120 [9], and the monovalent cation-selective channel, transient receptor potential channel type M5 (TRPM5) [10]. Genetic deletion of each of these proteins attenuates preferences that rodents display for LCUFAs [6, 7, 9, 10].

4.1. Taste cell signaling

Basic criteria for a fat taste, defined by Richard Mattes (see Table 1 [4]), include a defined class of effective stimuli, and unique transduction mechanisms. Gilbertson and colleagues reported that the application of LCUFAs with two or more double bonds in the *cis* configuration (e.g., 18:2 FA, 20:4 FA), but not saturated fatty acids or LCUFAs with their double bonds in the *trans* configuration, to isolated rat fungiform papillae induced a prolonged stimulus-induced depolarization via the inhibition of DRK channels [90]. Later studies revealed that taste cells contain a variety of subtypes of DRK channels, with a particularly high expression of mRNA for the Kv1.5 subtype in the anterior tongue in rat [102]. Together, the evidence might represent one component of intracellular transduction for fat taste signaling. Sweet and sour tastes, however, also reduce K⁺ current and depolarize taste cells in mudpuppy and hamster [103, 104]. Therefore, DRK responses might not be unique to fatty acids, but might represent a convergent downstream mechanism involved in taste more generally. Potential differences between species, however, are possible and should be considered when interpreting the results.

The Besnard group reported that application of LCUFAs to isolated mouse taste cells containing CD36 induced a large and rapid rise in Ca²⁺ that included the production of inositol triphosphate, which triggers the release of Ca²⁺ from intracellular pools, and suggests a G-protein coupled receptor-mediated event [105]. CD36 are not G-protein coupled, however, raising the question of how they transduce a signal that includes the activation of phospholipase c pathways (i.e., inositol triphosphate production)[10]. An influx of extracellular Ca²⁺ in response to fatty acid signaling is also thought to occur by phosphorylation of Src-protein-tyrosine kinases (Src-PTK), which opens store-operated calcium channels (SOCs) [105]. The C-terminal tail of CD36 is proposed to interact with Src-PTK, thus, providing a mechanism that couples the interactions of LCUFAs with CD36 on the taste cell membrane to subsequent intracellular signaling events [106]. Furthermore, stromal interaction molecule 1 (STIM1), which is a Ca²⁺ sensor involved in the activation of SOCs, might be critical for the opening of SOCs in taste cells in response to LCUFAs [107].

Preferences for LCUFAs were abolished in STIM-1 knockout mice, which suggests the importance for STIM-1 in the signaling cascade that drives preferences for fatty acids. Interestingly, intracellular signaling cascades initiated by LCUFAs *in vitro* in isolated mouse circumvallate taste cells were met with the release of the monoamine neurotransmitters, 5-hydroxytryptamine and noradrenalin, into the culture medium [105]. This event was dependent on the opening of SOC channels and required STIM-1 [107], a result that highlights possible neurotransmitter signals by which taste bud cells communicate fat taste to afferent gustatory nerves and the brainstem. Other possible signaling molecules involved in fat taste cannot be excluded [5], including glucagon-like peptide-1 (GLP-1), which is expressed by circumvallate papillae, and recently shown to be important for small quantities of LCUFAs to reinforce taste responses to sucrose in mice [108]. GPR120 was also implicated in the actions of GLP-1, which included the release of GLP-1 by circumvallate papillae into the culture medium in the presence of a specific GPR120 agonist, GSK137647A [108]. Furthermore, mice lacking GLP-1 receptors failed to elicit preferences for low concentrations of LCUFAs in a 12-hour two-bottle choice test. Additional investigations, however, will be critical to definitively identify the specific neurotransmitter(s) that link taste cell activation by fatty acids with neural signaling to the brainstem by gustatory afferents.

4.2. CD36

CD36 has gained considerable experimental attention as a plausible candidate for the elusive fat receptor(s) [101]. CD36 is a multifunctional glycoprotein that is present in a variety of tissues in the body, including the tongue [101]. Immunoreactivity for CD36 was found in rodent and human circumvallate and foliate papillae, with high concentrations of CD36 localized to the apical side of taste buds in lingual epithelium [6, 8, 109]. Recent studies utilized RNAi to selectively reduce the expression of CD36 on the tongue in obesity-prone Osborne-Mendel and obesity-resistant S5B/Pl rats [110]. This local genetic modification reduced preferences for the most preferred concentration of 18:2 FA in the 48-hour two-bottle choice test.

Besnard and colleagues reported that application of 18:2 FA to the tongue of anaesthetized mice and rats with surgical ligation of the esophagus, which minimizes the chances for post-ingestive influences, induced increases in pancreaticobiliary secretion and protein content [6]. In contrast to wild-type, mice lacking CD36 receptors failed to exhibit both, enhanced gastric secretions following application of 18:2 FA to the tongue, and preferences for LCUFAs. Notably, secretion mostly occurred in wild-type with application of LCUFAs (i.e., 18:1 FA, 18:2 FA, 18:3 FA), but was absent for saturated fats (18:0 FA) and short-chain saturated fatty acids (caprylic acid, 10:0 FA). A lack of effect for saturated fats under these conditions resembles our recent data showing that, in contrast to most 18 carbon unsaturated fatty acids tested, sham feeding 18:0 FA also failed to elicit endocannabinoid accumulation in the rat jejunum [46]. Together, the results suggest selectivity of cephalic-phase responses for specific fatty acids in rodents, which provides further evidence for a receptor-mediated signaling event unique to LCUFAs. Limitations of esophageal ligation to isolate taste, however, include the possibility of disrupting neurotransmission carried by branches of the vagus nerve, which run along the esophagus below the diaphragm and have recently been

shown to be required for oral exposure of dietary fats to initiate gut endocannabinoid signaling [38]. Thus, the sham-feeding model in behaving rodents, which does not carry the same risk of modifying vagal neurotransmission, might be beneficial for investigating related cephalic-phase responses (see Figure 1).

4.3. TRPM5

TRPM5 are activated by elevated levels of intracellular Ca^{2+} , which results in depolarization of the cell membrane [5], and recent studies suggest that TRPM5 are important for fat taste [10]. Supporting this role for TRPM5, application of the TRPM5 channel blocker, triphenylphosphine oxide, to taste bud cells from mouse fungiform and circumvallate papillae blunted their activation by 18:2 FA [10]. Calcium increases and inward current induced by 18:2 FA were also both reduced in taste bud cells harvested from TRPM5 knockout mice when compared to wild-type controls. Furthermore, TRPM5 knockout mice failed to display preferences for 18:2 FA or soybean oil in the two-bottle choice test when compared to controls [10, 111]. TRPM5 are also important for detection of other primary tastes [111] and are likely involved in a more general role in downstream signaling mechanisms of taste transduction.

4.4. GPR40 and GPR120

The G-protein coupled receptors, GPR40 and GPR120, are located on type 1 and type 2 taste bud cells, and mice lacking these receptors exhibit reduced preferences for LCUFAs and nerve (i.e., chorda tympani and glossopharyngeal) responses to lingual application of LCUFAs [9]. Furthermore, LCUFAs activate Ca^{2+} signaling *in vitro* in HEK 293T cells transfected with mouse or human GPR40 or GPR120 [94]. Sclafani and colleagues, however, recently reported that GPR40 and GPR120 might not be necessary for the cephalic component of fat preference [112]. Mice lacking GPR40, GPR120, or double knockouts, exhibited similar preferences to wild-type mice for intralipid versus water in a two-bottle choice test [112]. In contrast to oral exposure, knockout mice trained in the conditioned flavor preference paradigm to drink a non-nutritive sweet flavor (conditioned stimulus +) paired with intragastric infusion of intralipid, and another non-nutritive sweet flavor (conditioned stimulus -) paired with intragastric infusion of water, exhibited reduced intakes and preferences for the CS+ solution in subsequent one-bottle intake or two-bottle choice tests when compared to wild-type mice. Studies from another group showed that non fatty acid agonists of GPR40 and GPR120 initiated glossopharyngeal nerve activation in wild-type, but not mice lacking GPR40 or GPR120; however, these same agonists failed to elicit any discernable preferences in two-bottle choice tests in wild-type [113]. Collectively, these studies suggest that GPR40 and GPR120 are not necessary—but are possibly sufficient—for preferences associated with tasting dietary lipids, and their importance might lie in the post-oral regulation of preference [112].

5. Obesity and dietary fat detection

Recent studies in humans and rodents suggest that the oral detection and perception of dietary fats might be dysregulated under conditions of obesity. Several laboratories report that obese rats [97] and mice [98], when compared to lean, require higher concentrations of

nutritive fats (i.e., corn oil or 18:2 FA) in order to effectively engage in consumption. The data could be interpreted to suggest a dysregulation in the sensory perception of fats in DIO rodents; however, the authors' measurements of intakes do not reflect taste thresholds as psychophysically defined [114]. Thus, changes in, for example, the motivation to consume under conditions of DIO, rather than changes in detection thresholds, is possible and should be noted. Nonetheless, Ca^{2+} signaling in isolated taste cells from mice fed a high fat diet for four weeks were diminished in the presence of LCUFAs, when compared to control mice fed a standard chow [98]. Stewart and colleagues reported a similar phenotype in obese humans when compared to DIO rodents. Obese men required higher concentrations of 18:1 FA for detection, when compared to lean subjects [99], and detection thresholds for 18:1 FA positively correlated with BMI [115]. Furthermore, subjects ranked as hyposensitive to 18:1 FA, when compared to hypersensitive subjects, reported increased energy intake, and reductions in the ability to distinguish minor changes in the fat content of custard [115]. Hypersensitivity to 18:1 FA, in contrast, was associated with lower BMI, lower total energy consumption, and increases in the detection of custards with lower fat content [73].

Recent evidence suggests that the quantity of fats consumed might be a predictor of sensitivity to fatty acids [116]. Lean or overweight/obese human subjects (mean BMI 23 and 28, respectively) were maintained for four weeks on a high-fat (greater than 45% of caloric intake) or low-fat diet (less than 20% of caloric intake) and assessed for taste sensitivities to 18:1 FA. All subjects maintained on the low fat diet exhibited increased sensitivities to 18:1 FA when compared to baseline measurements obtained prior to the onset of testing. Lean subjects maintained on the high-fat diet exhibited reduced sensitivities to 18:1 FA, while overweight/obese subjects did not show any changes. The authors suggest that obese subjects might already be "adapted" to chronically consuming a high-fat diet, and thus, do not show any further differences in their oral perception of fatty acids following four weeks on a regimented high-fat diet. Lean or overweight/obese subjects, however, did not differ in dietary intake and detection of 18:1 FA at baseline before experimental intervention, raising the question of the generalization of the experimental data to normal feeding conditions. Furthermore, other studies in human also revealed no correlation between the quantity of fat normally consumed by the test subjects and their detection thresholds for 18:1 FA [71].

5.1. CD36 expression in obesity

Recent studies in human and rodent suggest that the expression or function of the proposed fatty acid receptor, CD36, might be dysregulated in obesity. Pepino and colleagues reported that obese human subjects (mostly African American women included in the study) homozygous for a common single nucleotide polymorphism in the CD36 gene (rs1761667 A-allele), which associates with reduced CD36 expression, displayed eight-fold increases in the detection threshold for 18:1 FA and triolein versus obese carriers of the CD36 rs1761667 G-allele [70]. Moreover, Wistar rats maintained on a high-fat diet for eight weeks displayed reduced CD36 mRNA levels in circumvallate papillae and reduced number of cells containing immunoreactivity for CD36 protein, when compared to rats maintained on a standard chow [117]. Other studies found that mice fed ad-libitum a no-fat diet had comparable levels of CD36 mRNA in circumvallate papillae when compared to fasted controls; however, mice fed ad-libitum a high-fat (30% w/w) or a low-fat diet (0.5%)

exhibited reductions in levels of CD36 mRNA, with no changes to GPR120 mRNA in any condition [118]. Furthermore, application of soybean oil to the tongue in anaesthetized mice fed ad-libitum a standard chow resulted in reductions in CD36 mRNA and protein expression within one hour after exposure. The authors of this study suggested that small amounts of fat are sufficient to decrease CD36 gene expression and might contribute to a reduction in the oral detection of dietary fats. In support of this hypothesis, mice maintained on a high-fat diet, when compared to controls, displayed impairments in the detection and preference for low concentrations of rapeseed oil in a two-bottle choice test. Furthermore, DIO mice failed to exhibit the normal reduction in CD36 protein [98] that occurs within one hour after refeeding in fasted control mice maintained on a standard chow [98, 118], and have impaired calcium signaling in isolated CD36-positive circumvallate papillae [98]. CD36 protein expression in taste bud cells of fasted mice fed a high-fat diet for four or 23 weeks, however, were not different when compared to those maintained on a standard chow. Thus, given the similar expression of CD36 protein in diet-induced obese (DIO) and lean mice under control conditions, coupled with the previously discussed data in humans in which no correlation was found between the quantity of fats normally eaten and the oral detection of 18:1 FA [71], it is preliminary to conclude that a dysregulation in CD36 expression in obesity contributes to altered fat detection, and further studies will be required.

6. Humans and cephalic-phase lipid release

Cephalic-phase biochemical responses in the periphery are widely reported and viewed as anticipatory signals for the arrival of nutrients [119, 120]. Early work in humans from the 1950's found that consuming a meal after direct loading of the highly fat soluble vitamin A into the intestine led to a rapid increase in circulating vitamin A content, which is an indirect indicator of elevated triacylglycerol levels [121]. This effect was reproduced by the authors in a modified sham-feeding paradigm in humans that involves mastication of a palatable food, then expectoration without swallowing. The results suggest a cephalic mechanism for the rise in circulating triacylglycerol levels. Moreover, pretreatment with the muscarinic acetylcholine receptor antagonist, atropine, delayed the rise in plasma vitamin A levels after a meal, which suggests that cholinergic signals carried by the efferent vagus nerve are required for the response [122].

Richard Mattes further explored the role for fat taste in driving increases in circulating triacylglycerol levels in the modified sham-feeding paradigm in humans [123]. Oral exposure to fats, despite wearing nose plugs to eliminate olfactory cues, led to increases in serum triacylglycerol levels that were maintained at all time points, while odor-only exposure increased triacylglycerol levels at four hours only. Additional studies showed that tasting butter, but not fat substitutes (i.e., Olestra), elicited a rise in serum triacylglycerol levels, a result that highlights the ability for gustatory signals from dietary fats, but not non-nutritive fats, to initiate changes in circulating triacylglycerols [124]. Interestingly, tasting non-fat cream cheese triggered increases in serum triacylglycerol levels; however, the effect was more pronounced with full-fat cream cheese [125]. Further studies revealed that oral fat exposure for as little as 10 seconds led to an initial spike in triacylglycerol levels [125], then a more prolonged postprandial rise at later time points [125, 126]. Thus, it is possible that a portion of lipids from a high-fat meal are retained in enterocytes and mobilized following

subsequent meals, leading to a rapid initial spike in serum triacylglycerols (i.e., second meal effect) [127]. Other investigations in humans support this possibility. For these studies, subjects were provided a high-fat meal, then five hours later, given water or glucose, followed by biopsies of jejunum mucosa via esophageal-gastro-duodenoscopy [128]. Subjects that drank water retained significantly more lipid droplets in enterocytes and displayed lower levels of circulating triacylglycerols than those that consumed glucose, a result that further supports the possibility that lipids are stored in enterocytes and rapidly released by a cephalic mechanism after a subsequent meal. It would be of interest, however, to perform the same test in subjects after tasting dietary fats in order to evaluate the selectivity and magnitude of the response to different macronutrients.

Our recent work suggests that tasting a fatty liquid meal drives endocannabinoid signaling at cannabinoid CB₁Rs in the rat jejunum through efferent vagal neurotransmission, and this peripheral signal is required to maintain intake and preferences for dietary fats [38, 46]. It is possible that this cephalic-phase signal initiated by fats might serve, in addition to the promotion of feeding, a broader role in lipid processing. Several studies support this hypothesis. Pharmacological inhibition of endocannabinoid degradation in mice with the non-selective inhibitor, isopropyl dodecylfluorophosphonate, induced hypertriglyceridemia through a mechanism that included disrupting apolipoprotein-E-mediated clearance of plasma triacylglycerols [129]. This effect was blocked with the CB₁R antagonist, AM251, and absent in CB₁R or apolipoprotein-E knockout mice. Furthermore, inhibiting CB₁Rs in DIO rats with rimonabant significantly reduced levels of circulating triacylglycerols and non-esterified fatty acids, and triacylglycerol content in skeletal muscle [130]. A similar response was found in weight-stable baboons: rimonabant treatment significantly increased turnover of free fatty acids and triacylglycerol-rich lipoproteins in circulation [131]. Endocannabinoid mechanisms in peripheral organs, such as the liver [132, 133] or adipose tissue [134–137], are important for energy storage. The localization of fat-induced endocannabinoid signaling in the gut [15, 46], however, is suggestive of an anticipatory signal that might regulate the processing of dietary lipids, and their release into, or clearance from, circulation.

7. Conclusions and future directions

Studies from the past two decades have substantially expanded our understanding of the physiological underpinnings of dietary fat detection in mammals, and support fat taste as a primary taste quality. Many knowledge gaps, however, remain. The evidence suggests that gustation, texture, and olfaction are all to varying degrees involved in the perception of fatty foods, with gustatory signals from free fatty acids as a likely irreplaceable component. This diverse set of characteristics contributes to the difficulty in defining fat as a primary taste quality, and at the same time, highlights the unique nature of these molecules and our possibly dogmatic interpretation of what constitutes a basic taste quality. Some of the more immediate key elements that need to be addressed in terms of adequately defining a specific fat taste are 1) a more definitive identification of specific “fat receptor(s)” that interact with free fatty acids in the oral cavity of humans and other mammals, 2) characterization of key intracellular mechanisms that transduce the receptor-mediated signal from fatty acids into an electrical current, 3) the specific neurotransmitter(s) that communicate signaling from taste

bud cells to the afferent gustatory nerves, 4) how fat taste is represented in forebrain structures, 5) how specific signaling mechanisms in the periphery, including the endocannabinoid system, contribute to fat taste and preference, and more generally, 6) how these systems are dysregulated in obesity.

Acknowledgments

This manuscript is based on work presented during the 2013 Annual Meeting of the Society for the Study of Ingestive Behavior, July 30–August 3, 2013. The SSIB meeting was made possible in part by generous unrestricted donations from Ajinomoto North America, Inc, The Coca-Cola Company, GlaxoSmithKline Consumer Health Care, PepsiCo, Inc, Research Diets, Inc, Advanced Targeting Systems, Elsevier, Lafayette-Campden Neuroscience, Sable Systems International, Senomyx, and TSE. Funding from the National Institute on Drug Abuse of the National Institutes of Health is gratefully acknowledged: grant K99 DA034009 to NVD.

References

1. Small DM. Flavor is in the brain. *Physiology & Behavior*. 2012; 107:540–52. [PubMed: 22542991]
2. Breslin PA, Spector AC. Mammalian taste perception. *Current Biology*. 2008; 18:R148–55. [PubMed: 18302913]
3. Mattes RD. Is there a fatty acid taste? *Annu Rev Nutr*. 2009; 29:305–27. [PubMed: 19400700]
4. Mattes RD. Accumulating evidence supports a taste component for free fatty acids in humans. *Physiology & Behavior*. 2011; 104:624–31. [PubMed: 21557960]
5. Roper SD. Taste buds as peripheral chemosensory processors. *Seminars in Cell & Developmental Biology*. 2013; 24:71–9. [PubMed: 23261954]
6. Laugerette F, Passilly-Degrace P, Patris B, Niot I, Febbraio M, Montmayeur JP, et al. CD36 involvement in orosensory detection of dietary lipids, spontaneous fat preference, and digestive secretions. *The Journal of Clinical Investigation*. 2005; 115:3177–84. [PubMed: 16276419]
7. Sclafani A, Ackroff K, Abumrad NA. CD36 gene deletion reduces fat preference and intake but not post-oral fat conditioning in mice. *Am J Physiol Regul Integr Comp Physiol*. 2007; 293:R1823–32. [PubMed: 17804586]
8. Fukuwatari T, Kawada T, Tsuruta M, Hiraoka T, Iwanaga T, Sugimoto E, et al. Expression of the putative membrane fatty acid transporter (FAT) in taste buds of the circumvallate papillae in rats. *FEBS letters*. 1997; 414:461–4. [PubMed: 9315741]
9. Cartoni C, Yasumatsu K, Ohkuri T, Shigemura N, Yoshida R, Godinot N, et al. Taste preference for fatty acids is mediated by GPR40 and GPR120. *J Neurosci*. 2010; 30:8376–82. [PubMed: 20573884]
10. Liu P, Shah BP, Croasdell S, Gilbertson TA. Transient receptor potential channel type M5 is essential for fat taste. *J Neurosci*. 2011; 31:8634–42. [PubMed: 21653867]
11. Travers SP, Nicklas K. Taste bud distribution in the rat pharynx and larynx. *Anat Rec*. 1990; 227:373–9. [PubMed: 2372140]
12. Vandenbeuch A, Clapp TR, Kinnamon SC. Amiloride-sensitive channels in type I fungiform taste cells in mouse. *BMC Neuroscience*. 2008; 9:1. [PubMed: 18171468]
13. Kinnamon SC, Finger TE. A taste for ATP: neurotransmission in taste buds. *Frontiers in cellular neuroscience*. 2013; 7:264. [PubMed: 24385952]
14. Hamilton RB, Norgren R. Central projections of gustatory nerves in the rat. *The Journal of Comparative Neurology*. 1984; 222:560–77. [PubMed: 6199385]
15. DiPatrizio NV, Piomelli D. The thrifty lipids: endocannabinoids and the neural control of energy conservation. *Trends Neurosci*. 2012; 35:403–11. [PubMed: 22622030]
16. Grill HJ, Hayes MR. Hindbrain neurons as an essential hub in the neuroanatomically distributed control of energy balance. *Cell Metab*. 2012; 16:296–309. [PubMed: 22902836]
17. Yu T, Shah BP, Hansen DR, Park-York M, Gilbertson TA. Activation of oral trigeminal neurons by fatty acids is dependent upon intracellular calcium. *European Journal of Physiology*. 2012; 464:227–37. [PubMed: 22644615]

18. Pittman DW, Contreras RJ. Responses of single lingual nerve fibers to thermal and chemical stimulation. *Brain Research*. 1998; 790:224–35. [PubMed: 9593906]
19. Stratford JM, Curtis KS, Contreras RJ. Chorda tympani nerve transection alters linoleic acid taste discrimination by male and female rats. *Physiology & Behavior*. 2006; 89:311–9. [PubMed: 16963089]
20. Pittman D, Crawley ME, Corbin CH, Smith KR. Chorda tympani nerve transection impairs the gustatory detection of free fatty acids in male and female rats. *Brain Research*. 2007; 1151:74–83. [PubMed: 17428454]
21. Gaillard D, Laugerette F, Darcel N, El-Yassimi A, Passilly-Degrace P, Hichami A, et al. The gustatory pathway is involved in CD36-mediated orosensory perception of long-chain fatty acids in the mouse. *FASEB J*. 2008; 22:1458–68. [PubMed: 18162488]
22. Treesukosol Y, Blonde GD, Jiang E, Gonzalez D, Smith JC, Spector AC. Necessity of the glossopharyngeal nerve in the maintenance of normal intake and ingestive bout size of corn oil by rats. *Am J Physiol Regul Integr Comp Physiol*. 2010; 299:R1050–8. [PubMed: 20668233]
23. DiLeone RJ, Taylor JR, Picciotto MR. The drive to eat: comparisons and distinctions between mechanisms of food reward and drug addiction. *Nature Neuroscience*. 2012; 15:1330–5. [PubMed: 23007187]
24. Kenny PJ. Common cellular and molecular mechanisms in obesity and drug addiction. *Nature Reviews Neuroscience*. 2011; 12:638–51. [PubMed: 22011680]
25. Volkow ND, Wang GJ, Fowler JS, Tomasi D, Baler R. Food and Drug Reward: Overlapping Circuits in Human Obesity and Addiction. *Curr Top Behav Neurosci*. 2011
26. Avena NM, Bocarsly ME, Hoebel BG. Animal models of sugar and fat bingeing: relationship to food addiction and increased body weight. *Methods in Molecular Biology*. 2012; 829:351–65. [PubMed: 22231826]
27. Drewnowski A. Food perceptions and preferences of obese adults: a multidimensional approach. *Int J Obes*. 1985; 9:201–12. [PubMed: 4055227]
28. White MA, Whisenhunt BL, Williamson DA, Greenway FL, Netemeyer RG. Development and validation of the food-craving inventory. *Obesity Research*. 2002; 10:107–14. [PubMed: 11836456]
29. Pepino MY, Finkbeiner S, Mennella JA. Similarities in food cravings and mood states between obese women and women who smoke tobacco. *Obesity (Silver Spring)*. 2009; 17:1158–63. [PubMed: 19247281]
30. Miller WC, Lindeman AK, Wallace J, Niederpruem M. Diet composition, energy intake, and exercise in relation to body fat in men and women. *Am J Clin Nutr*. 1990; 52:426–30. [PubMed: 2393005]
31. Piomelli D. The molecular logic of endocannabinoid signalling. *Nature Reviews Neuroscience*. 2003; 4:873–84. [PubMed: 14595399]
32. DiPatrizio NV, Simansky KJ. Activating parabrachial cannabinoid CB1 receptors selectively stimulates feeding of palatable foods in rats. *J Neurosci*. 2008; 28:9702–9. [PubMed: 18815256]
33. DiPatrizio NV, Simansky KJ. Inhibiting parabrachial fatty acid amide hydrolase activity selectively increases the intake of palatable food via cannabinoid CB1 receptors. *Am J Physiol Regul Integr Comp Physiol*. 2008; 295:R1409–14. [PubMed: 18768763]
34. Piomelli D. A fatty gut feeling. *Trends in Endocrinology and Metabolism*. 2013; 24:332–41. [PubMed: 23567058]
35. Ferreira JG, Tellez LA, Ren X, Yeckel CW, de Araujo IE. Regulation of fat intake in the absence of flavour signalling. *The Journal of Physiology*. 2012; 590:953–72. [PubMed: 22219333]
36. Ackroff, K.; Sclafani, A. Oral and Postoral Determinants of Dietary Fat Appetite. In: Montmayeur, JP.; le Coutre, J., editors. *Fat Detection: Taste, Texture, and Post Ingestive Effects*. Boca Raton (FL): 2010.
37. Greenberg D, Smith GP. The controls of fat intake. *Psychosom Med*. 1996; 58:559–69. [PubMed: 8948004]
38. DiPatrizio NV, Astarita G, Schwartz G, Li X, Piomelli D. Endocannabinoid signal in the gut controls dietary fat intake. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108:12904–8. [PubMed: 21730161]

39. Liang NC, Hajnal A, Norgren R. Sham feeding corn oil increases accumbens dopamine in the rat. *Am J Physiol Regul Integr Comp Physiol*. 2006; 291:R1236–9. [PubMed: 16763080]
40. Smith KS, Berridge KC, Aldridge JW. Disentangling pleasure from incentive salience and learning signals in brain reward circuitry. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108:E255–64. [PubMed: 21670308]
41. Berridge KC. 'Liking' and 'wanting' food rewards: brain substrates and roles in eating disorders. *Physiology & Behavior*. 2009; 97:537–50. [PubMed: 19336238]
42. Adachi SI, Endo Y, Mizushige T, Tsuzuki S, Matsumura S, Inoue K, et al. Increased Levels of Extracellular Dopamine in the Nucleus Accumbens and Amygdala of Rats by Ingesting a Low Concentration of a Long-Chain Fatty Acid. *Bioscience, Biotechnology, and Biochemistry*. 2013
43. Suzuki A, Yamane T, Imaizumi M, Fushiki T. Integration of orosensory and postingestive stimuli for the control of excessive fat intake in mice. *Nutrition*. 2003; 19:36–40. [PubMed: 12507637]
44. Lorenz D, Nardi P, Smith GP. Atropine methyl nitrate inhibits sham feeding in the rat. *Pharmacology, Biochemistry, and Behavior*. 1978; 8:405–7.
45. Pradhan SN, Roth T. Comparative behavioral effects of several anticholinergic agents in rats. *Psychopharmacologia*. 1968; 12:358–66. [PubMed: 5652217]
46. DiPatrizio NV, Joslin A, Jung KM, Piomelli D. Endocannabinoid signaling in the gut mediates preference for dietary unsaturated fats. *FASEB J*. 2013; 27:2513–20. [PubMed: 23463697]
47. Dore R, Valenza M, Wang X, Rice KC, Sabino V, Cottone P. The inverse agonist of CB receptor SR141716 blocks compulsive eating of palatable food. *Addict Biol*. 2013 epub, ahead of print.
48. Mindell S, Smith GP, Greenberg D. Corn oil and mineral oil stimulate sham feeding in rats. *Physiology & Behavior*. 1990; 48:283–7. [PubMed: 2255732]
49. Nissenbaum JW, Sclafani A. Sham-feeding response of rats to Polycose and sucrose. *Neurosci Biobehav Rev*. 1987; 11:215–22. [PubMed: 3614789]
50. Berthoud HR. The vagus nerve, food intake and obesity. *Regul Pept*. 2008; 149:15–25. [PubMed: 18482776]
51. Dockray GJ. Enteroendocrine cell signalling via the vagus nerve. *Current Opinion in Pharmacology*. 2013 epub ahead of print.
52. Sykaras AG, Demenis C, Case RM, McLaughlin JT, Smith CP. Duodenal enteroendocrine I-cells contain mRNA transcripts encoding key endocannabinoid and fatty acid receptors. *PLoS ONE*. 2012; 7:e42373. [PubMed: 22876318]
53. Senin LL, Al-Massadi O, Folguiera C, Pardo M, Barja-Fernandez S, Roca-Rivada A, et al. The gastric CB1 receptor modulates ghrelin production through the mTOR pathway to regulate food intake. *PLoS ONE*. 2013; 8:e80339. [PubMed: 24303008]
54. Mattes RD. Oral detection of short-, medium-, and long-chain free fatty acids in humans. *Chem Senses*. 2009; 34:145–50. [PubMed: 19091695]
55. Ramirez I. Role of olfaction in starch and oil preference. *Am J Physiol*. 1993; 265:R1404–9. [PubMed: 8285284]
56. Fukuwatari T, Shibata K, Iguchi K, Saeki T, Iwata A, Tani K, et al. Role of gustation in the recognition of oleate and triolein in anosmic rats. *Physiology & Behavior*. 2003; 78:579–83. [PubMed: 12782211]
57. Takeda M, Sawano S, Imaizumi M, Fushiki T. Preference for corn oil in olfactory-blocked mice in the conditioned place preference test and the two-bottle choice test. *Life Sciences*. 2001; 69:847–54. [PubMed: 11487096]
58. Gilbertson, TA.; Yu, T.; Shah, BP. Gustatory Mechanisms for Fat Detection. In: Montmayeur, JP.; le Coutre, J., editors. *Fat Detection: Taste, Texture, and Post Ingestive Effects*. Boca Raton (FL): 2010.
59. Pittman, DW. Role of the Gustatory System in Fatty Acid Detection in Rats. In: Montmayeur, JP.; le Coutre, J., editors. *Fat Detection: Taste, Texture, and Post Ingestive Effects*. Boca Raton (FL): 2010.
60. Stratford, JM.; Contreras, RJ. Peripheral Gustatory Processing of Free Fatty Acids. In: Montmayeur, JP.; le Coutre, J., editors. *Fat Detection: Taste, Texture, and Post Ingestive Effects*. Boca Raton (FL): 2010.

61. Smith, JC. Orosensory Factors in Fat Detection. In: Montmayeur, JP.; le Coutre, J., editors. Fat Detection: Taste, Texture, and Post Ingestive Effects. Boca Raton (FL): 2010.
62. Mattes, RD. Fat Taste in Humans: Is It a Primary?. In: Montmayeur, JP.; le Coutre, J., editors. Fat Detection: Taste, Texture, and Post Ingestive Effects. Boca Raton (FL): 2010.
63. Smith GP, Greenberg D. Orosensory stimulation of feeding by oils in preweanling and adult rats. *Brain Research Bulletin*. 1991; 27:379–82. [PubMed: 1959033]
64. Yoneda T, Saitou K, Mizushige T, Matsumura S, Manabe Y, Tsuzuki S, et al. The palatability of corn oil and linoleic acid to mice as measured by short-term two-bottle choice and licking tests. *Physiology & Behavior*. 2007; 91:304–9. [PubMed: 17459430]
65. Hamosh M, Scow RO. Lingual lipase and its role in the digestion of dietary lipid. *The Journal of Clinical Investigation*. 1973; 52:88–95. [PubMed: 4682389]
66. Hand AR. The fine structure of von Ebner's gland of the rat. *J Cell Biol*. 1970; 44:340–53. [PubMed: 5411078]
67. Hamosh M, Burns WA. Lipolytic activity of human lingual glands (Ebner). *Lab Invest*. 1977; 37:603–8. [PubMed: 599904]
68. Hamosh M. Lingual and gastric lipases. *Nutrition*. 1990; 6:421–8. [PubMed: 2134569]
69. Kawai T, Fushiki T. Importance of lipolysis in oral cavity for orosensory detection of fat. *Am J Physiol Regul Integr Comp Physiol*. 2003; 285:R447–54. [PubMed: 12702486]
70. Pepino MY, Love-Gregory L, Klein S, Abumrad NA. The fatty acid translocase gene CD36 and lingual lipase influence oral sensitivity to fat in obese subjects. *J Lipid Res*. 2012; 53:561–6. [PubMed: 22210925]
71. Kulkarni B, Mattes R. Evidence for presence of nonesterified fatty acids as potential gustatory signaling molecules in humans. *Chem Senses*. 2013; 38:119–27. [PubMed: 23250958]
72. Laffargue A, de Kochko A, Dussert S. Development of solid-phase extraction and methylation procedures to analyse free fatty acids in lipid-rich seeds. *Plant Physiology and Biochemistry*. 2007; 45:250–7. [PubMed: 17360190]
73. Stewart JE, Feinle-Bisset C, Golding M, Delahunty C, Clifton PM, Keast RS. Oral sensitivity to fatty acids, food consumption and BMI in human subjects. *Br J Nutr*. 2010; 104:145–52. [PubMed: 20196892]
74. Tucker RM, Laguna L, Quinn R, Mattes RD. The effect of short, daily oral exposure on non-esterified fatty acid sensitivity. *Chemosensory Perception*. 2013; 6:78–85.
75. Grabenhorst F, Rolls ET. The representation of oral fat texture in the human somatosensory cortex. *Human Brain Mapping*. 2013
76. De Araujo IE, Rolls ET. Representation in the human brain of food texture and oral fat. *J Neurosci*. 2004; 24:3086–93. [PubMed: 15044548]
77. Gilbertson TA, Liu L, York DA, Bray GA. Dietary fat preferences are inversely correlated with peripheral gustatory fatty acid sensitivity. *Annals of the New York Academy of Sciences*. 1998; 855:165–8. [PubMed: 9929599]
78. Blasbalg TL, Hibbeln JR, Ramsden CE, Majchrzak SF, Rawlings RR. Changes in consumption of omega-3 and omega-6 fatty acids in the United States during the 20th century. *Am J Clin Nutr*. 2011; 93:950–62. [PubMed: 21367944]
79. Moreau, RA. Corn Oil. *Vegetable Oils in Food Technology*. Wiley-Blackwell; 2011. p. 273-89.
80. Yoneda T, Saitou K, Asano H, Mizushige T, Matsumura S, Eguchi A, et al. Assessing palatability of long-chain fatty acids from the licking behavior of BALB/c mice. *Physiology & Behavior*. 2009; 96:735–41. [PubMed: 19385029]
81. Alvheim AR, Torstensen BE, Lin YH, Lillefosse HH, Lock EJ, Madsen L, et al. Dietary linoleic acid elevates endogenous 2-arachidonoylglycerol and anandamide in Atlantic salmon (*Salmo salar* L.) and mice, and induces weight gain and inflammation in mice. *Br J Nutr*. 2012:1–10.
82. Dayton S, Hashimoto S, Dixon W, Pearce ML. Composition of lipids in human serum and adipose tissue during prolonged feeding of a diet high in unsaturated fat. *J Lipid Res*. 1966; 7:103–11. [PubMed: 5900208]
83. Krishnan S, Cooper JA. Effect of dietary fatty acid composition on substrate utilization and body weight maintenance in humans. *European journal of nutrition*. 2013

84. Guyenet SJ, Schwartz MW. Clinical review: Regulation of food intake, energy balance, and body fat mass: implications for the pathogenesis and treatment of obesity. *J Clin Endocrinol Metab.* 2012; 97:745–55. [PubMed: 22238401]
85. Tam J, Vemuri VK, Liu J, Batkai S, Mukhopadhyay B, Godlewski G, et al. Peripheral CB1 cannabinoid receptor blockade improves cardiometabolic risk in mouse models of obesity. *The Journal of Clinical Investigation.* 2010; 120:2953–66. [PubMed: 20664173]
86. Randall PA, Vemuri VK, Segovia KN, Torres EF, Hosmer S, Nunes EJ, et al. The novel cannabinoid CB1 antagonist AM6545 suppresses food intake and food-reinforced behavior. *Pharmacology, Biochemistry, and Behavior.* 2010; 97:179–84.
87. Cluny NL, Vemuri VK, Chambers AP, Limebeer CL, Bedard H, Wood JT, et al. A novel peripherally restricted cannabinoid receptor antagonist, AM6545, reduces food intake and body weight, but does not cause malaise, in rodents. *British Journal of Pharmacology.* 2011; 161:629–42. [PubMed: 20880401]
88. LoVerme J, Duranti A, Tontini A, Spadoni G, Mor M, Rivara S, et al. Synthesis and characterization of a peripherally restricted CB1 cannabinoid antagonist, URB447, that reduces feeding and body-weight gain in mice. *Bioorg Med Chem Lett.* 2009; 19:639–43. [PubMed: 19128970]
89. Gilbertson TA, Khan NA. Cell signaling mechanisms of oro-gustatory detection of dietary fat: advances and challenges. *Progress in Lipid Research.* 2013 epub, ahead of print.
90. Gilbertson TA, Fontenot DT, Liu L, Zhang H, Monroe WT. Fatty acid modulation of K⁺ channels in taste receptor cells: gustatory cues for dietary fat. *Am J Physiol.* 1997; 272:C1203–10. [PubMed: 9142845]
91. Chale-Rush A, Burgess JR, Mattes RD. Evidence for human orosensory (taste?) sensitivity to free fatty acids. *Chem Senses.* 2007; 32:423–31. [PubMed: 17361006]
92. Chale-Rush A, Burgess JR, Mattes RD. Multiple routes of chemosensitivity to free fatty acids in humans. *American journal of Physiology.* 2007; 292:G1206–12. [PubMed: 17234892]
93. Pittman DW, Smith KR, Crawley ME, Corbin CH, Hansen DR, Watson KJ, et al. Orosensory detection of fatty acids by obesity-prone and obesity-resistant rats: strain and sex differences. *Chem Senses.* 2008; 33:449–60. [PubMed: 18372387]
94. Galindo MM, Voigt N, Stein J, van Lengerich J, Raguse JD, Hofmann T, et al. G protein-coupled receptors in human fat taste perception. *Chem Senses.* 2012; 37:123–39. [PubMed: 21868624]
95. Yoshida R, Ohkuri T, Jyotaki M, Yasuo T, Horio N, Yasumatsu K, et al. Endocannabinoids selectively enhance sweet taste. *Proceedings of the National Academy of Sciences of the United States of America.* 2010; 107:935–9. [PubMed: 20080779]
96. Matias I, Gatta-Cherifi B, Tabarin A, Clark S, Leste-Lasserre T, Marsicano G, et al. Endocannabinoids measurement in human saliva as potential biomarker of obesity. *PLoS ONE.* 2012; 7:e42399. [PubMed: 22860123]
97. Shin AC, Zheng H, Pistell PJ, Berthoud HR. Roux-en-Y gastric bypass surgery changes food reward in rats. *International Journal of Obesity (2005).* 2011; 35:642–51. [PubMed: 20805826]
98. Chevrot M, Bernard A, Ancel D, Buttet M, Martin C, Abdoul-Azize S, et al. Obesity alters the gustatory perception of lipids in the mouse: plausible involvement of lingual CD36. *J Lipid Res.* 2013; 54:2485–94. [PubMed: 23840049]
99. Stewart JE, Seimon RV, Otto B, Keast RS, Clifton PM, Feinle-Bisset C. Marked differences in gustatory and gastrointestinal sensitivity to oleic acid between lean and obese men. *Am J Clin Nutr.* 2011; 93:703–11. [PubMed: 21310831]
100. Little TJ, Feinle-Bisset C. Effects of dietary fat on appetite and energy intake in health and obesity--oral and gastrointestinal sensory contributions. *Physiology & Behavior.* 2011; 104:613–20. [PubMed: 21596051]
101. Degrace-Passilly P, Besnard P. CD36 and taste of fat. *Curr Opin Clin Nutr Metab Care.* 2012; 15:107–11. [PubMed: 22248592]
102. Liu L, Hansen DR, Kim I, Gilbertson TA. Expression and characterization of delayed rectifying K⁺ channels in anterior rat taste buds. *American Journal of Physiology.* 2005; 289:C868–80. [PubMed: 15930148]

103. Kinnamon SC, Dionne VE, Beam KG. Apical localization of K⁺ channels in taste cells provides the basis for sour taste transduction. *Proceedings of the National Academy of Sciences of the United States of America*. 1988; 85:7023–7. [PubMed: 2457924]
104. Cummings TA, Daniels C, Kinnamon SC. Sweet taste transduction in hamster: sweeteners and cyclic nucleotides depolarize taste cells by reducing a K⁺ current. *Journal of Neurophysiology*. 1996; 75:1256–63. [PubMed: 8867134]
105. El-Yassimi A, Hichami A, Besnard P, Khan NA. Linoleic acid induces calcium signaling, Src kinase phosphorylation, and neurotransmitter release in mouse CD36-positive gustatory cells. *J Biol Chem*. 2008; 283:12949–59. [PubMed: 18321850]
106. Martin C, Chevrot M, Poirier H, Passilly-Degrace P, Niot I, Besnard P. CD36 as a lipid sensor. *Physiology & Behavior*. 2011; 105:36–42. [PubMed: 21354192]
107. Dramane G, Abdoul-Azize S, Hichami A, Vogtle T, Akpona S, Chouabe C, et al. STIM1 regulates calcium signaling in taste bud cells and preference for fat in mice. *The Journal of Clinical Investigation*. 2012; 122:2267–82. [PubMed: 22546859]
108. Martin C, Passilly-Degrace P, Chevrot M, Ancel D, Sparks SM, Drucker DJ, et al. Lipid-mediated release of GLP-1 by mouse taste buds from circumvallate papillae: putative involvement of GPR120 and impact on taste sensitivity. *J Lipid Res*. 2012; 53:2256–65. [PubMed: 22904345]
109. Simons PJ, Kummer JA, Luiken JJ, Boon L. Apical CD36 immunolocalization in human and porcine taste buds from circumvallate and foliate papillae. *Acta Histochemica*. 2011; 113:839–43. [PubMed: 20950842]
110. Chen CS, Bench EM, Allerton TD, Schreiber AL, Arceneaux KP 3rd, Primeaux SD. Preference for linoleic acid in obesity-prone and obesity-resistant rats is attenuated by the reduction of CD36 on the tongue. *Am J Physiol Regul Integr Comp Physiol*. 2013 epub ahead of print.
111. Sclafani A, Zukerman S, Glendinning JI, Margolskee RF. Fat and carbohydrate preferences in mice: the contribution of alpha-gustducin and Trpm5 taste-signaling proteins. *Am J Physiol Regul Integr Comp Physiol*. 2007; 293:R1504–13. [PubMed: 17652359]
112. Sclafani A, Zukerman S, Ackroff K. GPR40 and GPR120 fatty acid sensors are critical for post-oral but not oral mediation of fat preferences in the mouse. *Am J Physiol Regul Integr Comp Physiol*. 2013 epub, ahead of print.
113. Godinot N, Yasumatsu K, Barcos ME, Pineau N, Ledda M, Viton F, et al. Activation of tongue-expressed GPR40 and GPR120 by non caloric agonists is not sufficient to drive preference in mice. *Neuroscience*. 2013; 250:20–30. [PubMed: 23831422]
114. Treesukosol Y, Lyall V, Heck GL, DeSimone JA, Spector AC. A psychophysical and electrophysiological analysis of salt taste in Trpv1 null mice. *Am J Physiol Regul Integr Comp Physiol*. 2007; 292:R1799–809. [PubMed: 17234959]
115. Stewart JE, Newman LP, Keast RS. Oral sensitivity to oleic acid is associated with fat intake and body mass index. *Clin Nutr*. 2011; 30:838–44. [PubMed: 21757270]
116. Stewart JE, Keast RS. Recent fat intake modulates fat taste sensitivity in lean and overweight subjects. *International Journal of Obesity (2005)*. 2012; 36:834–42. [PubMed: 21829156]
117. Zhang XJ, Zhou LH, Ban X, Liu DX, Jiang W, Liu XM. Decreased expression of CD36 in circumvallate taste buds of high-fat diet induced obese rats. *Acta Histochemica*. 2011; 113:663–7. [PubMed: 20926122]
118. Martin C, Passilly-Degrace P, Gaillard D, Merlin JF, Chevrot M, Besnard P. The lipid-sensor candidates CD36 and GPR120 are differentially regulated by dietary lipids in mouse taste buds: impact on spontaneous fat preference. *PLoS ONE*. 2011; 6:e24014. [PubMed: 21901153]
119. Berthoud HR, Jeanrenaud B. Sham feeding-induced cephalic phase insulin release in the rat. *Am J Physiol*. 1982; 242:E280–5. [PubMed: 7039361]
120. Vahl TP, Drazen DL, Seeley RJ, D'Alessio DA, Woods SC. Meal-anticipatory glucagon-like peptide-1 secretion in rats. *Endocrinology*. 2010; 151:569–75. [PubMed: 19915164]
121. Mendeloff AI. The effects of eating and of sham feeding upon the absorption of vitamin A palmitate in man. *The Journal of Clinical Investigation*. 1954; 33:1015–21. [PubMed: 13174658]
122. Robertson MD, Mason AO, Frayn KN. Timing of vagal stimulation affects postprandial lipid metabolism in humans. *Am J Clin Nutr*. 2002; 76:71–7. [PubMed: 12081818]

123. Mattes RD. The taste of fat elevates postprandial triacylglycerol. *Physiology & Behavior*. 2001; 74:343–8. [PubMed: 11714498]
124. Mattes RD. Oral exposure to butter, but not fat replacers elevates postprandial triacylglycerol concentration in humans. *J Nutr*. 2001; 131:1491–6. [PubMed: 11340105]
125. Mattes RD. Brief oral stimulation, but especially oral fat exposure, elevates serum triglycerides in humans. *American Journal of Physiology*. 2009; 296:G365–71. [PubMed: 19074638]
126. Mattes RD. Oral fat exposure increases the first phase triacylglycerol concentration due to release of stored lipid in humans. *J Nutr*. 2002; 132:3656–62. [PubMed: 12468603]
127. Chavez-Jauregui RN, Mattes RD, Parks EJ. Dynamics of fat absorption and effect of sham feeding on postprandial lipema. *Gastroenterology*. 2010; 139:1538–48. [PubMed: 20493191]
128. Robertson MD, Parkes M, Warren BF, Ferguson DJ, Jackson KG, Jewell DP, et al. Mobilisation of enterocyte fat stores by oral glucose in humans. *Gut*. 2003; 52:834–9. [PubMed: 12740339]
129. Ruby MA, Nomura DK, Hudak CS, Mangravite LM, Chiu S, Casida JE, et al. Overactive endocannabinoid signaling impairs apolipoprotein E-mediated clearance of triglyceride-rich lipoproteins. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105:14561–6. [PubMed: 18794527]
130. Cota D, Sandoval DA, Olivieri M, Prodi E, D'Alessio DA, Woods SC, et al. Food intake-independent effects of CB1 antagonism on glucose and lipid metabolism. *Obesity (Silver Spring)*. 2009; 17:1641–5. [PubMed: 19325539]
131. Vaidyanathan V, Bastarrachea RA, Higgins PB, Voruganti VS, Kamath S, Dipatrizio NV, et al. Selective cannabinoid-1 receptor blockade benefits fatty acid and triglyceride metabolism significantly in weight-stable nonhuman primates. *Am J Physiol Endocrinol Metab*. 2012; 303:E624–34. [PubMed: 22761159]
132. Kola B, Hubina E, Tucci SA, Kirkham TC, Garcia EA, Mitchell SE, et al. Cannabinoids and ghrelin have both central and peripheral metabolic and cardiac effects via AMP-activated protein kinase. *J Biol Chem*. 2005; 280:25196–201. [PubMed: 15899896]
133. Haas MJ, Mazza AD, Wong NC, Mooradian AD. Inhibition of apolipoprotein a-I gene expression by obesity-associated endocannabinoids. *Obesity (Silver Spring)*. 2012; 20:721–9. [PubMed: 22016100]
134. Gary-Bobo M, Elachouri G, Scatton B, Le Fur G, Oury-Donat F, Bensaid M. The cannabinoid CB1 receptor antagonist rimonabant (SR141716) inhibits cell proliferation and increases markers of adipocyte maturation in cultured mouse 3T3 F442A preadipocytes. *Molecular Pharmacology*. 2006; 69:471–8. [PubMed: 16282221]
135. Bouaboula M, Hilaiet S, Marchand J, Fajas L, Le Fur G, Casellas P. Anandamide induced PPARgamma transcriptional activation and 3T3-L1 preadipocyte differentiation. *European Journal of Pharmacology*. 2005; 517:174–81. [PubMed: 15987634]
136. Karaliota S, Siafaka-Kapadai A, Gontinou C, Psarra K, Mavri-Vavayanni M. Anandamide increases the differentiation of rat adipocytes and causes PPARgamma and CB1 receptor upregulation. *Obesity (Silver Spring)*. 2009; 17:1830–8. [PubMed: 19543211]
137. Cota D, Marsicano G, Tschop M, Grubler Y, Flachskamm C, Schubert M, et al. The endogenous cannabinoid system affects energy balance via central orexigenic drive and peripheral lipogenesis. *The Journal of Clinical Investigation*. 2003; 112:423–31. [PubMed: 12897210]

Highlights

- Fat taste is not universally recognized as a primary taste quality
- Recent evidence suggests potential fat receptors in the oral cavity of mammals
- Gustatory signals from fats possibly drive intake and preference for fatty foods
- The endocannabinoid system plays a key role in dietary fat intake and preference

Figure 1.

The sham feeding rat. Sham feeding isolates the orosensory component of feeding by eliminating its post-ingestive consequences. The sham feeding protocol is based on the methods of Gerard P. Smith [37], and recently adapted for our studies examining cephalic-phase changes in endocannabinoid signaling [38, 46]. In the sham feeding paradigm, liquid diets are consumed from sipper tubes, then drain out of a reversible gastric cannulae that is surgically implanted into the stomach (see a and b). Liquids drain through a tube that is threaded into the cannulae, and placed through a longitudinal opening in the floor of the cage that allows for free movement of the animal during testing. Fluids are collected in a vessel placed below the cage, which can be weighed to determine that all consumed liquids properly drain through the gastric cannulae.

Table 1

Elements of a taste. Minimal elements of a primary taste quality, as recently extended and defined by Richard Mattes [4].

Minimal elements of a primary taste quality
1) Provide some adaptive advantage
2) Have a defined class of effective stimuli
3) Have unique transduction mechanisms
4) Initiate peripheral signals conveyed by gustatory nerves that are decoded in gustatory centers
5) Be perceptible and unique
6) Evoke a functional physiological and/or behavioral response