Published in final edited form as:

Behav Neurosci. 2014 June ; 128(3): 360–370. doi:10.1037/a0035892.

Basic Sleep and Circadian Science as Building Blocks for Behavioral Interventions: A Translational Approach for Mood Disorders

Lauren D. Asarnow, Adriane M. Soehner, and Allison G. Harvey

University of California, Berkeley

Abstract

Sleep and circadian functioning has been of particular interest to researchers focused on improving treatments for psychiatric illness. The goal of the present paper is to highlight the exciting research that utilizes basic sleep and circadian science as building blocks for intervention in the mood disorders. The reviewed evidence suggests that the sleep and circadian systems are 1) disrupted in the mood disorders and linked to symptoms, 2) open systems that can be modified, 3) the focus of interventions which have been developed to effectively treat sleep disturbance within mood disorders, and 4) intimately linked with mood, such that improvements in sleep are associated with improvements in mood. Although, significant positive treatment effects are evident, more research is needed to fill the gap in our basic understanding of the relationship between sleep and mood.

Keywords

sleep; circadian; mood disorders; cognitive-behavior therapy for insomnia; interpersonal social rhythms therapy; light therapy; sleep deprivation

The World Health Organization's study on the global burden of disease lists mental illness as the leading source of disability, accounting for nearly 40% of all medical disability (World Health Organization, 2002). The associated economic burden is estimated at \$317 billion annually (Kessler et al., 2008; Insel, 2008). An analysis of mortality in 8 states indicates that individuals with serious mental illness die 13 to 32 years earlier than those without mental illness (Colton & Manderscheid, 2006).

Accordingly, there is a need for new and improved treatments for psychiatric illness (Insel, 2009). Sleep and circadian functioning has been of particular interest to researchers focused on improving treatments for psychiatric illness. Across psychiatric disorders it is clear that sleep and circadian systems are often disrupted (Benca, Obermeyer, Thisted & Gillin, 1992; Harvey, 2008a). Indeed, the accumulated evidence indicates that sleep and circadian systems are implicated in the cause and/or maintenance of psychiatric disorders (e.g. Buysse et al., 1997; Jackson, Cavanagh & Scott, 2003; Harvey, 2008a; NIH, 2005). This evidence dispels a previously widely held assumption that sleep disturbance is secondary to, or an

epiphenomenon of, the so-called “primary” psychiatric disorder (Harvey, 2008a). Moreover, as will be discussed later in the present review, the sleep and circadian systems are modifiable.

The goal of this paper is to highlight the exciting research that utilizes basic sleep and circadian science as building blocks for interventions. Although we focus on the mood disorders, there are similar sets of findings across a wide range of psychiatric disorders including schizophrenia (Freeman, Pugh, Vorontsova, & Southgate, 2009; Myers, Startup, & Freeman, 2011) and post traumatic stress disorder (Germain et al., 2012). Perhaps not surprisingly then, the NIMH research domain criteria also highlight sleep and circadian disturbance as a key domain for research on psychiatric illness going forward (Insel et al., 2010).

Sleep and Circadian Disruption in the Mood Disorders

The mood disorders are a broad class of psychiatric disorders characterized by mood episodes, specifically depression and mania. Major Depressive Disorder (MDD) and Bipolar Disorder (BD), are two disorders that fall collectively into the mood disorder category. MDD and BD are amongst the most severe psychiatric illnesses. Estimates of the lifetime rates of suicide attempts of persons with bipolar and unipolar depression are 29.2% and 15.9% respectively (Chen & Dilsaver, 1996). Moreover, MDD is the leading global cause of disability (Insel & Charney, 2003).

MDD is characterized by depressive episodes, which are defined as periods of two weeks or longer with down or depressed mood and/or loss of pleasure and/or interest in daily activities. These mood symptoms are also accompanied by appetite changes, sleep disturbance, feelings of worthlessness or excessive guilt, psychomotor retardation or agitation, difficulty concentrating or making decisions, or suicidal ideation. Bipolar Disorder (BD) is characterized by episodes of mania or hypomania and depression. Manic episodes are periods of elevated, expansive or irritable mood lasting at least one week, or less if resulting in hospitalization. Hypomania is a less severe form of mania that can last 4 or more days. The manic or hypomanic symptoms can also include inflated self-esteem or grandiosity, decreased need for sleep (e.g., one feels rested after only 3 hours of sleep), more talking than is usual or pressured speech, racing thoughts, increased distractibility, increase in goal-directed activity (socially, at work or school, or sexually) or psychomotor agitation.

Disturbances of the sleep-wake cycle are a hallmark feature of the mood disorders, occurring in both mania and depression. These sleep disturbances arise from abnormalities in sleep-wake generating and/or timing (circadian) mechanisms. Though these disturbances are common in both MDD and BD, their presentation varies between mood disorders. Among patients with MDD, approximately 50–52% report insomnia, 27 – 30% report hypersomnia and 50% report early waking during depressive episodes (Korszun et al., 2004). During periods of remission, estimates are that between 7 and 55% continue to report sleep and circadian disturbances (Nierenberg et al., 2010). Among patients with BD, in a sample of 59 patients evidence indicates that 100% report insomnia (Winokur, Clayton, & Reich, 1969) and 78% report hypersomnia (Detre, Himmelijoch, & Swartzburg, 1972).

during depressive episodes. During manic episodes, insomnia occurs in 24–34% of patients (Winokur & Tanna, 1969). In the period between depressive and manic/hypomanic episodes, referred to as the *interepisode period* of BD, insomnia occurs in 55% (Harvey, Schmidt, Scarnà, Semler & Goodwin, 2005) and hypersomnia occurs in 25% of cases (Kaplan, Gruber, Eidelman, Talbot & Harvey, 2011). Among patients with BD and MDD, sleep disturbance is complicated and understudied.

Sleep and circadian disturbances observed among patients with mood disorders are of critical importance for at least 2 reasons. First, sleep is vital for intact emotion and mood regulation (Horne, 1985; Gruber et al., 2007; Dinges et al., 1997). Research in healthy adults has indicated that neural circuitry crucial for emotion regulation is destabilized following a night of sleep deprivation (Yoo et al. 2007; Sotres-Bayon, Bush, & LeDoux, 2004). Further, a study of naturalistic sleep loss in medical residents reported that less sleep predicted increased negative affect following a goal-thwarting event and diminished positive emotions following a goal-enhancing event (Zohar, Tzischinsky, Epstein & Lavie, 2005). Second, the adverse effects of sleep loss on cognitive functioning have been clearly demonstrated (e.g., Van Dongen, Maislin, Mullington & Dinges, 2003). Of particular importance within the mood disorders, sleep disturbance has detrimental effects on tests of cognitive flexibility and decision-making (Walker, Liston, Hobson & Stickgold, 2002). Also, sleep loss disrupts the encoding and later retention for positive emotional memory (Phelps, 2004). Hence individuals with mood disorders, who already have documented deficits in emotion-regulation (Gross & Muñoz, 1995) and cognitive processing (see Gotlib & Joormann, 2010 for review), may experience even more adverse effects from sleep disturbance than healthy populations.

Moreover, the sleep and circadian disruptions observed in MDD and BD have been identified as early indicators of the disorder and/or poor prognosis. Unipolar depressed patients with disturbed sleep have significantly worse clinical outcomes, attrition rates, and response to treatment than good sleepers (Dew et al., 1997; Thase et al., 1997). Additionally, sleep disturbances (measured via self-report and polysomnography) are associated with increased suicidal ideation, as well as slower and lower rates of remission from depression (Buysse et al., 1997; Thase, 1998). Further, two epidemiologic studies have shown that sleep problems identified 47% of the new cases of depression in the following year (Ford & Kamerow, 1989; Eaton, Badawi & Melton, 1995). Sleep disturbance has been found to be the most common prodromal symptom of manic episodes and the 6th most common prodromal symptom of depressive episodes (Jackson et al., 2003). In BD, disturbed sleep appears to be a predictor of worsening symptoms (see Harvey, 2008b for review). Indeed, in inter-episode BD shortened sleep duration predicted worsened depressive symptoms at a 6-month follow-up (Perlman, Johnson & Mellman, 2006). Similarly, interepisode sleep symptoms are correlated with future manic and depressive symptoms and impairment in BD (Eidelman, Talbot, Gruber, Hairston & Harvey, 2010; Kaplan et al., 2011).

Together, the clinical data point to the likely importance of identifying points of intervention to improve sleep and circadian functioning for patients with mood disorders. However, one particular challenge in the study of the sleep and circadian systems is the difficulty isolating

whether the dysfunction arises within the sleep system, the circadian system, or both systems, a problem we now address.

Identifying Targets for Intervention in the Sleep and Circadian Systems

The sleep/wake cycle is regulated by an interaction between two opponent processes (Borbély & Wirz-Justice, 1982). Process S regulates sleep-wake system through homeostatic self-modulation. This process regulates the duration and structure of sleep based on prior sleep and wakefulness. Specifically, sleep pressure increases during extended wakefulness and dissipates during early sleep. Sleep homeostasis influences sleep propensity; that is, sleep homeostasis results in an increased tendency to sleep when a person has been sleep-deprived, and a decreased tendency to sleep after having had a substantial amount of sleep. Process C regulates the circadian clock, arising from the endogenous pacemaker in the hypothalamic suprachiasmatic nuclei (SCN) (Reppert & Weaver, 2002). Circadian regulation of sleep is characterized by a relatively low sleep propensity during waking hours and an increase in sleep propensity during night hours. The maximal circadian propensity for sleep occurs in the early morning hours (typically 3:00 AM – 5:00 AM), near the time of peak melatonin secretion and core body temperature nadir. At the molecular level, intrinsically rhythmic cells within the SCN generate rhythmicity via an autoregulatory transcription-translation feedback loop regulating expression of circadian genes. The process by which the pacemaker is set to a 24-hour period and kept in appropriate phase with seasonally shifting day length is called entrainment, which occurs via *zeitgebers*. *Zeitgeber* is German for ‘time giver’; these are environmental cues that support entrainment. In sum, the sleep and circadian processes work in concert to facilitate consolidated nighttime sleep in humans.

However, while the sleep and circadian processes are interrelated, they are also independent (Borbély & Wirz-Justice, 1982). The independent actions of Process C and Process S can be most clearly isolated using laboratory-based manipulations. Forced desynchrony (FD) protocols are currently the gold standard method for distinguishing the influences of Process S and Process C. Specifically, FD protocols hold participants to a non-24 hour day outside of the range of entrainment of the biological clock. This forces the endogenous clock to free-run to its intrinsic 24–25 hours period, thus decoupling the circadian clock from the build up of homeostatic sleep drive (Hanneman, 2001). Such experimental manipulations revealed that the sleep and circadian processes can be unpaired. Moreover, this work indicates that the phase relationship between the sleep and circadian systems during a 24-hour day is crucial for enabling consolidated periods of sleep and wakefulness. Disruption to the phase relationship between these systems results in poorly consolidated sleep and daytime fatigue or sleepiness.

From an intervention perspective, it is particularly exciting that process C and S are *open systems* susceptible to modification. The endogenous period generated in the SCN is close to, but generally not equal to, 24 hours. As a result, entrainment to a 24-hour day is achieved through a combination of internal cues from the sleep homeostat (Process S) and exogenous inputs (*zeitgebers*). The phase, period and amplitude of the internal clock can be adjusted via changes to the timing/regularity of *zeitgebers* and manipulations of the sleep homeostat.

Process S contributes to entrainment primarily through modulation of the sleep homeostat. When sleep deprivation occurs, sleep pressure builds, thereby causing the individual to feel sleepy earlier than is typical for their circadian clock. Likewise, if an individual takes an afternoon nap, sleep pressure decreases, resulting in the individual being less tired at their usual bedtime. Through changes in sleep timing, the sleep homeostat can affect regulation of the circadian system. Experimental evidence supports this link between the two processes. In hamsters, sleep deprivation was shown to rapidly reset the circadian clock and alter gene expression within the circadian system (Antle & Mistleberger, 2000). Sleep loss also appears to modulate the response of the hamster circadian system to light (Mistleberger, Landry & Marchant, 1997; Challet, Turek, Laute, & Van Reeth, 2001). In humans, a study of 10 free-running patients demonstrated that endogenous circadian rhythm (measured via core body temperature) is intimately linked to sleep propensity; the increase in circadian sleep tendency that occurs as the night progresses counteracts the decrease in sleep propensity associated with accumulated sleep and allows humans to maintain a consolidated 8 hour sleep episode (Dijk & Czeisler, 1994).

With regard to Process C, entrainment occurs via exogenous inputs or zeitgebers, which are environmental events that can affect the phase and period of the clock. Cycles of alternating light and darkness are one of the most powerful exogenous inputs of circadian rhythms in humans (Roenneberg & Foster, 1997; Czeisler & Wright, 1999). In mammals, there are specialized photoreceptive ganglion cells whose exclusive purpose is believed to be entrainment of the circadian rhythm via light intensity measurement (Klerman et al., 2002). Light is detected by ocular photoreceptors and information regarding light intensity is then transmitted to the brain via the retinohypothalamic tract. This tract terminates in the suprachiasmatic nuclei (SCN), lying just above the optic chiasm. Retinal light exposure early in the night will delay the timing of the clock, while light exposure later in the night and early in the morning will advance the timing of the clock (Czeisler et al. 1989; Johnson, 1990). Humans are highly responsive to the phase-delaying effects of light during the early biological night. According to Zeitzer, Dijk, Kornauer, Brown and Czeisler (2000), even small changes in ordinary light exposure during the late evening hours can significantly affect both plasma melatonin concentrations and the entrained phase of the human circadian pacemaker.

Nonphotic cues can also exert a powerful influence on the SCN. Social cues, such as the timing of social interactions, meals and exercise, are a dominant nonphotic cue (Mistleberger, Antle, Glass & Miller, 2000). The best evidence for social entrainment in humans is from a small sample of subjects with total blindness (inability to perceive light of any intensity) who synchronized to the 24 h day, or to near-24 h sleep-wake schedules under laboratory conditions with regular social interactions (Mistleberger & Skene, 2005). In healthy adults, higher daily regularity in social rhythms (social interactions, mealtimes, etc., occurring at about the same time each day) is associated with a biomarker of stronger circadian function (deeper nocturnal body temperature trough) and better subjective sleep quality (Monk, Petrie, Hayes & Kupfer, 1994). Work with animal models complements and extends findings in humans. In the absence of photic cues, female rats living in isolation show a greater delay in their circadian relative to those living in groups (Cambras, Chiesa, Araujo &

Díez-Noguera, 2004), suggesting that cohabitation helps to stabilize circadian rhythmicity. In a study by Sulzman, Fuller and Moore-Ede (1977), only light-dark cycles and cycles of food availability were shown to be entraining agents in a sample of monkeys. Moreover, feeding period restriction has been shown to modulate the expression of peripheral circadian clock genes (Jang & Lee, 2012). Objective body temperature data has confirmed marked differences in the effects of varied restricted feeding schedules on circadian rhythms (Nelson & Halberg, 1986). Exercise and activity can also serve as exogenous inputs that entrain the circadian system, with several studies reporting that increased nocturnal physical activity can phase delay human circadian rhythms (Buxton et al., 1997; Van Reeth et al., 1994; Edgar & Dement, 1991).

As we move to the next section and think more about the interaction between the sleep and circadian systems, we need to be cognizant of the difficulty in truly differentiating between the sleep and circadian processes. Though they are independent processes they are also overlapping and difficult to tease apart.

How do the sleep and circadian systems become dysregulated in the mood disorders?

Popular early theories linking sleep and circadian rhythms to mood episodes proposed that, in mood disorders, the endogenous circadian pacemaker becomes misaligned with sleep timing (Frank, Kupfer, Ehlers & Monk, 1994; Kripke et al., 1994; Wehr, Wirz-Justice, Goodwin, Duncan & Gillin, 1979) and sleep disturbance serves as a psychobiological marker of mood disorder (Kupfer, 1976; Benca et al., 1992). In the past 30 years, researchers have begun to elaborate upon early hypotheses to better understand how the sleep and circadian systems become dysregulated in mood disorders. Below is a summary of a selection of the most well investigated sleep and circadian disturbances and what we know about why these sleep and circadian disturbances develop in the mood disorders. As will become evident, there are many open questions and domains for future research.

Insomnia is highly prevalent in the mood disorders. Insomnia is defined as difficulty initiating or maintaining sleep or non-restorative sleep for at least 1 month that causes significant impairment or distress in social, occupational, or other areas of functioning. Estimates from large epidemiological studies observed a 50–75% comorbidity rate between insomnia and other psychiatric illnesses (Ford & Kamerow, 1989; Lichstein, Wilson & Johnson, 2000). Despite limited information concerning insomnia's underlying pathophysiology, there is a general agreement that insomnia is perpetuated by the interplay of cognitive and behavioral mechanisms, which serve to alter homeostatic and circadian regulation. For example, daytime napping or spending extra time in bed in pursuit of elusive, unpredictable sleep may only serve to interfere with the body's homeostatic mechanisms that operate automatically to increase sleep drive in the face of increasing periods of wakefulness (i.e., sleep debt). Alternately, the habit of remaining in bed well beyond the normal rising time following a poor night's sleep may disrupt the body's circadian system by delaying circadian phase (Edinger & Wohlgemuth, 1999; Webb, 1988). Indeed, experiments in which bedtime (clock time at which participants went to sleep) was manipulated to occur at different phases of the circadian rhythm demonstrated that an

uninterrupted 8 hour sleep episode can only be achieved when sleep is initiated approximately 6 hours before the core body temperature minimum (which is linked to the bodies' circadian rhythm; Dijk & Czeisler, 1994). Moreover, there is likely to be a cyclical influence of sleep disturbance and mood disorder, with worsening sleep problems leading to a decline in mood as well as the maintenance of daytime distress and symptoms and, conversely, daytime distress and symptoms worsening the sleep problems (Harvey, 2008a).

While insomnia has been established as a core symptom across mood disorders, much less is known about hypersomnia in mood disorders. Across diagnostic nosologies, such as the DSM-IV, ICD-10 and ICSD-2, hypersomnia is broadly defined by a combination of prolonged nighttime sleep episodes, increased nighttime wakefulness, frequent daytime napping, and excessive daytime sleepiness (Kaplan & Harvey, 2009). Epidemiologic data suggest hypersomnia is present in 10–40% of patients with various mood disorders (Baldwin & Papakostas, 2006). Despite its prevalence, very little research has been done describing how hypersomnia develops in the mood disorders. Billiard, Dolenc, Aldaz, Ondze and Besset (1994) compared individuals with hypersomnia with and without mood disorders. While excessive sleepiness and prolonged sleep length were evident in a subset of individuals with mood disorders, for others the reports of hypersomnia were more closely associated with illness-related lethargy. Billiard et al. (1994) further noted that many individuals in the mood disorders group, as compared to the idiopathic hypersomnia group, spent a good deal of time in bed without any actual sleep. Excessive time in bed or long sleep could stymie build up of homeostatic sleep pressure and destabilize sleep timing, contributing to disrupted sleep regulation. Thus, akin to insomnia, it is hypothesized that hypersomnia disrupts both the homeostatic and circadian systems (Kaplan & Harvey, 2009).

Inadequate, irregular, or inappropriate timing of exposure to light and dark is hypothesized to play a role in the mood disorders (Boivin, 2000). For example, seasonal affective disorder, a depressive episode occurring in winter or fall when light exposure is lowest and spontaneously remits in spring or summer when light exposure is higher (Rosenthal et al., 1984). Moreover, studies have shown that insomnia complaints and depressed mood are associated with low light exposure (Kripke et al., 1994; Wallace-Guy et al., 2002). In addition, MDD and BD are commonly accompanied by circadian phase shifts such that exposure to light decreases (Emens, Lewy, Kinzie, Arntz & Rough, 2009; Liebenluft, Feldman-Naim, Turner, Schwartz & Wehr, 1995). Researchers have also postulated that an increased sensitivity to light exposure may be a marker of BD (Lewy, Wehr, Goodwin, Newsome & Rosenthal, 1981; Phelps, 2008) although subsequent research has failed to replicate this finding (Nurnberger et al., 2000).

Reduced physical activity and unstable social rhythms are also implicated in sleep and circadian disruptions within the mood disorders (Frank et al., 1994). Indeed, lower activity levels that are characteristic of the mood disorders are hypothesized as contributors to sleep disturbance (e.g., Harvey et al. 2005). Similarly, depression and and interepisode BD have been associated with lower social support (Moos, Cronkite & Moos, 1998; Eidelman et al., 2010) and more social strain (Franks et al. 1992). Two studies highlight that the onset of manic episodes are associated with life events that involve disruption of social rhythms (Malkoff-Schwartz, et al., 1998; Malkoff-Schwartz, et al., 2000).

Taken together, the sleep and circadian disruptions observed in mood disorders point to specific deficiencies in entrainment.

How does the basic science inform intervention?

Because sleep and circadian processes are open systems, disturbances in the sleep and circadian systems are modifiable. In the section that follows, several treatment breakthroughs are described that draw directly from the basic science provided by sleep and circadian researchers.

Cognitive Behavioral Therapy for Insomnia (CBT-I)

Cognitive Behavioral Therapy (CBT) was originally designed to target problematic thoughts and behaviors, which are characteristic of individuals with psychiatric disorders (Beck, 1964). Its effectiveness in altering negative thought and behavior patterns has been demonstrated across psychiatric disorders (see Beck, 2005 for a review).

Cognitive Behavioral Therapy for Insomnia (CBT-I) is a modification of classic CBT. CBT-I aims to change sleep habits and scheduling factors, as well as misconceptions about sleep and insomnia, that perpetuate sleep difficulties. CBT-I is a multi-component treatment. It is a short treatment, typically unfolding over 2 to 8 50-minute sessions, delivered individually or in groups. As demonstrated in Table 1, the treatment components of this intervention are directly drawn from basic science research.

CBT-I is currently the recommended first-line treatment for insomnia. Though hypnotic medications are effective for the short-term management of insomnia, there is limited evidence about sustained efficacy with long-term use (Krystal, 2009). In contrast, a review of CBT-I outcomes by the Standards of Practice Committee of the American Academy of Sleep Medicine found that CBT-I is highly effective in achieving both short-term improvements in sleep and sustained gains over long-term follow-up (Morin et al., 2006). Comparing short-term effects of pharmacological treatments for insomnia and CBT-I, a meta-analysis of 21 RCTs concluded that treatment effects for the two treatment modalities were generally comparable, though CBT-I is more effective in decreasing time to initiate sleep (Smith et al., 2002). Another study by Sivertsen et al. (2006) compared CBT-I, zopiclone and placebo. The results clearly favored CBT-I, which resulted in improved short and long-term functioning relative to zopiclone.

As insomnia is a pervasive problem across psychiatric conditions (Lichstein, Wilson, & Johnson, 2000; Ohayon & Roth, 2001), adaptations of CBT-I have been increasingly evaluated in the context of comorbid psychiatric illness in an effort to improve outcomes. Consistent with the growing evidence that insomnia and comorbid psychiatric disorders mutually maintain one another, accruing evidence suggests that CBT-I results in improvements not only in insomnia symptoms, but also psychiatric symptoms (Smith, Huang, & Manber, 2005). Recent studies have explored the application of CBT-I to MDD and BD.

In a sample of patients with MDD prescribed escitalopram, Manber et al. (2008) randomized half to CBT-I while the other half received placebo psychotherapy. Relative to placebo psychotherapy, the addition of CBT-I to pharmacologic anti-depressant treatment resulted in significantly higher rates of remission from both insomnia (7.7% vs. 50%) and depression (33.3% vs. 61.5%). In a later study, Manber et al. (2011) evaluated whether response to CBT-I treatment differed in patients with severe versus mild depression, comorbid with insomnia. Results indicated that improvement in insomnia, perceived energy, productivity, self-esteem, and overall treatment satisfaction did not differ by depression severity. Moreover, following CBT-I patients with more severe depressive symptoms experienced significant reductions in suicidal ideation and depression severity.

As discussed earlier, sleep complaints are common across all phases of BD (Jackson et al., 2003). Importantly, insomnia persists between episodes in 55% of patients (Harvey et al., 2005). In a recent case series, Kaplan and Harvey (2013) reported that CBT-I has a positive impact on sleep in patients with inter-episode BD and comorbid insomnia. Regularizing bedtimes and rise times was often sufficient to bring about improvements in sleep. This finding indicates that circadian disruptions in BD may be a key factor maintaining comorbid insomnia. Moreover, when changes in mood and daytime sleepiness were carefully monitored, CBT-I was found to be a safe and efficacious treatment for insomnia in BD.

Interpersonal and Social Rhythm Therapy (IPSRT)

The social zeitgeber hypothesis proposes that major life events (e.g., loss of a significant relationship or a job) not only have psychological salience, but also destabilize key zeitgebers, such as daily activities and light exposure (Frank et al., 1997). Social Rhythm Therapy is a psychotherapy based on the social zeitgeber hypothesis designed to increase stability in social rhythms. In the treatment of BD, Social Rhythm Therapy is typically integrated with principles from interpersonal psychotherapy in a treatment known as Interpersonal and Social Rhythm Therapy (IPSRT; E. Frank, et al., 1994).

IPSRT has 3 main goals 1) to help patients with BD stabilize their daily routines, 2) to assist patients in noticing the bi-directional relationship between mood and interpersonal interactions and 3) to ameliorate interpersonal difficulties (Frank, 2007; Miklowitz, 2006). The beginning of IPSRT involves the tracking of daily sleep timing, activities, social interactions and mood. After several weeks of tracking their daily activities, patients begin to see the connections between their sleep and daily activities and mood. The therapist and patient collaboratively create a plan to stabilize the patients' social schedule and sleep timing in order to regularize the circadian system.

The effectiveness of IPSRT for BD has been demonstrated in several studies (Frank, et al., 1997; Frank et al., 2005; Miklowitz, et al., 2007). A study by Frank et al. (1997) randomly assigned patients with BD to medication, active clinical management (a conventional clinic approach of medication combined with counseling), or IPSRT. When compared with active clinical management, patients receiving IPSRT experienced greater stability in daily routines and more time prior to recurrences of mood episodes. IPSRT was most effective in delaying mood episodes in patients who succeeded in stabilizing their daily routines. Patients who received IPSRT were also more likely to remain well through a 2-year follow-up (Frank et

al., 2005). This effect was mediated by the substantially increased regularity of social routines among subjects receiving IPSRT. Another study (Miklowitz et al., 2007) compared a combined treatment of IPSRT and Family Focused Therapy (FFT) to brief psychoeducational crisis management in a medicated sample of patients with BD. During one year, patients who received IPSRT combined with FFT experienced longer periods of remission.

Taken together, the literature suggests that psychosocial intervention designed to improve biological rhythm regularity, such as IPSRT, has beneficial effects on various psychiatric symptoms associated with a range of disorders (Malkoff-Schwartz et al., 2000).

Light and Dark Therapy

In 1984, Rosenthal and colleagues published a case series on recurrent depression with a seasonal pattern (seasonal affective disorder). The experimental treatment of a sample of these depressed patients with artificial bright light resulted in remission from depression (Rosenthal et al., 1984). The researchers ability to modify the circadian system through the manipulation of a powerful zeitgeber, light, laid the groundwork for the development and refinement of light therapy.

Light therapy attempts to reset the phase of the endogenous circadian clock relative to the light-dark cycle and/or increase its amplitude. The goals of light therapy include 1) synchronizing the sleep-wake cycle with the subjective night, 2) shifting the circadian clock to facilitate sleep at a desired time of day/night, and 3) advancing the circadian clock to attain indirect effects on mood (Wirz-Justice, Benedetti & Terman, 2009; Shirani & Louis, 2009). Light treatment involves the patient sitting in front of on a light therapy unit at a prescribed distance yielding an intensity of light ranging from 2,500–10,000 lux (Terman, Amira, Terman & Ross, 1996). The optimal dose and timing of bright light therapy is a matter of ongoing interest (Anderson, Glod, Dai, Cao, & Lockley, 2009), particularly in individuals who are light sensitive (Liebenluft et al., 1995). Effective timing of light administration is best predicted by an individuals' circadian timing (Terman, Terman, Lo, & Cooper, 2001). The patient is not required to look directly at the unit, indeed most patients read or eat while undergoing light exposure. Side effects associated with light therapy may include eyestrain, headache, nausea, and agitation. As some patients are resistant to sitting in front of a light box, clinicians often use management of natural light for similar therapeutic effects. Managing exposure to light in the evening can help patients down regulate sufficiently and allow the biology governing sleep onset to take over (Wallace-Guy et al., 2002). For example, turning down lights, turning off television, computers and cell phones an hour before bedtime can be a simple but powerful intervention for some patients.

While bright light exposure is a first line treatment for depression with a seasonal pattern, a few studies have also demonstrated effectiveness in non-seasonal depression (Even, Schröder, Friedman, & Rouillon, 2008; Golden, et al., 2005). A meta-analysis of placebo-controlled bright light studies found support for the use of bright light treatment in non-seasonal MDD and in seasonal affective disorder, with effect sizes of 0.53 and 0.84, respectively (Golden et al., 2005). Interestingly, bright light therapy delivered as an adjunct to antidepressant pharmacotherapy for non-seasonal depression was not effective (Golden et

al., 2005). Limited trials of light therapy have also demonstrated the effectiveness of light therapy in post-partum depression (Oren, et al., 2002) and chronic fatigue syndrome (Terman, Levine, Terman, & Doherty, 1998). Light therapy has also been associated with reduced suicidal ideation in 45% of patients with seasonal affective disorder (Lam, Tam, Shiah, Yatham & Zis, 2000). Notably, the presence of pre-treatment hypersomnia has been associated with favorable bright light treatment outcome (Terman et al., 1996; Avery et al., 1991)

There is a dearth of research investigating light therapy in patients with BD, as some reports have suggested that individuals with BD may be hypersensitive to the antidepressant effects of light. Indeed, hypomania or mania can occur as a potential adverse effect of light exposure in BD (Tuunainen, Kripke & Ende, 2004). For example, in a sample of women with bipolar depression, Sit, Wisner, Hanusa, Stull & Terman (2007) examined dosing of bright light, varying intensity and frequency of exposure across 2-week intervals. Three out of 4 of the women developed mixed mood states (including irritability, elevated energy and racing thoughts), while one entered a sustained period of remission. Thus, researchers have investigated light restriction (or dark therapy) as a treatment for mania (Wehr et al., 1998). Barbini et al. (2005) randomly allocated bipolar patients in a manic episode to 14 hours of darkness over three consecutive days or treatment as usual. Patients who received dark therapy exhibited a greater decrease in manic symptoms relative to the treatment as usual group. In a case report of two severely manic patients who were asked to enter a dark room for a daytime nap, both patients were asleep within 'seconds following eye closure' (Van Sweden, 1986). This may indicate that 'reduced need for sleep' experienced during mania may be associated with light hypersensitivity rather than a truly reduced sleep need. The few studies of dark therapy that exist are promising, and more research is needed in this domain for patients with BD.

Sleep Deprivation

Thirty years ago, the first experimental sleep deprivation study in a depressed patient with severe insomnia revealed an unexpected and paradoxical improvement in depression the following day (Pflug & Tölle 1971). The rapid overnight remission of severe depression following sleep deprivation remains one of the most striking phenomena in psychiatry, though depressive symptoms generally return after recovery sleep (Wu & Bunney, 1990).

Sleep deprivation, or wake therapy, generally consists of total sleep deprivation for one night or for the second half of one night (referred to as partial sleep deprivation). Because depressive symptoms rapidly return after subsequent sleep, sleep deprivation is not widely used alone (Wirz-Justice & Van den Hoofdakker, 1999). A combination of sleep deprivation (or partial sleep deprivation) with other treatments, including pharmacotherapy and light therapy, shows promise in achieving a rapid and maintained therapeutic response (Benedetti et al., 1997; Barbini, Colombo, Barbini, Campori & Smeraldi, 1999; Colombo, Benedetti, Barbini, Campori & Smeraldi, 2000). In order to verify that exposure to light during the night is not the active mechanism of sleep deprivation, researchers investigated whether sleep deprivation was also effective in the dark (Wehr, Rosenthal, Sack, & Gillin, 1985). Indeed, although patients' experienced much more difficulty staying awake, sleep

deprivation in the dark was also effective. Although the mechanisms are not well-understood, one theory posits that the antidepressant effects of sleep deprivation may be due to an enhancement of process S, and the relapse of depression following recovery sleep to return to baseline levels of the abnormal S process (Borbely & Wirz-Justice, 1982). In support of this hypothesis, a recent investigation reported that selectively deprived slow wave sleep (thus enhancing process S without disrupting total sleep time) in a small sample of unmedicated depressed patients led to an acute reduction in depressive symptomatology (Landsness, Goldstein, Peterson, Tononi & Benca, 2011).

Sleep deprivation in BD has potential benefits and risks. There is some evidence that, following partial sleep deprivation, bipolar depressed patients show a greater decrease in negative affect than recurrent unipolar depressed patients (Szuba, Baxter, Fairbanks, Guze, Schwartz, 1991; Barbini et al 1998). However, a risk when treating bipolar depression is a switch from depression into mania. In patients with BD, a close relationship has been observed between sleep loss and the onset of mania (e.g. Barbini et al., 1998). Indeed, an investigation of 3 cycles of sleep deprivation (alone or in combination with heterogeneous medications) in a sample of 206 patients with bipolar depression observed a 4.85% switch rate into mania and a 5.83% switch rate into hypomania (Colombo et al. 1999). Although, these percentages are low and comparable to those observed with antidepressant drug treatments, these results highlight the importance of careful monitoring of the sleep and wake patterns of patients with BD.

There are few studies investigating the consequences of sleep deprivation on the circadian system. One study found that after repeated partial sleep deprivation, the circadian rhythm of melatonin was delayed by 45 minutes (Lo et al., 2012). In total sleep deprivation conditions, because the circadian process is independent from the sleep process, circadian rhythms appear to modulate the timing but not the amount of accumulated total sleep rebound (Mistlberger et al., 1983). The circadian effects of sleep deprivation will be important to consider and manage in future research, particularly if this treatment becomes more widely used.

Overall, sleep deprivation has been described as the most rapid antidepressant available today. Data on hundreds of depressed patients of all diagnostic subcategories show marked improvement within hours in approximately 60% of patients (see Wirz-Justice et al., 2005 for review). Lack of antidepressant response to the first dose sleep deprivation does not mean that the patient will not respond to later doses of sleep deprivations (Fahndrich, 1988; Gordijn et al., 1995). However, the question of how to sustain the remarkable antidepressant response that occurs after sleep deprivation in some depressed patients remains an open domain for further investigation.

Conclusions

We have reviewed evidence suggesting that the sleep and circadian systems are 1) disrupted in the mood disorders and linked to symptoms, 2) open systems that can be modified, 3) the focus of interventions which have been developed to effectively treat sleep disturbance within mood disorders, and 4) intimately linked with mood, such that improvements in sleep

are associated with improvements in mood. Treatment of sleep and circadian disturbances within mood disorders are an exciting domain of research. Although, significant positive treatment effects are evident, more research is needed. Significant gaps include 1) our basic understanding of the relationship between sleep and mood, 2) the modification of existing treatments for sleep disturbance in mood disorders for children, adolescents, and the elderly, 3) improving existing treatments and/or developing novel treatments, and 4) dissemination research focusing on improving access to these efficacious treatments.

Acknowledgments

This research was supported by the National Science Foundation DGE 1106400 and the National Institute of Mental Health Grants R34 MH080958 and T32MH089919.

References

- Anderson JL, Glod CA, Dai J, Cao Y, Lockley SW. Lux vs. wavelength in light treatment of Seasonal Affective Disorder. *Acta Psychiatrica Scandinavica*. 2009; 120(3):203–212. [PubMed: 19207131]
- Antle MC, Mistlberger RE. Circadian clock resetting by sleep deprivation without exercise in the Syrian hamster. *The Journal of Neuroscience*. 2000; 20(24):9326–9332. [PubMed: 11125012]
- Avery DH, Khan A, Dager SR, Cohen S, Cox GB, Dunner DL. Morning or evening bright light treatment of winter depression? The significance of hypersomnia. *Biological Psychiatry*. 1991; 29(2):117–126. [PubMed: 1995083]
- Baldwin DS, Papakostas GI. Symptoms of fatigue and sleepiness in major depressive disorder. *The Journal of Clinical Psychiatry*. 2005; 67:9–15. [PubMed: 16848671]
- Barbini B, Benedetti F, Colombo C, Dotoli D, Bernasconi A, Cigala-Fulgosi M, Smeraldi E. Dark therapy for mania: a pilot study. *Bipolar Disorders*. 2005; 7(1):98–101. [PubMed: 15654938]
- Barbini B, Colombo C, Benedetti F, Campori E, Bellodi L, Smeraldi E. The unipolar–bipolar dichotomy and the response to sleep deprivation. *Psychiatry Research*. 1998; 79(1):43–50. [PubMed: 9676825]
- Beck AT. Thinking and depression: II. Theory and therapy. *Archives of General Psychiatry*. 1964; 10(6):561–571. [PubMed: 14159256]
- Beck AT. The current state of cognitive therapy: a 40-year retrospective. *Archives of General Psychiatry*. 2005; 62(9):953–959. [PubMed: 16143727]
- Benca RM, Obermeyer WH, Thisted RA, Gillin JC. Sleep and psychiatric disorders: a meta-analysis. *Archives of General Psychiatry*. 1992; 49(8):651–668. [PubMed: 1386215]
- Benedetti F, Barbini B, Lucca A, Campori E, Colombo C, Smeraldi E. Sleep deprivation hastens the antidepressant action of fluoxetine. *European Archives of Psychiatry and Clinical Neuroscience*. 1997; 247(2):100–103. [PubMed: 9177956]
- Benedetti F, Colombo C, Barbini B, Campori E, Smeraldi E. Ongoing lithium treatment prevents relapse after total sleep deprivation. *Journal of Clinical Psychopharmacology*. 1999; 19(3):240–245. [PubMed: 10350030]
- Benedetti F, Serretti A, Colombo C, Campori E, Barbini B, di Bella D, Smeraldi E. Influence of a functional polymorphism within the promoter of the serotonin transporter gene on the effects of total sleep deprivation in bipolar depression. *American Journal of Psychiatry*. 1999; 156(9):1450–1452. [PubMed: 10484962]
- Billiard M, Dolenc L, Aldaz C, Ondze B, Besset A. Hypersomnia associated with mood disorders: a new perspective. *Journal of Psychosomatic Research*. 1994; 38:41–47. [PubMed: 7799250]
- Boivin DB. Influence of sleep–wake and circadian rhythm disturbances in psychiatric disorders. *Journal of Psychiatry and Neuroscience*. 2000; 25(5):446–458. [PubMed: 11109296]
- Borbély AA, Wirz-Justice A. Sleep, sleep deprivation and depression: A hypothesis derived from a model of sleep regulation. *Human Neurobiology*. 1982; 1(3):205–210. [PubMed: 7185793]

- Buxton OM, Frank SA, L'Hermite-Baleriaux M, Leproult R, Turek FW, Van Cauter E. Roles of intensity and duration of nocturnal exercise in causing phase delays of human circadian rhythms. *American Journal of Physiology*. 1997; 273:536–542.
- Buyse DJ, Frank E, Lowe KK, Cherry CR, Kupfer DJ. Electroencephalographic sleep correlates of episode and vulnerability to recurrence in depression. *Biological Psychiatry*. 1997; 41(4):406–418. [PubMed: 9034535]
- Cambras T, Chiesa J, Araujo J, Díez-Noguera A. Effects of photoperiod on rat motor activity rhythm at the lower limit of entrainment. *Journal of Biological Rhythms*. 2004; 19(3):216–225. [PubMed: 15155008]
- Challet E, Turek FW, Laute MA, Van Reeth O. Sleep deprivation decreases phase-shift responses of circadian rhythms to light in the mouse: role of serotonergic and metabolic signals. *Brain Research*. 2001; 909(1):81–91. [PubMed: 11478924]
- Chen YW, Dilsaver SC. Lifetime rates of suicide attempts among subjects with bipolar and unipolar disorders relative to subjects with other Axis I disorders. *Biological Psychiatry*. 1996; 39(10):896–899. [PubMed: 8860192]
- Colombo C, Benedetti F, Barbini B, Campori E, Smeraldi E. Rate of switch from depression into mania after therapeutic sleep deprivation in bipolar depression. *Psychiatry Research*. 1999; 86(3):267–270. [PubMed: 10482346]
- Colton CW, Manderscheid RW. Congruencies in increased mortality rates, years of potential life lost, and causes of death among public mental health clients in eight states. *Preventing Chronic Disease*. 2006; 3(2):1–14.
- Czeisler CA, Kronauer RE, Allan JS, Duffy JF, Jewett ME, Brown EN, Ronda JM. Bright light induction of strong (type 0) resetting of the human circadian pacemaker. *Science*. 1989; 244(4910):1328–1333. [PubMed: 2734611]
- Czeisler CA, Wright KP. Influence of light on circadian rhythmicity in humans. *Lung Biology in Health and Disease*. 1999; 133:149–180.
- Detre T, Himmelijoch J, Swartzburg M. Hypersomnia and manic-depressive disease. *American Journal of Psychiatry*. 1972; 128(10):1303–1305. [PubMed: 4335278]
- Dew MA, Reynolds CF III, Houck PR, Hall M, Buysse DJ, Frank E, Kupfer DJ. Temporal profiles of the course of depression during treatment: predictors of pathways toward recovery in the elderly. *Archives of General Psychiatry*. 1997; 54(11):1016–1024. [PubMed: 9366658]
- Dijk DJ, Czeisler CA. Paradoxical timing of the circadian rhythm of sleep propensity serves to consolidate sleep and wakefulness in humans. *Neuroscience Letters*. 1994; 166(1):63–68. [PubMed: 8190360]
- Dinges DF, Pack F, Williams K, Gillen KA, Powell JW, Ott GE, Pack AI. Cumulative sleepiness, mood disturbance and psychomotor vigilance performance decrements during a week of sleep restricted to 4–5 hours per night. *Sleep: Journal of Sleep Research & Sleep Medicine*. 1997; 20(4):267–277.
- Eaton WW, Badawi M, Melton B. Prodromes and precursors: epidemiologic data for primary prevention of disorders with slow onset. *The American Journal of Psychiatry*. 1995; 152(7):967–972.
- Edgar DM, Dement WC. Regularly scheduled voluntary exercise synchronizes the mouse circadian clock. *American Journal of Physiology*. 1991; 261:928–931.
- Edinger JD, Wohlgemuth WK. The significance and management of persistent primary insomnia: the past, present and future of behavioral insomnia therapies. *Sleep Medicine Reviews*. 1999; 3(2):101–118. [PubMed: 15310480]
- Eidelman P, Talbot LS, Gruber J, Hairston I, Harvey AG. Sleep architecture as correlate and predictor of symptoms and impairment in interepisode bipolar disorder: taking on the challenge of medication effects. *Journal of Sleep Research*. 2010; 19(4):516–524. [PubMed: 20408930]
- Emens J, Lewy A, Kinzie JM, Arntz D, Rough J. Circadian misalignment in major depressive disorder. *Psychiatry Research*. 2009; 168(3):259–261. [PubMed: 19524304]
- Even C, Schröder CM, Friedman S, Rouillon F. Efficacy of light therapy in nonseasonal depression: a systematic review. *Journal of Affective Disorders*. 2008; 108(1):11–23. [PubMed: 17950467]

- Fähndrich, E. *Klinische Diagnostik und Therapie der Depression*. Regensburg: Roderer Verlag; 1988. *Chronobiologie der Depression: therapeutische und theoretische Aspekte des Schlafentzuges*; p. 126-141.
- Ford DE, Kamerow DB. Epidemiologic study of sleep disturbances and psychiatric disorders. *JAMA: The Journal of the American Medical Association*. 1989; 262(11):1479-1484. [PubMed: 2769898]
- Frank, E. *Treating bipolar disorder: A clinician's guide to interpersonal and social rhythm therapy*. Guilford Press; 2007.
- Frank E, Hlastala S, Ritenour A, Houck P, Tu XM, Monk TH, Kupfer DJ. Inducing lifestyle regularity in recovering bipolar disorder patients: results from the maintenance therapies in bipolar disorder protocol. *Biological Psychiatry*. 1997; 41(12):1165-1173. [PubMed: 9171907]
- Frank E, Kupfer DJ, Ehlers CL, Monk TH. Interpersonal and social rhythm therapy for bipolar disorder: integrating interpersonal and behavioral approaches. *Behaviour Therapist*. 1994; 17:143-143.
- Frank E, Kupfer DJ, Thase ME, Mallinger AG, Swartz HA, Fagiolini AM, Monk T. Two-year outcomes for interpersonal and social rhythm therapy in individuals with bipolar I disorder. *Archives of General Psychiatry*. 2005; 62(9):996-1004. [PubMed: 16143731]
- Franks P, Shields C, Campbell T, McDaniel S, Harp J, Botelho RJ. Association of social relationships with depressive symptoms: Testing an alternative to social support. *Journal of Family Psychology*. 1992; 6(1):49-59.
- Freeman D, Pugh K, Vorontsova N, Southgate L. Insomnia and paranoia. *Schizophrenia Research*. 2009; 108(1):280-284. [PubMed: 19097752]
- Germain A, Richardson R, Moul DE, Mammen O, Haas G, Forman SD, Nofzinger EA. Placebo-controlled comparison of prazosin and cognitive-behavioral treatments for sleep disturbances in US military veterans. *Journal of Psychosomatic Research*. 2012; 72(2):89-96. [PubMed: 22281448]
- Golden RN, Gaynes BN, Ekstrom RD, Hamer RM, Jacobsen FM, Suppes T, Nemeroff CB. The efficacy of light therapy in the treatment of mood disorders: a review and meta-analysis of the evidence. *American Journal of Psychiatry*. 2005; 162(4):656-662. [PubMed: 15800134]
- Gordijn MCM, Beersma DGM, Bouhuys AL, Korte HJ, Van den Hoofdakker RH. A longitudinal study of sleep deprivation responses in depression: the variability is highly related to diurnal mood variability. *Acta Neuropsychiatry*. 1995; 7:58-60.
- Gotlib IH, Joormann J. Cognition and depression: current status and future directions. *Annual Review of Clinical Psychology*. 2010; 27(6):285-312.
- Gross JJ, Muñoz RF. Emotion regulation and mental health. *Clinical Psychology: Science and Practice*. 1995; 2(2):151-164.
- Gruber S, Rathgeber K, Bräunig P, Gauggel S. Stability and course of neuropsychological deficits in manic and depressed bipolar patients compared to patients with major depression. *Journal of Affective Disorders*. 2007; 104(1):61-71. [PubMed: 17360041]
- Hanneman SK. Measuring circadian temperature rhythm. *Biological Research for Nursing*. 2001; 2(4):236-248. [PubMed: 11876463]
- Harvey AG. Insomnia, psychiatric disorders, and the transdiagnostic perspective. *Current Directions in Psychological Science*. 2008a; 17(5):299-303.
- Harvey A. Sleep and circadian rhythms in bipolar disorder: seeking synchrony, harmony, and regulation. *American Journal of Psychiatry*. 2008b; 165(7):820-829. [PubMed: 18519522]
- Harvey AG, Schmidt DA, Scarnà A, Semler CN, Goodwin GM. Sleep-related functioning in euthymic patients with bipolar disorder, patients with insomnia, and subjects without sleep problems. *American Journal of Psychiatry*. 2005; 162(1):50-57. [PubMed: 15625201]
- Horne JA. Sleep function, with particular reference to sleep deprivation. *Annals of Clinical Research*. 1985; 17(5):199-208. [PubMed: 3909914]
- Insel T. Assessing the economic costs of serious mental illness. *American Journal of Psychiatry*. 2008; 165(6):663-665. [PubMed: 18519528]
- Insel TR. Translating scientific opportunity into public health impact: a strategic plan for research on mental illness. *Archives of General Psychiatry*. 2009; 66(2):128-133. [PubMed: 19188534]

- Insel TR, Charney DS. Research on major depression. *JAMA: The Journal of the American Medical Association*. 2003; 289(23):3167–3168. [PubMed: 12813123]
- Insel TR, Cuthbert BN, Garvey MA, Heinssen RK, Pine DS, Quinn KJ, Wang PS. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*. 2010; 167(7):748–751. [PubMed: 20595427]
- Jackson A, Cavanagh J, Scott J. A systematic review of manic and depressive prodromes. *Journal of Affective Disorders*. 2003; 74(3):209–217. [PubMed: 12738039]
- Jang H, Lee G. Feed period restriction alters the expression of peripheral circadian rhythm genes without changing body weight in mice. *PLoS One*. 2012; 7:49993.
- Johnson, CH. *An Atlas of Phase Response Curves for Circadian and Circatidal Rhythms*. Department of Biology, Vanderbilt University; Nashville, TN, USA: 1990.
- Kaplan KA, Gruber J, Eidelman P, Talbot LS, Harvey AG. Hypersomnia in inter-episode bipolar disorder: Does it have prognostic significance? *Journal of Affective Disorders*. 2011; 132(3):438–444. [PubMed: 21489637]
- Kaplan KA, Harvey AG. Hypersomnia across mood disorders: a review and synthesis. *Sleep Medicine Reviews*. 2009; 13(4):275–285. [PubMed: 19269201]
- Kaplan KA, Harvey AG. Behavioral Treatment of Insomnia in Bipolar Disorder. *American Journal of Psychiatry*. 2013 In press.
- Kessler RC, Heeringa S, Lakoma MD, Petukhova M, Rupp AE, Schoenbaum M, Zaslavsky AM. The individual-level and societal-level effects of mental disorders on earnings in the United States: Results from the National Comorbidity Survey Replication. *The American Journal of Psychiatry*. 2008; 165(6):703–711. [PubMed: 18463104]
- Klerman EB, Shanahan TL, Brotman DJ, Rimmer DW, Emens JS, Rizzo J3, Czeisler CA. Photic resetting of the human circadian pacemaker in the absence of conscious vision. *Journal of Biological Rhythms*. 2002; 17(6):548–555. [PubMed: 12465888]
- Korszun A, Moskvina V, Brewster S, Craddock N, Ferrero F, Gill M, McGuffin P. Familiality of symptom dimensions in depression. *Archives of General Psychiatry*. 2004; 61(5):468–474. [PubMed: 15123491]
- Kripke DF, Juarez S, Cole RJ, Ancoli-Israel S, Hauri PJ, Wisbey JA, Gruen W. Adult illumination exposures and some correlations with symptoms. *Evolution of Circadian Clock*. 1994:349–360.
- Kupfer DJ. REM latency: a psychobiologic marker for primary depressive disease. *Biological Psychiatry*. 1976; 11(2):159–174. [PubMed: 183839]
- Landsness EC, Goldstein MR, Peterson MJ, Tononi G, Benca RM. Antidepressant effects of selective slow wave sleep deprivation in major depression: a high-density EEG investigation. *Journal of Psychiatric Research*. 2011; 45(8):1019–1026. [PubMed: 21397252]
- Lam RW, Tam EM, Shiah IS, Yatham LN, Zis AP. Effects of light therapy on suicidal ideation in patients with winter depression. *Journal of Clinical Psychiatry*. 2000; 61(1):30–32. [PubMed: 10695643]
- Leibenluft E, Feldman-Naim S, Turner EH, Schwartz PJ, Wehr TA. Salivary and plasma measures of dim light melatonin onset (DLMO) in patients with rapid cycling bipolar disorder. *Biological Psychiatry*. 1996; 40(8):731–735. [PubMed: 8894065]
- Lewy A, Wehr T, Goodwin F, Newsome D, Rosenthal N. Manic-depressive patients may be supersensitive to light. *The Lancet*. 1981; 317(8216):383–384.
- Lichstein KL, Wilson NM, Johnson CT. Psychological treatment of secondary insomnia. *Psychology and Aging*. 2000; 15(2):232–240. [PubMed: 10879578]
- Lo JC, Groeger JA, Santhi N, Arbon EL, Lazar AS, Hasan S, Dijk DJ. Effects of partial and acute total sleep deprivation on performance across cognitive domains, individuals and circadian phase. *PLoS One*. 2012; 7(9):e45987. [PubMed: 23029352]
- Malkoff-Schwartz S, Frank E, Anderson BP, Hlastala SA, Luther JF, Sherrill JT, Kupfer DJ. Social rhythm disruption and stressful life events in the onset of bipolar and unipolar episodes. *Psychological Medicine*. 2000; 30(05):1005–1016. [PubMed: 12027038]
- Malkoff-Schwartz S, Frank E, Anderson B, Sherrill JT, Siegel L, Patterson D, Kupfer DJ. Stressful life events and social rhythm disruption in the onset of manic and depressive bipolar episodes: a

- preliminary investigation. *Archives of General Psychiatry*. 1998; 55(8):702–707. [PubMed: 9707380]
- Manber R, Bernert RA, Suh S, Nowakowski S, Siebern AT, Ong JC. CBT for insomnia in patients with high and low depressive symptom severity: adherence and clinical outcomes. *Journal of Clinical Sleep Medicine: JCSM: Official Publication of the American Academy of Sleep Medicine*. 2011; 7(6):645–652. [PubMed: 22171204]
- Manber R, Edinger JD, Gress JL, San Pedro-Salcedo MG, Kuo TF, Kalista T. Cognitive behavioral therapy for insomnia enhances depression outcome in patients with comorbid major depressive disorder and insomnia. *Sleep*. 2008; 31:489–495. [PubMed: 18457236]
- Miklowitz DJ. A review of evidence-based psychosocial interventions for bipolar disorder. *Journal of Clinical Psychiatry*. 2006; 67(11):28–33. [PubMed: 17029494]
- Miklowitz DJ, Otto MW, Frank E, Reilly-Harrington NA, Wisniewski SR, Kogan JN, Sachs GS. Psychosocial treatments for bipolar depression: a 1-year randomized trial from the Systematic Treatment Enhancement Program. *Archives of General Psychiatry*. 2007; 64(4):419–426. [PubMed: 17404119]
- Mistlberger RE, Antle MC, Glass JD, Miller JD. Behavioral and serotonergic regulation of circadian rhythms. *Biological Rhythm Research*. 2000; 31(3):240–283.
- Mistlberger RE, Bergmann BM, Waldenar W, Rechtschaffen A. Recovery sleep following sleep deprivation in intact and suprachiasmatic nuclei-lesioned rats. *Sleep: Journal of Sleep Research & Sleep Medicine*. 1983; 6(3):217–233.
- Mistlberger RE, Landry GJ, Marchant EG. Sleep deprivation can attenuate light-induced phase shifts of circadian rhythms in hamsters. *Neuroscience Letters*. 1997; 238(1):5–8. [PubMed: 9464641]
- Mistlberger RE, Skene DJ. Nonphotic entrainment in humans? *Journal of Biological Rhythms*. 2005; 20(4):339–352. [PubMed: 16077153]
- Monk TH, Petrie SR, Hayes AJ, Kupfer DJ. Regularity of daily life in relation to personality, age, gender, sleep quality and circadian rhythms. *Journal of Sleep Research*. 1994; 3(4):196–205. [PubMed: 10607126]
- Morin CM, Bootzin RR, Buysse DJ, Edinger JD, Espie CA, Lichstein KL. Psychological and behavioral treatment of insomnia: update of the recent evidence (1998–2004). *Sleep*. 2006; 29(11):1398–1414. [PubMed: 17162986]
- Moos RH, Cronkite RC, Moos BS. Family and extrafamily resources and the 10-year course of treated depression. *Journal of Abnormal Psychology*. 1998; 107(3):450–460. [PubMed: 9715580]
- Myers E, Startup H, Freeman D. Cognitive behavioural treatment of insomnia in individuals with persistent persecutory delusions: a pilot trial. *Journal of Behavior Therapy and Experimental Psychiatry*. 2011; 42(3):330–336. [PubMed: 21367359]
- Nelson W, Halberg F. Meal-timing, circadian rhythms and the life span of mice. *The Journal of Nutrition*. 1986; 116:2244–2253. [PubMed: 3794831]
- Nierenberg AA, Husain MM, Trivedi MH, Fava M, Warden D, Wisniewski SR, Rush AJ. Residual symptoms after remission of major depressive disorder with citalopram and risk of relapse: a STAR* D report. *Psychological Medicine*. 2010; 40(01):41–50. [PubMed: 19460188]
- NIH. National Institutes of Health State-of-the-Science Conference Statement on Manifestations and Management of Chronic Insomnia in Adults (final statement). *Sleep*. 2005; 28:1049–1057. [PubMed: 16268373]
- Nurnberger JI Jr, Adkins S, Lahiri DK, Mayeda A, Hu K, Lewy A, Davis-Singh D. Melatonin suppression by light in euthymic bipolar and unipolar patients. *Archives of General Psychiatry*. 2000; 57(6):572–579. [PubMed: 10839335]
- Oren DA, Wisner KL, Spinelli M, Epperson CN, Peindl KS, Terman JS, Terman M. An open trial of morning light therapy for treatment of antepartum depression. *American Journal of Psychiatry*. 2002; 159(4):666–669. [PubMed: 11925310]
- Perlman CA, Johnson SL, Mellman TA. The prospective impact of sleep duration on depression and mania. *Bipolar disorders*. 2006; 8(3):271–274. [PubMed: 16696829]
- Pflug B, Tölle R. Therapie endogener Depressionen durch Schlafentzug. *Nervenarzt*. 1971; 42(117):1124.

- Phelps EA. Human emotion and memory: interactions of the amygdala and hippocampal complex. *Current Opinion in Neurobiology*. 2004; 14(2):198–202. [PubMed: 15082325]
- Phelps J. Dark therapy for bipolar disorder using amber lenses for blue light blockade. *Medical Hypotheses*. 2008; 70(2):224–229. [PubMed: 17637502]
- Reppert SM, Weaver DR. Coordination of circadian timing in mammals. *Nature*. 2002; 418(6901):935–941. [PubMed: 12198538]
- Roenneberg T, Foster RG. Twilight times: light and the circadian system. *Photochemistry and Photobiology*. 1997; 66(5):549–561. [PubMed: 9383985]
- Rosenthal NE, Sack DA, Gillin JC, Lewy AJ, Goodwin FK, Davenport Y, Wehr TA. Seasonal affective disorder: a description of the syndrome and preliminary findings with light therapy. *Archives of General Psychiatry*. 1984; 41(1):72–80. [PubMed: 6581756]
- Shirani A, Louis EKS. Illuminating rationale and uses for light therapy. *Journal of clinical sleep medicine: JCSM: Official Publication of the American Academy of Sleep Medicine*. 2009; 5(2):155–163. [PubMed: 19968050]
- Sit D, Wisner KL, Hanusa BH, Stull S, Terman M. Light therapy for bipolar disorder: a case series in women. *Bipolar Disorders*. 2007; 9(8):918–927. [PubMed: 18076544]
- Sivertsen B, Omvik S, Pallesen S, Bjorvatn B, Havik OE, Kvale G, Nordhus IH. Cognitive behavioral therapy vs zopiclone for treatment of chronic primary insomnia in older adults. *JAMA: the Journal of the American Medical Association*. 2006; 295(24):2851–2858. [PubMed: 16804151]
- Smith MT, Perlis ML, Park A, Smith MS, Pennington J, Giles DE, Buysse DJ. Comparative meta-analysis of pharmacotherapy and behavior therapy for persistent insomnia. *American Journal of Psychiatry*. 2002; 159(1):5–11. [PubMed: 11772681]
- Sotres-Bayon F, Bush DE, LeDoux JE. Emotional perseveration: an update on prefrontal-amygdala interactions in fear extinction. *Learning & Memory*. 2004; 11(5):525–535. [PubMed: 15466303]
- Sulzman FM, Fuller CA, Moore-Ede MC. Feeding time synchronizes primate circadian rhythms. *Physiology & Behavior*. 1977; 18(5):775–779. [PubMed: 410048]
- Szuba MP, Baxter LR Jr, Fairbanks LA, Guze BH, Schwartz JM. Effects of partial sleep deprivation on the diurnal variation of mood and motor activity in major depression. *Biological Psychiatry*. 1991; 30(8):817–829. [PubMed: 1751624]
- Terman M, Amira L, Terman JS, Ross DC. Predictors of response and nonresponse to light treatment for winter depression. *American Journal of Psychiatry*. 1996; 153(11):1423–1429. [PubMed: 8890675]
- Terman M, Levine SM, Terman JS, Doherty S. Chronic fatigue syndrome and seasonal affective disorder: comorbidity, diagnostic overlap, and implications for treatment. *The American Journal of Medicine*. 1998; 105(3):115S–124S. [PubMed: 9790493]
- Terman JS, Terman M, Lo ES, Cooper TB. Circadian time of morning light administration and therapeutic response in winter depression. *Archives of General Psychiatry*. 2001; 58(1):69–75. [PubMed: 11146760]
- Thase ME. Depression, sleep, and antidepressants. *The Journal of Clinical Psychiatry*. 1998; 59(4):55–65. [PubMed: 9554322]
- Thase ME, Buysse DJ, Frank E, Cherry CR, Cornes CL, Mallinger AG, Kupfer DJ. Which depressed patients will respond to interpersonal psychotherapy? The role of abnormal EEG sleep profiles. *American Journal of Psychiatry*. 1997; 154(4):502–509. [PubMed: 9090337]
- Tuunainen A, Kripke DF, Endo T. Light therapy for non-seasonal depression. *Cochrane Database Syst Rev*. 2004:2.
- Van Dongen HP, Maislin G, Mullington JM, Dinges DF. The cumulative cost of additional wakefulness: dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep*. 2003; 26(2):117–129. [PubMed: 12683469]
- Van Reeth O, Sturis J, Byrne MM, Blackman JD, L'Hermite-Baleriaux M, LeProult R, Van Cauter E. Nocturnal exercise phase delays circadian rhythms of melatonin and thyrotropin secretion in normal men. *American Journal of Physiology-Endocrinology and Metabolism*. 1994; 266(6):E964–E974.

- Van Sweden B. Disturbed vigilance in mania. *Biological Psychiatry*. 1986; 21(3):311–313. [PubMed: 3947712]
- Walker MP, Liston C, Hobson JA, Stickgold R. Cognitive flexibility across the sleep–wake cycle: REM-sleep enhancement of anagram problem solving. *Cognitive Brain Research*. 2002; 14(3): 317–324. [PubMed: 12421655]
- Wallace-Guy GM, Kripke DF, Jean-Louis G, Langer RD, Elliott JA, Tuunainen A. Evening light exposure: Implications for sleep and depression. *Journal of the American Geriatrics Society*. 2002; 50(4):738–739. [PubMed: 11982677]
- Webb WB. An objective behavioral model of sleep. *Sleep: Journal of Sleep Research & Sleep Medicine*. 1988; 11(5):488–496.
- Wehr TA, Rosenthal NE, Sack DA, Gillin JC. Antidepressant effects of sleep deprivation in bright and dim light. *Acta Psychiatrica Scandinavica*. 1985; 72(2):161–165. [PubMed: 4050508]
- Wehr TA, Turner EH, Shimada JM, Lowe CH, Barker C, Leibenluft E. Treatment of a rapidly cycling bipolar patient by using extended bed rest and darkness to stabilize the timing and duration of sleep. *Biological Psychiatry*. 1998; 43(11):822–828. [PubMed: 9611672]
- Wehr TA, Wirz-Justice A, Goodwin FK, Duncan W, Gillin JC. Phase advance of the circadian sleep–wake cycle as an antidepressant. *Science*. 1979; 206(4419):710–713. [PubMed: 227056]
- Weiland-Fiedler P, Erickson K, Waldeck T, Luckenbaugh DA, Pike D, Bonne O, Neumeister A. Evidence for continuing neuropsychological impairments in depression. *Journal of Affective Disorders*. 2004; 82(2):253–258. [PubMed: 15488254]
- Winokur, G.; Clayton, P.J.; Reich, T. Manic depressive illness. St Louis: CV Mosby; 1969.
- Winokur G, Tanna VL. Possible role of X-linked dominant factor in manic depressive disease. *Diseases of the Nervous System*. 1969; 30(2):89–94. [PubMed: 4975420]
- Wirz-Justice A, Benedetti F, Berger M, Lam RW, Martiny K, Terman M, Wu JC. Chronotherapeutics (light and wake therapy) in affective disorders. *Psychological Medicine*. 2005; 35(07):939–944. [PubMed: 16045060]
- Wirz-Justice, A.; Benedetti, F.; Terman, M. Chronotherapeutics for affective disorders. Basel: Karger; 2009.
- Wirz-Justice A, Van den Hoofdakker RH. Sleep deprivation in depression: what do we know, where do we go? *Biological Psychiatry*. 1999; 46(4):445–453. [PubMed: 10459393]
- World Health Organization. World health report 2002. Reducing risks, promoting healthy life. Vol. 2002. Geneva: WHO; 2003.
- Wu JC, Bunney WE. The biological basis of an antidepressant response to sleep deprivation and relapse: review and hypothesis. *The American Journal of Psychiatry*. 1990; 147(1):14–21. [PubMed: 2403471]
- Yoo SS, Gujar N, Hu P, Jolesz FA, Walker MP. The human emotional brain without sleep—a prefrontal amygdala disconnect. *Current Biology*. 2007; 17(20):R877–R878. [PubMed: 17956744]
- Zeitler JM, Dijk DJ, Kronauer RE, Brown EN, Czeisler CA. Sensitivity of the human circadian pacemaker to nocturnal light: melatonin phase resetting and suppression. *The Journal of Physiology*. 2000; 526(3):695–702. [PubMed: 10922269]
- Zohar D, Tzischinsky O, Epstein R, Lavie P. The effects of sleep loss on medical residents' emotional reactions to work events: a cognitive-energy model. *Sleep: Journal of Sleep and Sleep Disorders Research*. 2005; 28(1):47–54.

Table 1

How Behavioral Change Components of CBT-I Map onto Basic Sleep and Circadian Science.

Behavior Changes Targeted with CBT-I	How Maps Onto Basic Science
Go to bed only when sleepy	Allows for adequate build-up of sleep drive prior to bedtime (Process S)
Wind down before bed	Decreases hyperarousal, and when done regularly, conditions the body to prepare for sleep (Process C and Process S)
Getting out of bed during nighttime awakenings	Conditioning bed as cue for sleep
Restricted time in bed	Sleep restriction increases sleep drive (Process S)
Regular risetime	Entrainment of circadian phase (Process C); If a poor night of sleep, allows for greater build-up of sleep drive the following day (Process S)
Sunlight or bright light on awakening	Entrainment of circadian phase (Process C)
No naps	Serves to increase sleep drive (Process S)