- Danish adults: the Copenhagen allergy study. *J Allergy Clin Immunol* 2003;111:847-53.
- 6 Wickens KL, Crane J, Kemp TJ, Lewis SJ, D'Souza WJ, Sawyer GM, et al. Family size, infections, and asthma prevalence in New Zealand children. *Epidemiology* 1999;10:699-705.
 - 7 Strachan DP, Taylor EM, Carpenter RG. Family structure, neonatal infection, and hay fever in adolescence. *Arch Dis Child* 1996;74:422-6.
 - 8 Forastiere F, Agabiti N, Corbo GM, Dell'Orco V, Porta D, Pistelli R, et al. Socioeconomic status, number of siblings, and respiratory infections in early life as determinants of atopy in children. *Epidemiology* 1997;8:566-70.
 - 9 Nafstad P, Magnus P, Jaakkola JJ. Early respiratory infections and childhood asthma. *Pediatrics* 2000;106:E38.
 - 10 Matricardi PM, Rosmini F, Ferrigno L, Nisini R, Rapicetta M, Chionne P, et al. Cross sectional retrospective study of prevalence of atopy among Italian military students with antibodies against hepatitis A virus. *BMJ* 1997;314:999-1003.
 - 11 Ponsonby AL, Couper D, Dwyer T, Carmichael A, Kemp A. Relationship between early life respiratory illness, family size over time, and the development of asthma and hay fever: a seven year follow up study. *Thorax* 1999;54:664-9.
 - 12 McKeever TM, Lewis SA, Smith C, Collins J, Heatlie H, Frischer M, et al. Early exposure to infections and antibiotics and the incidence of allergic disease: a birth cohort study with the West Midlands general practice research database. *J Allergy Clin Immunol* 2002;109:43-50.
 - 13 Olsen J, Melbye M, Olsen SF, Sørensen TI, Aaby P, Andersen AM, et al. The Danish national birth cohort—its background, structure and aim. *Scand J Public Health* 2001;29:300-7.
 - 14 Benn CS, Benfeldt E, Andersen PK, Olesen AB, Melbye M, Björkstén B. Atopic dermatitis in young children: diagnostic criteria for use in epidemiological studies based on telephone interviews. *Acta Derm Venereol* 2003;83:347-50.
 - 15 Matricardi PM, Ronchetti R. Are infections protecting from atopy? *Curr Opin Allergy Clin Immunol* 2001;1:413-9.
 - 16 Strannegård Ö, Strannegård IL. The causes of the increasing prevalence of allergy: is atopy a microbial deprivation disorder? *Allergy* 2001;56:91-102.
 - 17 Kalliomäki M, Isolauri E. Role of intestinal flora in the development of allergy. *Curr Opin Allergy Clin Immunol* 2003;3:15-20.
 - 18 Björkstén B. The intrauterine and postnatal environments. *J Allergy Clin Immunol* 1999;104:1119-27.
 - 19 Rystedt I, Strannegård IL, Strannegård Ö. Recurrent viral infections in patients with past or present atopic dermatitis. *Br J Dermatol* 1986;114:575-82.
 - 20 Böhme M, Lannerö E, Wickman M, Nordvall SL, Wahlgren CF. Atopic dermatitis and concomitant disease patterns in children up to two years of age. *Acta Derm Venereol* 2002;82:98-103.
- (Accepted 5 March 2004)
- doi 10.1136/bmj.38069.512245.FE

Commentary: the defence of dirt

Geoff Watts

28 New End
Square, London
NW3 1LS

Geoff Watts
science editor, *BMJ*

Correspondence to:
geoff@scileg.
freeserve.co.uk

As the Duchess of Windsor once remarked, one can never be too thin or too rich—wisely, she did not add too clean. The hygiene hypothesis aims to explain why some people have allergies and some do not, and why the prevalence of allergic disorders has been increasing over the past century. It suggests that the modern obsession with cleanliness may be counter-productive; in childhood, at least, it may encourage the development of allergic disorders. Benn and colleagues have attempted to untangle the link between infections in infancy and atopic disease.¹

Formalised in the late 1980s, the hypothesis seemed to fly in the face of common sense. After all, hadn't the hygiene movement ushered in a dramatic reduction in infectious disease? This initial scepticism was compounded by the suspicion that infections might actually provoke allergy. Despite these doubts, the evidence kept piling up.² In epidemiological studies, factors such as large household size, a poor standard of living, not using antibiotics, and early exposure to farm animals all emerged as protective.

The original attempt to explain the immunological basis of the hygiene hypothesis invoked a loss of balance between two sets of the body's immune cells—the Th1 and Th2 lymphocytes.³ A reduced exposure to childhood infection, it was argued, meant a low level of Th1 activity in the body, so bringing about a relative excess of Th2 activity—and a consequent tendency to allergy. This neat explanation began to fall apart when it was appreciated that Th1 mediated conditions, such as type 1 diabetes and inflammatory bowel disease, were also on the increase, and in the same regions of the world.

Clearly, any reinterpretation of the hygiene hypothesis had to take account of this parallel increase in conditions associated with Th1 and Th2. A recent attempt manages to do just this.⁴ It suggests that the effect of hygiene is to diminish the body's production of a third group of players in the immune drama—the regulatory T cells. It is a shortage of these that results in the emer-

gence of allergy. The driving force in this case, goes the argument, is our freedom not from pathogenic microbes but from a group of organisms including mycobacteria, lactobacilli, and helminth worms. These "old friends," as Rook describes them,⁴ have lived with us for countless generations. In adapting to their more or less benign presence, the immune system has learned not to over-react. It exercises this self suppression by generating regulatory T cells.

In the absence of its "old friends" the system produces fewer of these cells. The consequence is a state of relatively uncontrolled effector T cell exuberance and, depending presumably on genetic factors of some kind, a predisposition to allergy or to more serious autoimmune conditions such as inflammatory bowel disease and diabetes. The findings reported by Benn and colleagues,¹ which refer only to "clinically apparent" infections, are entirely consistent with this view of the "old friends" being the organisms responsible for the protection against allergy.

Does this version of the hygiene hypothesis suggest a method of vaccination against allergy based on stimulating the body's production of regulatory T cells? It does, and preliminary experiments are already under way.⁵ Immunology's love affair with dirt is blossoming and may yet bear fruit.

Competing interests: None declared.

1 Benn CS, Melbye M, Wohlfahrt J, Björkstén B, Aaby P. Cohort study of the sibling effect, infectious diseases, and risk of atopic dermatitis during the first 18 months of life. *BMJ* 2004 doi:10.1136/bmj.38069.512245.FE.

2 Bach JF. The effect of infections on susceptibility to autoimmune and allergic diseases. *N Engl J Med* 2002;347:911-20.

3 Strachan DP. Family size, infection and atopy: the first decade of the "hygiene hypothesis". *Thorax* 2000;55(suppl 1):S2-10.

4 Rook GAW, Adams V, Hunt J, Palmer R, Martinelli R, Brunet LR. Mycobacteria and other environmental organisms as immunomodulators for immunoregulatory disorders. *Springer Semin Immunopathol* 2004;25: 237-55.

5 Arkwright PD, David TJ. Intradermal administration of a killed Mycobacterium vaccae suspension (SRL 172) is associated with improvement in atopic dermatitis in children with moderate-to-severe disease. *J Allergy Clin Immunol* 2001;107:531-4.

doi 10.1136/bmj.38075.565822.55