Published in final edited form as:

Exerc Sport Sci Rev. 2011 October ; 39(4): 161–166. doi:10.1097/JES.0b013e3182297439.

The Mechanistic Connection Between the Trunk, Knee, and Anterior Cruciate Ligament Injury

Timothy E. Hewett^{1,2,3,4} and Gregory D. Myer^{1,2,3,5}

¹The Ohio State University Sports Health and Performance Institute, Departments of Physiology and Cell Biology, Orthopaedic Surgery, Family Medicine, and Biomedical Engineering, The Ohio State University, Columbus

²Cincinnati Children's Hospital Medical Center, Division of Sports Medicine, Cincinnati, Ohio

³The University of Cincinnati College of Medicine, Departments of Pediatrics and Orthopaedic Surgery, Cincinnati, OH

⁴Department of Rehabilitation Sciences, College of Allied Health Sciences, The University of Kentucky, Lexington, KY

⁵Rocky Mountain University of Health Professions, Departments of Athletic Training, Sports Orthopaedics, and Pediatric Science, Provo, UT

Abstract

Neuromuscular control of the trunk and knee predicts anterior cruciate ligament injury risk with high sensitivity and specificity. These predictors are linked, as lateral trunk positioning creates high knee abduction torque (load). The hypotheses explored are that lateral trunk motion increases load and that neuromuscular training that increases trunk control will decrease load.

Keywords

knee injury; high risk biomechanics; ACL injury prevention; neuromuscular control; female sports injuries; identification of knee injury risk factors

INTRODUCTION: THE MECHANISTIC CONNECTION BETWEEN THE TRUNK AND THE KNEE

The goal of this perspective for progress is to lead to advances in the understanding of the mechanisms and prevention of anterior cruciate ligament (ACL) injuries in female athletes, who are at a 2- to 10-fold increased risk of ACL injury than male athletes. The overall scientific objectives are to determine the mechanisms by which female athletes become more susceptible to ACL injury and to optimize the effectiveness of interventions designed to prevent ACL injuries. Specifically, this perspective is directed toward understanding and

Copyright © 2011 by the American College of Sports Medicine.

Address for correspondence: Timothy E. Hewett, Ph.D., The Sports Medicine Biodynamics Center and Human Performance Laboratory, 3333 Burnet Ave., Cincinnati, Ohio, 45206 (tim.hewett@cchmc.org).

exploring the following hypotheses: 1) Does decreased neuromuscular control of the trunk (*i.e.*, active stabilization of lateral trunk motion) underlie increased loading of the knee joint in female athletes? 2) Does this increased loading occur through both biomechanical (GRFv driven) and neuromuscular (hip adductor driven) mechanisms? 3) Does neuromuscular training (*i.e.*, training focused to increase active stabilization and control during dynamic movements) of the trunk decrease abduction load at the knee?

Analysis of neuromuscular control of the trunk may be a useful new perspective for the determination of the mechanisms underlying increased risk for ACL injury and for the optimization of risk screening and intervention programs. The preponderance of prior research has focused on the resultant symptoms, knee motions, and loads, as risk factors, as opposed to the potential underlying root of the problem, neuromuscular control of the trunk. The causes of risky knee loads and motions are not likely in the knee, rather it is probable that high-risk knee loads are the resultant symptoms to reduced control of body posture and trunk accelerations. A narrow perspective, which focuses on knee motions and loads may not only lead to the misinterpretation of underlying injury mechanism, but it may prohibit the application of optimal intervention, as the determination of knee load levels cannot be incorporated readily into clinic or field-based screening programs and can be difficult to communicate to athletes. The valgus moment at the knee is not visible, and the relevant differences in the valgus angle are in the range of five to six degrees, which hardly is quantifiable by the naked eye. However, relative trunk control, or increased trunk motion, is not only a likely major underlying factor; it is more visible and can be explained readily to athletes and athletic trainers (Figs. 1 and 2). Hence, examination of neuromuscular control of the trunk results in measurable variables that likely are closer to the higher level control mechanisms.

A series of studies have developed the logical basis behind the hypotheses delineated above. 1) Hewett *et al.* (27) demonstrated that the mechanism of noncontact ACL injuries in female athletes, as observed on video, includes lateral trunk motion with the body shifted over one leg, which was associated with high knee abduction or medial knee collapse (30,33,48). Lateral perturbation of the trunk is another common component of the mechanism (5,32). Hence, increased lateral sway of the trunk may underlie medial collapse of the knee joint in female athletes. 2) Zazulak *et al.* (56) demonstrated that trunk displacement and coronal plane knee load both predict ACL injury risk in female athletes with high sensitivity and specificity and accuracy (91%) (20). Therefore, increased knee loading occurs through neuromuscular and both biomechanical mechanisms related to increased trunk inertia and motion. 3) Finally, both initial (25) and confirmatory investigation (45,50) have demonstrated that neuromuscular control of the trunk and lower extremity can be improved with neuromuscular training. Neuromuscular control of the trunk has the potential to decrease abduction load at the knee and to decrease ACL injury risk (17,40–42,45). With these concepts in mind, we will focus the remainder of the “Perspectives for Progress” review on the scientific justification (*e.g.*, trunk and knee are coupled mechanically and dynamically via the ground reaction force (GRF) lever arm) and the evidence of a correlation from movement analysis studies (Fig. 2).

Prophylactic intervention for ACL injury could prevent a significant percentage of the 100,000 to 250,000 injuries that occur each year in the United States (17,53). The President's council on physical fitness recommends that women, both young and old, remain active to maintain optimal health. Reduction of female injury rates from five times to equal male athletes' potentially would allow female athletes annually to continue the health benefits of sports participation and avoid the long-term complications of osteoarthritis, which occurs with a 10 to 100-fold greater incidence in ACL-injured than in uninjured athletes (34).

ACL Injury Rate Differences and Long-Term Sequelae in Female Athletes

Adolescent and mature female athletes who participate in pivoting and jumping sports experience ACL injuries at a 2- to 10-fold greater rate than male athletes participating in the same high-risk cutting and landing sports (1). The combination of this greater susceptibility and a 10-fold increase in the female sports population since the inception (June 23, 1972) of Title IX has resulted in a dramatic increase in the number of ACL injuries in female athletes. In the United States, 100,000–250,000 ACL injuries occur each year (17). The costs may exceed \$650 million annually in female varsity athletics alone (19). Neuromuscular control deficits at the hip and trunk may follow maturation (Fig. 1) and may contribute to decreased active neuromuscular control of the lower extremity that may lead to increased knee abduction loads and strain on the knee ligaments (3,18,20,37). The costs associated with ACL injury do not end with the ACL reconstruction and rehabilitation. There is a strong association between ACL injury and development of posttraumatic knee osteoarthritis at a relatively young age, which also occurs with much greater incidence in female athletes than male athletes (34). It is estimated that between 50% and 100% of women with an ACL injury will show significant pain, functional limitations, and radiographic signs of knee osteoarthritis within 12–20 yr of the index injury (34). Prediction of who is at risk and how and when their risk increases has potential to prevent unnecessary injury and to significantly improve outcomes after injury.

Coronal Plane Components of the ACL Injury Mechanism in Female Athletes

Trunk displacement and coronal plane knee load both predict ACL injury risk in female athletes. Knee abduction loads and neuromuscular control of the trunk both predict ACL injury risk with high sensitivity and specificity. Knee abduction load predicted ACL injury risk with 78% sensitivity and 73% specificity (20). A logistic regression model that incorporated lateral trunk motion predicted ACL injury risk in female athletes with 83% sensitivity and 76% specificity but did not predict either general knee or specifically ACL injury risk in male athletes (56). The mechanism of ACL injury may differ in female and male athletes, especially with respect to the dynamic positioning of the knee, as female athletes demonstrate greater valgus collapse of the lower extremity primarily in the coronal plane (32). Most ACL injuries in female athletes occur by noncontact mechanisms during landing and lateral pivoting (48). The mechanism of noncontact ACL injuries as observed on video has several common components in female athletes: high knee abduction, lateral trunk motion with the body shifted over one leg and the plantar surface of the foot fixed flat on the playing surface, displaced away from the trunk and low knee flexion (30,32,48).

Unanticipated perturbation of the trunk is another common component of the mechanism (5,33).

Trunk motion can influence knee abduction load through mechanical and neuromuscular mechanisms (Figs. 1 and 2) (22). For example, if the trunk moves laterally (relative to the stance limb), the GRF vector (GRFv) may move laterally and have a greater lever arm relative to the knee joint center. This directly will increase the potential for knee abduction loading in combination with increased inertial acceleration of the trunk or thigh segments during dynamic movement. The resultant knee load simultaneously increases in magnitude of GRFv because of unicompartamental (*i.e.*, lateral compartment loads from knee abduction stance) knee joint loading (22,43). In response to the lateral trunk motion and increased knee load, it is essential to counteract with increasing reactive hip adductor torque to maintain upright stance and dissipate lower extremity forces. Increased hip adductor torque can increase knee abduction moments, which are predictors of peak GRF (25) and an important determinant of joint load. Knee abduction torque places knee ligaments in the high slope (load) segment of their force-length curve and elicits knee pain in female athletes (52). Video evidence of ACL injuries shows that the female trunk usually moves lateral toward the ACL injured limb as the knee abducts (Fig. 2) (26). Ultimately, when the trunk leans to one side, the hip adductors are activated to right the pelvis and trunk to maintain stance; however, this strategy of movement, during dynamic high-load scenarios seems to initiate a viscous mechanical cycle, which underlies the inciting mechanism of knee (ACL and patellofemoral pain disorder) injury in female sports.

Mechanical Mechanisms Related to Lateral Trunk Motion and Knee Load in Female Athletes

Trunk position and knee external abduction moment (load) may be linked mechanically, as lateral positioning of the trunk can create abduction loads at the knee (Figs. 2 and 3) (54). In the coronal plane, applying static equilibrium mechanics and neglecting the inertia of the body segments between hip and ground, if the GRF passes lateral to the center of the head of the femur, an external hip abduction torque results (54). This external moment is balanced by a hip adductor generated torque of equal magnitude. There is a strong association ($r = 0.62\text{--}0.77$) between lateral trunk lean and both hip adductor torque and coronal plane knee valgus stress during pathological gait (13,49). Interestingly, even alteration of arm position relative to the centerline of the body can increase the external knee abduction load by 29%–60% (13,49). At the low knee flexion angles that are present during ACL injuries, the ACL, rather than the MCL, can be the primary restraint to knee abduction loads (55). Knee load and ACL injury may be outcomes that result from an unstable, collapsing lower extremity column under axial load (GRF) passing through the lateral knee compartment.

Neuromuscular Trunk Control Mechanisms may Increase Knee Load in Female Athletes

Trunk stability is related to the ability of the hip to control the trunk in response to forces generated from distal body segments as well as from unexpected perturbations (23,36). Deficits in neuromuscular control of the trunk during cutting and landing may lead to uncontrolled lateral trunk motion that may increase knee abduction motion and torque through mechanical (lateral GRF motion) and neuromuscular (increased hip adductor

torque) mechanisms (18,20). Insufficient neuromuscular control of the trunk may increase strain on the ACL and lead to injury via either one or both of these mechanisms (3,18,22,27). Neuromuscular control of the hip is required to control coronal plane trunk and pelvis motion. An external hip abduction moment created by the GRF moving lateral to the center of the femoral head is counterbalanced internally by hip adductor torque (22).

The Underlying Mechanisms V Growth in the Absence of Sufficient Power?

Following the growth spurt that occurs during maturation, female subjects have very different neuromuscular control profiles than male subjects. Recent published studies have observed that male subjects experience a significant increase in neuromuscular strength and coordination as skeletal growth and maturation progresses, a so-called neuromuscular spurt not often observed in female subjects (21,41). As bone length and body mass increases, male subjects also demonstrate greater neuromuscular control of the knee joint than female subjects, allowing them to better absorb loads (Fig. 1). In lay terms, growth results in larger machines in both sexes, but as male subjects mature, they adapt with disproportionately more muscle “horsepower” to match the control demands of their larger machine. Female subjects do not show similar adaptations. In the absence of neuromuscular adaptations in strength and muscle recruitment, female knees are exposed to greater GRFs and high external knee abduction moments (load), particularly in landing, pivoting, and deceleration sports. Collectively, these studies have driven the central hypothesis that increased bone length and body mass, in the absence of matching adaptations in neuromuscular strength and coordinated muscle recruitment, expose the female knee to greater load and likelihood of ACL injury (Fig. 2) (20–22).

Female subjects activate the hip musculature differently than male subjects in response to sudden loading (12). Women adduct the hip more than men during both low- and high-intensity activities. They begin descent in a more abducted knee position and remain in a more abducted alignment relative to men throughout a squat motion or during landing (16). Female subjects also demonstrate more hip adduction than male subjects during cutting (39). Increased hip adduction during dynamic motion and decreased hip muscle abductor strength and recruitment can increase knee load and injury risk (16,20). Ipsilateral trunk lean is a sign of weak hip abductors as it moves the center of mass closer to the stance limb to reduce demand on the weak abductors (51). During single leg landing and cutting, the entire body mass must be balanced over one lower extremity. Because the trunk comprises greater than half of the body’s mass, lateral trunk motion increases GRF and load (51). The internal response is an equal and opposite increase in the counterbalancing hip adductor torque (54). The resulting increase in relative hip adductor-abductor torque ratio then likely increases knee load.

Based on the current literature, it is hypothesized that increased lateral trunk motion, change in direction of the GRF velocity ($\dot{GRF}_v$) and hip adductor torque increase knee load of female athletes (22). Deficits in neuromuscular control of the trunk may contribute to lower extremity joint instability and injury (24). Landing and cutting require high levels of neuromuscular control to maintain stability and performance (35). Dynamic stability of the knee is dependent on accurate sensory input and appropriate motor responses to rapid

changes in body position during cutting and landing (20,24). Neuromuscular control of the hip, trunk, and knee is based on feedback control. The position and load of each segment is used to modify the descending movement commands (11). Impaired control of the hip and trunk can increase lower extremity injury. For example, abdominal muscle fatigue contributes to hamstrings injuries (7). Subjects with ankle sprains had a delay in onset of gluteus maximus and medius activation (2). Female, but not male, athletes who experienced ankle injury had greater body sway before injury than uninjured controls (4).

Effects of Neuromuscular Training in Female Athletes

Movement biomechanics and lower extremity strength can be altered in female subjects with neuromuscular training (25). Neuromuscular power (*i.e.*, rate of muscular recruitment and force generation, as evidenced by vertical jump height) can increase within 6 wk of training and may result in decreases in peak impact forces and knee abduction torques (25).

Observed changes in female subjects may be greater than those in male subjects as their baseline neuromuscular performance levels are lower (31). If neuromuscular training can decrease ACL injury risk, it is likely that the mechanisms underlying increased risk are neuromuscular in nature (19). ACL injury risk may be reduced in trained female athletes during landing and cutting (15). Elite female athletes show reductions in ACL injuries with neuromuscular training (47). These prospective studies indicate that neuromuscular training has the potential to decrease ACL injury rates in female subjects.

Neuromuscular control of the trunk and lower extremity can be improved with neuromuscular training (20,25,40). Neuromuscular training may increase coronal plane trunk and hip control in female subjects (25,31). For example, during a drop-jump, a two-footed plyometric activity, post training results showed that lower extremity valgus was reduced at the hip. Conversely, during a single-leg landing task, the most significant modifications may occur at the knee (44). Therefore, the effects of training in the coronal plane are likely to be movement task specific. These results indicate that resultant load is controlled primarily by hip torque during a two-foot landing but more by GRFv during single leg tasks. Increased coronal plane control at both the hip and trunk may be necessary to reduce ACL injury risk (6,38). Lower extremity coronal joint motions and torques linked to increased ACL injury risk often are correlated, indicating that control of knee load may require synergistic and antagonistic contribution from the trunk, hip, and knee (20). Perturbation-enhanced training may increase trunk control and decrease knee abduction load in female subjects (29,45).

Female subjects tend to utilize greater coronal plane control rather than a sagittal plane control strategy for the lower extremity (8,10). They tend to utilize a “hip strategy” for single leg control and balance during landing and cutting (9,16). For example, coronal plane excursions are greater and more rapid at the hip and knee during walking in female subjects (29). The knee functions optimally as a sagittal plane hinge, not a coronal plane hinge or ball and socket joint, as the large muscles of the lower extremity that control coronal plane trunk, hip, and knee motion or torque absorb and dissipate force most effectively and efficiently in the sagittal, rather than coronal plane. Women also demonstrate less active joint stiffness than men (28).

Future Directions to Improve Trunk Control Reduce Dangerous Knee Loads and Prevent ACL Injury

So what are the next steps in the decipherment of this hidden, chaotic, and destructive process? First, the newly developed interventions to be tested must target all planes of motion. Future study designs must be rigorous and must employ randomized, controlled trials, with placebo or sham treatments in place. Sham treatments must be in place to control for bias, confounding, unnecessary error, and potentially erroneous misrepresentation of treatment effects. Moreover, result interpretation may be the most common and greatest offender and must be avoided in investigations focused on the effects of intervention on risk factors to female athletes.

SUMMARY AND CONCLUSIONS

This Perspective for Progress has provided a strong, evidence-based rationale for the development of effective “core-based” interventions to decrease ACL injury risk in high-risk female athletes. Increased hip abductor muscle recruitment and strength likely has a direct effect on the knee abduction loading of the ACL during cutting and landing. Although ACL injuries likely occur too quickly (<100 milliseconds) for reflexive muscular activation (>100 milliseconds), athletes can adopt preparatory muscle recruitment and movement patterns that reduce the probability of injuries caused by unexpected perturbations (32,52). The current evidence indicate that decreased neuromuscular control of the trunk leads to increased joint load (knee abduction moment) via two mechanisms (lateral motion of the GRF and reactive hip adduction torque) and results in increased ACL injury risk in female athletes (Figs. 1–3). One would theorize that knee load (the measurable symptom that increases ACL injury risk) will be decreased by neuromuscular training designed to increase coronal plane trunk stability (22). Further evidence supports the contention that increased load will be decreased by neuromuscular training that increases coronal plane control of trunk via decreased

GRFv and hip adductor strength and recruitment (22,40,41,45). Knowledge of these mechanisms also may be used to categorize female athletes into high-, moderate-, and low-risk groups to allow for targeted training to underlying mechanics of load, which may maximize the effectiveness of neuromuscular training programs aimed to reduce injury risk in female athletes.

Acknowledgments

This work was supported in part by NIH/NIAMS R01 AR049735, R01 AR05563 and R01 AR056259 (T.E.H.). We wish to dedicate this Perspective for Progress to our eight sisters and to all the millions of sisters who have been given vast opportunities for the healthy physical activity of sports participation since the passage of Title IX of the Education Assistance Act over the last 4 decades.

References

1. Arendt E, Dick R. Knee injury patterns among men and women in collegiate basketball and soccer. NCAA data and review of literature. *Am. J. Sports Med.* 1995; 23:694–701. [PubMed: 8600737]
2. Beckman SM, Buchanan TS. Ankle inversion injury and hypermobility: effect on hip and ankle muscle electromyography onset latency. *Arch. Phys. Med. Rehabil.* 1995; 76:1138–1143. [PubMed: 8540791]

3. Bendjaballah MZ, Shirazi-Adl A, Zukor DJ. Finite element analysis of human knee joint in varus-valgus. *Clin. Biomech.* 1997; 12:139–148.
4. Beynnon BD, Renstrom PA, Haugh L, Uh BS, Barker H. A prospective, randomized clinical investigation of the treatment of first-time ankle sprains. *Am. J. Sports Med.* 2006; 34:1401–1412. [PubMed: 16801691]
5. Boden BP, Torg JS, Knowles SB, Hewett TE. Video analysis of anterior cruciate ligament injury: abnormalities in hip and ankle kinematics. *Am. J. Sports Med.* 2009; 37:252–259. [PubMed: 19182110]
6. Chaudhari AM, Andriacchi TP. The mechanical consequences of dynamic frontal limb alignment for non-contact ACL injury. *J. Biomech.* 2006; 39:330–338. [PubMed: 16321635]
7. Devlin L. Recurrent posterior thigh symptoms detrimental to performance in rugby union: predisposing factors. *Sports Med.* 2000; 29:273–287. [PubMed: 10783902]
8. Ford KR, Myer GD, Hewett TE. Valgus knee motion during landing in high school female and male basketball players. *Med. Sci. Sports Exerc.* 2003; 35:1745–1750. [PubMed: 14523314]
9. Ford KR, Myer GD, Smith RL, Vianello RM, Seiwert SL, Hewett TE. A comparison of dynamic coronal plane excursion between matched male and female athletes when performing single leg landings. *Clin. Biomech.* 2006; 21:33–40.
10. Ford KR, Myer GD, Toms HE, Hewett TE. Gender differences in the kinematics of unanticipated cutting in young athletes. *Med. Sci. Sports.* 2005; 37:124–129.
11. Gandevia SC, McCloskey DI, Burke D. Kinaesthetic signals and muscle contraction. *Trends Neurosci.* 1992; 15:62–65. [PubMed: 1374964]
12. Granata KP, Orishimo KF, Sanford AH. Trunk muscle coactivation in preparation for sudden load. *J. Electromyogr. Kinesiol.* 2001; 11:247–254. [PubMed: 11532595]
13. Gupta RT, Vankoski S, Novak RA, Dias LS. Trunk kinematics and the influence on valgus knee stress in persons with high sacral level myelomeningocele. *J. Pediatr. Orthop.* 2005; 25:89–94. [PubMed: 15614067]
14. Hewett, TE.; Biro, FM.; McLean, SG.; Van den Bogert, AJ. Book Identifying Female Athletes at High Risk for ACL Injury. National Institutes of Health; 2003. Identifying female athletes at high risk for ACL injury.
15. Hewett TE, Ford KR, Myer GD. Anterior cruciate ligament injuries in female athletes: part 2, a meta-analysis of neuromuscular interventions aimed at injury prevention. *Am. J. Sports Med.* 2006; 34:490–498. [PubMed: 16382007]
16. Hewett TE, Ford KR, Myer GD, Wanstrath K, Scheper M. Gender differences in hip adduction motion and torque during a single leg agility maneuver. *J. Orthop. Res.* 2006; 24:416–421. [PubMed: 16479597]
17. Hewett TE, Lindenfeld TN, Riccobene JV, Noyes FR. The effect of neuromuscular training on the incidence of knee injury in female athletes. A prospective study. *Am. J. Sports Med.* 1999; 27:699–706. [PubMed: 10569353]
18. Hewett TE, Myer GD, Ford KR. Reducing knee and anterior cruciate ligament injuries among female athletes: a systematic review of neuromuscular training interventions. *J. Knee Surg.* 2005; 18:82–88. [PubMed: 15742602]
19. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: part 1, mechanisms and risk factors. *Am. J. Sports Med.* 2006; 34:299–311. [PubMed: 16423913]
20. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am. J. Sports Med.* 2005; 33:492–501. [PubMed: 15722287]
21. Hewett TE, Myer GD, Ford KR, Slaughterbeck JR. Preparticipation physical exam using a box drop vertical jump test in young athletes: the effects of puberty and sex. *Clin. J. Sport Med.* 2006; 16:298–304. [PubMed: 16858212]
22. Hewett, TE.; Myer, GD.; Ford, KR.; Powers, S. Book Neuromuscular Intervention Targeted to Mechanisms of ACL Load in Female Athletes. National Institutes of Health; 2008. Neuromuscular intervention targeted to mechanisms of ACL load in female athletes.

23. Hewett TE, Myer GD, Zazulak BT. Hamstrings to quadriceps peak torque ratios diverge between sexes with increasing isokinetic angular velocity. *J. Sci. Med. Sport.* 2008; 11:452–459. [PubMed: 17875402]
24. Hewett TE, Paterno MV, Myer GD. Strategies for enhancing proprioception and neuromuscular control of the knee. *Clin. Orthop.* 2002; 402:76–94. [PubMed: 12218474]
25. Hewett TE, Stroupe AL, Nance TA, Noyes FR. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *Am. J. Sports Med.* 1996; 24:765–773. [PubMed: 8947398]
26. Hewett TETJ, Boden BP. Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in female athletes: lateral trunk and knee abduction motion are combined components of the injury mechanism. *Br. J. Sports Med.* 2009; 43:417–422. [PubMed: 19372088]
27. Hewett TE, Torg JS, Boden BP. Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in female athletes: lateral trunk and knee abduction motion are combined components of the injury mechanism. *Br. J. Sports Med.* 2009; 43:417–422. [PubMed: 19372088]
28. Hurd WJ, Chmielewski TL, Axe MJ, Davis IM, Snyder-Mackler L. Differences in normal and perturbed walking kinematics between male and female athletes. *Clin. Biomech.* 2004; 19:465–472.
29. Hurd WJ, Chmielewski TL, Snyder-Mackler L. Perturbation-enhanced neuromuscular training alters muscle activity in female athletes. *Knee Surg. Sports Traumatol. Arthrosc.* 2006; 14:60–69. [PubMed: 15937713]
30. Ireland ML. The female ACL: why is it more prone to injury? *Orthop. Clin. North Am.* 2002; 33:637–651. [PubMed: 12528906]
31. Kraemer WJ, Mazzetti SA, Nindl BC, et al. Effect of resistance training on women's strength/power and occupational performances. *Med. Sci. Sports Exerc.* 2001; 33:1011–1025. [PubMed: 11404668]
32. Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am. J. Sports Med.* 2007; 35(3):359–367. [PubMed: 17092928]
33. Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am. J. Sports Med.* 2007; 35:359–367. [PubMed: 17092928]
34. Lohmander LS, Ostengren A, Englund M, Roos H. High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis. Rheum.* 2004; 50:3145–3152. [PubMed: 15476248]
35. Loram ID, Lakie M. Human balancing of an inverted pendulum: position control by small, ballistic-like, throw and catch movements. *J. Physiol. (Lond.).* 2002; 540:1111–1124. [PubMed: 11986396]
36. MacKinnon CD, Winter DA. Control of whole body balance in the frontal plane during human walking. *J. Biomech.* 1993; 26:633–644. [PubMed: 8514809]
37. Markolf KL, Burchfield DM, Shapiro MM, Shepard MF, Finerman GA, Slaughterbeck JL. Combined knee loading states that generate high anterior cruciate ligament forces. *J. Orthop. Res.* 1995; 13:930–935. [PubMed: 8544031]
38. McLean SG, Huang X, Su A, van den Bogert AJ. Sagittal plane biomechanics cannot injure the ACL during sidestep cutting. *Clin. Biomech.* 2004; 19:828–838.
39. McLean SG, Su A, van den Bogert AJ. Development and validation of a 3-D model to predict knee joint loading during dynamic movement. *J. Biol. Chem.* 2003; 125:864–874.
40. Myer GD, Brent JL, Ford KR, Hewett TE. A pilot study to determine the effect of trunk and hip focused neuromuscular training on hip and knee isokinetic strength. *Br. J. Sports Med.* 2008; 42:614–619. [PubMed: 18308886]
41. Myer GD, Chu DA, Brent JL, Hewett TE. Trunk and hip control neuromuscular training for the prevention of knee joint injury. *Clin. Sports Med.* 2008; 27:425–448. ix. [PubMed: 18503876]

42. Myer GD, Ford KR, Brent JL, Hewett TE. Differential neuromuscular training effects on ACL injury risk factors in “high-risk” versus “low-risk” athletes. *BMC Musculoskelet. Disord.* 2007; 8:1–7. [PubMed: 17204151]
43. Myer GD, Ford KR, Hewett TE. The effects of gender on quadriceps muscle activation strategies during a maneuver that mimics a high ACL injury risk position. *J. Electromyogr. Kinesiol.* 2005; 15:181–189. [PubMed: 15664147]
44. Myer GD, Ford KR, McLean SG, Hewett TE. The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics. *Am. J. Sports Med.* 2006; 34:490–498. [PubMed: 16382007]
45. Myer GD, Ford KR, Palumbo JP, Hewett TE. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *J. Strength Cond. Res.* 2005; 19:51–60. [PubMed: 15705045]
46. Myer GD, Ford KR, Paterno MV, Nick TG, Hewett TE. The effects of generalized joint laxity on risk of anterior cruciate ligament injury in young female athletes. *Am. J. Sports Med.* 2008; 36(6): 1073–1080. [PubMed: 18326833]
47. Myklebust G, Engebretsen L, Braekken IH, Skjølberg A, Olsen OE, Bahr R. Prevention of anterior cruciate ligament injuries in female team handball players: a prospective intervention study over three seasons. *Clin. J. Sport Med.* 2003; 13:71–78. [PubMed: 12629423]
48. Olsen OE, Myklebust G, Engebretsen L, Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am. J. Sports Med.* 2004; 32:1002–1012. [PubMed: 15150050]
49. Ounpuu S, Thomson JD, Davis RB, DeLuca PA. An examination of the knee function during gait in children with myelomeningocele. *J. Pediatr. Orthop.* 2000; 20:629–635. [PubMed: 11008743]
50. Paterno MV, Myer GD, Ford KR, Hewett TE. Neuromuscular training improves single-limb stability in young female athletes. *J. Orthop. Sports Phys. Ther.* 2004; 34:305–317. [PubMed: 15233392]
51. Perry, J. *Gait Analysis: Normal and Pathological Function.* Thorofare (NJ): SLACK Inc.; 1992.
52. Pope MH, Johnson RJ, Brown DW. The role of the musculature in injuries to the medial collateral ligament. *J. Bone Joint Surg.* 1979; 61A:398–402. [PubMed: 429411]
53. Toth AP, Cordasco FA. Anterior cruciate ligament injuries in the female athlete. *J. Gend. Specif. Med.* 2001; 4:25–34. [PubMed: 11727468]
54. Winter, DA. *Biomechanics and Motor Control of Human Movement.* New York (NY): John Wiley & Sons, Inc.; 2005.
55. Wu G, Siegler S, Allard P, et al. ISB recommendation on definitions of joint coordinate system of various joints for the reporting of human joint motion — part I: ankle, hip, spine. *International Society of Biomechanics. J. Biomech.* 2002; 35:543–548. [PubMed: 11934426]
56. Zazulak BT, Hewett TE, Reeves NP, Goldberg B, Cholewicki J. Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. *Am. J. Sports Med.* 2007; 35:1123–1130. [PubMed: 17468378]

Figure 1.

Theory linking growth, NM adaptation, trunk NMC, "dynamic valgus," and knee joint load to anterior cruciate ligament (ACL) injury risk (14,46).

Figure 2. Mechanical model linking lateral trunk motion to change in direction of the ground reaction force velocity (GRFv), hip adductor torque, and knee load during cutting and landing ACL injuries.

Figure 3. Conceptual model of lateral trunk motion leading to increased Δ GRFv and hip adductor torque and knee load. This will result in clustered load subgroups of female athletes.