Published in final edited form as:

Angew Chem Int Ed Engl. 2014 September 22; 53(39): 10443–10447. doi:10.1002/anie.201404917.

Total Synthesis of ¹²-Prostaglandin J₃, a Highly Potent and Selective Antileukemic Agent**

K. C. Nicolaou,

Department of Chemistry, BioScience Research Collaborative, Rice University, 6100 Main Street, Houston, TX 77005 (USA), kcn@rice.edu

Philipp Heretsch,

Department of Chemistry, BioScience Research Collaborative, Rice University, 6100 Main Street, Houston, TX 77005 (USA)

Abdelatif ElMarrouni,

Department of Chemistry, BioScience Research Collaborative, Rice University, 6100 Main Street, Houston, TX 77005 (USA)

Christopher R. H. Hale,

Department of Chemistry, BioScience Research Collaborative, Rice University, 6100 Main Street, Houston, TX 77005 (USA)

Kiran K. Pulukuri,

Department of Chemistry, BioScience Research Collaborative, Rice University, 6100 Main Street, Houston, TX 77005 (USA)

Avinash K. Kudva,

Department of Veterinary and Biomedical Sciences, The Pennsylvania State University, University Park, PA 16802 (USA)

Vivek Narayan, and

Department of Veterinary and Biomedical Sciences, The Pennsylvania State University, University Park, PA 16802 (USA)

K. Sandeep Prabhu

Department of Veterinary and Biomedical Sciences, The Pennsylvania State University, University Park, PA 16802 (USA)

Abstract

**Financial support was provided by The Cancer Prevention Research Institute of Texas, The Welch Foundation, Rice University, and the National Institutes of Health (to K.S.P.; R01 CA175576). Fellowships to A.E. (Fundación Alfonso Martín Escudero, Postdoctoral Fellowship) and C.R.H.H. (NSF Graduate Research Fellowship) are gratefully acknowledged. We thank Drs. Lawrence B. Alemany and Quinn Kleerekoper for NMR-spectroscopic assistance and Dr. Christopher L. Pennington for mass-spectrometric assistance. We also thank Professor Robert F. Paulson for valuable discussions.

© Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Correspondence to: K. C. Nicolaou.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201404917>.

A catalytic asymmetric total synthesis of the potent and selective antileukemic 12 -prostaglandin J_3 (12 -PGJ₃) is described. The convergent synthesis proceeded through intermediates **2** and **3**, constructed enantioselectively from readily available starting materials and coupled through an aldol reaction followed by dehydration to afford stereoselectively the cyclopentenone alkylidene structural motif of the molecule.

Keywords

asymmetric catalysis; C–H activation; leukemia; natural products; total synthesis

Recent reports described 12 -prostaglandin J_3 (12 -PGJ₃, **1**, Figure 1) as a potent and selective ablator of leukemia stem cells in vitro and in vivo.^[1,2] In view of the increasing interest in cancer stem cells as drivers for growth, perpetuation, recurrence and drug resistance in various types of cancer,^[3] 12 -PGJ₃ may serve as an important tool to decipher cancer biology and a lead compound for drug discovery and development. Naturally formed from ω -3 eicosapentaenoic acid [(5Z,8Z,11Z,14Z,17Z)-5,8,11,14,17-icosapentaenoic acid, EPA],^[1,2,4] this secondary metabolite was isolated in minute amounts and characterized by UV spectroscopic and mass spectrometric methods.^[1,2] The impressive in vitro potency and selectivity of 12 -PGJ₃ against chronic myelogenous leukemia (CML) stem cells (IC₅₀ = 12 nM) and its ability to effectively cure this form of leukemia in a mouse model,^[1,2] coupled with its scarcity, prompted us to undertake its total synthesis with the intention of rendering it readily available for thorough biological investigations and full structural characterization.

The molecular structure of 12 -PGJ₃ (**1**) is characterized by its 2-alkylidene cyclopentenone structural motif, and while stable by itself it could, in principle, undergo conjugate nucleophilic attack by thiols, particularly at the endocyclic double bond^[4,5] (i.e. C9, prostaglandin numbering). This property may or may not be involved in the mechanism of action of this intriguing molecule. The *E*-configuration of the 12 -olefinic bond was presumed on the basis of thermodynamic stability. Figure 1 summarizes, in retrosynthetic format, the devised strategy for a catalytic asymmetric total synthesis of 12 -PGJ₃ (**1**). Thus, disconnection of the 12 -olefinic bond through an aldol/dehydration process led to advanced intermediates cyclopentenone **2** and aldehyde **3**. These intermediates were then traced back to building blocks **4** and **5** (for **2**) and **6** and **7** (for **3**) through a Wittig olefination and an asymmetric Mukaiyama aldol reaction, respectively. The latter was intended to secure the configuration of the C15 asymmetric center, while a palladium-catalyzed asymmetric Tsuji–Troost alkylation reaction was reserved to install, using as precursor cyclopentene ester fragment **4**, the other remote stereocenter (C8) in its proper configuration. The required oxygenation at C11 was to be achieved through a C–H activation/functionalization process at the stage of building block **4**. While this strategy includes similar features to that employed by the Kobayashi group^[6] to synthesize a related compound (12 -PGJ₂),^[6,7] it differs from it in significant ways, including the catalytic asymmetric Tsuji–Troost reaction to install the C8 stereocenter, the C–H activation/oxygenation of C11, and the application of the catalytic asymmetric Mukaiyama aldol reaction to install the C15 stereocenter.

Scheme 1 summarizes the construction of advanced intermediate **2** starting from commercially available 2-cyclopentenone (**8**) via intermediates **9** and **10**. Thus, employing modified literature procedures,^[8] **8** was first converted to cyclopentenol acetate **9** (DIBAL-H reduction followed by acetylation, 62% overall yield; similar results were obtained with LiAlH₄ or NaBH₄/CeCl₃ as reducing agents, see SI), and thence to dimethyl ester **10** through a palladium-catalyzed Tsuji–Trost asymmetric coupling reaction^[9] with dimethyl malonate (DMM) in the presence of catalytic amounts of [η³-C₃H₅PdCl]₂ and (S,S)-DACH-phenyl Trost ligand and Cs₂CO₃ (71% yield, 97% ee^[10]). Monodecarboxylation of the latter (KI, 130 °C) then furnished intermediate **4** in 94% yield. The desired C–H activation/functionalization of **4** to its enone counterpart **11** required considerable experimentation and led to the rhodium-catalyzed procedure^[11] requiring Rh₂(cap)₄ and *t*BuOOH as an oxidant (48% yield) as shown in Scheme 1. Other methods attempted to carry out this transformation (i.e. **4**→**11**) included the use of Mn(OAc)₃ and *t*BuOOH^[12] (see Table 1 and Scheme 2) or bleach and *t*BuOOH,^[13] both of which furnished the desired product, albeit in lower yields and with more cumbersome protocols. At this juncture we had the opportunity to explore briefly the scope of the rhodium-catalyzed and other oxygenations of substituted cyclopentenones such as **4** and related compounds and discovered an interesting and potentially useful regioselectivity effect. Thus, and as shown in Table 1, while methyl ester **4** and its Weinreb amide sibling **4a** reacted with *t*BuOOH and Rh₂(cap)₄ catalyst to afford the disubstituted enones **11** and **11a**, respectively (through intermediate peroxide species **4e**, Scheme 2), the corresponding reduced substrates **4b** and **4c** behaved differently, leading instead to the trisubstituted enones **11b** and **11c**, respectively, in which the enone functionality had undergone transposition towards the side chain (see Table 1 and Scheme 2). This structural shift may be attributed to electronic effects and the radical nature of the oxygenation reactions as depicted mechanistically in Scheme 2. Thus, reaction of Rh₂(cap)₄ with *t*BuOOH produces *t*BuOO[•] (see Scheme 2a),^[9] initiating the reactions shown in Scheme 2b and 2c. Substrates **4** and **4a** carry an electron-withdrawing group at the β-position from the trisubstituted carbon on the cyclopentene ring that can exert a destabilizing effect on the corresponding tertiary radical at that position, thereby favoring formation of the corresponding secondary radical **4d**^[14] which leads to products **11** and **11a** via intermediate peroxide **4e** (see Scheme 2b). On the other hand, substrates **4b** and **4c** are not suffering from this predicament and, therefore, lead to the more stable tertiary radical **4f** which undergoes rapid rearrangement to the less hindered secondary radical species **4g**, whose facile reaction with *t*BuOO[•] leads to products **11b** and **11c** through the corresponding peroxide (see Scheme 2c). A similar mechanistic rationale may be applicable to the manganese-catalyzed processes shown in Table 1. These observations could be expanded to wider explorations in search of new synthetic technologies.

Returning to Scheme 1 and the synthesis of advanced intermediate **2**, enone **11** was further elaborated to allylic alcohol **12** by Luche reduction^[15] (NaBH₄, CeCl₃, 95% yield, ca. 10:1 dr, inconsequential) and then to hydroxy aldehyde **13** through DIBAL-H reduction (2.2 equiv., 95% yield) in preparation for the pending Wittig olefination reaction. The latter proceeded smoothly by generating the ylid from {5-[(4-methoxybenzyl)oxy]pentyl}-triphenyl-phosphonium iodide [IPh₃P(CH₂)₅OPMB; for preparation see SI] and NaHMDS at –78 °C, and reacting it at low temperature (–78→25 °C) with hydroxy aldehyde **13**,

furnishing, after oxidation with PCC, enone **2**^[6] in 73% overall yield and good stereoselectivity (*Z* : *E* 10:1, by ¹H-NMR spectroscopic analysis). An alternative, slightly higher yielding, albeit longer, sequence for the conversion of allylic alcohol **12** to enone **2** was also carried out as shown in Scheme 1. Thus, **12** was silylated (TBSCl, imid.) to afford the expected TBS ether, which was subjected to DIBAL-H reduction to give the corresponding aldehyde, whose reaction with the ylid generated from IPh₃P(CH₂)₅OPMB and NaHMDS as before furnished bis-olefin **14** (*Z* : *E* 10:1) in 85% overall yield. Desilylation of the latter (TBAF, 91% yield) followed by PCC oxidation then led to enone **2** in 93% yield.

Scheme 3 depicts the catalytic asymmetric synthesis of the required aldehyde fragment **3**. Thus, commercially available 3-hexynol (**16**) was oxidized with DMP to give hex-3-ynal (**17**)^[16] (95% yield, crude, 95 purity by ¹H NMR-analysis). Due to its rather labile nature, the latter was immediately, and without further purification, subjected to enantioselective Mukaiyama aldol reaction employing the trimethylsilyl acetal (**18**)^[17] of benzyl acetate and (*R*)-NOBIN Mukaiyama catalyst (**19**)^[18] (5 mol%) to furnish, upon sequential aqueous work-up and exposure to TBAF, hydroxybenzyl ester **20** (72% yield, 95% ee by ¹⁹F-NMR spectroscopic analysis of the corresponding Mosher esters). The absolute configuration of **20** was confirmed by full Mosher ester analysis (see SI). The hydroxy group of the latter intermediate was then protected as a TBS ether (TBSCl, imid., 88% yield), and the resulting acetylenic compound **21** was selectively reduced with Lindlar catalyst in the presence of quinoline under a hydrogen atmosphere to afford exclusively *Z*-olefin **22** in 99% yield. Selective reduction of the benzyl ester moiety of intermediate **22** with DIBAL-H then produced the targeted aldehyde fragment **3** in 89% yield. An alternative catalytic asymmetric route to aldehyde **3** was also developed employing as a key step a Keck allylation^[19] of 3-*tert*-butyl dimethylsilyloxy propanal with tri-*n*-butylstannane in the presence of catalytic amounts of (*S*)-BINOL^[20] to provide, after silylation and ozonolysis, known aldehyde **23**^[21] in high enantiomeric excess (95% ee, Scheme 3). Wittig olefination of **23** then furnished **24** (87% yield, *Z* : *E* ca. 8:1, chromatographically separable), which was desilylated (py•HBr₃, 63% yield) and oxidized (DMP, 99% yield) to afford aldehyde **3** via hydroxy compound **25** as shown in Scheme 3.

With both fragments, cyclopentenone **2** and aldehyde **3**, readily available in their correct enantiomeric forms, their union through the planned aldol reaction and further elaboration of the product to ¹²-PGJ₃ (**1**) became the next task. Thus, and as shown in Scheme 4, generation of the enolate of **2** at -78 °C with LDA, followed by addition of **3** yielded the expected aldol product **26** as a mixture of C13 epimers (ca. 3:1 dr) in 79% yield.^[6] While the stereochemical outcome of the aldol reaction at C13 was inconsequential, the exclusive stereoselectivity with regards to the C12 stereocenter as a result of steric control was welcome. Mesylation of the C13 diastereomeric mixture **27** (MsCl, Et₃N) followed by treatment of the resulting mesylates (as a mixture or individually) with Al₂O₃^[6] (as purchased, presumably containing traces of water) produced exclusively the *E*-¹²-configurational isomer **28**^[22] in 62% overall yield for the two steps. It should be noted that the presence of moisture in the Al₂O₃ is essential for the success of this reaction as employment of anhydrous Al₂O₃ (obtained by heating commercially available reagent at

400 °C under vacuum) under the same conditions led to double elimination from mesylate TBS ether **27** to afford pentaene **28b** (53% yield) as shown in Scheme 4. All that now remained to reach the coveted ¹²-PGJ₃ (**1**) was the elaboration of the C1 functional group to the desired carboxylic acid moiety and the removal of the TBS group. The former task was most efficiently accomplished through the sequential use of DDQ (to remove the PMB group, **29**, 87% yield), PCC (to convert the resulting hydroxy group to the aldehyde, **30**, 91% yield), and NaClO₂ (Pinnick oxidation,^[6,23] to oxidize the latter moiety to the carboxylic acid group, **31**, 95% yield). Finally, desilylation of the C15 hydroxy group of the resulting precursor (**31**) with aqueous HF furnished ¹²-PGJ₃ (**1**) in 92% yield. ¹²-PGJ₃ methyl ester (**32**) was also prepared by treatment with trimethylsilyldiazomethane (93% yield) for the purposes of more convenient characterization and biological evaluation. Synthetic ¹²-PGJ₃ (**1**) proved identical with the authentic substance by HPLC, UV spectroscopy and mass spectrometry and was found to be equipotent against chronic myelogenous leukemia (CML) stem cells. ¹H NMR spectroscopic analysis of a small authentic sample confirmed its identity to the synthetic material.^[24]

The described chemistry renders ¹²-prostaglandin J₃ (¹²-PGJ₃, **1**) readily available for thorough biological and pharmacological investigations and opens the way for analogue design, synthesis, and biological evaluation of this important ω-3 series of eicosanoids. It also provides inspiration and the foundation for method development in the field of regioselective C–H activation/functionalization of appropriately substituted olefinic substrates to produce useful chemical building blocks. Further research in both areas of investigation is currently in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Hegde S, Kaushal N, Ravindra KC, Chiaro C, Hafer KT, Gandhi UH, Thompson JT, van den Heuvel JP, Kennett MJ, Hankey P, Paulson RF, Prabhu KS. *Blood*. 2011; 118:6909–6919. [PubMed: 21967980]
2. Kudva AK, Kaushal N, Mohinta S, Kennett MJ, August A, Paulson RF, Prabhu KS. *PLOS One*. 2013; 8:e80622. [PubMed: 24312486]
3. Williams SA, Anderson WC, Santaguida MT, Dylla SJ. *Lab. Invest*. 2013; 93:970–982. [PubMed: 23917877]
4. a) Brooks JD, Milne GL, Yin H, Sanchez SC, Porter NA, Morrow JD. *J. Biol. Chem*. 2008; 283:12043–12055. [PubMed: 18263929] b) Lefils-Lacourtablaise J, Socorro M, Gélœn A, Daira P, Debarb C, Loizon E, Guichadant M, Dominguez Z, Vidal H, Lagarde M, Bernoud-Hubac N. *PLOS One*. 2013; 8:e63997. [PubMed: 23734181]
5. Suzuki M, Mori M, Niwa T, Hirata R, Furuta K, Ishikawa T, Noyori R. *J. Am. Chem. Soc*. 1997; 119:2376–2385.
6. a) Acharya HP, Kobayashi Y. *Tetrahedron Lett*. 2004; 45:1199–1202. b) Acharya HP, Kobayashi Y. *Tetrahedron*. 2006; 62:3329–3343.
7. Das S, Chandrasekhar S, Yadav JS, Grée R. *Chem. Rev*. 2007; 107:3286–3337. [PubMed: 17590055]
8. Jacquet O, Bergholz T, Magnier-Bouvier C, Mellah M, Guillot J-C, Fiaud R. *Tetrahedron*. 2010; 66:222–226.

9. a) Trost BM, Bunt RC. *Angew. Chem. Int. Ed. Engl.* 1996; 35:99–102. *Angew. Chem.* **1996**, 108, 70–73. b) Miyazaki T, Yokoshima S, Simizu S, Osada H, Tokuyama H, Fukuyama T. *Org. Lett.* 2007; 9:4737–4740. [PubMed: 17935340]
10. The ee of this compound was determined by optical rotation comparison with that of the known enantiomer, see ref. [9b].
11. Catino AJ, Forslund RE, Doyle MP. *J. Am. Chem. Soc.* 2004; 126:13622–13623. [PubMed: 15493912]
12. Shing TKM, Yeung Y-Y, Su PL. *Org. Lett.* 2006; 8:3149–3151. [PubMed: 16805574]
13. Marwah P, Marwah A, Lardy HA. *Green Chem.* 2004; 6:570–577.
14. Newhouse T, Baran PS. *Angew. Chem. Int. Ed.* 2011; 50:3362–3374. *Angew. Chem.* 2011; 123:3422–3435. and references therein.
15. Luche JL. *J. Am. Chem. Soc.* 1978; 100:2226–2227.
16. Wavrin L, Viala J. *Synthesis.* 2002:326–330.
17. Kiyooka, S-i; Matsumoto, S.; Shibata, T.; Shinozaki, K-i. *Tetrahedron.* 2010; 66:1806–1816.
18. Carreira EM, Singer RA, Lee W. *J. Am. Chem. Soc.* 1994; 116:8837–8838.
19. Keck GE, Boden EP. *Tetrahedron Lett.* 1984; 25:265–268.
20. Kurosu M, Lorca M. *Synlett.* 2005:1109–1112.
21. White JD, Blakemore PR, Browder CC, Hong J, Lincoln CM, Nagorny PA, Robarge LA, Wardrop DJ. *J. Am. Chem. Soc.* 2001; 123:8593–8595. [PubMed: 11525667]
22. The E- ¹²-geometry was assigned on the basis of the observed H13 chemical shift: $\delta = 6.60$ ppm; see Bundy GL, Morton DR, Peterson DC, Nishizawa EE, Miller WL. *J. Med. Chem.* 1983; 26:790–799. [PubMed: 6854581]
23. Bal BS, Childers WE Jr, Pinnick HW. *Tetrahedron.* 1981; 37:2091–2096.
24. Due to its natural scarcity ¹²-PGJ₃ was isolated through a tedious process in only very small amounts (~20–40 μ g, impure). See Supporting Information for ¹H NMR spectroscopic comparison with a readily available synthetic sample.

Figure 1.
Retrosynthetic analysis of Δ^{12} -PGJ₃ (1).

Scheme 1.

Synthesis of cyclopentenone fragment **2**. Reagents and conditions: (a) DIBAL-H (1 M in CH₂Cl₂, 1.2 equiv.), CH₂Cl₂, 0 °C, 30 min; (b) Ac₂O (2.0 equiv.), Et₃N (2.5 equiv.), DMAP (0.1 equiv.), CH₂Cl₂, 0 → 25 °C, 18 h, 62% for two steps; (c) dimethyl malonate (3.0 equiv.), $[(\eta^3\text{-C}_3\text{H}_5)_2\text{PdCl}]_2$ (0.005 equiv.), (S,S)-DACH-phenyl Trost ligand (0.015 equiv.), Cs₂CO₃ (3.0 equiv.), CH₂Cl₂, 25 °C, 3 h, 71%; (d) KI (8.0 equiv.), DMI:H₂O (10:1), 130 °C, 12 h, 94%; (e) Rh₂(cap)₄ (0.01 equiv.), *t*BuOOH (5.0 equiv.), K₂CO₃ (0.5 equiv.), CH₂Cl₂, 1.5 h, 25 °C; then Rh₂(cap)₄ (0.01 equiv.), *t*BuOOH (5.0 equiv.), 1.5 h, 25 °C, 48%;

(f) $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (1.0 equiv.), NaBH_4 (1.0 equiv.), $-30\text{ }^\circ\text{C}$, 10 min, 95%; (g) DIBAL-H (1 M in CH_2Cl_2 , 2.2 equiv.), CH_2Cl_2 , $-78\text{ }^\circ\text{C}$, 45 min, 95%; (h) $\text{IPh}_3\text{P}(\text{CH}_2)_5\text{OPMB}$ (2.5 equiv.), NaHMDS (1 M in THF, 3.0 equiv.), THF, $-78 \rightarrow 25\text{ }^\circ\text{C}$, 18 h, 79%; (i) PCC (2.0 equiv.), CH_2Cl_2 , $25\text{ }^\circ\text{C}$, 3 h, 93%; (j) TBSCl (1.5 equiv.), imid. (3.0 equiv.), CH_2Cl_2 , $0 \rightarrow 25\text{ }^\circ\text{C}$, 15 min; (k) DIBAL-H (1 M in CH_2Cl_2 , 1.1 equiv.), CH_2Cl_2 , $-78\text{ }^\circ\text{C}$, 45 min; (l) $\text{IPh}_3\text{P}(\text{CH}_2)_5\text{OPMB}$ (1.5 equiv.), NaHMDS (1 M in THF, 2.0 equiv.), THF, $-78 \rightarrow 25\text{ }^\circ\text{C}$, 6 h, 85% for three steps; (m) TBAF (1 M in THF, 1.2 equiv.), THF, $0 \rightarrow 25\text{ }^\circ\text{C}$, 5 h, 91%.

DACH-phenyl Trost ligand: 1,2-diaminocyclohexane-*N,N'*-bis(2'-diphenylphosphinobenzoyl); DIBAL-H: diisobutylaluminum hydride; DMAP: 4-dimethylaminopyridine; DMI: 1,3-dimethyl-2-imidazolidinone; DMM: dimethyl malonate; imid.: imidazole; NaHMDS: sodium *bis*(trimethylsilyl)amide; PCC: pyridinium chlorochromate; PMB: *para*-methoxybenzyl; $\text{Rh}_2(\text{cap})_4$: dirhodium tetracaprolactamate; TBAF: tetrabutylammonium fluoride; TBS: *tert*-butyldimethylsilyl; THF: tetrahydrofuran.

Scheme 2.

Mechanistic rationale for the regioselective formation of enones **11**, **11a** and transposed enones **11b** and **11c** (see Table 1).

Scheme 3.

Synthesis of aldehyde fragment **3**. Reagents and conditions: (a) DMP (1.3 equiv.), CH_2Cl_2 , $0 \rightarrow 25^\circ\text{C}$, 1.5 h, 95% crude; (b) benzyl acetate trimethylsilyl acetal (**18**, 2.0 equiv.), (R)-NOBIN Mukaiyama catalyst (**19**, 0.05 equiv.), Et_2O , $-78 \rightarrow -15^\circ\text{C}$, 4 h; then aq. workup; then TBAF (1 M in THF, 4.0 equiv.), THF, 30 min, 72%; (c) TBSCl (2.0 equiv.), imid. (3.0 equiv.), CH_2Cl_2 , 25°C , 3 h, 88%; (d) Pd (5% on CaCO_3/Pb , 0.1 equiv.), quinoline (1.0 equiv.), EtOAc, 25°C , 30 min, 99%; (e) DIBAL-H (1 M in CH_2Cl_2 , 1.3 equiv.), CH_2Cl_2 , $-78 \rightarrow -25^\circ\text{C}$, 1 h, 89%; (f) BrPh_3PPr (2.0 equiv.), NaHMDS (2.0 equiv.), 0°C , 30 min;

then **23**, $-78 \rightarrow 25\text{ }^{\circ}\text{C}$, THF, 2 h, 87%; g) py^*HBr_3 (0.1 equiv.), MeOH, $-10\text{ }^{\circ}\text{C}$, 1.25 h, 63%; h) DMP (1.3 equiv.), CH_2Cl_2 , $0 \rightarrow 25\text{ }^{\circ}\text{C}$, 2 h, 99%. DMP: 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1*H*)-one; NOBIN: 2-amino-2'-hydroxy-1,1'-binaphthyl; py: pyridine; TMS: trimethylsilyl.

Scheme 4.

Coupling and completion of the synthesis of 12 -PGJ₃ (**1**). Reagents and conditions: (a) LDA (2.0 equiv.); then **2** (1.0 equiv.); then **3** (1.2 equiv.), THF, -78°C , 30 min, 79%; (b) MsCl (5.0 equiv.), Et₃N (10 equiv.), CH₂Cl₂, 0°C , 5 min; (c) Al₂O₃ (21 equiv.), CH₂Cl₂, 25°C , 8 h, 62% for two steps; (d) DDQ (1.5 equiv.), CH₂Cl₂:H₂O (16:1), 0°C , 45 min; (e) PCC (2.0 equiv.), CH₂Cl₂, 25°C , 2 h; (f) NaClO₂ (1.5 equiv.), NaH₂PO₄ (1.5 equiv.), 2-methyl-2-butene (10 equiv.), *t*BuOH:H₂O (4:3), 25°C , 30 min, 75% for three steps; (g) HF (50% aq, 50 equiv.), MeCN, 0°C , 45 min, 92%; (h) TMSCHN₂ (2 M in Et₂O), C₆H₆:

MeOH (3:2), 25 °C, 30 min, 93%; i) Al₂O₃ anhyd. (30 equiv.), CH₂Cl₂, 25 °C, 1 h, 53%.
DDQ: 2,3-dichloro-5,6-dicyano-1,4-benzoquinone; LDA: lithium diisopropylamide.

Table 1Regioselective C–H activation/oxygenation of substituted cyclopentenones.^[a]

Entry	Substrate	Catalyst	Product	Yield [%] ^[d]
1		Rh ₂ (cap) ₄ ^[b]		48
2		Mn(OAc) ₃ ^[c]		35
3		Rh ₂ (cap) ₄ ^[b]		41
4		Mn(OAc) ₃ ^[c]		37
5		Rh ₂ (cap) ₄ ^[b]		63
6		Mn(OAc) ₃ ^[c]		61
7		Rh ₂ (cap) ₄ ^[b]		45
8		Mn(OAc) ₃ ^[c]		39

^[a] Reactions were carried out on 1.0 mmol scale at 25 °C. Reagents and Conditions:^[b] Rh₂(cap)₄ (0.01 equiv.), *t*BuOOH (5.0 equiv.), K₂CO₃ (0.5 equiv.), CH₂Cl₂, 1.5 h; then Rh₂(cap)₄ (0.01 equiv.), *t*BuOOH (5.0 equiv.), 1.5 h;^[c] Mn(OAc)₃ (0.25 equiv.), *t*BuOOH (4.0 equiv.), MS Å3, EtOAc, 24 h.^[d] Yields refer to chromatographically isolated and spectroscopically pure products.