Published in final edited form as:

Sci Signal. ; 7(327): ra49. doi:10.1126/scisignal.2004732.

A PLC γ 1-Dependent, Force-Sensitive Signaling Network in the Myogenic Constriction of Cerebral Arteries

Albert L. Gonzales^{2,3}, Ying Yang², Michelle N. Sullivan², Lindsey Sanders², Fabrice Dabertrand³, David C. Hill-Eubanks³, Mark T. Nelson^{3,4}, and Scott Earley^{1,*}

¹Department of Pharmacology, University of Nevada School of Medicine, Reno, NV 89557–0318, USA

²Vascular Physiology Research Group, Department of Biomedical Sciences, Colorado State University, Fort Collins, CO 80523, USA

³Department of Pharmacology, University of Vermont, Burlington, VT 05405, USA

⁴Institute of Cardiovascular Sciences, University of Manchester, Manchester M13 9NT, UK

Abstract

Maintaining constant blood flow in the face of fluctuations in blood pressure is a critical autoregulatory feature of cerebral arteries. An increase in pressure within the artery lumen causes the vessel to constrict through depolarization and contraction of the encircling smooth muscle cells. This pressure-sensing mechanism involves activation of two types of transient receptor potential (TRP) channels: TRPC6 and TRPM4. We provide evidence that the activation of the γ 1 isoform of phospholipase C (PLC γ 1) is critical for pressure sensing in cerebral arteries. Inositol 1,4,5-trisphosphate (IP₃), generated by PLC γ 1 in response to pressure, sensitized IP₃ receptors (IP₃Rs) to Ca²⁺ influx mediated by the mechanosensitive TRPC6 channel, synergistically increasing IP₃R-mediated Ca²⁺ release to activate TRPM4 currents, leading to smooth muscle depolarization and constriction of isolated cerebral arteries. Proximity ligation assays demonstrated colocalization of PLC γ 1 and TRPC6 with TRPM4, suggesting the presence of a force-sensitive, local signaling network comprising PLC γ 1, TRPC6, TRPM4, and IP₃Rs. Src tyrosine kinase activity was necessary for stretch-induced TRPM4 activation and myogenic constriction, consistent with the ability of Src to activate PLC γ isoforms. We conclude that contraction of cerebral artery smooth muscle cells requires the integration of pressure-sensing signaling pathways and their convergence on IP₃Rs, which mediate localized Ca²⁺-dependent depolarization through the activation of TRPM4.

*Corresponding author. searley@medicine.nevada.edu.

Author contributions: A.L.G., Y.Y., M.N.S., L.S., F.D., and S.E. designed and performed the experiments; A.L.G., Y.Y., M.N.S., F.D., and S.E. analyzed the data; A.L.G., F.D., and S.E. constructed the figures; A.L.G., D.C.H.-E., M.T.N., and S.E. conceptualized and wrote the paper.

Competing interests: The authors declare that they have no competing interests.

INTRODUCTION

Localized increases in intraluminal pressure provoke vasoconstriction of small arteries and arterioles (1). This important part of the autoregulatory mechanism, first described by Bayliss and termed the vascular myogenic response (2), allows regional blood flow to remain essentially constant during transient changes in perfusion pressure (3, 4). Myogenic vasoconstriction results from pressure-induced depolarization of the vascular smooth muscle cell plasma membrane (5), which causes Ca^{2+} influx through voltage-dependent Ca^{2+} channels (VDCCs) (6), as well as increases in the Ca^{2+} sensitivity of the contractile apparatus (7–9). Longitudinal stretch depolarizes the plasma membrane of isolated arterial myocytes (10), demonstrating that these cells can directly detect mechanical strain within the vascular wall resulting from pressure increases. However, the underlying molecular mechanisms responsible for sensing stretch and eliciting membrane depolarization remain elusive.

In the simplest theoretical scheme to account for pressure-induced depolarization, stretch of the plasma membrane alters the activity of inherently mechanosensitive ion channels. In support of this possibility, various studies have implicated several members of the transient receptor potential (TRP) cation channel superfamily, including TRPP2 (11), TRPV2 (12), TRPC6 (13), and TRPM4 (14, 15), as well as the Ca^{2+} -activated Cl^- channel TME-M16A (16, 17), in pressure-induced membrane depolarization. TRPC6 (13) and TRPM4 channels appear to be indispensable for pressure-induced cerebral artery constriction (14, 18). TRPC6 is a Ca^{2+} -permeable, nonselective cation channel that is directly activated by the second messenger diacylglycerol (DAG) (19). Some studies have reported that TRPC6 channels are inherently mechanosensitive (20), whereas others have suggested that DAG generated by the activity of phospholipase C (PLC) is involved in mediating the response to stretch (21). Unlike TRPC6, TRPM4 is impermeant to Ca^{2+} , selective for monovalent cations, and activated by increased concentrations of intracellular Ca^{2+} (22, 23). In vascular smooth muscle cells, TRPM4 activity is stimulated by Ca^{2+} released from the sarcoplasmic reticulum (SR) through inositol 1,4,5-trisphosphate (IP_3) receptors (24, 25). Despite a report that TRPM4 channels are inherently mechanosensitive based on data obtained from heterologous expression systems (26), the mechanosensitivity of TRPM4 has not been clearly established.

Although TRPC6 and TRPM4 channels have been implicated in the myogenic response, their position in the pressure-responsive pathway and the mechanism by which their activities are integrated and regulated to achieve a graded response remain unclear. However, the activation of TRPC6 and TRPM4 by DAG and IP_3 , respectively, which are generated by cleavage of the membrane phospholipid phosphatidylinositol 4,5-bisphosphate [$\text{PtdIns}(4,5)\text{P}_2$] by PLC, suggests that PLC may be centrally involved in stretch-induced activation of both TRPM4 and TRPC6. Because PLC isoforms lack trans-membrane domains and thus are unlikely to respond directly to mechanical stimuli, the nature of the stress-sensing entity remains an open question. Several pathways can sense and signal stretch, including actin polymerization (9, 27, 28) and integrin-mediated (29) and G_q protein-coupled receptor-mediated signaling (21). It is posited that both PLC and TRPC6 play a role in this latter pathway, but details of this signaling cascade remain incompletely

characterized and certain key features of the model rely largely on data obtained from heterologous expression systems.

Here, we sought to determine how second messengers generated by PLC and interactions between TRPM4 and TRPC6 channels are linked to bring about pressure-induced membrane depolarization. We showed that generation of IP₃ by the γ 1 isoform of PLC (PLC γ 1) was essential for pressure-induced TRPM4 channel activity, smooth muscle membrane depolarization, and generation of myogenic tone. We also showed that PLC γ 1 activity alone was insufficient to stimulate TRPM4 currents in response to stretch and that Ca²⁺ influx through TRPC6 was also required, implying that TRPC6 transduces a mechanical stimulus to TRPM4. Finally, we demonstrated that TRPC6, but not TRPM4, is directly mechanosensitive and that the stress-sensing mechanism upstream of PLC γ 1 depended on the tyrosine kinase Src. Together, these findings describe a mechanism in which converging signaling pathways integrate the sensation of stretch in vascular smooth muscle cells to mediate pressure-induced membrane depolarization and myogenic vasoconstriction.

RESULTS

TRPM4 is not inherently mechanosensitive

Previous studies have indicated that TRPC6 and TRPM4 are critically involved in pressure-induced cerebral arterial myocyte membrane depolarization (13–15, 18), suggesting that these channels can be activated by mechanical stretch. We examined the mechanosensitivity of each channel by measuring changes in single-channel activity in response to membrane stretch induced by applying negative pressure (–20 mmHg) through the recording electrode in patch-clamp experiments using human embryonic kidney (HEK) cells expressing recombinant yellow fluorescent protein (YFP)-tagged TRPC6 or green fluorescent protein (GFP)-tagged TRPM4 (30). In agreement with previous studies (20, 21), the total open probability of TRPC6 channels increased nearly threefold in response to stretch of the plasma membrane (Fig. 1A). Current-voltage relationships recorded under both pressure conditions were linear with a slope conductance consistent with the previously reported unitary conductance of TRPC6 [~35 pS (19)]. Basal channel activity recorded from HEK cells expressing TRPM4 channels did not change when negative pressure was applied (Fig. 1B). The slope conductance of these currents was ~25 pS, similar to the reported unitary conductance of TRPM4 channels (23). These data indicate that heterologously expressed TRPC6 channels, but not TRPM4 channels, are mechanosensitive, under the conditions used for this study.

Phospholipase activity is necessary for stretch-induced TRPM4 activity in arterial myocytes

In native, freshly isolated cerebral artery smooth muscle cells, TRPM4 activity can be continuously recorded as transient inward cation currents (TICCs) (25). TICC activity increased after membrane stretch associated with cell swelling induced by an 80 mosmol/liter hypotonic challenge (Fig. 1C). We also examined the mechanosensitivity of TRPM4 in these cells using voltage-clamp in the amphotericin B-perforated configuration, and recorded TICCs when defined amounts of suction were applied to the plasma membrane by

the recording electrode. TICC activity increased in proportion to the amount of applied negative pressure (fig. S1A). Activation was maximal at an applied intrapipette pressure of -20 mmHg, and the half-maximal pressure of activation (P_{50}) was -12.8 mmHg (fig. S1, A and B). Stretch-activated increases in TICC activity returned to baseline after release of intrapipette pressure, indicating that current activation did not result from permanent damage to the plasma membrane (fig. S1, A and C). Basal and stretch-induced TICC activity was attenuated by the selective TRPM4 inhibitor 9-phenanthrol (15, 31, 32) (fig. S1D), demonstrating that currents originated from TRPM4 channels. These findings indicate that TRPM4 is activated by stretch of the plasma membrane in native cerebral artery smooth muscle cells, consistent with the channel's role in myogenic vasoconstriction.

TRPM4 lacks inherent mechanosensitivity, but endogenous TRPM4 channels are activated when stretch is applied to the plasma membrane of arterial myocytes, suggesting that the channel is indirectly activated by force-sensitive signaling pathways in these cells. TRPM4 activity requires high concentrations (~ 1 to 10 μM) of intracellular Ca^{2+} (22, 23), and in native smooth muscle cells, the channel is stimulated by Ca^{2+} released from proximal IP_3 receptors (IP_3Rs) (24, 25). IP_3 is generated by PLC-mediated hydrolysis of $\text{PtdIns}(4,5)\text{P}_2$, suggesting that tonic PLC activity is necessary for basal TICC activity. Consistent with this possibility, swelling-induced TICC activity was lost in the presence of U73122 (Fig. 1D). Furthermore, basal TRPM4 activity was eliminated by the PLC inhibitor U73122 and was restored by the membrane-permeant IP_3 analog Bt- IP_3 (fig. S1E). TICC activity was insensitive to U73343, an inactive analog of U73122 (fig. S2A). TICC activity stimulated by applying negative pressure to the plasma membrane was also attenuated after administration of U73122 (fig. S1F). Control experiments demonstrated that the frequency and amplitude of spontaneous transient outward currents (STOCs), activated by Ca^{2+} released from the SR through ryanodine receptors (33), were not altered by U73122 (fig. S2B), suggesting that the compound did not alter SR Ca^{2+} stores under these conditions. These findings indicate that tonic generation of IP_3 by PLC is necessary for basal and stretch-induced TICC activity in smooth muscle cells.

We assessed the consequences of disrupted stretch-induced, PLC-dependent activation of TRPM4 activity for myogenic constriction using isolated cerebral resistance arteries. In agreement with earlier findings (34), arteries pretreated with U73122 exhibited diminished pressure-dependent vasoconstriction compared with vessels treated with vehicle controls (fig. S3A). U73122 pretreatment also decreased vasoconstriction in response to the purinergic receptor vasoconstrictor agonist uridine 5'-triphosphate (UTP) (fig. S3B), but constriction in response to increased concentrations of extracellular KCl did not differ between groups (fig. S3C). These findings indicate that general PLC activity is necessary for both pressure- and agonist-induced vasoconstriction, and its blockade does not directly influence constriction in response to membrane depolarization.

PLC γ 1 regulates TRPM4 activity in arterial myocytes

Several PLC isoforms are present in vascular smooth muscle cells (35). We detected PLC γ 1, PLC γ 2, and PLC β 2 in native cerebral artery smooth muscle cells using reverse transcription polymerase chain reaction (RT-PCR) and immunocytochemistry (Fig. 2, A and B). PLC γ 1

and PLC γ 2 immunolabeling appeared at or near the plasma membrane, whereas PLC β 2 immunolabeling was detected as discrete puncta located primarily within the cytosol (Fig. 2B). TRPM4 immunolabeling was also concentrated at or near the plasma membrane (Fig. 2B). We used an in situ proximity ligation assay (36) to identify PLC isoforms that are closely associated with TRPM4 in cerebral myocytes. This technique produces positive immunolabeling only if the two targeted proteins are localized within 40 nm of each other. Dual immunolabeling for PLC γ 1 and TRPM4 produced significantly higher numbers of puncta per cell near the plasma membrane than were detected with dual labeling for TRPM4 and PLC γ 2 or PLC β 2 (Fig. 2C). These findings suggest that PLC γ 1 and TRPM4 are within 40 nm of each other near the plasma membrane in contractile smooth muscle cells, indicating that PLC γ 1 could be selectively involved in the regulation of TRPM4 activity.

Small interfering RNA (siRNA) was used to test the hypothesis that PLC γ 1 regulates TRPM4 activity in native cerebral artery smooth muscle cells. siRNA targeting PLC γ 1 or control (scrambled) siRNA was introduced into the arterial myocytes of intact cerebral arteries using a previously described reversible permeabilization technique (25) (fig. S4). TICC were recorded from arterial myocytes, enzymatically isolated from vessels treated with PLC γ 1 or control (nonsilencing) siRNA, whereas stretch was applied to the plasma membrane. Cells isolated from PLC γ 1 siRNA-treated vessels exhibited an about 73% decrease in stretch-induced TICC activity compared with cells from vessels treated with control siRNA (Fig. 2D). TICC activity in cells isolated from vessels treated with PLC γ 1 siRNA was partially restored by administration of Bt-IP $_3$ (Fig. 2D), indicating that diminished IP $_3$ production after PLC γ 1 silencing contributes to the loss of TRPM4 activity. These findings demonstrate that the activity of PLC γ 1 is necessary for the generation of TRPM4 currents in cerebral myocytes in response to stretch of the plasma membrane.

PLC γ 1 is essential for pressure-induced smooth muscle cell depolarization and the development of myogenic tone

Acute knockdown of TRPM4 largely eliminates pressure-induced smooth muscle cell depolarization and myogenic constriction of cerebral arteries (14, 15), and the current findings indicate that stretch-induced TRPM4 activity is dependent on the generation of IP $_3$ by PLC γ 1. Therefore, we examined the role of PLC γ 1 in pressure-induced smooth muscle membrane depolarization and vasoconstriction using siRNA-mediated silencing in intact cerebral arteries. Microelectrode recordings obtained from pressurized vessels demonstrated that the mean membrane potential of smooth muscle cells in control siRNA-treated arteries was -59.5 ± 6.6 mV at rest and was depolarized to -39.0 ± 3.0 mV by an increase in pressure to 80 mmHg. This response was significantly attenuated in PLC γ 1 siRNA-treated arteries, demonstrating a role for PLC γ 1 in pressure-induced depolarization of cerebral artery smooth muscle cells (Fig. 3, A and B). Moreover, myogenic tone was essentially abolished in PLC γ 1 siRNA-treated arteries (Fig. 3, C and D), demonstrating a requirement for PLC γ 1 in the myogenic response to pressure. In contrast, we found that agonist-induced vasoconstriction with UTP did not differ between PLC γ 1 siRNA- and control siRNA-treated vessels (Fig. 3E), indicating that PLC γ 1 activity is not involved in purinergic receptor-mediated vasoconstriction. Arterial constriction in response to increased extracellular K $^+$ ion concentration did not differ between PLC γ 1 siRNA- and control

siRNA-treated vessels (Fig. 3F), indicating that absence of PLC γ 1 does not directly impair membrane depolarization-dependent vasoconstrictor pathways. These findings show that PLC γ 1 is specifically involved in pressure-induced smooth muscle cell depolarization and is required for the development of myogenic tone. Considering the inhibitory effects of U73122 on UTP-mediated vasoconstriction (Fig. 2B), the absence of an effect of PLC γ 1 on the contractile response to UTP suggests that agonist-induced vasoconstrictor responses require the activity of a different PLC isoform (for example, PLC β). The close proximity of PLC γ 1 and TRPM4 (Fig. 2C) and our previous work showing that TRPM4 and IP $_3$ receptors are functionally coupled (24) suggest that TRPM4, PLC γ 1, and IP $_3$ receptors are part of a force-sensitive signaling network in vascular smooth muscle cells that mediates membrane potential depolarization in response to increases in intraluminal pressure.

Ca $^{2+}$ influx through TRPC6 is required for pressure-induced Ca $^{2+}$ waves and activation of TRPM4

Previous studies have shown that knocking down either TRPM4 or TRPC6 largely eliminates (~80 to 90%) the myogenic response in cerebral arteries, consistent with a serial mechanism (13, 14). However, we previously have shown that TRPM4 channels are not responsive to extracellular Ca $^{2+}$ influx, arguing against a direct link between TRPC6-mediated Ca $^{2+}$ influx and TRPM4 activation in the myogenic response. In particular, TRPM4 is activated by IP $_3$ R-mediated release of Ca $^{2+}$ from intracellular stores. An alternative mechanism that is compatible with these observations and a serial linkage between TRPC6 and TRPM4 is a Ca $^{2+}$ -induced Ca $^{2+}$ -release mechanism in which Ca $^{2+}$ influx through stretch-activated TRPC6 channels activates IP $_3$ Rs, promoting the release of Ca $^{2+}$ from intracellular stores that, in turn, stimulates TRPM4 activity. This mechanism predicts that stretch induces Ca $^{2+}$ influx and release from intracellular stores and that both events are required to activate TRPM4. We examined the role of TRPC6-mediated Ca $^{2+}$ influx in pressure-induced Ca $^{2+}$ waves in arteries and stretch-activated TRPM4 currents in freshly isolated myocytes. In intact pressurized arteries that were continuously exposed to the L-type Ca $^{2+}$ channel blocker nimodipine and the ryanodine receptor blocker tetracaine to isolate IP $_3$ R-mediated events, the percentage of vascular smooth muscle cells exhibiting spontaneous propagating Ca $^{2+}$ waves increased in a pressure-dependent manner (Fig. 4A). Preincubation with a TRPC6 antibody targeting an extracellular epitope eliminated these intraluminal pressure-induced Ca $^{2+}$ waves (Fig. 4, A and B) and blocked currents in TRPC6-expressing HEK 293 cells (fig. S6). In vessels preincubated with the inhibitory antibody, administration of the purinergic receptor agonist UTP restored Ca $^{2+}$ waves (Fig. 4, A and B), possibly through G $_q$ protein-coupled receptor-dependent activation of TRPC3 (37). These results suggest that increases in intraluminal pressure and stretch of the plasma membrane elicit Ca $^{2+}$ release through IP $_3$ Rs and that this response is governed by activation of plasma membrane TRPC6 channels.

To determine whether Ca $^{2+}$ influx through TRPC6 was necessary for the pressure-dependent, IP $_3$ R-mediated activation of TRPM4, we examined the role of TRPC6 in stretch-induced TICC activity. Internal SR Ca $^{2+}$ stores were maintained after acute removal of extracellular Ca $^{2+}$, accomplished by immediate (<1 to 2 min) exposure to zero extracellular Ca $^{2+}$ (fig. S5). Under these conditions, stimulated Ca $^{2+}$ release events from internal stores

are maintained (38). Removal of extracellular Ca^{2+} prevented stretch-induced increases in TRPM4 activity (Fig. 4B). In agreement with our previous findings (24, 25), depletion of SR Ca^{2+} stores with the sarco/endoplasmic reticulum Ca^{2+} -adenosine triphosphatase (ATPase) (SERCA) inhibitor cyclopiazonic acid essentially abolished both basal and stretch-induced TICC activity (Fig. 4C). These data indicate that Ca^{2+} influx and intact SR Ca^{2+} stores are necessary for stretch-induced increases in TRPM4 activity in native smooth muscle cells.

We next sought to establish that TRPC6, which is Ca^{2+} -permeable, activated by plasma membrane stretch (13, 20, 21) (see also Fig. 1A), and necessary for myogenic vasoconstriction of cerebral arteries (13), is the mechanosensitive ion channel involved in this response. Using in situ proximity ligation assay, we found that TRPC6 channels, but not large-conductance Ca^{2+} -activated K^{+} (BK_{Ca}) channels, are within 40 nm of TRPM4 channels (Fig. 4D). These data indicate that TRPC6 channels are positioned to provide the Ca^{2+} influx necessary for stretch-induced activation of TRPM4. Stretch-induced increases in TRPM4 activity were attenuated by GsMTx-4, a peptide derived from the venom of the Chilean rose hair tarantula that inhibits mechanosensitive ion channels, including TRPC6 (20) (Fig. 4E). GsMTX-4 blocked recombinant TRPC6 channels expressed in HEK 293 cells, but had no direct effects on heterologously expressed TRPM4 channels (fig. S6). These results indicate a functional linkage between TRPC6 and TRPM4. Stretch-activated TRPM4 currents in cerebral artery myocytes were also attenuated by an antibody that binds to an extracellular epitope of TRPC6 (Fig. 4E, middle), providing additional support for functional communication between TRPC6 and TRPM4. The anti-TRPC6 antibody blocked currents recorded from TRPC6-expressing HEK 293 cells, but not those recorded from HEK 293 cells transfected with TRPM4 (fig. S6), and did not affect currents recorded from cerebral artery myocytes when coad-ministered with its antigenic peptide (Fig. 4E), indicating that these effects were specific. Consistent with these observations and the results of in situ proximity ligation assay experiments, the anti-TRPC6 antibody immunolabeled the plasma membrane of native cerebral artery smooth muscle cells but failed to immunolabel these cells when the antibody was preincubated with the antigenic peptide (Fig. 4E, right). In addition, swelling-induced TICC activity was absent in cells that were preincubated with the TRPC6 inhibitory antibody (Fig. 4F). Collectively, these data indicate that Ca^{2+} influx through TRPC6 is necessary for stretch-induced IP_3R Ca^{2+} -mediated activation of TRPM4 in contractile cerebral artery smooth muscle cells.

AT₁ receptors and Src family tyrosine kinase activity contribute to stretch-induced TRPM4 activity in arterial myocytes

The dependence of the myogenic response on $\text{PLC}\gamma 1$ and the presumed absence of PLC mechanosensitivity prompted us to address the possible upstream involvement of the Src family of nonreceptor tyrosine kinases, which can signal through $\text{PLC}\gamma$ isoforms (35) and rapidly activate in response to mechanical stress (39). Consistent with a role for Src family tyrosine kinases in this pathway, we found that two structurally unrelated Src inhibitors, PP2 (Fig. 5A) or SKI-1 (fig. S7A), inhibited stretch-induced TRPM4-mediated currents. Administration of the Src kinase inhibitors PP2 (Fig. 5B) and SKI-1 (fig. S7B) nearly abolished predeveloped myogenic tone in isolated cerebral arteries. PP2 did not affect KCl-induced constriction (Fig. 5C). These data indicate that Src tyrosine kinase activity is

required for the development of myogenic tone. Previous work indicated that stimulation of angiotensin II (Ang II) receptor type 1 (AT₁R) initiates Src tyrosine kinase–dependent phosphorylation of PLC γ 1 in smooth muscle cells (40). In agreement with these results, administration of the AT₁R inhibitor losartan prevented stretch-induced increases in TRPM4 activity (Fig. 5D), suggesting a possible upstream role for this G_{q/11}-coupled receptor in the pressure-response pathway.

The findings presented here support a signaling pathway (Fig. 6) in which pressure induces PLC γ 1 activity through Src tyrosine kinase, likely with the involvement of the upstream G_{q/11}-coupled receptor AT₁R, leading to the generation of IP₃ and DAG. PLC γ 1-derived IP₃ sensitizes IP₃Rs located on the SR to the influx of Ca²⁺ through DAG-stimulated (fig. S8) and/or pressure-activated TRPC6 channels to generate localized Ca²⁺ nanodomains that activate TRPM4 channels located on the plasma membrane. The resulting influx of Na⁺ depolarizes the membrane to initiate Ca²⁺ influx through VDCCs and causes myocyte contraction.

DISCUSSION

Contraction of arterial myocytes in response to increased intraluminal pressure constitutes a basic intrinsic autoregulatory blood flow control mechanism. The ability of vascular smooth muscle cells to sense and appropriately respond to pressure is central to this critical response, but the underlying mechanisms are poorly understood. The findings of the current study incorporate previous studies implicating PLC (34, 41, 42), PKC (protein kinase C) (30, 43, 44), TRPC6 (13), and TRPM4 (14, 15, 18) as major players in this process and describe a signaling pathway responsible for the sensation of stretch and initiation of myogenic constriction. According to this model, stretch of the plasma membrane has a twofold effect: (i) AT₁R- and Src tyrosine kinase–dependent activation of PLC γ 1 and generation of IP₃, and (ii) stimulation of Ca²⁺ influx through TRPC6. IP₃ acts primarily by enhancing the sensitivity of IP₃Rs to local cytosolic Ca²⁺ (45); therefore, IP₃Rs play a key role in integrating these two signaling pathways. IP₃-mediated sensitization of IP₃Rs acts synergistically with TRPC6-mediated Ca²⁺ influx to promote Ca²⁺ release from IP₃Rs through a Ca²⁺-induced Ca²⁺-release mechanism, resulting in activation of proximal TRPM4 channels in response to increases in luminal pressure. Our previous observation that the TRPM4 inhibitor 9-phenanthrol relaxes vessels with tone independent of a change in pressure (15) suggests that this mechanism also operates under steady-state conditions, although it is possible that other pathways contribute.

PLC activity is necessary for the development of myogenic tone (34, 41, 42), such that pharmacological inhibition leads to hyperpolarization and disruption of sustained pressure-induced constriction of arteries (34, 41). Intracellular concentrations of endogenous IP₃ in smooth muscle cells are higher in myocytes isolated from vessels exposed to higher intraluminal pressures (46), suggesting a positive relationship between PLC activity and membrane stretch. In addition, Mufti *et al.* reported an increase in IP₃R-mediated Ca²⁺-release events in cerebral arteries in response to higher intraluminal pressure (47). These data suggest that pressure can stimulate PLC activity, thereby generating IP₃ and the subsequent release of Ca²⁺ through IP₃Rs. Yet, PLC is a common signaling enzyme with

multiple isoforms that contribute to different signaling pathways. PLC γ 1 is located proximal to TRPM4 and is a key element in stretch-induced, but not agonist-dependent, TRPM4 activation, suggesting that isoform specificity and microdomain localization can prevent crosstalk between different PLC subtypes.

PLC γ isoforms are downstream targets of Src tyrosine kinase activity (35), and previous studies have indicated that PLC γ 1 can be activated by Src tyrosine kinase-dependent phosphorylation after Ang II-induced AT $_1$ R stimulation in smooth muscle cells (40). These studies have advanced the concept that G $_q$ protein-coupled receptors, as direct mechanosensors (21, 48–50), contribute to myogenic vasoconstriction by activating depolarizing cation currents (21). Zou *et al.* demonstrated that agonist-independent, stretch-induced hypertrophy of cardiomyocytes is blocked by inhibition of AT $_1$ Rs but remains intact in cells isolated from angiotensinogen knockout mice (48). Furthermore, Mederos y Schnitzler and co-workers have shown that losartan diminishes swelling-induced increases in intracellular Ca $^{2+}$ in primary renal arterial smooth muscle cells and partially reverses the myogenic tone of isolated rat cerebral arteries (21). These authors further reported that all G $_q$ protein-coupled receptors tested in an exogenous expression system (HEK cells) confer responsiveness to osmotic swelling when coexpressed with TRPC6, and suggested that mechanosensitivity may be a general feature of G $_q$ protein-coupled receptors (21). Although our data also support a role for AT $_1$ R in the pressure-dependent activation of TRPC6 (and TRPM4) in cerebral arteries, we found that these effects did not extend to the P2Y receptor, another class of G $_q$ protein-coupled receptor. Specifically, whereas general inhibition of PLC eliminated the tone induced by the P2Y receptor agonist UTP (fig. S3), knockdown of PLC γ 1 in vessels did not alter UTP-induced tone (Fig. 3E). These data suggest that P2Y receptors are not involved in the activation of PLC γ 1, and thus, activation of this pathway does not appear to be a general property of G $_q$ protein-coupled receptors, in contrast to the results reported by Mederos y Schnitzler *et al.* (21). One possible explanation is that different PLC isoform(s) could couple to different G $_q$ protein-coupled receptors, a question that was not specifically addressed in this previous study. It has been suggested that the purinergic G $_q$ protein-coupled receptors P2Y $_4$ or P2Y $_6$, but not AT $_1$ Rs, are involved in generating myogenic tone in smaller penetrating parenchymal arterioles (51), implying that different mechanosensitive G $_q$ protein-coupled receptors are involved in myogenic tone in different vascular beds. Further investigation is needed to clarify the details of mechanotransduction mechanisms, particularly PLC isoform dependence, and assess the involvement of other PLC-dependent mechanosensitive pathways, such as those involving integrins (52, 53) or other extracellular matrix proteins.

TRPC6 clearly contributes to the generation of myogenic tone (13), but whether these channels are directly or indirectly mechanosensitive has been a matter of controversy. TRPC6-deficient mice have been reported to exhibit enhanced membrane depolarization, myogenic tone, and increased blood pressure (54), a phenotype that has been attributed to up-regulation of constitutively active, DAG-sensitive TRPC3 channels. In addition, an acute antisense strategy directed against TRPC6 in intact cerebral arteries, which prevents functional compensation by closely related channels, results in attenuation of pressure-dependent depolarization (13). These data implicate TRPC6 as a player in the pressure-

sensing mechanism but do not address whether this channel responds directly or indirectly to mechanical stretch. Various groups have characterized mechanosensitive channels in smooth muscle cells (55–57), including Park *et al.*, who have identified stretch-activated cation channels in native vascular smooth muscle cells with a unitary conductance of ~30 pS that are stimulated by a DAG analog and require PLC activity (58), properties that are reminiscent of TRPC channels. These observations are consistent with the possibility that DAG generated by pressure-induced increases in PLC γ 1 activity stimulate TRPC6 in cerebral artery smooth muscle, but do not preclude direct mechanosensitivity of the channel. In agreement with this possibility, Spassova *et al.* have reported that stretch-induced activation of recombinant TRPC6 is independent of PLC activity, consistent with the channel exhibiting direct mechanosensitivity (20).

Collectively, our results define a force-sensitive signaling module in arterial smooth muscle cells in which converging mechanosensitive inputs (namely, AT₁Rs and TRPC6) are integrated by PLC γ 1- and IP₃R-mediated Ca²⁺ release to link membrane stretch with membrane depolarization (TRPM4) to generate myogenic tone. Although our model assumes that TRPC6 is directly activated by stretch, it is capable of accommodating an indirect TRPC6 activation mechanism.

MATERIALS AND METHODS

General electrophysiological recordings

Currents were recorded using an Axopatch 200B amplifier equipped with an Axon CV 203BU headstage (Molecular Devices). Recording electrodes (2 to 3 megohms) were pulled, polished, and coated with wax to reduce capacitance. Currents were filtered at 1 kHz, digitized at 40 kHz, and stored for subsequent analysis. Clampex and Clampfit versions 10.2 (Molecular Devices) were used for data acquisition and analysis, respectively. All experiments were performed at room temperature (22°C).

HEK 293 culture and transfection

HEK 293 cells were cultured in 1× high-glucose Dulbecco's modified Eagle's medium (DMEM) (Gibco) supplemented with 10% fetal bovine serum (Gibco) and 0.5% penicillin-streptomycin (Gibco). Cells were incubated at 37°C with 6% CO₂ and subcultured when confluent using 0.25% trypsin-EDTA (Gibco), and media were changed every 2 to 3 days. HEK 293 cells were transfected with DNA encoding TRPC6-YFP or TRPM4-GFP using a Gene Pulser Xcell (Bio-Rad, 110-V 25-ms pulse) and transferred to culture flasks. Transfected cells were replated on coverslips 4 hours before patch-clamp experiments.

Electrophysiological recordings from TRPM4- and TRPC6-expressing HEK 293 cells

TRPM4 and TRPC6 single-channel activity was recorded using the cell-attached patch-clamp configuration. The patch pipette solution contained 80 mM Cs-aspartate, 30 mM CsCl, 1.5 mM MgCl₂, 10 mM tetraethylammonium (TEA), 60 mM D-mannitol, and 10 mM Hepes at pH 7.2 (CsOH). Bath solution contained 120 mM KCl, 1.5 mM MgCl₂, 10 mM Hepes, 50 mM D-mannitol, and 1 mM EGTA at pH 7.4 (KOH). Mechanical stimulation of the cell membrane was accomplished by applying suction (or negative pressure of –20

mmHg) through the recording pipette with a DPM-1B Pneumatic Transducer Tester (Fluke Biomedical).

Conventional whole-cell patch-clamp studies were conducted using nontransfected HEK 293 cells or cells transiently transfected with TRPC6-YFP or TRPM4-GFP. Transfected cells were identified by fluorescence. Cells were pretreated (20 min) with the TRPC6 inhibitor GsMTx-4 (5 μ M), a peptide derived from the venom of *Grammostola spatulata* (Chilean rose hair tarantula), or an anti-TRPC6 inhibitory antibody [1:250, Alomone (ALC-120)]. Macroscopic whole-cell currents from HEK 293 cells transfected with TRPM4-GFP were recorded in external bathing solution containing 146 mM NaCl, 5 mM CaCl₂, 10 mM Hepes, and 10 mM glucose (pH 7.4). The bathing solution was supplemented with the potassium channel blocker TEA (10 mM). The pipette solution contained 146 mM CsCl, 1 mM MgCl₂, 10 mM Hepes, and 0.1 mM CaCl₂. Macroscopic whole-cell currents from HEK 293 cells transfected with TRPC6-YFP were recorded in external bathing solution containing 150 mM NaCl, 0.08 mM KCl, 0.8 mM MgCl₂, 5.4 mM CaCl₂, 10 mM Hepes, and 10 mM glucose (pH 7.4). The pipette solution contained 130 mM CsOH, 110 mM aspartic acid, 1 mM MgCl₂, 10 mM EGTA, and 10 mM Hepes. After baseline currents were recorded, carbachol (10 μ M) was administered to induce receptor-mediated TRPC6 currents. For all experiments, cells were initially voltage-clamped at a membrane potential of 0 mV. Currents were measured during voltage ramps between -100 and +100 mV (1.6 s) repeated every 4 s.

Animals

Male Sprague-Dawley rats (250 to 350 g; Harlan) were used for these studies in accordance with protocols approved by the Institutional Animal Care and Use Committees of the Colorado State University. Animals were deeply anesthetized with pentobarbital sodium (50 mg, intraperitoneally) and euthanized by exsanguination. Cerebral and cerebellar arteries were dissected from brains and cleaned of connective tissue in cold Mops-buffered saline composed of 3 mM Mops (pH 7.4), 145 mM NaCl, 5 mM KCl, 1 mM MgSO₄, 2.5 mM CaCl₂, 1 mM KH₂PO₄, 0.02 mM EDTA, 2 mM pyruvate, 5 mM glucose, and 1% bovine serum albumin. Arteries were stored in Mops-buffered saline on ice before further manipulation.

Cerebral artery smooth muscle cell isolation

Vessels were washed in magnesium-based physiological saline solution (Mg-PSS) containing 5 mM KCl, 140 mM NaCl, 2 mM MgCl₂, 10 mM Hepes, and 10 mM glucose. Arteries were initially digested in papain (0.6 mg/ml) (Worthington) and dithioerythritol (1 mg/ml) in Mg-PSS at 37°C for 15 min, followed by a 15-min incubation at 37°C in type II collagenase (1.0 mg/ml) (Worthington) in Mg-PSS. The digested arteries were washed three times in ice-cold Mg-PSS solution and incubated on ice for 30 min. After this incubation period, vessels were triturated to liberate smooth muscle cells and stored in ice-cold Mg-PSS before use. Smooth muscle cells were studied within 6 hours after isolation.

Electrophysiological recordings from isolated smooth muscle cells

Isolated smooth muscle cells were placed in a recording chamber (Warner Instruments) and allowed to adhere to glass coverslips for 20 min at room temperature. Gigohm seals were obtained in Mg-PSS. Amphotericin B (40 μ M) was included in the pipette solution to perforate the membrane. Perforation was deemed acceptable if series resistance was less than 50 megohms. STOC and TICC activities were recorded in normal external bathing solution containing 134 mM NaCl, 6 mM KCl, 1 mM MgCl₂, 2 mM CaCl₂, 10 mM Hepes, and 10 mM glucose at pH 7.4 (NaOH). The pipette solution contained 110 mM K-aspartate, 1 mM MgCl₂, 30 mM KCl, 10 mM NaCl, 10 mM Hepes, and 5 μ M EGTA at pH 7.2 (NaOH). Additional external solutions include Ca²⁺-free external solution containing 134 mM NaCl, 6 mM KCl, 1 mM MgCl₂, 0.73 mM CaCl₂, 1 mM EGTA, 10 mM Hepes, and 10 mM glucose at pH 7.4 (NaOH). Where specified, a hyperosmotic solution was reached using 100 mM mannitol. A hypotonic challenge of 80 mosmol/liter was achieved by incubating and recording baseline activity in the above hypertonic solution (376 mosmol/liter) followed by exposure to the original recording solution (without mannitol, 296 mosmol/liter). In our recording solutions, the calculated reversal potentials for total monovalent cations were -1.8 and -30.6 mV for monovalent anions (Cl⁻). STOCs were defined as transient events greater than 10 pA (more than one BK_{Ca} channel), and the frequency was calculated by dividing the number of events by the time between the first and the last event. Defined amounts of stretch of the plasma membrane were achieved by applying negative pressure through the recording electrode using a DPM-1B Pneumatic Transducer Tester (Fluke Biomedical). Stretch-induced TICC activity was recorded after the administration of 9-phenanthrol (30 μ M), U73122 (10 μ M), Bt-IP₃ (10 μ M), Ang II (100 nM), losartan (1 μ M), PP2 (300 nM), GsMTx-4 (5 μ M), U73343 (10 μ M), or Src family kinase inhibitor-1 (SKI-1, 300 nM). TICC activity at -70 mV was quantified as the calculated sum of the open channel probability of multiple open states of 1.75 pA. This value was based on the reported unitary conductance of TRPM4 (25 pS). Channel open probability was calculated using the following equation:

$$NP_o = \sum_{j=1}^N \left(\frac{t_j \cdot j}{T} \right)$$

where t_j is the time spent in seconds, with $j = 1, 2, \dots, N$ channels open; N is the max number of channels observed; and T is the duration of measurement.

Isolated vessel experiments

Arteries were harvested and transferred to a vessel chamber (Living Systems Inc.). The proximal end of the vessel was cannulated with a glass micropipette and secured with monofilament thread. Blood was gently rinsed from the lumen, and the distal end of the vessel was cannulated and secured. Arteries were pressurized to 10 mmHg with PSS composed of 119 mM NaCl, 4.7 mM KCl, 1.8 mM CaCl₂, 1.2 mM MgSO₄, 24 mM NaHCO₃, 0.2 mM KH₂PO₄, 10.6 mM glucose, and 1.1 mM EDTA, and superfused (5 ml/min) with warmed (37°C) PSS aerated with a normoxic gas mixture (21% O₂, 6% CO₂, balance N₂). After a 15-min equilibration period, intraluminal pressure was slowly increased to 80 mmHg, vessels were stretched to remove bends, and pressure was reduced to 20

mmHg for an additional 15-min equilibration period. Inner diameter was continuously monitored using video microscopy and edge detection software (IonOptix). Arteries pressurized to 20 mmHg were exposed to isotonic PSS containing 60 mM KCl to assess viability of the preparation. Arteries were pressurized to 80 mmHg, and stable myogenic tone was allowed to develop before the administration of U73122 (10 μ M). To determine maximum (or passive) diameter, vessels were later superfused with Ca^{2+} -free PSS: 119 mM NaCl, 4.7 mM KCl, 1.2 mM MgSO_4 , 24 mM NaHCO_3 , 0.2 mM KH_2PO_4 , 10.6 mM glucose, 1.1 mM EDTA, 3 mM EGTA, and 0.01 mM diltiazem.

Ca^{2+} imaging experiments of isolated vessels

Arteries were harvested and incubated in the dark at room temperature in Mops-buffered saline containing 10 μ M fluo-4 AM (Invitrogen) and pluronic acid (2.5 μ g/ml) for 1.5 hours. Vessels were cannulated on a glass micropipette, pressurized, and superfused with warmed (37°C) modified PSS (125 mM NaCl, 3 mM KCl, 26 mM NaHCO_3 , 1.25 mM NaH_2PO_4 , 1 mM MgCl_2 , 4 mM D-glucose, and 2 mM CaCl_2), and aerated with a gas mixture consisting of 20% O_2 , 5% CO_2 , 75% N_2 to maintain pH (pH 7.4). IP_3R -mediated Ca^{2+} waves were isolated by continuously exposing vessels to the L-type Ca^{2+} channel blocker nimodipine (300 nM) and the ryanodine receptor blocker tetracaine (100 μ M). Images were acquired at 20 frames per second on a 138×138 - μ m field of view (512×512 pixels) using a Solamere confocal scanning unit (QLC 100), an electron-multiplying CCD (charge-coupled device) camera, and a 40 $\times$ water immersion objective. The fluorophore was excited by illumination with a 488-nm krypton/argon laser, and fluorescence emission was collected above 495 nm. Ca^{2+} images were acquired using Andor Revolution TL software (Andor Technology) and analyzed with custom software (SparkAn) created by A. D. Bonev (University of Vermont, Burlington, VT).

Reverse transcription polymerase chain reaction

After enzymatic dispersal of rat cerebral arteries, total RNA was extracted (RNeasy Protect Mini Kit, Qiagen Inc.) and first-strand complementary DNA (cDNA) was synthesized using an Omniscript Reverse Transcription Kit (Qiagen Inc.). RT-PCR was performed with the following PLC-specific QuantiTect Primers (Qiagen Inc.) spanning intron/exon boundaries for the following: PLC γ 1, QT00184100 (Rn_Plcg1_1_SG); PLC γ 2, QT01080338 (Rn_Plcg2_1_SG); and PLC β 2, QT00188062 (Rn_Plcb2_1_SG). PCR products were resolved on 2% agarose gels. These primers yielded PCR products of 84 base pairs (bp) (PLC γ 1), 108 bp (PLC γ 2), and 102 bp (PLC β 2). All PCRs included a template-free control. RT-PCR experiments were repeated using RNA extracted from at least three different animals.

Membrane staining and Mag-fluo-4 Ca^{2+} imaging

Cells were enzymatically dissociated from cerebral vessels as described above, and allowed to adhere to glass slides for 20 min at 4°C. Membrane-specific fluorescent staining was performed with ER-Tracker Red (5 μ g/ml) (Invitrogen), and SR Ca^{2+} stores were assessed using the low-affinity ($K_d = 22$ μ M) Ca^{2+} indicator Mag-fluo-4 AM (5 μ M, Invitrogen). Cells were incubated in ER-Tracker Red and Mag-fluo-4 AM in the patch-clamp recording

solution for 30 min at 37°C. Fluorescent images were obtained using a spinning disk confocal microscope (Andor). ER-Tracker Red and Mag-fluo-4 fluorescence were excited by illumination with 543- and 488-nm line, respectively. All images were acquired at 1024 × 1024 pixels and were analyzed using ImageJ [National Institutes of Health (NIH)].

Immunocytochemistry

Cells were enzymatically dissociated from freshly isolated and siRNA-treated vessels as described above. Cells were allowed to adhere to glass slides for 20 min at 4°C, then were fixed with 4% formaldehyde for 10 min, permeabilized with cold methanol (−80°C), and blocked with 2% bovine serum albumin. Cells were incubated in 2% bovine serum albumin blocking solution containing polyclonal primary antibodies overnight at 4°C. Cells were washed and incubated in appropriate fluorescent secondary antibodies for 2 hours at room temperature. Fluorescent images were obtained using a spinning disk confocal microscope (Andor) and a 100× oil immersion objective. Texas Red and fluorescein isothiocyanate (FITC) were excited by illumination with 543- and 488-nm line, respectively. All images were acquired at 1024 × 1024 pixels and were analyzed using a Volocity imaging software package (version 6.0, PerkinElmer Inc.) and ImageJ version 1.42q (NIH). Total fluorescence was determined using the mean fluorescence of the ROI encompassing the whole cell.

Proximity ligation assay

Colocalization of proteins was studied in freshly isolated cerebral artery myocytes using an in situ proximity ligation assay detection kit (Duolink, Olink Biosciences Inc.). Cells were enzymatically dissociated as described. Glass slides were pretreated with a blocking solution containing 1% bovine serum albumin and 1% fish gelatin in Mg-PSS for 10 min at room temperature. Cells were allowed to adhere to glass slides for 20 min at 4°C and fixed with 4% formaldehyde for 10 min at room temperature and 2 hours at 4°C. After subsequent washes in Mg-PSS, cells were permeabilized with cold methanol (−80°C) and incubated overnight in blocking solution containing paired primary antibodies. After incubation in primary antibodies, cells were washed in blocking solution followed by three 10-min washes in 10 ml of Duolink In Situ Wash Buffer A containing 10 mM tris, 150 mM NaCl, and 0.05% Tween 20 (pH 7.4). Cells were incubated in a humidified chamber at 37°C for 60 min with secondary anti-rabbit PLUS and anti-goat MINUS proximity ligation assay probes and later washed three times for 5 min with 10 ml of Wash Buffer A at room temperature. Cells were incubated in ligation ligase solution for 30 min at 37°C in a humidified chamber and later washed three times for 2 min with 10 ml of Wash Buffer A at room temperature. Last, cells were incubated in amplification polymerase solution for 100 min at 37°C in a humidified chamber and later washed twice for 2 min with 10 ml of Duolink In situ Wash Buffer B containing 200 mM tris and 100 mM NaCl (pH 7.5). Cells were further washed with a 1% Wash Buffer B for 1 min and mounted using Duolink In situ Mounting Medium containing DAPI nuclear stain. Fluorescent images were obtained using a spinning disk confocal microscope (Andor) and a 100× oil immersion objective. Generation of positive signal (bright puncta) occurs only when the two proximity ligation assay probes are in close proximity (<40 nm). Excitation of fluorescent puncta was achieved with a 543-nm laser, and autofluorescence of the cytosol was illuminated using a 488-nm laser. Images were analyzed in the Volocity imaging software package (version 6.0, PerkinElmer Inc.). To demonstrate

the specificity of the primary antibodies, we performed proximity ligation assay experiments using two antibodies (goat and rabbit) targeting different epitopes of the same protein. Under these conditions, positive proximity ligation assay puncta were observed as the two primary antibodies were bound in close proximity. The density of positive puncta per cell was determined using an automated object-finding protocol in the Velocity imaging software package.

Antibodies

The following primary antibodies were used: anti-PLC γ 1 (Santa Cruz, sc-426) for immunocytochemistry and proximity ligation assay experiments; anti-PLC γ 2 (Santa Cruz, sc-31751) for immunocytochemistry and proximity ligation assay experiments; anti-PLC β 2 (Santa Cruz, sc-206) for immunocytochemistry and proximity ligation assay experiments; anti-TRPC6 (Alomone, ACC-120) for immunocytochemistry, electrophysiology, and Ca²⁺ imaging experiments; anti-TRPC6 (Santa Cruz, sc-19197) for proximity ligation assay experiments; anti-TRPM4 (Abcam, ab63080) for immunocytochemistry and proximity ligation assay experiments; anti-TRPM4 (Santa Cruz, sc-67125) for immunocytochemistry and proximity ligation assay experiments; and anti-BK_{Ca} (Alomone, APC-021) for proximity ligation assay experiments. Secondary antibodies used were the following: sheep anti-rabbit Texas Red (Abcam, ab6793-1), bovine anti-rabbit FITC (Santa Cruz, sc-2365), donkey anti-goat Texas Red (Santa Cruz, sc-2783), and donkey anti-goat FITC (Abcam, ab6881).

RNA interference and reverse permeabilization

siRNAs directed against PLC γ 1 were used to decrease protein abundance in isolated cerebral arteries. siRNAs purchased from Qiagen [1027280 (AllStars Negative Control) and S101961939 (Rn_Plcg1_1)] were dissolved at a concentration of 20 μ M in siRNA suspension buffer. Control or PLC γ 1 siRNAs were introduced into intact cerebral arteries using a reversible permeabilization procedure. Arteries were permeabilized by first incubating segments for 20 min at 4°C in the following solution: 120 mM KCl, 2 mM MgCl₂, 10 mM EGTA, 5 mM Na₂ ATP, and 20 mM TES (pH 6.8). Arteries were then placed in a similar solution containing siRNA (40 nM) for 3 hours at 4°C and then transferred to a third siRNA-containing solution with increased MgCl₂ (10 mM) for 30 min at 4°C. Permeabilization was reversed by placing arteries in a Mops-buffered physiological siRNA-containing solution consisting of 140 mM NaCl, 5 mM KCl, 10 mM MgCl₂, 5 mM glucose, and 2 mM Mops (pH 7.1, 22°C) for 30 min at room temperature. Ca²⁺ was gradually increased in the latter solution from nominally Ca²⁺-free to 0.01, 0.1, and 1.8 mM over a 45-min period. After the reversible permeabilization procedures, arteries were cultured for 2 to 3 days in DMEM/F-12 culture medium supplemented with L-glutamine (2 mM) (Gibco) and 0.5% penicillin-streptomycin (Gibco).

Real-time RT-PCR

Arteries treated with control or PLC γ 1 siRNA were enzymatically dissociated, and RNA was immediately isolated, purified, and synthesized into cDNA as described above. PLC γ 1 mRNA abundance was measured using a real-time SYBR Green detection assay (Bio-Rad), QuantiTect (Qiagen Inc.) primers for PLC γ 1 [QT00184100 (Rn_Plcg1_1_SG)], and iQ5

Multicolor Real-Time PCR Detection System (Bio-Rad). Samples were normalized to β -actin [QT00193473 (NM_031144)], and cycling parameters were selected on the basis of the protocol for QuantiTect primer assays (Qiagen Inc.). The decrease in PLC γ 1 mRNA abundance was calculated according to the Pfaffl method (59).

Smooth muscle cell membrane potential

To measure smooth muscle cell membrane potential, cerebral arteries were isolated and pressurized to either 20 or 80 mmHg, and smooth muscle cells were impaled through the adventitia with glass intracellular microelectrodes (tip resistance 100 to 200 megohm). AWPI Intra 767 amplifier was used to record membrane potential (E_m). Analog output from the amplifier was recorded using IonWizard software (sample frequency 20 Hz). Criteria for acceptance of E_m recordings were (i) an abrupt negative deflection of potential as the microelectrode was advanced into a cell, (ii) stable membrane potential for at least 1 min, and (iii) an abrupt change in potential to about 0 mV after the electrode was retracted from the cell.

Calculations and statistics

All data are means $\pm$ SE. Values of n refer to the number of arteries for the isolated vessel experiments or the number of cells for immunocytochemistry, proximity ligation assay, and patch-clamp experiments. Unpaired Student's t tests were performed to detect differences in total PLC γ 1-dependent fluorescence from cells isolated from control compared to PLC γ 1 siRNA-treated vessels and to demonstrate differences in TRPM4: TRPC6 compared to TRPM4:BK Ca proximity ligation assay labeling. In addition, unpaired Student's t tests were performed to detect differences in TICC total open probability from cells isolated from control compared to PLC γ 1 siRNA-treated vessels and control compared to PLC γ 1 siRNA-treated vessels treated with Bt-IP $_3$. Paired Student's t tests were performed to determine differences in single-channel and macroscopic TRPC6-YFP currents in response to intrapipette pressure, GsMTx-4, and anti-TRPC6 inhibitory peptide. In addition, paired Student's t tests were performed to determine differences in TICC total open probability in response to Bt-IP $_3$ in cells isolated from PLC γ 1 siRNA-treated vessels. One-way analysis of variance (ANOVA) was performed to demonstrate differences in proximity ligation assay labeling for TRPM4:PLC γ 1, TRPM4:PLC γ 2, and TRPM4: PLC β 2, and the percentage of cells with Ca $^{2+}$ waves in Ca $^{2+}$ imaging experiments in pressurized vessels. One-way repeated-measures ANOVAs were performed to demonstrate differences in TICC total open probability in response to intrapipette pressures and after administration of U73122, Bt-IP $_3$, and CPA. Two-way repeated-measures ANOVAs were performed to demonstrate differences between TICC total open probability at different intrapipette pressures and in the presence or absence of 9-phenanthrol or extracellular Ca $^{2+}$; myogenic tone at different intraluminal pressures in the presence or absence of U73122; myogenic tone and membrane potential at different intraluminal pressures from control versus PLC γ 1 siRNA-treated vessels; and arterial tone at different concentrations of UTP from vessels treated with U73122. Individual groups were compared using a Student-Newman-Keuls post hoc test. P 0.05 was accepted as statistically significant for all experiments.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank M. M. Tamkun and the members of his laboratory for generating the TRPC6-YFP expression plasmid used for these studies; Z. Garcia and A. Simpson for technical assistance; and A. Howard for statistical assistance.

Funding: This work was supported by a Monfort Excellence Award from the Monfort Family Foundation (to S.E.), Fondation Leducq for the Transatlantic Network of Excellence on the Pathogenesis of Small Vessel Disease of the Brain (to M.T.N.), Totman Medical Research Trust (to M.T.N.), and the NIH [grants R01HL091905 (to S.E.), HL044455 (to M.T.N.), R37 DK053832 (to M.T.N.), 1P01HL095488 (to M.T.N.), T32 HL007594 (to A.L.G.), and F31HL094145 (to A.L.G.)].

REFERENCES AND NOTES

1. Davis MJ, Hill MA. Signaling mechanisms underlying the vascular myogenic response. *Physiol Rev.* 1999; 79:387–423. [PubMed: 10221985]
2. Bayliss WM. On the local reactions of the arterial wall to changes of internal pressure. *J Physiol.* 1902; 28:220–231. [PubMed: 16992618]
3. Björnberg J, Grände PO, Maspers M, Mellander S. Site of autoregulatory reactions in the vascular bed of cat skeletal muscle as determined with a new technique for segmental vascular resistance recordings. *Acta Physiol Scand.* 1988; 133:199–210. [PubMed: 3227915]
4. Järhult J, Mellander S. Autoregulation of capillary hydrostatic pressure in skeletal muscle during regional arterial hypo- and hypertension. *Acta Physiol Scand.* 1974; 91:32–41. [PubMed: 4835709]
5. Harder DR. Pressure-dependent membrane depolarization in cat middle cerebral artery. *Circ Res.* 1984; 55:197–202. [PubMed: 6744529]
6. Knot HJ, Nelson MT. Regulation of arterial diameter and wall $[Ca^{2+}]$ in cerebral arteries of rat by membrane potential and intravascular pressure. *J Physiol.* 1998; 508:199–209. [PubMed: 9490839]
7. VanBavel E, Wesselman JP, Spaan JA. Myogenic activation and calcium sensitivity of cannulated rat mesenteric small arteries. *Circ Res.* 1998; 82:210–220. [PubMed: 9468192]
8. Moreno-Domínguez A, Colinas O, El-Yazbi A, Walsh EJ, Hill MA, Walsh MP, Cole WC. Ca^{2+} sensitization due to myosin light chain phosphatase inhibition and cytoskeletal reorganization in the myogenic response of skeletal muscle resistance arteries. *J Physiol.* 2013; 591:1235–1250. [PubMed: 23230233]
9. Gokina NI, Park KM, McElroy-Yaggy K, Osol G. Effects of Rho kinase inhibition on cerebral artery myogenic tone and reactivity. *J Appl Physiol.* 2005; 98:1940–1948. [PubMed: 15626753]
10. Davis MJ, Donovitz JA, Hood JD. Stretch-activated single-channel and whole cell currents in vascular smooth muscle cells. *Am J Physiol Cell Physiol.* 1992; 262:C1083–C1088.
11. Sharif-Naeini R, Folgering JH, Bichet D, Duprat F, Lauritzen I, Arhatte M, Jodar M, Dedman A, Chatelain FC, Schulte U, Retailleau K, Loufrani L, Patel A, Sachs F, Delmas P, Peters DJ, Honoré E. Polycystin-1 and -2 dosage regulates pressure sensing. *Cell.* 2009; 139:587–596. [PubMed: 19879844]
12. Muraki K, Iwata Y, Katanosaka Y, Ito T, Ohya S, Shigekawa M, Imaizumi Y. TRPV2 is a component of osmotically sensitive cation channels in murine aortic myocytes. *Circ Res.* 2003; 93:829–838. [PubMed: 14512441]
13. Welsh DG, Morielli AD, Nelson MT, Brayden JE. Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circ Res.* 2002; 90:248–250. [PubMed: 11861411]
14. Earley S, Waldron BJ, Brayden JE. Critical role for transient receptor potential channel TRPM4 in myogenic constriction of cerebral arteries. *Circ Res.* 2004; 95:922–929. [PubMed: 15472118]
15. Gonzales AL, Garcia ZI, Amberg GC, Earley S. Pharmacological inhibition of TRPM4 hyperpolarizes vascular smooth muscle. *Am J Physiol Cell Physiol.* 2010; 299:C1195–C1202. [PubMed: 20826763]

16. Bulley S, Neeb ZP, Burris SK, Bannister JP, Thomas-Gatewood CM, Jangsangthong W, Jaggar JH. TMEM16A/ANO1 channels contribute to the myogenic response in cerebral arteries. *Circ Res.* 2012; 111:1027–1036. [PubMed: 22872152]
17. Thomas-Gatewood C, Neeb ZP, Bulley S, Adebisi A, Bannister JP, Leo MD, Jaggar JH. TMEM16A channels generate Ca^{2+} -activated Cl^- currents in cerebral artery smooth muscle cells. *Am J Physiol Heart Circ Physiol.* 2011; 301:H1819–H1827. [PubMed: 21856902]
18. Reading SA, Brayden JE. Central role of TRPM4 channels in cerebral blood flow regulation. *Stroke.* 2007; 38:2322–2328. [PubMed: 17585083]
19. Hofmann T, Obukhov AG, Schaefer M, Harteneck C, Gudermann T, Schultz G. Direct activation of human TRPC6 and TRPC3 channels by diacylglycerol. *Nature.* 1999; 397:259–263. [PubMed: 9930701]
20. Spassova MA, Hewavitharana T, Xu W, Soboloff J, Gill DL. A common mechanism underlies stretch activation and receptor activation of TRPC6 channels. *Proc Natl Acad Sci USA.* 2006; 103:16586–16591. [PubMed: 17056714]
21. Mederos y Schnitzler M, Storch U, Meibers S, Nurwakagari P, Breit A, Essin K, Gollasch M, Gudermann T. G_q -coupled receptors as mechanosensors mediating myogenic vasoconstriction. *EMBO J.* 2008; 27:3092–3103. [PubMed: 18987636]
22. Nilius B, Prenen J, Droogmans G, Voets T, Vennekens R, Freichel M, Wissenbach U, Flockerzi V. Voltage dependence of the Ca^{2+} -activated cation channel TRPM4. *J Biol Chem.* 2003; 278:30813–30820. [PubMed: 12799367]
23. Launay P, Fleig A, Perraud AL, Scharenberg AM, Penner R, Kinet JP. TRPM4 is a Ca^{2+} -activated nonselective cation channel mediating cell membrane depolarization. *Cell.* 2002; 109:397–407. [PubMed: 12015988]
24. Gonzales AL, Earley S. Endogenous cytosolic Ca^{2+} buffering is necessary for TRPM4 activity in cerebral artery smooth muscle cells. *Cell Calcium.* 2012; 51:82–93. [PubMed: 22153976]
25. Gonzales AL, Amberg GC, Earley S. Ca^{2+} release from the sarcoplasmic reticulum is required for sustained TRPM4 activity in cerebral artery smooth muscle cells. *Am J Physiol Cell Physiol.* 2010; 299:C279–C288. [PubMed: 20427713]
26. Morita H, Honda A, Inoue R, Ito Y, Abe K, Nelson MT, Brayden JE. Membrane stretch-induced activation of a TRPM4-like nonselective cation channel in cerebral artery myocytes. *J Pharmacol Sci.* 2007; 103:417–426. [PubMed: 17420615]
27. Cipolla MJ, Osol G. Vascular smooth muscle actin cytoskeleton in cerebral artery forced dilatation. *Stroke.* 1998; 29:1223–1228. [PubMed: 9626298]
28. Cipolla MJ, Gokina NI, Osol G. Pressure-induced actin polymerization in vascular smooth muscle as a mechanism underlying myogenic behavior. *FASEB J.* 2002; 16:72–76. [PubMed: 11772938]
29. Martinez-Lemus LA, Crow T, Davis MJ, Meininger GA. $\alpha_v\beta_3$ - and $\alpha_5\beta_1$ -integrin blockade inhibits myogenic constriction of skeletal muscle resistance arterioles. *Am J Physiol Heart Circ Physiol.* 2005; 289:H322–H329. [PubMed: 15722407]
30. Crnich R, Amberg GC, Leo MD, Gonzales AL, Tamkun MM, Jaggar JH, Earley S. Vasoconstriction resulting from dynamic membrane trafficking of TRPM4 in vascular smooth muscle cells. *Am J Physiol Cell Physiol.* 2010; 299:C682–C694. [PubMed: 20610768]
31. Simard C, Sallé L, Rouet R, Guinamard R. Transient receptor potential melastatin 4 inhibitor 9-phenanthrol abolishes arrhythmias induced by hypoxia and re-oxygenation in mouse ventricle. *Br J Pharmacol.* 2012; 165:2354–2364. [PubMed: 22014185]
32. Grand T, Demion M, Norez C, Mettey Y, Launay P, Becq F, Bois P, Guinamard R. 9-Phenanthrol inhibits human TRPM4 but not TRPM5 cationic channels. *Br J Pharmacol.* 2008; 153:1697–1705. [PubMed: 18297105]
33. Nelson MT, Cheng H, Rubart M, Santana LF, Bonev AD, Knot HJ, Lederer WJ. Relaxation of arterial smooth muscle by calcium sparks. *Science.* 1995; 270:633–637. [PubMed: 7570021]
34. Osol G, Laher I, Kelley M. Myogenic tone is coupled to phospholipase C and G protein activation in small cerebral arteries. *Am J Physiol.* 1993; 265:H415–H420. [PubMed: 8342661]
35. Rebecchi MJ, Pentyla SN. Structure, function, and control of phosphoinositide-specific phospholipase C. *Physiol Rev.* 2000; 80:1291–1335. [PubMed: 11015615]

36. Gustafsdottir SM, Schallmeiner E, Fredriksson S, Gullberg M, Söderberg O, Jarvius M, Jarvius J, Howell M, Landegren U. Proximity ligation assays for sensitive and specific protein analyses. *Anal Biochem.* 2005; 345:2–9. [PubMed: 15950911]
37. Reading SA, Earley S, Waldron BJ, Welsh DG, Brayden JE. TRPC3 mediates pyrimidine receptor-induced depolarization of cerebral arteries. *Am J Physiol Heart Circ Physiol.* 2005; 288:H2055–H2061. [PubMed: 15604128]
38. Rainbow RD, Macmillan D, McCarron JG. The sarcoplasmic reticulum Ca^{2+} store arrangement in vascular smooth muscle. *Cell Calcium.* 2009; 46:313–322. [PubMed: 19836074]
39. Na S, Collin O, Chowdhury F, Tay B, Ouyang M, Wang Y, Wang N. Rapid signal transduction in living cells is a unique feature of mechanotransduction. *Proc Natl Acad Sci USA.* 2008; 105:6626–6631. [PubMed: 18456839]
40. Haendeler J, Yin G, Hojo Y, Saito Y, Melaragno M, Yan C, Sharma VK, Heller M, Aebbersold R, Berk BC. GIT1 mediates Src-dependent activation of phospholipase $\text{C}\gamma$ by angiotensin II and epidermal growth factor. *J Biol Chem.* 2003; 278:49936–49944. [PubMed: 14523024]
41. Jarajapu YP, Knot HJ. Role of phospholipase C in development of myogenic tone in rat posterior cerebral arteries. *Am J Physiol Heart Circ Physiol.* 2002; 283:H2234–H2238. [PubMed: 12388230]
42. Jarajapu YP, Guberski DL, Grant MB, Knot HJ. Myogenic tone and reactivity of cerebral arteries in type II diabetic BBZDR/Wor rat. *Eur J Pharmacol.* 2008; 579:298–307. [PubMed: 18036520]
43. Earley S, Straub SV, Brayden JE. Protein kinase C regulates vascular myogenic tone through activation of TRPM4. *Am J Physiol Heart Circ Physiol.* 2007; 292:H2613–H2622. [PubMed: 17293488]
44. Osol G, Laher I, Cipolla M. Protein kinase C modulates basal myogenic tone in resistance arteries from the cerebral circulation. *Circ Res.* 1991; 68:359–367. [PubMed: 1991343]
45. Foskett JK, White C, Cheung KH, Mak DO. Inositol trisphosphate receptor Ca^{2+} release channels. *Physiol Rev.* 2007; 87:593–658. [PubMed: 17429043]
46. Narayanan J, Imig M, Roman RJ, Harder DR. Pressurization of isolated renal arteries increases inositol trisphosphate and diacylglycerol. *Am J Physiol.* 1994; 266:H1840–H1845. [PubMed: 8203583]
47. Mufti RE, Brett SE, Tran CH, Abd El-Rahman R, Anfinogenova Y, El-Yazbi A, Cole WC, Jones PP, Chen SR, Welsh DG. Intravascular pressure augments cerebral arterial constriction by inducing voltage-insensitive Ca^{2+} waves. *J Physiol.* 2010; 588:3983–4005. [PubMed: 20736418]
48. Zou Y, Akazawa H, Qin Y, Sano M, Takano H, Minamino T, Makita N, Iwanaga K, Zhu W, Kudoh S, Toko H, Tamura K, Kihara M, Nagai T, Fukamizu A, Umemura S, Iiri T, Fujita T, Komuro I. Mechanical stress activates angiotensin II type 1 receptor without the involvement of angiotensin II. *Nat Cell Biol.* 2004; 6:499–506. [PubMed: 15146194]
49. Storch U, Mederos y Schnitzler M, Gudermann T. G protein-mediated stretch reception. *Am J Physiol Heart Circ Physiol.* 2012; 302:H1241–H1249. [PubMed: 22227128]
50. Mederos y Schnitzler M, Storch U, Gudermann T. AT_1 receptors as mechanosensors. *Curr Opin Pharmacol.* 2011; 11:112–116. [PubMed: 21147033]
51. Brayden JE, Li Y, Tavares MJ. Purinergic receptors regulate myogenic tone in cerebral parenchymal arterioles. *J Cereb Blood Flow Metab.* 2013; 33:293–299. [PubMed: 23168530]
52. Schnapp LM, Goswami S, Rienzi N, Kotliansky VE, Gotwals P, Schachter EN. Integrins inhibit angiotensin II-induced contraction in rat aortic rings. *Regul Pept.* 1998; 77:177–183. [PubMed: 9809813]
53. Jones NP, Peak J, Brader S, Eccles SA, Katan M. $\text{PLC}\gamma 1$ is essential for early events in integrin signalling required for cell motility. *J Cell Sci.* 2005; 118:2695–2706. [PubMed: 15944397]
54. Dietrich A, Mederos M, Schnitzler Y, Gollasch M, Gross V, Storch U, Dubrovskaya G, Obst M, Yildirim E, Salanova B, Kalwa H, Essin K, Pinkenburg O, Luft FC, Gudermann T, Birnbaumer L. Increased vascular smooth muscle contractility in *TRPC6*^{−/−} mice. *Mol Cell Biol.* 2005; 25:6980–6989. [PubMed: 16055711]
55. Kirber MT, Walsh JV Jr, Singer JJ. Stretch-activated ion channels in smooth muscle: A mechanism for the initiation of stretch-induced contraction. *Pflügers Arch.* 1988; 412:339–345. [PubMed: 2459658]

56. Davis MJ, Meininger GA, Zawieja DC. Stretch-induced increases in intracellular calcium of isolated vascular smooth muscle cells. *Am J Physiol.* 1992; 263:H1292–H1299. [PubMed: 1329564]
57. Wu X, Davis MJ. Characterization of stretch-activated cation current in coronary smooth muscle cells. *Am J Physiol Heart Circ Physiol.* 2001; 280:H1751–H1761. [PubMed: 11247789]
58. Park KS, Kim Y, Lee YH, Earm YE, Ho WK. Mechanosensitive cation channels in arterial smooth muscle cells are activated by diacylglycerol and inhibited by phospholipase C inhibitor. *Circ Res.* 2003; 93:557–564. [PubMed: 12946944]
59. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 2001; 29:e45. [PubMed: 11328886]

Fig. 1. TRPM4 is not intrinsically mechanosensitive, and requires PLC and TRPC6 activity for cell swelling-induced activity

(A) Left: Representative on-cell patch-clamp recording of changes in single-channel activity in a HEK 293 cell transfected with TRPC6-YFP as negative pressure (-20 mmHg) was applied to the patch pipette. Middle: Single-channel current-to-voltage (*I/V*) amplitude relationship plotted for currents recorded with and without negative pressure ($n = 4$ to 7 cells in three experiments). Right: Summary of total open probability (NP_o) at the indicated intrapipette pressures at a holding potential of +70 mV ($n = 5$ cells in three transfection experiments; $*P < 0.05$). (B) Left: Representative on-cell patch-clamp recording of changes in single-channel activity in a HEK 293 cell transfected with TRPM4-GFP as negative pressure (-20 mmHg) was applied to the patch pipette. Middle: *I/V* relationship at the indicated intrapipette pressures ($n = 4$ to 5 cells in three transfection experiments). Right: Summary NP_o at the indicated intrapipette pressures at a holding potential of +70 mV ($n = 5$ cells in three transfection experiments; $*P < 0.05$). (C) Representative recording and summary of TICC activity after 80 mosmol/liter osmotic challenge ($n = 5$ cells from three animals; $*P < 0.05$ compared to control, $\#P < 0.05$ compared to hypotonic). (D) Representative recording and summary of the effects of the PLC inhibitor U73122 on swelling-induced TICC activity ($n = 5$ cells from three animals; $*P < 0.05$ compared to control).

Fig. 2. PLCγ1 colocalizes with TRPM4 in cerebral artery smooth muscle cells

(A) PLCγ1, PLCγ2, and PLCβ2 detected by RT-PCR in RNA extracted from whole cerebral arteries. Representative of data obtained from at least three animals. NT, no template control. (B) Immunofluorescent images of labeling for PLCγ1, PLCγ2, or PLCβ2 (green) and TRPM4 (red) in freshly isolated cerebral artery smooth muscle cells. Scale bar, 10 μm. Representative of 15 cells from three animals. (C) Immunofluorescent images and summary of proximity ligation assay (PLA) labeling for PLCγ1:TRPM4 (top), PLCγ2:TRPM4 (middle), and PLCβ2:TRPM4 (bottom). Autofluorescence of cytosol is green, and the occurrences of protein colocalization are visualized as bright fluorescent puncta (red, insert). Representative of data obtained from at least three animals ($n = 15$ cells from three animals; $*P = 0.05$ compared to PLCγ1:TRPM4). Scale bars, 10 and 1 μm (insert). (D) Representative recordings and summary of TICC activity in cells isolated from arteries treated with control (top) and PLCγ1 siRNA (bottom). Bt-IP₃ was administered to cells isolated from arteries treated with PLCγ1 siRNA as indicated ($n = 6$ to 11 cells from five animals; $*P = 0.05$ compared to control siRNA, $^{\#}P = 0.05$ compared to PLCγ1 siRNA baseline).

Fig. 3. PLC γ 1 siRNA attenuates pressure-induced membrane depolarization and vasoconstriction

(A) Representative recordings of smooth muscle membrane potential in cerebral arteries treated with either control (top) or PLC γ 1 (bottom) siRNA at the indicated intraluminal pressures. (B) Summary data demonstrating the effect of PLC γ 1 siRNA on cerebral arterial myocyte resting membrane potential (E_m) ($n = 5$ arteries from four animals; $*P = 0.05$ compared to control siRNA at 20 mmHg, $\#P = 0.05$ compared to control siRNA at 80 mmHg). (C) Recordings demonstrating the effects of increasing intraluminal pressure on luminal diameter from arteries treated with either control (top) or PLC γ 1 (bottom) siRNA. (D) Summary data showing the effects of PLC γ 1 siRNA treatment on myogenic tone ($n = 6$

to 7 arteries from four animals; $*P = 0.05$). (E) Summary data demonstrating the effect of increasing concentrations of UTP on luminal diameter from arteries treated with either control or PLC γ 1 siRNA ($n = 5$ arteries from four animals). (F) Summary data of the vasoconstriction in response to increased extracellular [KCl] from arteries treated with either control or PLC γ 1 siRNA ($n = 7$ arteries from four animals).

Fig. 4. Ca^{2+} influx through TRPC6 is required for stretch-induced activation of TRPM4 in arterial myocytes

(A) Representative images and recordings demonstrating the effects of increasing intraluminal pressure from 30 to 70 mmHg on Ca^{2+} waves in fluo-4 AM-loaded arteries preincubated with TRPC6 inhibitory antibody. Images are composed of compressed image stacks illustrating the number of cells (grayscale) and number of cells with Ca^{2+} waves (color topography). Scale bars, 5 μm . Spontaneous Ca^{2+} signals were detected as fractional fluorescence (F/F_0) from 15 regions of interest (ROIs) marked as colored squares. Summary of the effects of the TRPC6 inhibitory antibody on pressure- or UTP-induced Ca^{2+} waves of intraluminal pressures ($n = 3$ to 4 vessels from three animals; $*P < 0.05$ compared to control at 30 mmHg, $\#P < 0.05$ compared to control at 70 mmHg, and $\dagger P < 0.05$ compared to anti-TRPC6 at 70 mmHg). (B) Representative recordings and summary of stretch-induced TICC activity in the presence (top) and absence (bottom) of extracellular Ca^{2+} ($n = 7$ cells from four animals; $*P < 0.05$ compared to control at 0 mmHg). (C) Representative recordings and summary data of stretch-dependent TICC activity in the presence of the SERCA pump blocker cyclopiazonic acid (CPA) ($n = 5$ cells from three animals; $*P < 0.05$ compared to

control at 0 mmHg). **(D)** Immunofluorescent images and summary of PLA labeling for TRPM4:TRPC6 and TRPM4:BK_{Ca} ($n = 15$ cells from three animals; $*P = 0.05$ compared to TRPM4:TRPC6). Autofluorescence of cytosol is green, and the areas of protein colocalization are visualized as bright fluorescent puncta (red, insert). Scale bars, 10 or 1 μm (insert). **(E)** Representative recordings and summary of stretch-dependent TICC activity from myocytes preincubated with GsMTx-4 (top), inhibitory TRPC6 antibody (middle), or the inhibitory TRPC6 antibody and antigenic peptide (bottom). Immunofluorescent images (right) of TRPC6 labeling in the absence (top) and presence of antigenic peptide (bottom). DAPI (4',6-diamidino-2-phenylindole)-stained cell nuclei are blue. Scale bars, 10 μm ($n = 15$ cells from three animals). Summary of the effects of GsMTx-4, TRPC6 inhibitory antibody, and antigenic peptide on stretch-dependent TICC activity ($n = 5$ cells from three animals; $*P = 0.05$ compared to control at 0 mmHg). **(F)** Representative recording and summary of swelling-induced TICC activity in myocytes preincubated with an inhibitory TRPC6 antibody ($n = 4$ cells from three animals).

Fig. 5. Src family tyrosine kinase activity is necessary for stretch-induced TRPM4 activity in arterial myocytes

(A) Representative recording and summary of the effects of the Src family kinase inhibitor PP2 on stretch-induced TICC activity [$n = 5$ cells from three animals; $*P = 0.05$ compared to control (0 mmHg), $^{\#}P = 0.05$ compared to control at -20 mmHg]. (B) Representative recording of the effects of repeated administration of PP2 on the luminal diameter of a cerebral artery with predeveloped myogenic tone (80 mmHg) ($n = 3$ to 5 arteries from four animals; $*P = 0.05$ compared to control). (C) Summary data indicating the effects of PP2 administration on vasoconstriction in response to increased extracellular [KCl] ($n = 3$ vessels from three animals). (D) Representative recording and summary of the effects of losartan on stretch-induced TICC activity ($n = 5$ cells from four animals; $*P = 0.05$ compared to control at 0 mmHg, $^{\#}P = 0.05$ compared to control at -20 mmHg).

Fig. 6. Proposed mechanism of stretch-dependent activation of TRPM4

Convergence of stretch-sensing pathways in cerebral artery myocytes includes (i) the activation of AT₁R, Src tyrosine kinase, and PLCγ1 in the generation of IP₃, and (ii) Ca²⁺ influx through TRPC6 channels onto IP₃R to activated TRPM4 channels that are responsible for pressure-induced depolarization of the plasma membrane. PM, plasma membrane; V_m, change in membrane potential.