CORRESPONDENCE

Iatrogenic Nerve Injuries Prevalence, Diagnosis and Treatment

by Prof. Dr. med. Gregor Antoniadis, Prof. Dr. med. Thomas Kretschmer,
Dr. med. Maria Teresa Pedro, PD Dr. med. Ralph W. König,
Dr. med. Christian Heinen, Prof. Dr. med. Hans-Peter Richter in issue 16/2014

Early Diagnosis and Treatment

Congratulations to the authors for choosing this important topic (1). As shown in their review, regeneration after nerve reconstruction is by no means guaranteed (1), in particular not in older patients (2). The most common nerve injury – damage to the median nerve during carpal tunnel surgery (1) – often occurs in middle-aged/ older patients where the prognosis for regeneration is not always favorable. Full recovery of motor function is the exception rather than the rule. Therefore, tendon transfers should already be considered at the time of nerve reconstruction (3). Especially proximal nerve damages with long regeneration distances have a poor prognosis, e.g. injury to the radial nerve associated with osteosynthesis of fractures of the humerus. The various tendon transfers to restore motor function are detailed in the AWMF guidelines for the treatment of peripheral nerve injuries, a publication rightfully cited by the authors of the review.

Although ideally a microsurgeon should be called in cases of intraoperative nerve transection, this is often not practical. We believe that most iatrogenic nerve injuries go unnoticed by the surgeon. On the other hand, it is not uncommon that patients present with unsatisfactory nerve repair where a suture was created with unsuitable suture material under inadequate magnification (microscope/ loupes) without microsurgical technique. As the authors point out in their review, secondary reconstruction by a microsurgeon is certainly more advisable than attempting to perform a nerve suture without the necessary infrastructure and/or adequate microsurgical training.

We agree with the authors' conclusion that iatrogenic nerve injuries should be diagnosed early and adequately treated by experienced microsurgeons. In addition, these surgeons should also be capable of performing tendon transfers to spare these patients from having to undergo multiple surgical procedures.

DOI: 10.3238/arztebl.2014.0602a

REFERENCES

1. Antoniadis G, Kretschmer T, Pedro MT, König RW, Heinen CPG, Richter HP: Iatrogenic neurological damage—prevalence, diagnosis and treatment. *Dtsch Arztebl Int* 2014; 111: 273–9.
2. Stang F, Stollwerck P, Prommersberger KJ, van Schoonhoven J: Posterior interosseus nerve vs. medial cutaneous nerve of the forearm: differences in digital nerve reconstruction. *Arch Orthop Trauma Surg* 2013; 133: 875–80.
3. Prommersberger KJ, van Schoonhoven J: Motorische Ersatzoperationen an der oberen Extremität. *Oper Orthop Traumatol* 2013; 25: 320.

Prof. Dr. med. Frank Unglaub

Medizinische Fakultät Mannheim der Universität Heidelberg,
Mannheim
Abteilung Handchirurgie, Vulpus Klinik,
Bad Rappenau

Prof. Dr. med. Peter Hahn

Abteilung Handchirurgie, Vulpus Klinik,
Bad Rappenau

Dr. med. Christian K. Spies

Abteilung Handchirurgie, Vulpus Klinik,
Bad Rappenau
christian.spies@vulpuslinik.de

Further Locations Prone to Iatrogenic Nerve Injuries

According to the review article (1), iatrogenic nerve injuries can be caused by transection, crushing and compression, penetration, and twisting by screws. Among the procedures associated with iatrogenic nerve injury, herniotomy is described as a major abdominal surgical procedure. This view certainly differs from the general perception, as inguinal hernia repair is a procedure which should typically be performed on a day surgery basis, according to the Social Security Code (SGB). On the other hand, the review does not mention other abdominal and vascular surgical procedures at all.

Indisputably, the best evaluated data material is available for the 90 000 thyroid surgeries performed in Germany each year. A recent multicenter study reports an incidence of transient and permanent recurrent laryngeal nerve palsy of 2–4%. Since patients undergoing thyroid surgery are typically seen by an ENT specialist before and after the operation, these data are unquestionably valid. Thus, the incidence of nerve injury during thyroid surgery is certainly much higher than the rate stated in Table 1.

Another intriguing aspect is related to the approximately 35 000 carotid artery surgeries performed in Germany each year. Peripheral nerve injury is one of the risks associated with the surgical treatment of carotid artery stenosis. Most commonly, the hypoglossal nerve is injured. The evaluation of peripheral nerve injuries is especially relevant as there is ongoing controversy whether carotid artery stenosis should be treated surgically or with a stent.

Even though the authors did apparently not attempt surgical repair of such nerve injuries, they should at least have been mentioned them in their discussion.

DOI: 10.3238/arztebl.2014.0602b

REFERENCES

1. Antoniadis G, Kretschmer T, Pedro MT, König RW, Heinen CPG, Richter HP: Iatrogenic neurological damage—prevalence, diagnosis and treatment. *Dtsch Arztebl Int* 2014; 111: 273–9.

Prof. Dr. med. Heiner H. Wenk

Zentrum für Chirurgie – Allgemein-, Gefäß- und Visceralchirurgie
Klinikum Bremen-Nord, Bremen
a.g.chirurgie@klinikum-bremen-nord.de

Early Surgical Intervention Is the Key

At the Department of Plastic, Reconstructive & Aesthetic Surgery, Innsbruck Medical University, a total of 23 patients (16 female, 7 male, mean age: 52.7 years) were treated for iatrogenic femoral nerve injuries between 2000 and 2010. In our patient population the main causes of nerve injury were the same as those reported by Antoniadis et al. (1). Preoperatively, the primary focus was on taking the patient's history and performing a physical examination, including the collection of information about sensory and motor losses. Here, the clinical-interdisciplinary, neurological and neuro-ultrasonographic examination proved to be valuable. Ultrasonography, performed by an experienced radiologist, can determine the exact location of the damage. Nerve regeneration, which may occur with time, can be monitored by means of electromyography and nerve conduction velocity tests, combined with a wandering Tinel's sign.

Piza-Katzer et al. (2) demonstrated the advantages of early surgical revision in cases with persistent loss of nerve function without clinical signs of improvement. Thus, we decided to perform surgical revisions in affected patients already after 3 months if no improvement of the clinical signs and symptoms and no wandering of Tinel's sign was observed. With this approach, the time between nerve injury and surgical revision was reduced from approx. 12 months to 6.4 months between 2008 and 2010. The time between the nerve injury and the referral to our hospital accounted for approx. 3 months of this period.

In summary, we can only agree with the authors of the article. Early surgical intervention can help to shorten these patients' suffering, and clinical regeneration can be achieved much earlier. In addition, the progressive muscle atrophy is stopped, resulting in faster rehabilitation. Once again we would like to point out that the mere suspicion of a nerve injury should trigger the patient's immediate referral to an expert to ensure that the rapid intervention approach is actually practiced.

DOI: 10.3238/arztebl.2014.0603a

REFERENCES

1. Antoniadis G, Kretschmer T, Pedro MT, König RW, Heinen CPG, Richter HP: Iatrogenic neurological damage—prevalence, diagnosis and treatment. *Dtsch Arztebl Int* 2014; 111: 273–9.
2. Piza-Katzer H, Schöller T, Stichelberger M: Iatrogenic femoral nerve lesions. *Handchir Mikrochir plast Chir* 2009; 41: 230–7.

Dr. med. univ. Gabriel Djedovic

PD Dr. med. Ulrich M. Rieger

Klinik für Plastische und Ästhetische,
Wiederherstellungs- und Handchirurgie
Frankfurt am Main
gabriel.djedovic@me.com

Dr. med. Eva-Maria Baur

Universitätsklinik für Plastische,
Rekonstruktive und Ästhetische Chirurgie,
Medizinische Universität Innsbruck

Tendon Transfers

In their review, the authors highlight a major and serious complication of surgical procedures. They clearly show that with proper diagnosis and treatment the problems experienced by affected patients can be significantly reduced (1). Regrettably, the article does not mention tendon transfers as an important option for (early) restoration of motor function.

In many cases, satisfactory reinnervation after injury and nerve reconstruction cannot be achieved, especially in older patients, as the ability of peripheral nerves to regenerate significantly declines with advancing age (2, 3). In this situation, functional reconstruction by means of tendon transfer significantly improves the outcome for the affected patients. This surgical technique functionally replace muscles which have lost their motor function with intact muscles or tendons. By transferring these muscles, the lost motor function can often be partially or completely restored.

Good examples are the Bridle procedure to treat foot drop after peroneal injury and tendon transfer surgery to restore finger and wrist extension following radial nerve palsy (4). Here—as the authors point out—an intensive physiotherapy program is essential to prevent contractures which may make it impossible to use tendon transfers. At times, an approach combining nerve reconstruction with tendon transfer may be useful.

These important options for the treatment of traumatic or iatrogenic injury to peripheral nerves of the extremities should always be taken into account when deciding on the further management of these patients.

DOI: 10.3238/arztebl.2014.0603b

REFERENCES

1. Antoniadis G, Kretschmer T, Pedro MT, König RW, Heinen CPG, Richter HP: Iatrogenic neurological damage—prevalence, diagnosis and treatment. *Dtsch Arztebl Int* 2014; 111: 273–9.
2. Brüser P: Motorische Ersatzoperationen an der oberen Extremität. *Dtsch Arztebl* 1999; 96: A-1134–9.
3. Rosén B, Lundborg G: The long term recovery curve in adults after median or ulnar nerve repair: a reference interval. *J Hand Surg Br* 2001; 26:196–200.
4. Steinau HU, Tofaute A, Huellmann K, et al.: Tendon transfers for drop foot correction: long-term results including quality of life assessment, and dynamometric and peobarographic measurements. *Arch Orthop Trauma Surg* 2011; 131: 903–10.

Dr. med. Tobias Hirsch

Prof. Dr. med. Marcus Lehnardt

Klinik für Plastische Chirurgie und Schwerbrandverletzte,
Handchirurgiezentrum, Operatives Referenzzentrum für Gliedmaßen-tumore
BG Universitätskliniken Bergmannsheil, Ruhr Universität Bochum
tobias.hirsch@rub.de

In Reply:

In our paper (1), we summarized the iatrogenic nerve injuries treated surgically in our department. We saw several patients with recurrent laryngeal nerve palsy after surgery (thyroid or anterior cervical spine surgery) at our outpatient clinic. In some of these patients, the palsies disappeared spontaneously. In cases of complete transection of this thin nerve, it is impossible for us to perform a nerve reconstruction. Since the ends of the severed nerve retract, the chances to discover them are slim. We recommend that in these cases reconstructive procedures should be performed by ETN specialists.

Iatrogenic hypoglossal nerve injuries require treatment by a peripheral nerve surgeon. We have not had this type of lesion in our patient population. If no reinnervation can be demonstrated clinically or by electromyography, surgical exposure and nerve reconstruction is indicated, unless the nerve was clearly severed during the operation. In these cases, primary end-to-end suture or early secondary surgical treatment should then be performed within 3 weeks (2).

We believe that in cases with iatrogenic nerve injury a reconstruction of the nerve should be performed by a surgeon specialized in peripheral nerve surgery. This is the governing principle that peripheral nerve surgeons should always follow. However, if the damage to the nerve is extensive or autologous grafting cannot be performed for various reasons, the option of a nerve transfer should be considered (3). Various nerve transfers can be performed, including

- the Oberlin transfer (coaptation of a fascicle of the ulnar nerve to the biceps branch of the musculocutaneous nerve) at the upper arm (4)
- a transfer of the triceps branch of the radial nerve to the motor portion of the axillary nerve in the shoulder region (3)
- a transfer of motor branches of the median nerve to the radial nerve branches for hand and finger extension at the proximal forearm (5, 6) or
- the diversion of the final path of the anterior interosseous nerve to the motor portion of ulnar nerve at the distal lower arm (7).

We believe that tendon transfers are indicated in cases where reinnervation was unsuccessful. The primary aim should always be to restore the continuity of the nerve to maintain the original situation. In our opinion, muscle and tendon transfers are a substitute and should not be regarded as more than this (2).

Ultimately, it would be best to prevent iatrogenic nerve injuries from happening in the first place. However, if they still occur after surgery, they should be detected and addressed. The patient requires treatment in a center specialized in peripheral nerve surgery within the timeframes detailed in our article.

DOI: 10.3238/arztebl.2014.0604

REFERENCES

1. Antoniadis G, Kretschmer T, Pedro MT, König RW, Heinen CPG, Richter HP: Iatrogenic neurological damage—prevalence, diagnosis and treatment. *Dtsch Arztebl Int* 2014; 111: 273–9.
2. Kretschmer T, Antoniadis G, Assmus H: *Nerven Chirurgie*. Heidelberg: Springer Verlag 2014.
3. Yang LJ-S, Chang KE-C, Chung KC: A systematic review of nerve transfer and nerve repair for the treatment of adult upper brachial plexus injury. *Neurosurgery* 2012; 71: 417–29.
4. Teboul F, Kakkar R, Ameer N, Beaulieu JY, Oberlin C: Transfer of fascicles from the ulnar nerve to the nerve to the biceps in the treatment of upper brachial plexus palsy. *J Bone Joint Surg Am* 2004; 86: 1485–90.
5. Davidge KM, Yee A, Kahn LC, Mackinnon SE: Median to radial nerve transfers for restoration of wrist, finger, and thumb extension. *J Hand Surg Am* 2013; 38: 1812–27.
6. Ray WZ, Mackinnon SE: Clinical outcomes following median to radial nerve transfers. *JHS* 2011; 36A: 202–8.
7. Flores LP: Distal anterior interosseous nerve transfer to the deep ulnar nerve and end-to-side suture of the superficial ulnar nerve to the third common palmar digital nerve for treatment of high ulnar nerve injuries: experience in five cases. *Arq Neuropsiquiatr* 2011; 69: 519–24.

Prof. Dr. med. Gregor Antoniadis

Dr. med. Maria T. Pedro

Neurochirurgische Klinik der Universität Ulm,
Bezirkskrankenhaus Günzburg
gregor.antoniadis@uni-ulm.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.