Published in final edited form as:

Cochrane Database Syst Rev. 2010 ; 2010(4): . doi:10.1002/14651858.CD008490.

Eletriptan for acute migraine headaches in adults

Sheena Derry¹, R Andrew Moore¹, and Henry J McQuay¹

¹Pain Research and Nuffield Department of Anaesthetics, University of Oxford, Oxford, UK

Abstract

This is the protocol for a review and there is no abstract. The objectives are as follows:

The objective of the review will be to determine the efficacy and tolerability of eletriptan compared to placebo and other active interventions in the treatment of acute migraine headaches in adults.

BACK GROUND

Description of the condition

Migraine is a common, disabling headache disorder, affecting about 12% of Western populations, and with considerable social and economic impact. It is more prevalent in women than men (on the order of 18% versus 6% 1-year prevalence), and in the age range 30 to 50 years (Hazard 2009; Lipton 2007; Moens 2007). The International Headache Society (IHS) classifies two major subtypes. Migraine without aura is the most common, and usually more disabling, subtype. It is characterised by attacks lasting 4 to 72 hours that are typically of moderate to severe pain intensity, unilateral, pulsating, aggravated by normal physical activity and associated with nausea and/or photophobia and phonophobia. Migraine with aura is characterised by reversible focal neurological symptoms that develop over a period of 5 to 20 minutes and last for less than 60 minutes, followed by headache with the features of migraine without aura. In some cases the headache may lack migrainous features or be absent altogether (IHS 2004).

A recent large prevalence study in the US found that over half of migraineurs had severe impairment or required bed rest during attacks. Despite this high level of disability and a strong desire for successful treatment, only a proportion of migraine sufferers seek professional advice for the treatment of attacks. The majority were not taking any preventive medication, although one-third met guideline criteria for offering or considering it. Nearly

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Contact address: Maura Moore, Pain Research and Nuffield Department of Anaesthetics, University of Oxford, West Wing (Level 6), John Radcliffe Hospital, Oxford, Oxfordshire, OX3 9DU, UK. maura.moore@pru.ox.ac.uk.

Editorial group: Cochrane Pain, Palliative and Supportive Care Group.

Publication status and date: New, published in Issue 4, 2010.

CONTRIBUTIONS OF AUTHORS: SD and RAM wrote the protocol and will carry out searches for studies. SD and RAM will carry out data extraction and analyses. HJM will act as arbitrator. All authors will be involved with writing the final review.

DECLARATIONS OF INTEREST: All review authors have received research support from charities, government and industry sources at various times. RAM and HJM have consulted for various pharmaceutical companies, including Pfizer Ltd, the manufacturer of eletriptan. RAM and HJM have received lecture fees from pharmaceutical companies related to analgesics and other healthcare interventions. Pfizer Ltd. supplied data from relevant clinical trials, but were not in any way involved in this review.

all (98%) migraineurs used acute treatments for attacks, with 49% using over-the-counter (OTC) medication only, 20% using prescription medication, and 29% using both. OTC medication included aspirin, other non-steroidal anti-inflammatory drugs (NSAIDs), paracetamol (acetaminophen) and paracetamol with caffeine (Bigal 2008; Diamond 2007; Lipton 2007). Similar findings have been reported from other large studies in France and Germany (Lucas 2006; Radtke 2009).

The significant impact of migraine with regard to pain, disability, social functioning, quality of relationships, emotional well-being and general health (Edmeads 1993; Osterhaus 1994; Solomon 1997) results in a huge burden for the individual, health services and society (Clarke 1996; Ferrari 1998; Hazard 2009; Hu 1999; Solomon 1997). The annual US economic burden relating to migraine, including missed days of work and lost productivity, is US\$14 billion (Hu 1999). Thus successful treatment of acute migraine attacks not only benefits patients by reducing their disability and improving health-related quality of life, but also reduces the need for healthcare resources and increases economic productivity (Jhingran 1996; Lofland 1999).

Description of the intervention

The symptomatic treatment of migraine advanced significantly with the development of the triptan class of drugs, of which sumatriptan was the first. Eletriptan (trade name Relpax) is a second-generation triptan, available as 20 mg and 40 mg oral tablets. The maximum daily dose is 80 mg. It is available by prescription only in most parts of the world. In England in 2008 there were almost 12,000 prescriptions for eletriptan in primary care, 10,500 of which were for the 40 mg dose (PCA 2009).

In order to establish whether eletriptan is an effective analgesic at a specified dose in acute migraine attacks, it is necessary to study its effects in circumstances that permit detection of pain relief. Such studies are carried out in individuals with established pain of moderate to severe intensity, using single doses of the interventions. Participants who experience an inadequate response with either placebo or active treatment are permitted to use rescue medication, and the intervention is considered to have failed in those individuals. In clinical practice, however, individuals would not normally wait until pain is of at least moderate severity, and may take a second dose of medication if the first dose does not provide adequate relief. Once analgesic efficacy is established in studies using single doses in established pain, further studies may investigate different treatment strategies and patient preferences. These are likely to include treating the migraine attack early while pain is mild, and using a low dose initially, with a second dose if response is inadequate.

How the intervention might work

Eletriptan is a 5-HT₁ agonist, selectively targeting the 5-HT (serotonin) 1B and 1D receptors. It has three putative mechanisms of therapeutic action (Ferrari 2002):

- vasoconstriction of dilated meningeal blood vessels;
- inhibition of the release of vasoactive neuropeptides from perivascular trigeminal sensory neurons;

- reduction of pain signal transmission in the trigeminal dorsal horn.

It is used for acute treatment, having no efficacy in preventing future attacks.

Why it is important to do this review

An earlier Cochrane review of eletriptan for acute migraine was withdrawn from publication (Smith 2007). It is important to reassess and re-analyse the data from that review together with the newer evidence now available. This is one of a series of reviews planned for acute treatments for migraine attacks.

OBJECTIVES

The objective of the review will be to determine the efficacy and tolerability of eletriptan compared to placebo and other active interventions in the treatment of acute migraine headaches in adults.

METHODS

Criteria for considering studies for this review

Types of studies—Randomised, double-blind, placebo-controlled studies using eletriptan to treat a discrete migraine headache episode will be included. Studies must have a minimum of 10 participants per treatment arm and report dichotomous data for at least one of the outcomes specified below. Studies reporting treatment of consecutive headache episodes will be accepted if outcomes for the first, or each, episode are reported separately. Cross-over studies will be accepted if there is adequate washout (48 hours) between treatments.

Types of participants—Studies must include adults (at least 18 years of age) with migraine. The diagnosis of migraine specified by the International Headache Society (IHS 1988; IHS 2004) will be used, although other definitions will be considered if they conform in general to IHS diagnostic criteria. There will be no restrictions on migraine frequency, duration or type (with or without aura). Participants taking stable prophylactic therapy to reduce migraine frequency will be accepted; details on the prophylactic therapy prescribed or allowed will be provided in the ‘Characteristics of included studies’ table.

Types of interventions—Included studies must use either a single dose of eletriptan to treat a discrete migraine headache episode when pain is of moderate to severe intensity, or investigate different dosing strategies and/or timing of the first dose in relation to headache intensity. There will be no restriction on dose, or route of administration, provided the medication is self-administered.

A placebo comparator is essential to demonstrate that eletriptan is effective in this condition. Active-controlled trials without a placebo will be considered as secondary evidence. Studies to demonstrate prophylactic efficacy in reducing the number or frequency of migraine headaches will not be included.

Types of outcome measures

Primary outcomes: The choice of main outcome measures for this review was made by taking into consideration scientific rigour, availability of data and patient preferences (Lipton 1999).

Patients with acute migraine headaches have rated complete pain relief, no headache recurrence, rapid onset of pain relief, and no side effects as the four most important outcomes (Lipton 1999).

In view of these patient preferences, and in line with the guidelines for controlled trials of drugs in migraine issued by the IHS (IHS 2000), the main outcomes to be considered will be:

- Pain-free at 2 hours, without the use of rescue medication;
- Reduction in headache pain ('headache relief') at 1 and 2 hours (pain reduced from moderate or severe to none or mild without the use of rescue medication);
- Sustained pain-free over 24 hours (pain-free within 2 hours, with no use of rescue medication or recurrence within 24 hours);
- Sustained pain reduction over 24 hours (headache relief at 2 hours, sustained for 24 hours, with no use of rescue medication or a second dose of study medication).

Pain intensity or pain relief should be measured by the patient (not the investigator or carer). Pain measures accepted for the primary outcomes will be:

- Pain intensity (PI): 4-point categorical scale, with wording equivalent to none, mild, moderate and severe; or 100 mm VAS;
- Pain relief (PR): 5-point categorical scale, with wording equivalent to none, a little, some, a lot, complete; or 100 mm VAS.

Only data obtained directly from the patient will be considered.

Secondary outcomes: Secondary outcomes considered will include:

- Participants with any adverse event over 24 hours post dose;
- Participants with particular adverse events over 24 hours post dose;
- Withdrawals due to adverse events over 24 hours post dose;
- Relief of headache-associated symptoms;
- Functional disability.

Search methods for identification of studies

Electronic searches—The following databases will be searched:

- Cochrane CENTRAL;
- MEDLINE (via OVID);

- EMBASE (via OVID);
- Oxford Pain Relief Database (Jadad 1996a).

See Appendix 1 for the search strategy for MEDLINE (via OVID). This will be modified for searching the other databases. There will be no language restrictions.

Searching other resources—Reference lists of retrieved studies and review articles will be searched for additional studies. Grey literature and abstracts will not be searched. Clinical trial reports made available to the authors of the original review by the manufacturer of eletriptan (Pfizer Ltd.) will be included. In addition, the authors of the present review will make a new written request for information about both published and unpublished data from Pfizer.

Data collection and analysis

Selection of studies—Two review authors will independently carry out the searches and select studies for inclusion. Titles and abstracts of all studies identified by electronic searches will be viewed on screen, and any that clearly do not satisfy inclusion criteria will be excluded. Full copies of the remaining studies will be read to identify those suitable for inclusion. Disagreements will be settled by discussion with a third review author.

Data extraction and management—Two review authors will independently extract data from included studies using a standard data extraction form. Disagreements will be settled by discussion with a third review author. Data will be entered into RevMan 5.0 by one author.

Assessment of risk of bias in included studies—Methodological quality will be assessed using the Oxford Quality Score (Jadad 1996b).

The scale is used as follows:

- Is the study randomised? If yes, give one point.
- Is the randomisation procedure reported and is it appropriate? If yes, add one point; if no, deduct one point.
- Is the study double-blind? If yes, add one point.
- Is the double-blind method reported and is it appropriate? If yes, add one point; if no, deduct one point.
- Are the reasons for patient withdrawals and dropouts described? If yes, add one point.

The scores for each study will be reported in the 'Characteristics of included studies' table.

A risk of bias table will also be completed, using assessments of randomisation, allocation concealment and blinding.

Measures of treatment effect—Relative risk (or ‘risk ratio’, RR) will be used to establish statistical difference. Numbers needed to treat (NNT) and pooled percentages will be used as absolute measures of benefit or harm.

The following terms will be used to describe adverse outcomes in terms of harm or prevention of harm:

- When significantly fewer adverse outcomes occur with eletriptan than with control (placebo or active) we will use the term the number needed to treat to prevent one event (NNTp).
- When significantly more adverse outcomes occur with eletriptan compared with control (placebo or active) we will use the term the number needed to harm or cause one event (NNH).

Unit of analysis issues—We will accept randomisation to individual patient only.

Dealing with missing data—The most likely source of missing data is in cross-over studies. Where this is an issue only first-period data will be used. In all cases (cross-over or parallel-group) where there are substantial missing data we will comment on this and perform sensitivity analysis if possible.

Assessment of heterogeneity—Heterogeneity of studies will be assessed visually (L’Abbe 1987).

Data synthesis—Studies using a single dose of eletriptan in established pain of at least moderate intensity will be analysed separately from studies in which medication is taken before pain is well established or in which a second dose of medication is permitted.

Effect sizes will be calculated and data combined for analysis only for comparisons and outcomes where there were at least two studies and 200 participants (Moore 1998). Relative risk of benefit or harm will be calculated with 95% confidence intervals (CIs) using a fixed-effect model (Morris 1995). NNT, NNTp and NNH with 95% CIs will be calculated using the pooled number of events by the method of Cook and Sackett (Cook 1995). A statistically significant difference from control will be assumed when the 95% CI of the relative risk of benefit or harm does not include the number one.

Significant differences between NNT, NNTp and NNH for different doses of active treatment, or between groups in the sensitivity analyses, will be determined using the z test (Tramer 1997). Data from comparisons and outcomes with only one study or fewer than 200 participants will be described in summary tables where appropriate for information and comparison, but will not be analysed quantitatively.

Subgroup analysis and investigation of heterogeneity—Issues for subgroup analysis are dose, formulation, and route of administration.

Sensitivity analysis—Sensitivity analysis is planned for study quality (Oxford Quality Score of 2 versus 3 or more), and for migraine type (with aura versus without aura). A minimum of two studies and 200 participants must be available for any sensitivity analysis.

Acknowledgments

SOURCES OF SUPPORT

Internal sources

- Oxford Pain Research funds, UK.

External sources

- NHS Cochrane Collaboration Programme Grant Scheme, UK.
- NIHR Biomedical Research Centre Programme, UK.

Appendix 1. Search strategy for MEDLINE (via OVID)

1. Serotonin Agonists/ OR Tryptamines/
2. (eletriptan OR Relpax).mp.
3. 1 OR 2
4. Headache/ OR exp Headache Disorders/ OR exp Migraine Disorders/
5. (headach* OR migrain* OR cephalgi* OR cephalalgi*).mp.
6. 4 OR 5
7. randomized controlled trial.pt.
8. controlled clinical trial.pt.
9. randomized.ab.
10. placebo.ab.
11. drug therapy.fs.
12. randomly.ab.
13. trial.ab.
14. groups.ab.
15. OR/7-14
16. 3 AND 6 AND 15

HISTORY

Protocol first published: Issue 4, 2010

NOTES

The review proposed in this protocol will replace a previously withdrawn review: Smith LA, Oldman A, McQuay HJ, Moore RA. Eletriptan for acute migraine. Cochrane Database of

Systematic Reviews 2007, Issue 1. Art. No.: CD003224. DOI: 10.1002/14651858.CD003224.pub2 (Smith 2007; originally published in 2001, Issue 3; withdrawn in 2007, Issue 1). That review was withdrawn at the request of the Cochrane Funding Arbitration Panel after a change in the The Cochrane Collaboration's commercial sponsorship policy. The original review fulfilled the criteria required by the Collaboration at the time it was submitted, and the source, nature and terms of funding were discussed openly and fully. This funding included an absolute right to publish for the authors, with no interference from the funding source. There was never any suggestion that the results of the review were mistaken, and those results have been confirmed by other, subsequent systematic reviews by other authors. Nevertheless, the Cochrane Pain, Palliative and Supportive Care Review Group (PaPaS) agreed to withdraw the review and seek a new team of authors to update, revise, and republish the review.

Staff at PaPaS' editorial office sought approval from the Cochrane Funding Arbitration Panel before registering the present title and another previously withdrawn title ('Rizatriptan for acute migraine headaches in adults') to new teams of authors. The Panel's reply was as follows:

Response to Funding Arbiter Referral - Ref 191208/038:

Your inquiry was reviewed by the Funding Arbitration panel. The panel recommends that the reviews go forward providing:

They are different reviews (not funded by industry) and in compliance with our policy and the financial ties of all authors (such as consultancies) be disclosed in the review.

Additional references

- Bigal 2008 . Bigal ME, Serrano D, Reed M, Lipton RB. Chronic migraine in the population: burden, diagnosis, and satisfaction with treatment. *Neurology*. 2008; 71(8):559–66. DOI: 10.1212/01.wnl.0000323925.29520.e7. [PubMed: 18711108]
- Clarke 1996 . Clarke CE, MacMillan L, Sondhi S, Wells NEJ. Economic and social impact of migraine. *Quarterly Journal of Medicine*. 1996; 89(1):77–84.
- Cook 1995 . Cook RJ, Sackett DL. The number needed to treat: a clinically useful measure of treatment effect. *BMJ*. 1995; 310(6977):452–4. [PubMed: 7873954]
- Diamond 2007 . Diamond S, Bigal ME, Silberstein S, Loder E, Reed M, Lipton RB. Patterns of diagnosis and acute and preventive treatment for migraine in the United States: results from the American Migraine Prevalence and Prevention study. *Headache*. 2007; 47(3):355–63. DOI: 10.1111/j.1526-4610.2006.00631.x. [PubMed: 17371352]
- Edmeads 1993 . Edmeads J, Findlay H, Tugwell P, Pryse-Phillips W, Nelson RF, Murray TJ. Impact of migraine and tension-type headache on life-style, consulting behaviour, and medication use: a Canadian population survey. *Canadian Journal of Neurological Sciences*. 1993; 20(2):131–7. [PubMed: 8334575]
- Ferrari 1998 . Ferrari MD. The economic burden of migraine to society. *Pharmacoeconomics*. 1998; 13(6):667–76. [PubMed: 10179702]
- Ferrari 2002 . Ferrari MD, Goadsby PJ, Roon KI, Lipton RB. Triptans (serotonin, 5-HT_{1B/1D} agonists) in migraine: detailed results and methods of a meta-analysis of 53 trials. *Cephalalgia*. 2002; 22(8):633–58. [PubMed: 12383060]
- Hazard 2009 . Hazard E, Munakata J, Bigal ME, Rupnow MF, Lipton RB. The burden of migraine in the United States: current and emerging perspectives on disease management and economic

- analysis. *Value in Health*. 2009; 12(1):55–64. DOI: 10.1111/j.1524-4733.2008.00404.x. [PubMed: 18671771]
- Hu 1999 . Hu XH, Markson LE, Lipton RB, Stewart WF, Berger ML. Burden of migraine in the United States: disability and economic costs. *Archives of Internal Medicine*. 1999; 159(8):813–8. [PubMed: 10219926]
- IHS 1988 . Headache Classification Committee of the International Headache Society. Classification and diagnostic criteria for headache disorders, cranial neuralgias and facial pain. *Cephalalgia*. 1998; 8(Suppl 7):1–96.
- IHS 2000 . International Headache Society Clinical Trials Subcommittee. Guidelines for controlled trials of drugs in migraine: second edition. *Cephalalgia*. 2000; 20(9):765–86. [PubMed: 11167908]
- IHS 2004 . Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders: 2nd edition. *Cephalalgia*. 2004; 24(Suppl 1): 1–160.
- Jadad 1996a . Jadad AR, Carroll D, Moore A, McQuay H. Developing a database of published reports of randomised clinical trials in pain research. *Pain*. 1996; 66(2-3):239–46. [PubMed: 8880846]
- Jadad 1996b . Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJM, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Controlled Clinical Trials*. 1996; 17(1):1–12. [PubMed: 8721797]
- Jhingran 1996 . Jhingran P, Cady RK, Rubino J, Miller D, Grice RB, Gutterman DL. Improvements in health-related quality of life with sumatriptan treatment for migraine. *Journal of Family Practice*. 1996; 42(1):36–42. [PubMed: 8537803]
- L'Abbe 1987 . L'Abbe KA, Detsky AS, O'Rourke K. Meta-analysis in clinical research. *Annals of Internal Medicine*. 1987; 107(2):224–33. [PubMed: 3300460]
- Lipton 1999 . Lipton RB, Stewart WF. Acute migraine therapy: do doctors understand what patients with migraine want from therapy? *Headache*. 1999; 39(Suppl 2):S20–S26.
- Lipton 2007 . Lipton RB, Bigal ME, Diamond M, Freitag F, Reed ML, AMPP Advisory Group. et al. Migraine prevalence, disease burden, and the need for preventive therapy. *Neurology*. 2007; 68(5):343–9. [PubMed: 17261680]
- Lofland 1999 . Lofland JH, Johnson NE, Batenhorst AS, Nash DB. Changes in resource use and outcomes for patients with migraine treated with sumatriptan: a managed care perspective. *Archives of Internal Medicine*. 1999; 159(8):857–63. [PubMed: 10219932]
- Lucas 2006 . Lucas C, Géraud G, Valade D, Chautard MH, Lantéri-Minet M. Recognition and therapeutic management of migraine in 2004, in France: results of FRAMIG 3, a French nationwide population-based survey. *Headache*. 2006; 46(5):715–25. DOI: 10.1111/j.1526-4610.2006.00430.x. [PubMed: 16643573]
- Moens 2007 . Moens G, Johannik K, Verbeek C, Bulters S. The prevalence and characteristics of migraine among the Belgian working population. *Acta Neurologica Belgica*. 2007; 107(3):84–90. [PubMed: 18072336]
- Moore 1998 . Moore RA, Gavaghan D, Tramer MR, Collins SL, McQuay HJ. Size is everything - large amounts of information are needed to overcome random effects in estimating direction and magnitude of treatment effects. *Pain*. 1998; 78(3):209–16. [PubMed: 9870574]
- Morris 1995 . Morris, JA.; Gardner, MJ. Calculating confidence intervals for relative risk, odds ratios and standardised ratios and rates. In: Gardner, MJ.; Altman, DG., editors. *Statistics with confidence - confidence intervals and statistical guidelines*. British Medical Journal; London: 1995. p. 50-63.
- Osterhaus 1994 . Osterhaus JT, Townsend RJ, Gandek B, Ware JE Jr. Measuring the functional status and well-being of patients with migraine headache. *Headache*. 1994; 34(6):337–43. [PubMed: 7928312]
- PCA 2009 . Anonymous. Prescription cost analysis, England 2008. The NHS Information Centre; 2009. ISBN: 978-1-84636-292-7

- Radtke 2009 . Radtke A, Neuhauser H. Prevalence and burden of headache and migraine in Germany. *Headache*. 2009; 49(1):79–89. DOI: 10.1111/j.1526-4610.2008.01263.x. [PubMed: 19125877]
- Smith 2007 . Smith LA, Oldman A, McQuay HJ, Moore RA. Eletriptan for acute migraine. *Cochrane Database of Systematic Reviews*. 2007; (1) Art. No.: CD003224. DOI: 10.1002/14651858.CD003224.pub2.
- Solomon 1997 . Solomon GD, Price KL. Burden of migraine. A review of its socioeconomic impact. *Pharmacoeconomics*. 1997; 11(Suppl 1):1–10. [PubMed: 10168038]
- Tramer 1997 . Tramèr MR, Reynolds DJM, Moore RA, McQuay HJ. Impact of covert duplicate results on meta-analysis: a case study. *BMJ*. 1997; 315(7109):635–40. [PubMed: 9310564]

**Indicates the major publication for the study*